Published in final edited form as:

J Phys Chem B. 2017 September 14; 121(36): 8437–8446. doi:10.1021/acs.jpcb.7b04318.

Role of Electrostatics in Protein-RNA Binding: The Global vs. the Local Energy Landscape

Zhaleh Ghaemi¹, Irisbel Guzman², David Gnutt¹, Zaida Luthey-Schulten^{1,3,4}, and Martin Gruebele^{1,3,*}

¹Department of Chemistry, University of Illinois, Urbana, Illinois 61801, USA

²Department of Biochemistry, University of Illinois, Urbana, Illinois 61801, USA

³Department of Physics, Center for the Physics of Living Cells, and Center for Biophysics and Quantitative Biology, University of Illinois, Urbana, IL 61801, USA

⁴Beckman Institute, University of Illinois, Urbana, IL 61801, USA

Abstract

U1A protein - stem loop 2 RNA association is a basic step in the assembly of the spliceosomal U1 small nuclear ribonucleoprotein. Long-range electrostatic interactions due to the positive charge of U1A are thought to provide high binding affinity for the negatively charged RNA. Short range interactions, such as hydrogen bonds and contacts between RNA bases and protein side chains, favor a specific binding site. Here, we propose that electrostatic interactions are as important as local contacts in biasing the protein-RNA energy landscape towards a specific binding site. We show by using molecular dynamics simulations that deletion of two long-range electrostatic interactions (K22Q and K50Q) leads to mutant-specific alternative RNA bound states. One of these states preserves many short-range interactions with aromatic residues in the original binding site, while the other one does not. We test the computational prediction with experimental temperature-jump kinetics using a tryptophan probe in the U1A-RNA binding site. The two mutants show the distinct predicted kinetic behaviors. Thus the stem loop 2 RNA has multiple binding sites on a rough RNA-protein binding landscape. We speculate that the rough protein-RNA binding landscape, when biased to different local minima by electrostatics, could be one way that protein-RNA interactions evolve towards new binding sites and novel function.

Graphical Abstract

*Corresponding author: mgruebel@illinois.edu, Tel: +1 217 333 6136.

Present address for D.G.: Lehrstuhl für Physikalische Chemie II, Ruhr-Universität Bochum, Universitätsstr. 150, D-44801 Bochum, Germany

The authors declare no competing financial interest.

SUPPORTING INFORMATION AVAILABLE

Additional theoretical and experimental figures are provided in supplementary information file. This information is available free of charge via the Internet at <http://pubs.acs.org/>.

INTRODUCTION

Interactions of RNA with ribonucleic acid-binding proteins are crucial for the maintenance of important structures and machinery in the cell, such as the ribosome and the spliceosome.¹⁻³ To enable these binding proteins to have a wide range of function, a variety of RNA binding domains have evolved that recognize diverse RNA sequences.⁴ The most abundant such domain in eukaryotes is the RNA recognition motif.² The structure in Figure 1A shows the RNA recognition motif located at the N-terminus of spliceosomal U1A protein, comprised of a $\beta_1\alpha_1\beta_2\beta_3\alpha_2\beta_4$ sequence, bound to stem loop 2 (SL2) RNA of the U1 small nuclear RNA.^{5,6} Not shown is a second recognition motif located at the C terminus of U1A protein, which does not bind RNA, because it is uncharged.⁷

The net charge of the N-terminal RNA-binding U1A protein is +7. Thus it binds to SL2 RNA with a sub-nanomolar dissociation constant (K_d).⁸ Surface plasmon resonance studies have shown that Lys→Gln (charge change +1→0) mutants with different long-range electrostatics have different contributions to K_d values.⁹ To predict these K_d values relative to the wild type, we recently developed a simple quantitative model based only on fluctuations within the bound complex and verified it experimentally.¹⁰ We also showed that short-range side chain dynamics of the free U1A protein are affected by Lys→Gln mutation,¹¹ indicating that local surface structure of the protein can be modulated by long-range electrostatics. Moreover, a fast (microsecond) phase has been detected in the early stages of U1A-RNA dissociation kinetics of the F56W pseudo wild type complex. This fast kinetic phase could indicate transient population of an additional binding site.¹²

It has been suggested that the binding mechanism of the U1A protein to SL2 RNA has two distinct 'lure' and 'lock' steps.^{13,14} During the 'lure' step, electrostatic interactions are believed to be crucial for attracting the RNA to the protein. Following by the 'lock' step, in which protein and RNA rearrange to occupy the native binding site.^{9,15,16} Here, we hypothesize that the long-range electrostatics are also important for the 'lock' step, by biasing a rough binding energy landscape towards a specific binding site.

We propose that different Lys→Gln mutations, despite being distant from the binding site, will not just change overall binding affinity, but actually result in RNA rearrangement on the protein surface and lead to alternative bound states. Furthermore, we suggest that these alternative bound states can be detected experimentally because of the aromatic residues in the main binding site (W56 in the extensively studied pseudo wild type) will interact differently with the bound RNA upon mutation.

To investigate these two specific hypotheses, we mutated two positively charged side chains Lys22 (K22) and Lys 50 (K50) in the U1A protein pseudo wild type (F56W). These particular residues were chosen for three reasons: (1) According to the crystallographic structures, they do not form any hydrogen bonds with the RNA.^{17,18} (2) K22 is relatively close to the RNA stem whereas K50 is relatively close to the RNA loop, thus affecting different areas of the protein-RNA interface (Figure 1A). (3) Both of these residues are conserved in mammals, *Drosophila* and plants^{19,20} (Figure S1), which suggest that K22 and K50 residues contribute to more than just a general positive charge of the U1A protein. We mutated the lysine residues to glutamine to eliminate the positive charge while keeping roughly the same side chain length.

We carried out multi-microsecond full-atom, explicitly solvated molecular dynamics (MD) simulations of the U1A-RNA complex at 298 K and 310 K. Analysis of the trajectories shows that the RNA repositions itself to alternative bound states on the protein surface, and that W56 at the active site interacts differently with RNA bases depending on mutation as well as temperature. Finally, we tested the computational predictions experimentally with circular dichroism spectroscopy, stopped flow kinetic and microsecond time-resolved temperature jump kinetics of the mutants, confirming that different mutants explore different local binding sites as a function of temperature, even though overall U1A-RNA affinity and protein structure is largely unaffected by the mutations

The U1A-RNA binding landscape is thus rugged, and specific long-range electrostatic interactions must be in place to select which local interactions will yield the binding site with the lowest free energy.

MATERIALS AND METHODS

Construction of the computational model systems

The structure of the U1A protein (residues 6–103) in complex with SL2 RNA was determined by X-ray crystallography (PDB ID: 4PKD).¹⁷ To simulate the N-terminal U1A protein (residues 1–102), Swiss-Model program^{21,22} was used to model residues 1 to 5, and the K22Q and K50Q mutations were introduced by psfgen of NAMD program.²³ The SL2-RNA sequence, was obtained by mutating the nucleotide residues at the 3' and 5' ends and addition of a Gua residue at the 5' end using the RNA composer program.²⁴ The first solvation layers were added with the Solvate²⁵ program, and the Solvate plugin of VMD program²⁶ was used to solvate the rest of the box with TIP3P water molecules.²⁷ After neutralizing the system by addition of K⁺ ions using the Ionize software,²⁸ 0.2 M KCl was added to the water box to reproduce the experimental condition. CHARMM36 force fields^{29,30} were used for protein and RNA and the determination of the protonation states was

done with program PropKa.³¹ A stepwise minimization and equilibration were performed as suggested by Eargle *et al.*³², with a 5 ns NPT equilibration, using a Langevin thermostat (damping coefficient of 5 ps⁻¹) and Langevin piston method^{33,34} (200 fs piston period and 100 fs piston decay). Periodic boundary condition and NPT ensemble were used for production runs. Long-range electrostatics was computed by particle-mesh Ewald algorithm³⁵, Lennard-Jones interactions were cut off at a distance of 1.2 nm and the time step was 1 fs. All simulations were performed with NAMD 2.11 package.²³

To achieve adequate sampling, simulations were performed on the XSEDE supercomputer in addition to the special-purpose supercomputer for molecular dynamics, Anton, at Pittsburgh Supercomputing Center.³⁶

Molecular dynamics simulations and structural analysis

To investigate the effects of electrostatic interactions on the U1A protein-RNA complex, we extensively simulated K22Q and K50Q mutant complexes in atomistic detail using MD simulations. A total of 1.88 μ s of simulation, with five independent replicas was performed for each complex. Both systems were simulated at two 298 K and 310 K temperatures, to mimic a temperature jump (T-jump) experiment that could validate the MD results. The list of simulations at different temperatures are summarized in Table S1.

We used the production runs to observe the conformational changes of the complexes at each and upon increasing the temperature.

Since the T-jump experiments show the fluorescence quenching from the Trp, we monitor the amino acids that can quench this signal by Dexter energy transfer.^{37–39} From each simulation, we obtained the CoM distance between the side chains of all residues which can quench the fluorescent signal of the probe (W56) and the Trp side chain. In Figure S2 all potentially effective residues (of both protein and RNA) according to³⁷ are presented. Among these residues, we considered those positioned up to 0.8 nm away from W56, particularly focusing on A6 that base stacks with W56 and C5 which is within 0.65 nm of W56.

Shortest paths and community analysis

A network is constructed on a set of nodes which are connected by edges. These nodes were defined as the C $_{\alpha}$ and one side chain atoms of all protein residues and Phosphate and side chain nitrogens of the RNA nucleotides. The edges are placed between pairs of nodes if any heavy atoms from the two corresponding residues are within 4.5 Å of each other for at least 50% of the trajectory. Each edge is weighted by the weight $W_{ij} = -\log(|C_{ij}|)$, where C_{ij} are the correlation value of the two nodes. We used generalized correlation based on mutual information²⁵ to calculate the correlation matrix for each mutant. A path length, D_{ij} , between the nodes i and j is the sum of the edge weights between the consecutive nodes (k, l) along the path: $D_{ij} = \sum_{k,j} W_{kl}$. The Floyd–Warshall algorithm was used to find the shortest distance D_{ij} between all pairs of nodes in the network. On the constructed network, we identified the communities within the complex. The community detection follows Girvan–Newman algorithm⁴⁰ and they represent the nodes that are densely interconnected with each other.

Protein expression and purification

The expression vector for the N-terminal domain of U1A was obtained from Nagai.⁴¹ Both mutants were generated by site-directed mutagenesis and confirmed by sequencing. The expression vectors contained a sequence coding for a hexaHis tag at the N-terminus of the protein to facilitate purification. Expression vectors for the mutant U1A proteins were transformed into *Escherichia coli* strain BL21DE3 (pLysS) competent cells. The cells were grown in LB medium, and protein expression was induced with 1 mM IPTG at OD₆₀₀=0.60. The cultures were grown for 5–6 hours after induction. The cells were pelleted, resuspended in 10 mL of lysis buffer (50 mM NaH₂PO₄, 300 mM NaCl, 20 mM imidazole, pH 7.6) and lysed by ultrasonication. The lysate was centrifuged at 10,000 rpm, the supernatant was loaded on a 1 mL Ni-NTA column, and the protein was eluted with elution buffer (50 mM NaH₂PO₄, 300 mM NaCl, 250 mM imidazole, pH 7.6). Eluted protein was dialyzed against storage buffer (10 mM potassium phosphate, 50 mM KCl) and concentrated by amicon filter MWCO 3,000. The concentration of each protein was determined using a BCA assay (Pierce). The histidine tag of all U1A mutants was removed by thrombin cleavage at a 400:1 ratio of protein to thrombin, with 1.79 units/μl of biotinylated thrombin on beads (Novagen). The reaction was carried out at room temperature overnight at protein concentrations up to 100 μM in 0.2 M KCl, buffered at pH 7.4 by 10 mM phosphate buffer. The beads were removed by filtering the solution through a 22 μm filter. The molecular weights of the purified proteins were confirmed by low-resolution electrospray ionization mass spectrometry and the purities of the proteins were assessed by SDS-PAGE.

Circular dichroism (CD)

Thermal stability of U1A protein mutants without RNA was measured within a J-715 spectropolarimeter equipped with a Peltier temperature control (Jasco Inc.) at 298, 310, and 320 K. A quartz cuvette (Starna Cells Inc.) with 200 mm path length was used to acquire CD spectra of 10 μM samples from 200–250 nm. For comparison with the literature¹¹ we used a solvent of 10 mM cacodylic acid, 50 mM NaCl at pH 7.4. Each CD spectrum in Figure S3 is an average of 50 spectra at 200 nm/min scan speed. The SI (Figure S4) compares pseudo wild type spectra under our simulated, stopped flow and temperature jump conditions at 0.2 M KCl.

Stopped flow kinetics experiments

U1A-SL2 RNA dissociation kinetics were monitored by jumping KCl concentration from 0.2 M to 1 M, and decreasing protein/RNA concentration from 1 to 0.5 μM. The fast mixing measurements were performed using a SX-20 stopped flow spectrometer (Applied Photophysics). The tryptophan was excited at 280 nm and fluorescence emission was monitored through a 350 nm interference filter (Applied Photophysics) with a 2 mm entrance and exit slit width. The W56 fluorescence intensity was monitored between 320–450 nm. The fast mixing time is approximately 3 ms. Fluorescence scans were collected in 10 s data files with 1000 data points (every 10 ms). During dissociation studies, a syringe with 1 μM U1A–RNA complex solution in binding buffer (10 mM potassium phosphate buffer, 0.2 M KCl, pH 7.4) was fast mixed with phosphate buffer (10 mM potassium phosphate buffer, 1.8 M KCl, pH 7.4) from another syringe. Upon mixing the protein:RNA

complex solution with the buffer in a 1:1 ratio, the final concentration of the U1A–RNA complex was 0.5 μ M. A final concentration of 1 M KCl was required to obtain an optimal dissociation fluorescence signal. For each sample, at least seven individual scans were averaged to give one data set. Three independent averages were used for data analysis.

Laser-induced temperature jump kinetics

Microsecond relaxation kinetics of the equilibrated complexes with SL2 RNA in binding buffer (10 mM potassium phosphate buffer, 0.2 M KCl, pH 7.4) were measured with a home-built temperature jump (T-jump) apparatus described elsewhere.¹² The SL2 RNA was heat-shocked before complexation to ensure correct secondary structure conformation and decrease significantly the dissociation constant (K_d). Only 20 μ M of U1A–SL2 RNA complex was utilized to avoid aggregation, but drive the equilibrium towards bound complex.

Laser temperature jumps of ~ 10 degrees were achieved using a Surelite III Q-switched Nd:YAG laser (Continuum Inc.) Raman-shifted to 1.9 μ m by passing the beam through a 1 m long tube with hydrogen gas pressurized to 300 psi. The beam was then passed through a 50 % beam splitter to allow the sample to be excited from two sides providing more uniform heating. The pre-jump equilibrium temperature was set using an automated temperature controller, model Lake Shore 330 (Lake Shore Cryotronics Inc.). The sample cell was made of fused silica tubing 3530S-100 (VibroCom) fused shut on one side. The fluorescence excitation path length was 0.3 mm. The sample was excited with a 80 MHz pulsed Ti:sapphire laser (KMLabs Inc.). The Ti:sapphire laser wavelength was 860 nm, which was frequency tripled with a third harmonic generator (CSK Optonics Inc.) to 287 nm. Tryptophan fluorescence was then guided from the sample by an optical light-guide (Oriol Corp.), passed through a B370 band-pass filter (Hoya Corp.) and collected by a photomultiplier R7400U-03 (Hamamatsu Corp.). The signal was then recorded and digitized every 100 ps by an oscilloscope DPO7254 (Tektronix Inc.) with 2.5 GHz bandwidth. The length of the time traces was 500 μ s and each trace contained many tryptophan fluorescence decays every 12.5 ns (80 MHz). The temperature jump was set to occur 153.75 s after the oscilloscope was triggered to start data collection to provide a pre-jump baseline. The fluorescence decay peak signal was usually 10–40 mV. The fluorescence lifetime decay analysis has been previously reported.¹¹

ϕ value analysis

We performed a ϕ value analysis ($\phi^\# = G^\#_{M-WT} / G_{M-WT}$) to study the effect of the mutations (M = K22Q or K50Q) on the transition state (indicated by #). For this analysis, the F56W pseudo wild type was used as reference wild type. G_{M-WT} was calculated with the free energies obtained from equilibrium experiments, electrophoretic mobility shift assay (EMSA), at 298 K for mutant and pseudo wild type (F56W) complexes.¹⁰ $G^\#_{M-WT}$ was calculated based on the dissociation times obtained from the stopped flow kinetics experiments described above.

RESULTS

Changes in long-range electrostatics bias the U1A-RNA energy landscape towards new temperature-sensitive binding sites

Microsecond-long (a total of 1.88 μ s) MD simulations were performed to test our hypothesis that changes in long-range electrostatic interactions stabilize different U1A-RNA conformation. Table S1 summarizes all the simulations at both 298 K and 310 K temperatures (see Materials and Methods for details). Previously, it was shown by MD simulations that, the U1A protein-RNA complex structure with F56W mutation (pseudo wild type) at 298 K, remains bound close to the known crystal structure.^{10,42} We also performed 1 μ s MD simulations for the pseudo wild type complex with five replicas at 298 K as a reference point. By analyzing our MD simulations of K22Q/F56W and K50Q/F56W mutant complexes (hereafter referred to as K22Q and K50Q for simplicity), we can provide structural details for the conformational changes occurred in 20% of our simulations as a function of temperature. RMSD values for about 0.5 μ s of frames from the MD simulations for each temperature, were used as input for the clustering of trajectories. The RMSD plots for all 1.8 μ s performed simulations are reported in Figures S5 and S6.

We show representative structures from a cluster analysis of the K22Q complex at 298 K (blue) in Figure 2A, and after simulated T-jump to 310 K (red) in Figure 2B, both overlaid on the pseudo wild type structure (gray).

At 298 K, the RNA-bound state maintains a similar binding conformation to the pseudo wild type, although the 3' end of the RNA loop is distorted. The distortion is visible in the middle left of Figure 2A, and results in increased RNA loop (residues 1–10) backbone RMSD with respect to the pseudo wild type structure (Figure 2E). After jumping to 310 K, the RNA moves counterclockwise to a new bound state on the surface of the protein, resulting in breakage of about 73% of hydrogen bonds (Table S2) and of the base stacking interactions at the crystallographic binding site, namely Y13-C5 and W56-A6. In addition, helix C, which is crucial for binding of the protein to the RNA^{16,43} changes its conformation from the initially 'open' to a more 'closed' state (black arrow in Figure 2B). These 'open' and 'closed' states have been described previously for the bare protein.¹¹ This conformational change at 310 K modifies the electrostatic potential of the complex to a more positive potential in the vicinity of the RNA binding site with respect to the electrostatic potential at 298 K, as shown in Figure 2C, D. The hydrogen bonds that stabilize the alternative RNA binding conformation are shown in Figure S7A, S7B.

The K50Q mutant behaves quite differently, as shown in Figure 3. We show the representative structures of the K50Q complex at 298 K (blue) in Figure 3A, and the simulated T-jump to 310 K (red) in Figure 3B, both are overlaid on the pseudo wild type structure (gray). Again, all structures were obtained from a cluster analysis of the trajectory frames in terms of RMSD.

For K50Q mutant, the RNA is unwrapped from the native U1A protein binding site at both temperatures (Figure 3A, E). Up to 53% of the native complex hydrogen bonds including the majority (78%) of the loop 3 hydrogen bonds network (Table S3) and base stacking

interactions are lost: the base of Y13 does not stack against C5, whereas the W56-A6 base stacking remains intact (Figure 3A, B). Figure S7C, S7D shows that the new RNA bound state forms hydrogen bonds with the protein. The lysine 50 mutation changes the electrostatic potential of loop 3 in the protein, which plays an important role in RNA binding and complex stability⁴⁴, to be more negative (Figure 3C, D).

The conformational changes of both mutant complexes can be quantified by monitoring the distance between the center of mass (CoM) of the backbone atoms of the RNA loop and the C α of W56 of U1A protein as a reaction coordinate. The reaction coordinate probability distributions for both mutant complexes, using all the 1.8 μ s simulation data, shown in Figure S5, is consistent with our description above.

Electrostatic mutations affect the correlated pathways within the complex

When protein and RNA move, the amplitude of fluctuations in certain regions grow and diminish together. Mutations can affect these correlations within the protein-RNA complex. The changes in correlated regions upon mutation were determined by analyzing the shortest correlated pathways within the complex. Essentially, the shortest correlated pathway follows the most correlated set of atoms from one region to another⁴⁵ (see Methods for details).

We focused on the changes of the pathways from the mutated residues to important protein sites such as W56, which base-stacks with the RNA. Figure 4A shows that for pseudo wild type complex, the shortest pathway from RNA stem, nucleotide U(-5), passes through the K22 side chain to connect to W56. After K22Q mutation, this pathway is largely rearranged and does not involve residue 22 (Figure 4B). In addition, the pathway connecting nucleotide U2 to W56 by passing through K50 is lost upon mutation, and there is no other pathway connecting these two parts of the complex. Thus side chains such as K22 and K50 are directly connected to local dynamics at the binding site.

The effects of the mutations on the correlations within the complex were determined by performing community analysis, which detects the atoms whose fluctuation amplitudes track one another. Specifically, we compared the communities containing K50 in pseudo WT and K50Q mutant complex. As shown in Figure 4C, the K50 community contains loop 3 residues and connects two opposite sides of the RNA loop, namely residues A1, U2, U3, G4 and C9, C10 and G11, together with parts of β 1. Upon K50Q mutation, most of the connections of the K50 to the RNA loop residues present in the pseudo WT are broken as a result of the RNA ‘unwrapping’ (Figure 4D). These results emphasize the global effects of the studied mutations on the overall complex, and how the correlated motions of protein and RNA are disrupted by the mutations.

MD simulations reveal short-range structural markers for different binding sites

We show that the above-mentioned large scale rearrangements of U1A and RNA binding caused by changes in long-range electrostatics are also reflected in changed local contacts of the binding site (containing W56 and Y13) with the RNA. W56 has been previously shown to capture the kinetics of U1A-RNA complex with T-jump experiment.¹² To quantify the short-range conformational changes, we mainly monitored the CoM distance of W56 side chain to 1) base C5 and 2) base A6 in the RNA (see Materials and Methods for details).

Additional protein residues for which their CoM distances to W56 change by increasing the temperature are shown in Figure S9.

For the K22Q complex at higher temperature, both base stacking interactions break (Figure 5A–D) so that W56–A6 and Y13–C5 CoM distances shift toward larger values. However, Figure 5E–H for K50Q complex shows that the W56–A6 distance, a conserved base stacking interaction in this protein family,^{18,41} does not change with increasing the temperature.

Based on the MD simulation results, we can thus say the following: two different mutants that disrupt long-range electrostatic interactions between U1A and SL2 RNA result in large, but different, changes in binding geometry. Population of the alternative bound state is temperature-sensitive for the K22Q mutant in the 298–310 K range, less so for the K50Q mutant. The global change of electrostatic environment manifests itself in the local environment of the W56 residue in the binding site: again, the larger shift is seen for K22Q, with the Trp binding relatively unaffected in K50Q with respect to the pseudo wild type.

To summarize the MD results, we predict that in a T-jump experiment, the K22Q complex will show a change in the fluorescent signal of W56, whereas the K50Q complex will not, for two reasons: 1) K22Q in Figure 2A switches from a mostly pseudo wild type binding geometry to a new ‘rolled counter-clockwise’ binding site only after a simulated T-jump, whereas K50Q is in the same new ‘unwrapped’ binding site at both temperatures. 2) The effect of temperature on the local Trp environment is greater for K22Q than K50Q, as shown in Figure 5. We now address this prediction experimentally.

Both mutants have similar secondary structure over the full temperature range

First, we checked the integrity of secondary structure of the mutants over the temperature range of interest (lowest temperature before T-jump at 298 K to highest temperature after T-jump at 318 K). The circular dichroism (CD) spectrum of each U1A mutant was measured at 298 K and 318 K and is shown in Figure S3 in comparison to literature data¹¹ for the pseudo wild type. The CD spectra for both mutants did not change significantly over the 298 to 318 K temperature range, indicating that a temperature jump does not introduce a large perturbation to the secondary structure of the mutants.

The CD spectra were similar in shape to the pseudo-wild type control protein (black curve in Figure S4), although the K22Q mutant had a slightly smaller and the K50Q mutant a slightly larger mean residue ellipticity.

Stopped flow kinetics show that the mutants do not greatly accelerate unbinding

To verify that the complete dissociation kinetics of the mutant complexes is similar to the previously reported pseudo wild type, stopped flow experiments were performed. A solution of the U1A protein–SL2 RNA complex was mixed with a buffered solution containing no U1A or SL2 RNA to change the KCl concentration from an initial concentration of 0.2 M KCl to a final concentration of 1 M KCl and the concentration of the U1A–SL2 RNA complex was reduced by a factor of 2 upon mixing. Under these conditions, the dissociation rate should dominate the relaxation process, so the observed rate is close to the dissociation rate.

A single phase was observed upon fast mixing for the dissociation of the complexes formed with K22Q and K50Q mutants shown in Figure 6. The fluorescence amplitude increased almost to the protein-only baseline for all of the positively charged mutants, indicating nearly full dissociation of the complex. Figure S10 shows the complex dissociation at the lower temperature of 298 K, which is slower as expected.

The dissociation time constant τ for the K22Q complex is 86 ± 2 ms, and for the K50Q complex is 110 ± 0.03 ms. The K22Q complex has a slightly higher binding affinity than the K50Q mutant complex,¹⁰ but its dissociation is slightly faster, indicating less destabilization of the transition state(s) relative to the pseudo wild type.

Temperature-jump measurements

Finally, we carried out microsecond T-jump experiments comparing the two mutants, to see if they show different behavior as predicted by MD. Our probe signal was the Trp fluorescence decay lifetime change χ normalized from 1 (pre-jump equilibrium) to 0 (post jump equilibrium). The protein-RNA complex solution in a 0.2 M KCl buffer was perturbed by ~ 10 K temperature jumps starting at 298 K or 308 K (See Materials and Methods and ref.¹² for more details of data analysis).

The K22Q mutant complex shows a Trp relaxation signal, with relaxation times of $\tau \approx 13$ and 27 μ s, and a doubling of the amplitude from low to high temperature (Figure 7A, B). This is consistent with a transition between the original binding site and a new binding site increasingly favored at higher temperature, as proposed by the MD simulations in Figure 2A, B. It is also consistent with the larger temperature sensitivity and local structural change around the W56 residue, as predicted by MD in Figure 5.

Quite a different result is obtained for the K50Q mutant complex (Figure 8). We did not observe a microsecond phase at all, as predicted by our MD simulation results. Based on the experimental data, either there is no conformational rearrangement of protein and RNA, or if there is, the Trp interaction with the RNA in the binding site is unaffected by the conformational change. The MD simulation results in Figures 3 and 5 are consistent with the latter interpretation: the K50Q mutant complex features a new binding site whose tryptophan local environment is temperature-independent.

DISCUSSION

It has been known that electrostatic interactions in U1A-SL2 RNA complex are important for affinity between the protein and RNA, whereas short-range interactions such as hydrogen bonds and base stacking interactions are thought to finalize the locking of the RNA into the native binding site. In this 'locking' step, electrostatic interactions were not thought to play a major role mainly based on the mutants K_{off} measurements.^{9,13}

Here we hypothesize that electrostatic interactions are also crucial for biasing the U1A-RNA energy landscape to stabilize the native complex binding site over alternative binding sites. Our MD simulations indeed show that this is so, and that the nature of the alternative binding site upon electrostatic mutation differs between mutants. For K22Q mutant, complex

remained close to the original binding site at 298 K, but switched to a ‘rolled over’ bound state upon simulated T-jump to 310 K. This suggests the new bound state, more stable at higher temperature, is looser and more entropically favored. The observation agrees with the ‘tight-loose’ binding site previously postulated for pseudo-wild type complex.¹² This observation is also not unexpected because this mutant is known to have a lower binding affinity than the pseudo wild type.¹⁰ The same complex showed significant disruption of the Trp-RNA interactions in Figure 5B. Thus we predicted that a change in tryptophan fluorescence could be observed in an experimental T-jump.

In contrast, another mutant complex, K50Q-RNA, occupied a binding site different from pseudo wild type at both temperatures. That observation suggests that the alternate bound state may be enthalpically favored, so it is stable even at low temperature. The alternative bound conformation is different from the one observed in K22Q complex at 310 K, showing that different electrostatics lead to population of alternative binding sites. Also in contrast to K22Q complex, the Trp-RNA interaction of the K50Q complex is temperature-independent, as seen in Figure 5H. Thus the protein-RNA binding energy landscape is quite sensitive to electrostatics, which do play a role in binding site selection. Thus we predicted that no change in tryptophan fluorescence could be observed in an experimental T-jump.

Of course our MD simulations, although extensive, cannot conclusively prove these hypotheses, but they make a clear prediction: T-jumps of K22Q complex should show a change in Trp quenching because the binding site is temperature sensitive and the Trp-RNA interactions change a lot; whereas T-jumps of K50Q complex should show much less change in Trp quenching because the binding site is less temperature sensitive and the Trp-RNA interactions change less dramatically.

Our T-jump experiments verify this prediction based on MD simulation. The K22Q-RNA complex shows a fast (microsecond) rearrangement of the local Trp environment, whereas the K50Q-RNA complex does not. The simulations suggest that temperature-sensitive switching between two binding sites is responsible for the fast kinetics observed for K22Q complex, whereas a temperature-insensitive new binding site is responsible for the lack of fast kinetics observed for K50Q-RNA, although the experiments cannot of course distinguish the microscopic reasons. The MD simulations and experiments nicely complement one another. Since the K22Q mutant complex populates a high-temperature alternative bound state which is thus entropically favored, this binding site is not as conformationally rigid as the native binding site.

The changes in local binding sites driven by changes in electrostatics have relatively little effect on the overall dissociation reaction, as measured by stopped flow experiments (Figure 6). Although both mutants bind somewhat more weakly than pseudo wild type¹⁰ the alternative bound states have similar barriers to dissociation. This observation shows that electrostatics can steer local binding, while having little effect on overall affinity. We checked this quantitatively by ϕ value analysis, as shown in Table 1 (see Materials and Methods for details). Both mutants have relatively large ϕ values, showing that many protein-RNA contacts have formed in the transition state. Although K50Q mutant forms the least stable complex, it is not the fastest to dissociate (Figure 9), probably because its

transition state still has 82 % of complex contacts formed (ϕ value in Table 1), vs. 70% in the K22Q mutant.

CONCLUSIONS

The combination of simulations and experimental data provides strong evidence to support our hypothesis that positively charged sites in U1A protein are important not just for strengthening the interaction between protein and RNA. They are also important for locking the RNA into its local binding site. Deletion of individual lysines allows other binding sites to come into play, sometimes only at higher temperature, when the alternative bound state is entropically favored, sometimes over a wide temperature range, when the alternative bound state is also enthalpically favored. Some of these states have base-side chain interactions similar to the original binding site, others disrupt the aromatic-RNA stacking interaction. Thus electrostatics can bias the rough energy landscape of RNA Recognition Motifs and their cognate RNAs towards different bound local minima. We speculate that the combination of a rugged landscape with long-range bias could facilitate evolution of new binding sites and functions upon mutations that alter electrostatics, rather than mutations in the binding sites themselves.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This work was supported by National Science Foundation [Grant MCB-1244570 to Z.L.-S and Z.G. and Graduate Research Fellowship DGE-1144245 to I.G.] and by the National Institutes of Health [Grant R01GM093318 to M.G. and I.G.]. M.G. held the James R. Eiszner Chair and Z.L.S held the William and Janet Lycan Chair in Chemistry while this work was carried out. Supercomputer time was provided by stampede-XSEDE [TG-MCA03S027] and Anton [PSCA15052]. Anton computer time was provided by the Pittsburgh Supercomputing Center (PSC) through Grant *R01GM116961* from the National Institutes of Health. The Anton machine at PSC was generously made available by D.E. Shaw Research.

References

1. Varani G, Nagai K. RNA recognition by RNP proteins during RNA processing. *Annu Rev Biophys Biomol Struct.* 1998; 27:407–445. [PubMed: 9646873]
2. Maris C, Dominguez C, Allain FH. The RNA recognition motif, a plastic RNA-binding platform to regulate post-transcriptional gene expression. *FEBS J.* 2005; 272(9):2118–2131. [PubMed: 15853797]
3. Nagai K, Oubridge C, Ito N, Avis J, Evans P. The RNP domain: a sequence-specific RNA-binding domain involved in processing and transport of RNA. *Trends Biochem Sci.* 1995; 20(6):235–240. [PubMed: 7543225]
4. Lunde BM, Moore C, Varani G. RNA-binding proteins: modular design for efficient function. *Nat Rev Mol Cell Biol.* 2007; 8(6):479–490. [PubMed: 17473849]
5. Scherly D, Boelens W, van Venrooij WJ, Dathan NA, Hamm J, Mattaj IW. Identification of the RNA binding segment of human U1 A protein and definition of its binding site on U1 snRNA. *EMBO J.* 1989; 8(13):4163–4170. [PubMed: 2531658]
6. Lutz-Freyermuth C, Query CC, Keene JD. Quantitative determination that one of two potential RNA-binding domains of the A protein component of the U1 small nuclear ribonucleoprotein complex binds with high affinity to stem-loop II of U1 RNA. *Proc Natl Acad Sci U S A.* 1990; 87(16):6393–6397. [PubMed: 1696729]

7. Lu JR, Hall KB. An RBD That Does Not Bind RNA - NMR Secondary Structure Determination and Biochemical-Properties of the C-Terminal RNA-Binding Domain from the Human U1A Protein. *J Mol Biol.* 1995; 247(4):739–752. [PubMed: 7723028]
8. Hall KB, Stump WT. Interaction of N-terminal domain of U1A protein with an RNA stem/loop. *Nucleic Acids Res.* 1992; 20(16):4283–4290. [PubMed: 1508720]
9. Law MJ, Linde ME, Chambers EJ, Oubridge C, Katsamba PS, Nilsson L, Haworth IS, Laird-Offringa Ia. The role of positively charged amino acids and electrostatic interactions in the complex of U1A protein and U1 hairpin II RNA. *Nucleic Acids Res.* 2006; 34(1):275–285. [PubMed: 16407334]
10. Ghaemi Z, Guzman I, Baek J, Gruebele M, Luthey-Schulten Z. Estimation of Relative Protein-RNA Binding Strengths from Fluctuations in the Bound State. *J Chem Theory Comp.* 2016; 12:4593–4599.
11. Guzman I, Ghaemi Z, Baranger AM, Luthey-Schulten Za, Gruebele M. Native Conformational Dynamics of the Spliceosomal U1A Protein. *J Phys Chem B.* 2015; 119:3651–3661. [PubMed: 25659967]
12. Anunciado D, Dhar A, Gruebele M, Baranger AM. Multistep kinetics of the U1A- SL2 RNA complex dissociation. *J Mol Biol.* 2011; 408(5):896–908. [PubMed: 21419778]
13. Katsamba PS, Myszkowski DG, Laird-Offringa IA. Two functionally distinct steps mediate high affinity binding of U1A protein to U1 hairpin II RNA. *J Biol Chem.* 2001; 276(24):21476–21481. [PubMed: 11297556]
14. Plantinga MJ, Korennykh AV, Piccirilli JA, Correll CC. Electrostatic interactions guide the active site face of a structure-specific ribonuclease to its RNA substrate. *Biochemistry.* 2008; 47(34): 8912–8918. [PubMed: 18672906]
15. Law MJ, Chambers EJ, Katsamba PS, Haworth IS, Laird-Offringa Ia. Kinetic analysis of the role of the tyrosine 13, phenylalanine 56 and glutamine 54 network in the U1A/U1 hairpin II interaction. *Nucleic Acids Res.* 2005; 33(9):2917–2928. [PubMed: 15914668]
16. Law MJ, Lee DS, Lee CS, Anglim PP, Haworth IS, Laird-Offringa Ia. The role of the C-terminal helix of U1A protein in the interaction with U1hpII RNA. *Nucleic Acids Res.* 2013; 41(14):7092–7100. [PubMed: 23703211]
17. Kondo Y, Oubridge C, van Roon AMM, Nagai K. Crystal structure of human U1 snRNP, a small nuclear ribonucleoprotein particle, reveals the mechanism of 5' splice site recognition. *Elife.* 2015; 4:1–19.
18. Oubridge C, Ito N, Evans PR, Teo C-H, Nagai K. Crystal structure at 1.92 Å resolution of the RNA-binding domain of the U1A spliceosomal protein complexed with an RNA hairpin. *Nature.* 1994; 372(6505):432–438. [PubMed: 7984237]
19. Simpson GG, Clark GP, Rothnie HM, Boelens W, van Venrooij W, Brown JW. Molecular characterization of the spliceosomal proteins U1A and U2B" from higher plants. *EMBO J.* 1995; 14(18):4540–4550. [PubMed: 7556097]
20. Flickinger TW, Salz HK. The *Drosophila* sex determination gene *snf* encodes a nuclear protein with sequence and functional similarity to the mammalian U1A snRNP protein. *Genes Dev.* 1994; 8(8):914–925. [PubMed: 7926776]
21. Kiefer F, Arnold K, Künzli M, Bordoli L, Schwede T. The SWISS-MODEL Repository and associated resources. *Nucleic Acids Res.* 2009; 37(suppl 1):D387–D392. [PubMed: 18931379]
22. Arnold K, Bordoli L, Kopp J, Schwede T. The SWISS-MODEL workspace: a web-based environment for protein structure homology modelling. *Bioinformatics.* 2006; 22(2):195–201. [PubMed: 16301204]
23. Phillips JC, Braun R, Wang W, Gumbart J, Tajkhorshid E, Villa E, Chipot C, Skeel RD, Kalé L, Schulten K. Scalable molecular dynamics with NAMD. *J Comput Chem.* 2005; 26(16):1781–1802. [PubMed: 16222654]
24. Popenda M, Szachniuk M, Antczak M, Purzycka KJ, Lukasiak P, Bartol N, Blazewicz J, Adamiak RW. Automated 3D structure composition for large RNAs. *Nucleic Acids Res.* 2012; 40(14):e112. [PubMed: 22539264]
25. Lange OF, Grubmüller H. Generalized correlation for biomolecular dynamics. *Proteins.* 2006; 62(4):1053–1061. [PubMed: 16355416]

26. Humphrey W, Dalke A, Schulten K. {VMD} -- {V}isual {M}olecular {D}ynamics. *J Mol Graph*. 1996; 14:33–38. [PubMed: 8744570]
27. Jorgensen WL, Madura JD. Solvation and Conformation of Methanol in Water. *J Am Chem Soc*. 1983; 105:1407–1413.
28. Baraëff A, , Roberts E. Ionize 1.6.0 University of Illinois at Urbana- Champaign; 2005
29. Denning EJ, Priyakumar UD, Nilsson L, Mackerell AD. Impact of 2'-hydroxyl sampling on the conformational properties of RNA: Update of the CHARMM all-atom additive force field for RNA. *J Chem Theory Comput*. 2011; 32(9):1929–1943.
30. Best RB, Zhu X, Shim J, Lopes PEM, Mittal J, Feig M, Mackerell AD. Optimization of the additive CHARMM all-atom protein force field targeting improved sampling of the backbone ϕ ψ and side-chain $\chi(1)$ and $\chi(2)$ dihedral angles. *J Chem Theory Comput*. 2012; 8(9):3257–3273. [PubMed: 23341755]
31. Rostkowski M, Olsson MHM, Sndergaard CR, Jensen JH. Graphical analysis of pH-dependent properties of proteins predicted using PROPKA. *BMC Struct Biol*. 2011; 11(1):6. [PubMed: 21269479]
32. Eargle J, Black Aa, Sethi A, Trabuco LG, Luthey-Schulten Z. Dynamics of Recognition between tRNA and elongation factor Tu. *J Mol Biol*. 2008; 377(5):1382–1405. [PubMed: 18336835]
33. Feller SE, Zhang Y, Pastor RW, Brooks BR. Constant pressure molecular dynamics simulation: The Langevin piston method. *J Chem Phys*. 1995; 103(11):4613–4621.
34. Martyna GJ, Tobias DJ, Klein ML. Constant pressure molecular dynamics algorithms. *J Chem Phys*. 1994; 101(5)
35. Darden T, York D, Pedersen L. Particle mesh Ewald: An $N \cdot \log(N)$ method for Ewald sums in large systems. *J Chem Phys*. 1993; 98(12):10089–10092.
36. Shaw DE, , Dror RO, , Salmon JK, , Grossman JP, , Mackenzie KM, , Bank JA, , Young C, , Deneroff MM, , Batson B, , Bowers KJ, , et al. Millisecond-scale Molecular Dynamics Simulations on Anton. Proceedings of the Conference on High Performance Computing Networking, Storage and Analysis; SC '09; New York, NY, USA: ACM; 2009 39: 139:11
37. Chen Y, Barkley MD. Toward Understanding Tryptophan Fluorescence in Proteins. *Biochemistry*. 1998; 37:9976–9982. [PubMed: 9665702]
38. Callis PR, Burgess BK. Tryptophan Fluorescence Shifts in Proteins from Hybrid Simulations : An Electrostatic Approach. *J Phys Chem B*. 1997; 5647(97):9429–9432.
39. Vivian JT, Callis PR. Mechanisms of tryptophan fluorescence shifts in proteins. *Biophys J*. 2001; 80(5):2093–2109. [PubMed: 11325713]
40. Girvan M, Newman MEJ. Community structure in social and biological networks. *Proc Natl Acad Sci U S A*. 2002; 99(12):7821–7826. [PubMed: 12060727]
41. Nagai K, Oubridge C, Jessen TH, Li J, Evans PR. Crystal structure of the RNA-binding domain of the U1 small nuclear ribonucleoprotein A. *Nature*. 1990; 348(6301):515–520. [PubMed: 2147232]
42. Kormos BL, Pieniazek SN, Beveridge DL, Baranger AM. U1A Protein-Stem Loop 2 RNA Recognition: Prediction of Structural Differences from Protein Mutations. 2011; 95(9):591–606.
43. Kurisaki I, Takayanagi M, Nagaoka M. Combined mechanism of conformational selection and induced fit in U1A-RNA molecular recognition. *Biochemistry*. 2014; 53(22):3646–3657. [PubMed: 24828852]
44. Katsamba PS, Bayramyan M, Haworth IS, Myszkowski DG, Laird-Offringa Ia. Complex role of the beta 2-beta 3 loop in the interaction of U1A with U1 hairpin II RNA. *J Biol Chem*. 2002; 277(36): 33267–33274. [PubMed: 12082087]
45. Sethi A, Eargle J, Black Aa, Luthey-Schulten Z. Dynamical networks in tRNA:protein complexes. *Proc Natl Acad Sci*. 2009; 106(16):6620–6625. [PubMed: 19351898]

Figure 1.

(A) Structure of the U1A-SL2 RNA complex: sites of mutation, K22 and K50 together with the probe W56 are shown in licorice representation. Sequences of B) SL2-RNA and C) pseudo wild type (F56W) U1A protein, with the positions of K22 and K50 mutation sites colored in red. The ribonucleoprotein (RNP) functional units (RNP1, RNP2) that contribute to the base stacking interactions with the RNA are colored in green.

Figure 2.

K22Q mutant complex conformations: (A) After 0.57 μ s production run at 298 K, the RNA (blue) and protein (light blue) structures do not show a large conformational change from the representative pseudo wild type structure (gray). (B) After the T-jump to 310 K, the RNA (red) finds a new binding site relative to the RNA in the pseudo wild type complex (gray). The black arrow shows that the helix C of the protein at 310 K (light red) rearranges relative to wild type protein (gray). (C) Electrostatic potential map (± 5 k_BT/e isosurfaces) of representative structures of the complexes at 298 K and (D) 310 K together. The corresponding RNA structure (magenta) is shown relative to pseudo wild type (yellow). The green circle highlights the main difference of the potentials of both temperatures. (E) RMSD of the protein and RNA loop backbone nucleotides at both temperatures with respect to the pseudo wild type structure.

Figure 3.

K50Q mutant complex conformations: (A) Even at 298 K, the RNA (blue) unwraps from the protein surface (light blue) and binds in a new conformation, when compared to the representative pseudo wild type structure (gray). (B) After T-jump to 310 K, the RNA (red) also occupies a similar new site on the protein (light red), overlaid on the pseudo wild type structure (gray). In his mutant helix C does not rearrange greatly (black arrow)(C) Electrostatic potential map (± 5 k_BT/e isosurfaces) of representative structures of the complexes at 298 K and (D) 310 K together. The corresponding RNA structure (magenta) is shown relative to pseudo wild type (yellow). The blue circle highlights the main difference of the potentials of both temperatures. (E) The RMSD of the protein and RNA loop backbone nucleotides at both temperatures with respect to the pseudo wild type structure.

Figure 4.

(A) The shortest paths connecting different parts of the RNA to the center of the pseudo wild type protein pass through K22 (orange path) or K50 (green path) residues. (B) These pathways are either completely eliminated as a result of K50Q mutation, or completely rearranged for the K22Q mutation. (C) The pseudo wild type community containing K50 shows extensive connections of loop3 and both sides of the RNA loop, (D) In the K50Q mutant complex, the interconnections between the loop 3 and RNA loop weakens and does not include the unwrapped region of the RNA.

Figure 5.

A, B) For K22Q mutant complex, after the T-jump, the crucial base stacking interactions between the W56-A6 and Y13-C5 are lost. (C, D) The probability distribution of the CoM distance between the side chain of C5, A6 and W56 (probe) at 298 K and 310 K temperatures. (E, F) For K50Q mutant complex, the W56-A6 base stacking interaction remains intact after the T-jump, with the probabilities of both base stacking interactions shown in G,H panels. The green histograms represent the same distances from the F56W simulation, shown as a reference.

Figure 6.

Stopped flow experiments of mutant-RNA complex dissociation at 308 K (Figure S10 shows 298 K data). The initial protein and RNA concentration was 1 μ M. Mixing with high salt buffer switched the KCl concentration from 0.2 M to 1 M to increase electrostatic screening and favor full dissociation (see Materials and Methods for solvent conditions). (A) K22Q and (B) K50Q. The dissociation times τ are both on the order of ~ 0.1 s, similar to the pseudo wild type measured value of 233 ± 2 ms at the same temperature (inset in panel B).¹²

Figure 7.

T-jump of 20 M 1:1 K22Q mutant complex in low salt (0.2 M KCl) favors bound complex, and shows a fast relaxation with time constant τ (A) Jump from 298 K, and (B) Jump from 308 K. The data was logarithmically binned. We assign the signal to a change in the local Trp environment as the complex samples a new binding site at higher temperature.

Figure 8.

T-jump experiment of 20 M 1:1 K50Q-RNA mutant complex starting at 298 K (blue) and 308 K (red). No signal is observed in either case, so the Trp environment is preserved over the temperature range probed.

Figure 9.

Schematic of the free energy landscape for the full dissociation step of U1A-SL2 RNA complex dissociation: Free energies were obtained from the K_d and ϕ value analysis for the complex and transition state, respectively. The positively charged mutants are colored green, the pseudo wild type (with F56W in the active site) is colored black.

Table 1Parameters from ϕ value analysis

Protein	G_{M-WT} (kJ/mole)	$G^{\#}_{M-WT}$ (kJ/mole)	$G (G^{\#}_{M-WT} - G_{M-WT})$ (kJ/mole)	$\phi^{\#}$
K22Q	6.7	4.6	-2.1 (faster)	0.70
K50Q	7.1	5.9	-1.2	0.82