

Published in final edited form as:

J Pediatr Orthop. 2010 March ; 30(2): 106–109. doi:10.1097/BPO.0b013e3181d076a3.

Prevalence of Vitamin D Insufficiency in African American Children with Forearm Fractures: A Preliminary Study

Leticia Manning Ryan, MD^{1,2}, Cinzia Brandoli, PhD³, Robert J Freishtat, MD,MPH^{1,4}, Joseph L Wright, MD,MPH^{1,2,5}, Laura Tosi, MD³, and James M Chamberlain, MD^{1,2}

Children's National Medical Center George Washington University School of Medicine and Health Sciences Washington, DC

¹Division of Emergency Medicine

²Research Center for Clinical and Community Research

³Division of Orthopaedics and Sports Medicine

⁴Research Center for Genetic Medicine

⁵Child Health Advocacy Institute

Abstract

Objectives—Forearm fractures account for a significant proportion of childhood injuries and appear to be increasing in incidence. Poor vitamin D status increases overall fracture risk in infants with rickets and adults with osteoporosis. Children with vitamin D insufficiency (serum 25-hydroxy vitamin D level < 20 ng/mL) have decreased bone mineral density (BMD) compared to children with normal vitamin D status. The relationship between vitamin D status and childhood forearm fracture has not been investigated.

Methods—This prospective study is enrolling African American children, ages 5–9 years, with a forearm fracture. Bone health evaluation includes measurement of serum 25-hydroxy vitamin D level and BMD by dual energy xray absorptiometry (DXA) scan. Univariable analyses are used to test the associations between fracture status and the independent variables serum vitamin D level and BMD.

Results—Vitamin D levels are available for seventeen cases. The mean (\pm SD) 25-hydroxy vitamin D level is 20.1 (\pm 7.3) ng/mL with a range of 10–38 ng/mL. The mean of this group is at the cutpoint for vitamin D insufficiency. Ten cases (59%) were vitamin D insufficient. DXA scan results for these patients are consistent with normal bony mineralization for age.

Conclusions—A significant proportion of African American children with fractures in our study are vitamin D insufficient. Analysis of serum 25-hydroxy vitamin D levels and BMD in additional cases and controls will determine the significance of these findings.

Clinical Relevance—Vitamin D insufficiency may play a previously unrecognized role in childhood fractures. Strong consideration should be given to routine vitamin D testing in African American children with forearm fractures.

Corresponding Author: Leticia Manning Ryan, MD Children's National Medical Center Division of Emergency Medicine 111 Michigan Avenue, NW Washington, DC 20010 Phone: 202-476-4177 Fax: 202-476-3573 lryan@cnmc.org.

The authors have no conflicts of interest related to this article to disclose.

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Keywords

Bone Health; Fractures; Vitamin D

Introduction

Approximately half of all children will fracture a bone during childhood; of those, nearly 40% will sustain at least one additional future fracture.¹ Skeletal fractures account for a significant proportion of childhood injuries^{2–6} and result in substantial costs and morbidity.^{7–9} Despite success in preventing other childhood injuries,¹⁰ fracture rates appear to be increasing.^{11–14}

Prior fracture is associated with an increased risk of subsequent fractures during growth.¹⁵ Thus, there appears to be a vulnerable subset of the population at higher risk for fractures. Pediatric fractures are often attributed to normal childhood development, which makes falls and injuries common;^{16,17} however, there is increasing evidence that such fractures may be related to deficient bone health. Specifically, recent studies suggest that elevated fracture risk in otherwise well children may be a function of lower bone mineralization associated with both environmental factors, including poor nutrition and physical inactivity^{18–22} and genetics.²³

Pediatric studies evaluating the role of bone health in childhood fractures have been primarily conducted in New Zealand and white children;^{19–21} yet, African American children may be at higher risk of bone health deficits, including vitamin D deficiency. Darker skin pigmentation is a risk factor for vitamin D deficiency;²⁴ additionally, African American children have significantly lower dietary intakes of vitamin D compared with other ethnic groups.²⁵ Limited published data reveal a high prevalence (up to 49%) of vitamin D insufficiency [serum 25-hydroxy vitamin D level < 20 ng/mL] in otherwise healthy African American children.²⁶

The relationship between low vitamin D levels and fracture risk has been studied extensively in infants^{27–32} but its significance in children and adolescents has not been investigated. Recently, we have initiated a study of bone health, including environmental and genetic analysis, in African-American children with an isolated forearm fracture to evaluate the relationship between bone health and pediatric forearm fractures. Our objective is to determine the prevalence of vitamin D insufficiency and abnormal bone mineral density in a sample of African American children with forearm fractures.

Materials and Methods

A convenience sample of patients who satisfied the inclusion and exclusion criteria listed in Table 1 was enrolled between December 2005 and December 2006. This prospective study was conducted at Children's National Medical Center (CNMC), an urban pediatric medical center in Washington, DC. It represents a pilot study focusing on the enrollment of forearm fracture patients who will later represent cases in a case-control study. This subsequent case-control study will compare these patients to age and race-matched fracture free controls.

Patients were recruited through the outpatient Orthopaedic Clinic and were studied in the CNMC General Clinical Research Center. This study was approved by the CNMC Institutional Review Board. All participants and/or their guardians provided informed consent and children 7–9 years of age provided assent.

Bone health evaluation includes measurement of serum 25-hydroxy vitamin D level and bone mineral density by dual energy x-ray absorptiometry (DXA) scan. Univariable analyses are used to test the associations between fracture status and the independent variables serum vitamin D

level and BMD. Data analysis was performed with SPSS Statistics 17.0 (SPSS Inc., Chicago, IL).

Results

During the study period, seventeen cases have been enrolled. The mean (\pm SD) case age is 7.3 (\pm 1.3) years; age range 5 to 9 years; 53% are male. None of the patients were observed to adhere to a cultural or religious practice of wearing all enveloping clothing to cover exposed skin.

The mean (\pm SD) 25-hydroxy vitamin D level is 20.1 (\pm 7.3) ng/mL with a range of 10–38 ng/mL. The mean of this group is at the cutpoint for vitamin D insufficiency [serum 25-hydroxy vitamin D level < 20 ng/mL]. Ten cases (59%) were vitamin D insufficient. The proportion of patients with vitamin D insufficiency by season of enrollment is summarized in Table 2.

DXA scan results for all patients are consistent with normal bony mineralization for age (z-score > -2). The mean (\pm SD) total body z-score for study participants is 0.73 (\pm 0.8) with a range of -1.0 to 2.3.

There was no radiographic evidence of osteopenia for any of the patients.

Discussion

Vitamin D is essential for calcium homeostasis and bone development/remodeling.^{33,34} Vitamin D deficiency has long been recognized as a cause of increased fracture risk associated with rickets in infants and osteomalacia in adults.^{33–35} This nutritional deficiency is more likely to occur in infants with darker skin, limited sun exposure, a history of breast feeding without vitamin D supplementation and poor dietary intake of vitamin D or calcium.^{24,36} This deficiency and its effects have been studied extensively in infants^{27–32} but its significance in children and adolescents has not been investigated.

There is an increasing awareness of a preclinical phase of vitamin D deficiency, i.e. vitamin D insufficiency,^{24,37} which increases the risk of osteoporotic fractures in adults.^{38,39} This term describes biochemical evidence of deficiency [serum 25-hydroxy vitamin D level < 20 ng/mL] without obvious clinical signs or symptoms, such as radiographic osteopenia.⁴⁰ There is growing evidence that vitamin D insufficiency is a prevalent health problem. Surveys of other at-risk populations have shown that a significant proportion have vitamin D insufficiency^{41–44} which is associated with diminished bone mineral content.⁴⁴

Vitamin D deficiency and/or insufficiency may also play an important but previously unrecognized role in fractures occurring in the pediatric age group. Few studies have investigated the effect of vitamin D intake, sun exposure, or vitamin D status on bone mineral density in children and young adults. However, in a 3-year prospective study of Finnish girls aged 9 to 15 years, girls with vitamin D deficiency had a 4% lower bone mineral density accumulation from baseline than did girls with normal vitamin D status.⁴⁵ In adults, each standard deviation decrease in bone mineral density approximately doubles fracture risk.⁴⁶

African American children are likely to be at higher risk of vitamin D deficiency and/or insufficiency. African American children have significantly lower daily intake of vitamin D compared with other racial/ethnic groups.²⁵ Adequate intake levels for vitamin D from food are met by less than half of African-American children.²⁵ The higher prevalence of lactose intolerance in this population may also contribute to nutritional deficiencies. In the United States, it is estimated that up to three fourths of African-Americans have symptoms of lactose

intolerance.⁴⁷ In addition, darker skin pigmentation is a risk factor for vitamin D deficiency.²⁴

In our pilot data, 59% of African American children with fractures were vitamin D insufficient. This prevalence is higher than baseline levels of vitamin D insufficiency reported in comparable populations. The effect of these insufficient levels on fracture risk in otherwise healthy children merits further careful evaluation.

Interestingly, our preliminary results showed normal BMD measurements and normal bone appearance on radiographs for forearm fracture patients with and without vitamin D insufficiency. Analysis of serum 25-hydroxy vitamin D levels and bone mineral density in additional cases and controls will determine the significance of these findings. Vitamin D insufficiency may play a previously unrecognized role in childhood fractures. Strong consideration should be given to routine vitamin D testing in African American children with forearm fractures.

Acknowledgments

On the title page, include a paragraph which lists all forms of support, including pharmaceutical and industry backing. If you have not received grants or other aid, write: "None of the authors received financial support for this study."

Paragraph summarizing forms of support/funding: This study is funded in part by the National Institutes of Health National Center for Research Resources (1K23 RR024467-01), Children's National Medical Center Bone Health Program, Children's National Medical Center General Clinical Research Center (5-MO1-RR-020359-02), Children's National Medical Center Board of Visitors and the DC-Baltimore Research Center on Child Health Disparities (5P20MD00165).

References

1. Jones IE, Williams SM, Dow N, et al. How many children remain fracture-free during growth? A longitudinal study of children and adolescents participating in the Dunedin Multidisciplinary Health and Development Study. *Osteoporos* 2002;13:990–995.
2. Waltzman ML, Shannon M, Bowen AP, Bailey MC. Monkeybar injuries: complications of play. *Pediatrics* 1999;103(5) Available at: <http://www.pediatrics.org/cgi/content/full/103/5/e58>.
3. Committee on Injury and Poison Prevention and Committee on Sports Medicine and Fitness. Trampolines at home, school, and recreational centers. *Pediatrics* 1999;103(5):1053–1056. [PubMed: 10224191]
4. Radelet MA, Lephart SM, Rubinstein EN, Myers JB. Survey of the injury rate for children in community sports. *Pediatrics* 2002;110(3) Available at: <http://www.pediatrics.org/cgi/content/full/110/3/e28>.
5. Lyons RA, Delahunty AM, Heaven M, et al. Incidence of childhood fractures in affluent and deprived areas: population based study. *BMJ* 2000;320:149. [PubMed: 10634734]
6. Stark AD, Bennet GC, Stone DH, et al. Association between childhood fractures and poverty: population based study. *BMJ* 2002;324:457. [PubMed: 11859047]
7. Galano GJ, Vitale MA, Kessler MK, et al. The most frequent traumatic orthopaedic injuries from a national pediatric inpatient population. *J Pediatr Orthop* 2005;25:39–44. [PubMed: 15614057]
8. Musgrave DS, Mendelson SA. Pediatric orthopedic trauma: principles in management. *Critical Care Medicine* 2002;30(11 Suppl):S431–43. [PubMed: 12528785]
9. Rodriguez-Merchan EC. Pediatric skeletal trauma: a review and historical perspective. *Clinical Orthopaedics & Related Research* 2005;432:8–13. [PubMed: 15738799]
10. Rivara FP, Grossman DC, Cummings P. Injury prevention. *N Engl J Med* 1997;337:543–548. [PubMed: 9262499]
11. Khosla S, Melton LJ, Dekutoski MB, et al. Incidence of childhood distal forearm fractures over 30 years: a population-based study. *JAMA* 2003;290:1479–1485. [PubMed: 13129988]
12. Landin LA. Fracture patterns in children. *Acta Othop Scand* 1983;54(suppl 202):1–109.

13. Kramhoft M, Bodtker S. Epidemiology of distal forearm fractures in Danish children. *Acta Orthop Scand* 1988;59:557–559. [PubMed: 3188862]
14. Jonsson B, Bengner U, Redlund-Johnell I, Johnell O. Forearm fractures in Malmo, Sweden: changes in the incidence occurring during the 1950s, 1980s, and 1990s. *Acta Orthop Scand* 1999;70:129–132. [PubMed: 10366911]
15. Goulding A, Jones IE, Williams SM, et al. First fracture is associated with increased risk of new fractures during growth. *J Pediatr* 2005;146:286–288. [PubMed: 15689927]
16. Plumert JM. Relations between children's overestimation of their physical abilities and accident proneness. *Developmental Psychology* 1995;31:866–876.
17. Schwebel DC. The role of impulsivity in children's estimation of physical ability: implications for children's unintentional injury risk. *Am J of Orthopsychiatry* 2004;74:584–588. [PubMed: 15554818]
18. Clark EM, Tobias JH, Ness AR, et al. Association between bone density and fractures in children: a systematic review and meta-analysis. *Pediatrics* 2006;117:e291–e297. [PubMed: 16452336]
19. Goulding A, Jones IE, Taylor RW, et al. Bone mineral density and body composition in boys with distal forearm fractures: a dual-energy x-ray absorptiometry study. *J Pediatr* 2001;139:509–515. [PubMed: 11598596]
20. Goulding A, Cannan R, Williams SM, et al. Bone mineral density in girls with forearm fractures. *J Bone Miner Res* 1998;13:143–148. [PubMed: 9443800]
21. Goulding A, Rockell JEP, Black RE, et al. Children who avoid drinking cow's milk are at increased risk for prepubertal bone fractures. *J Am Diet Assoc* 2004;104:250–253. [PubMed: 14760576]
22. Ma D, Jones G. The association between bone mineral density, metacarpal morphometry, and upper limb fractures in children: a population-based case-control study. *J Clin Endocrinol Metab* 2003;88:1486–1491. [PubMed: 12679427]
23. Fischer PR, Thacher TD, Pettifor JM, et al. Vitamin D receptor polymorphisms and nutritional rickets in Nigerian children. *J Bone Miner Res* 2000;15:2206–2210. [PubMed: 11092401]
24. Thomas MK, Demay MB. Vitamin D deficiency and disorders of vitamin D metabolism. *Endocrinol Metab Clin* 2000;29:611–627.
25. Moore CE, Murphy MM, Holick MF. Vitamin D intakes by children and adults in the United States differ among ethnic groups. *J Nutr* 2005;135:2478–2485. [PubMed: 16177216]
26. Rajakumar K, Fernstrom JD, Janosky JE, et al. Vitamin D insufficiency in preadolescent African-American children. *Clin Pediatr* 2005;44:683–692.
27. Pawley N, Bishop NJ. Prenatal and infant predictors of bone health: the influence of vitamin D. *Am J Clin Nutr* 2004;80(suppl):1748S–51S. [PubMed: 15585799]
28. Mylott BM, Kump T, Bolton ML, et al. Rickets in the dairy state. *WMJ* 2004;103:84–87. [PubMed: 15553572]
29. Bloom E, Klein EJ, Shushan D, et al. Variable presentations of rickets in children in the emergency department. *Ped Emerg Care* 2004;20:126–130.
30. Kreiter SR, Schwartz RP, Kirkman HN, et al. Nutritional rickets in African American breast-fed infants. *J Pediatr* 2000;137:153–157. [PubMed: 10931404]
31. Pettifor JM, Isdale JM, Sahakian J, et al. Diagnosis of subclinical rickets. *Arch Dis Child* 1980;55:155–157. [PubMed: 7377839]
32. Spence JT, Serwint JR. Secondary prevention of vitamin D-deficiency rickets. *Pediatrics* 2004;113:e70–72. [PubMed: 14702499]
33. Holick MF. High prevalence of vitamin D inadequacy and implications for health. *Mayo Clinic Proceedings* 2006;81:353–73. [PubMed: 16529140]
34. Holick MF. Vitamin D and bone health. *J Nutr* 1996;126:1159S–1164S. [PubMed: 8642450]
35. Lanou AJ, Berkow SE, Barnard ND. Calcium, dairy products, and bone health in children and young adults: a reevaluation of the evidence. *Pediatrics* 2005;115:736–43. [PubMed: 15741380]
36. Jacobsen ST, Hull CK, Crawford AH. Nutritional rickets. *Journal of Pediatric Orthopaedics* 1986;6(6):713–716. [PubMed: 3793894]
37. Allgrove J. Is nutritional rickets returning? *Arch Dis Child* 2004;89:699–701. [PubMed: 15269061]
38. LeBoff MS, Kohlmeier L, Hurwitz S, et al. Occult vitamin D deficiency in postmenopausal US women with acute hip fracture. *JAMA* 1999;281:1505–1511. [PubMed: 10227320]

39. Chapuy MC, Arlot ME, Duboeuf F, et al. Vitamin D3 and calcium to prevent hip fractures in elderly women. *N Engl J Med* 1992;327:1637–1642. [PubMed: 1331788]
40. Hanley DA, Davison KS. Vitamin D insufficiency in North America. *J Nutr* 2005;135:332–337. [PubMed: 15671237]
41. Al-Jurayyan NA, El-Desouki ME, Al-Herbish AS, et al. Nutritional rickets and osteomalacia in school children and adolescents. *Saudi Med J* 2002;23:182–185. [PubMed: 11938395]
42. Bachrach S, Fisher J, Parks JS. An outbreak of vitamin D deficiency rickets in a susceptible population. *Pediatrics* 1979;64:871–877. [PubMed: 574626]
43. Rucker D, Allan JA, Fick GH, et al. Vitamin D insufficiency in a population of healthy western Canadians. *CMAJ* 2002;166:1517–1524. [PubMed: 12074117]
44. Valimaki VV, Alfthan H, Lehmuskallio E, et al. Vitamin D status as a determinant of peak bone mass in young Finnish men. *J Clin Endocrinol Metab* 2004;89:76–80. [PubMed: 14715830]
45. Lehtonen-Veromaa MKM, Möttönen TT, Nuotio IO, et al. Vitamin D and attainment of peak bone mass among peripubertal Finnish girls: a 3-yr prospective study. *Am J Clin Nutr* 2002;76:1446–1453. [PubMed: 12450915]
46. Genant HK, Cooper C, Poor G, et al. Interim report and recommendations of the World Health Organization Task-Force for Osteoporosis. *Osteoporos Int* 1999;10:259–264. [PubMed: 10692972]
47. Byers KG, Savaiano DA. The myth of increased lactose intolerance in African-Americans. *Journal of the American College of Nutrition* 2005;24(6 Suppl):569S–73S. [PubMed: 16373956]

Table 1

Inclusion and exclusion criteria for study participants

Inclusion Criteria	Exclusion Criteria
<ul style="list-style-type: none"> • African American/black race per parental designation • Age 5–9 • Orthopedic Clinic patient • Isolated and radiographically demonstrated forearm fracture (radius, ulna, or both) • Parent/guardian fluent in English language 	<ul style="list-style-type: none"> • Current underlying bone mineralization disorder (osteomalacia, osteogenesis imperfecta) • Current or prior use of anti-epileptic medication

Table 2

Proportion of enrolled patients with vitamin D insufficiency by season of enrollment

Season of patient enrollment	Proportion of enrolled patients with vitamin D insufficiency (%)
Winter	1/1 (100%)
Spring	6/9 (66.7%)
Summer	1/2 (50%)
Fall	2/5 (40%)