

Published in final edited form as:

J Phys Chem B. 2010 April 15; 114(14): 4896–4903. doi:10.1021/jp9095375.

Reaction Rates and Mechanism of the Ascorbic Acid Oxidation by Molecular Oxygen Facilitated by Cu(II)-Containing Amyloid- β Complexes and Aggregates

Dianlu Jiang, Xiangjun Li, Lin Liu, Gargey B. Yagnik, and Feimeng Zhou *

Department of Chemistry and Biochemistry, California State University, Los Angeles, Los Angeles, California 90032

Abstract

A forefront of the research on Alzheimer's disease (AD) is the interaction of amyloid beta ($A\beta$) peptides with redox metal ions (e.g., Cu(II), Fe(III) and Fe(II)) and the biological relevance of the $A\beta$ -metal complexes to neuronal cell loss and homeostasis of essential metals and other cellular species. This work is concerned with the kinetic and mechanistic studies of the ascorbic acid oxidation reaction by molecular oxygen that is facilitated by Cu(II) complexes with $A\beta$ (1–16), $A\beta$ (1–42), and aggregates of $A\beta$ (1–42). The reaction rate was found to linearly increase with the concentrations of $A\beta$ -Cu(II) and dissolved oxygen, and be invariant with high ascorbic acid concentrations. The rate constants were measured to be 117.2 ± 15.4 and $15.8 \pm 2.8 \text{ M}^{-1}\text{s}^{-1}$ at low ($<100 \mu\text{M}$) and high AA concentrations, respectively. Unlike free Cu(II), in the presence of AA, $A\beta$ -Cu(II) complexes facilitate the reduction of oxygen by producing H_2O_2 as a major product. Such a conclusion is drawn on the basis that the reaction stoichiometry between AA and O_2 is 1:1 when $A\beta$ concentration is kept at a much greater value than that of Cu(II). A mechanism is proposed for the AA oxidation in which the oxidation states of the copper center in the $A\beta$ complex alternates between 2+ and 1+. The catalytic activity of Cu(II) towards O_2 reduction was found to decrease in the order of free Cu(II) $>$ $A\beta$ (1–16)-Cu(II) $>$ $A\beta$ (1–42)-Cu(II) $>$ Cu(II) complexed by the $A\beta$ oligomer/fibril mixture $>$ Cu(II) in $A\beta$ fibrils. The finding that Cu(II) in oligomeric and fibrous $A\beta$ aggregates possesses considerable activity towards H_2O_2 generation is particularly significant, since in senile plaques of AD patients the co-existing copper and $A\beta$ aggregates have been suggested to inflict oxidative stress through the production of reactive oxygen species (ROS). Although Cu(II) bound to oligomeric and fibrous $A\beta$ aggregates is less effective than free Cu(II) and the monomeric $A\beta$ -Cu(II) complex in producing ROS, in vivo the Cu(II)-containing $A\beta$ oligomers and fibrils might be more biologically relevant given their stronger association with cell membranes and the closer proximity of ROS to cell membranes.

Introduction


The pathological hallmarks of Alzheimer's disease (AD) include the formation of senile plaques composed of mainly amyloid- β ($A\beta$) variants of 40–43 residues, a dramatic loss of neuronal cells, and extensive oxidative stress.^{1–3} More intriguing is the fact that inside the senile plaques high levels of metals are accumulated.^{4–6} Consequently, recent years have seen an enormous interest in the interaction between metal and $A\beta$ and the possible roles of the metal- $A\beta$ complexes in the AD pathogenesis.^{7–19} Although it is still not clear how extensive oxidative stress in AD-afflicted brain occurs, it is becoming evident that the redox active metals (e.g., Cu(II) and Fe(III)) are involved in the production of reactive oxygen species (ROS) and

*Corresponding author. Phone: 323-343-2390. Fax: 323-343-6490. fzhou@calstatela.edu.

the aggravation of oxidative stress.^{3,19–21} Hence, in order to probe the role of metals and A β , a key issue is to understand the redox properties of the complexes of metals with A β and possible redox reactions of these complexes with biological species. Our group,^{16,22} among others,^{14,15} have systematically elucidated the redox properties of A β complexes with copper and iron. Findings from these efforts have improved our understanding of the metal-initiated production of ROS.

It has been well documented that A β binds copper with its hydrophilic domain (residues 1–16) through the three histidine residues at positions 6, 13, and 14.⁷ Other binding sites proposed for the Cu(II) binding include the amine and carboxylate groups of aspartic acid at the N-terminus, the tyrosine residue at position 10 and other carboxylates in the hydrophilic domain (For a comprehensive review please see Ref. 7). At least two binding modes with varying binding affinities have been proposed, and investigations on the detailed binding modes have continued.^{8,9,13,23–26} The binding (affinity) constants reported thus far have ranged from micromolar to attomolar,^{10,26–31} with most values lying between submicromolar and nanomolar.^{10,26,29,31} Such a large variation has been suggested to be a result of the differences in the concentrations of A β and Cu(II), buffer systems, A β sample preparations, and even data interpretations. The impact of these differences on the measurement of the binding affinity and stoichiometry between Cu(II) and A β has been reviewed and rationalized by recent reviews.^{7,13} Despite these conflicting results, it is generally agreed that Cu(II) binds to A β strongly and the resultant complexes are possibly related to oxidative stress (either as a prooxidant or an antioxidant or a dual role of prooxidant and antioxidant).^{12–19}

In vitro, the A β -Cu(II) complex, in the presence of an electron donor (reductant), has been shown to reduce O₂ to produce H₂O₂.^{15,32} Recently, we successfully measured the redox potential of the complex and suggested that thermodynamically a number of biological species are capable of reducing A β -Cu(II) to A β -Cu(I).¹⁶ A β -Cu(I) in turn can reduce O₂ and generate H₂O₂. Overall, the net effect is that A β -Cu(II) facilitates the oxidation of a biological species and the reduction of molecular oxygen, generating H₂O₂ as a ROS, as shown by reaction (3).



where Red and Ox represent the reduced and oxidized forms of a biological species, respectively. For reductant that can receive two electrons (e.g., ascorbic acid), reaction (3) becomes $\text{Red} + \text{O}_2 + 2\text{H}^+ \rightarrow \text{O}_x + \text{H}_2\text{O}_2$ or the stoichiometry between Red and O₂ is 1:1. Under normal circumstance without A β -Cu(I), reaction (3) would proceed very slowly, imposing no significant effect on the normal functions of cellular species. In contrast, in the presence of A β -Cu(I), reaction (3) may be dramatically accelerated, and both the biological species and molecular O₂ would be depleted, adversely impacting many biological processes.^{16,33}

The mixture of A β and Cu(II) (in many studies containing both complexed and free Cu(II)) has been reported to accelerate the oxidation of catechol and dopamine by H₂O₂.^{34,35} In this aspect, the mixture was proposed to act like an enzyme (or catalyst) for the oxidation reactions of these species. Ascorbic acid (AA), present as ascorbate anion at physiological pH, is a broad spectrum antioxidant (reductant) that acts against many ROS, and is present abundantly in

central nerve system (ranging from 500 μM in cerebrospinal fluid to 10 mM in neurons).³⁶ It has been reported that AD patients have a significantly lower antioxidant capacity in cerebral fluid than control subjects.³⁷ On the basis of the redox potentials of AA and the $\text{A}\beta\text{-Cu(II)}$ complex,¹⁶ its reaction with $\text{A}\beta\text{-Cu(II)}$ is thermodynamically favored. As such, its oxidation facilitated by $\text{A}\beta\text{-Cu(II)}$, similar to that shown in reaction (3), ought to occur. Given its biological importance and the implication of $\text{A}\beta\text{-Cu(II)}$ to the pathogenesis of AD, the study of the oxidation kinetics should provide insight into the role of $\text{A}\beta\text{-Cu(II)}$ in the development of AD.

Oxygen is also abundant in brain and is vital for a variety of brain functions.²⁰ It has been assumed that the complex formed between Cu(II) and $\text{A}\beta(1\text{--}42)$ participates in the reduction of molecular oxygen to hydroxyl radicals through the route $\text{O}_2 \rightarrow \text{O}_2^{\cdot-} \rightarrow \text{H}_2\text{O}_2 \rightarrow \text{OH}^{\cdot}$.¹² A recent theoretical paper by Hewitt and Rauk shows that the conversion of O_2 to H_2O_2 in the presence of the reduced form of $\text{A}\beta\text{-Cu(II)}$ (i.e., $\text{A}\beta\text{-Cu(I)}$) does not have to proceed via the superoxide $\text{O}_2^{\cdot-}$ intermediate.³² Results recently obtained by Baruch-Suchodolsky and Fischer¹² and Faller's group¹⁴ have suggested that $\text{A}\beta$ peptides (especially $\text{A}\beta(1\text{--}40)$ and $\text{A}\beta(1\text{--}42)$) may serve as scavengers of OH^{\cdot} . In the work by Baruch-Suchodolsky and Fischer, the $\text{A}\beta\text{-Cu(I)}$ complex was formed between $\text{A}\beta$ and Cu(I) without first reducing $\text{A}\beta\text{-Cu(II)}$ through a reductant.^{12,38} Interestingly, Guilloreau et al. showed that the amount of OH^{\cdot} generated by $\text{A}\beta(25\text{--}35)$, the segment that does not contain the Cu(II) binding domain but possesses the easily oxidizable methionine residue at position 35, is much greater than that produced by $\text{A}\beta(1\text{--}16)$ or $\text{A}\beta(1\text{--}42)$.¹⁴ This observation suggests that Cu(II) complexed by $\text{A}\beta(1\text{--}16)$ or $\text{A}\beta(1\text{--}42)$ is less capable than free Cu(II) of generating OH^{\cdot} (i.e., more capable of inhibiting OH^{\cdot} generation). Although many studies have focused on the production of H_2O_2 from O_2 ,^{16,32} it is not certain whether H_2O_2 produced with the aid of the $\text{A}\beta\text{-Cu(I)}$ complex can further decompose to OH^{\cdot} .

The oxidation kinetics of AA has been extensively investigated employing both free Cu(II) or Cu(II) complexes.^{39–41} Depending on the ligands coordinating the Cu(II) center, the kinetics and mechanisms of the AA oxidation vary significantly. For example, when Cu(II) is coordinated by poly-4-vinylpyridine, its catalytic activity toward the AA oxidation increases by four orders of magnitude.⁴² The copper(II)-catalyzed AA oxidation, in the presence or absence of strong ligands, is generally believed to proceed by an alternating reduction-oxidation mechanism with the formation of an intermediate of cuprous (Cu(I)) species.^{42,43} Nonetheless, AA oxidation catalyzed by histidine (His) ligated copper proceeds via a different mechanism in which a ternary complex of $\text{O}_2\text{-(His-Cu)-AA}$ is formed.⁴⁰ Furthermore, although in the presence of an electron donor and $\text{A}\beta\text{-Cu(II)}$ H_2O_2 has been detected and the reaction has been proposed to proceed along the pathway outlined by reactions (1)–(2), it is not clear if H_2O_2 is the exclusive product or if other ROS are also generated. To elucidate the potential involvement of $\text{A}\beta\text{-Cu(II)}$ in the AD pathogenesis, it is necessary to further investigate the mechanism of oxygen reduction facilitated by the $\text{A}\beta\text{-Cu(II)}$ complex in a milieu similar to biological conditions.

To the best of our knowledge, the rate of the $\text{A}\beta\text{-Cu(II)}$ -facilitated AA oxidation by O_2 has not been determined and the dependence of this rate on the concentrations of $\text{A}\beta$, Cu(II) , AA, and O_2 has not been thoroughly assessed. Furthermore, although it has been shown that $\text{A}\beta$ fibrils bind to Cu(II) as strongly as $\text{A}\beta$ monomers,²⁵ the ability of the Cu(I) -containing fibrils in scavenging OH^{\cdot} appears to be diminished with respect to that of the complex between $\text{A}\beta$ monomer and Cu(I) .¹² We are therefore also interested in whether the aggregation of $\text{A}\beta$ in the presence of Cu(II) may affect the rate of H_2O_2 production. In this work, we investigated the effect of $\text{A}\beta\text{-Cu(II)}$ on AA oxidation by molecular oxygen, the oxidation reaction kinetics and the stoichiometry of the reaction. The differences in reactivity and mechanism between free Cu(II) and $\text{A}\beta\text{-Cu(II)}$ are compared and discussed. Such a study sheds light onto the

possible role of A β as a neuroprotective agent or a toxin, an ambiguity that has long been under debate.^{12,44,45} We also compared the activity of A β -Cu(II) in monomeric and aggregate forms in facilitating the O₂ reduction to gain a better understanding of their relative toxicities in terms of their abilities to deplete biological species and to produce hydrogen peroxide. With the rate constant known, we should be able to learn how rapidly important biological redox species are consumed in the A β -Cu(II)-facilitated O₂ reduction reaction.

MATERIALS AND METHODS

Materials

Lyophilized A β (1–16) (DAEFRHDSGYEVHHQK) was synthesized by Genemed Synthesis (San Antonio, TX), and purified in-house using HPLC. A β (1–42) (DAEFRHDSGYEVHHQKLVFFAEDVGSNKGAIIGLMVGGVVV) was purchased from American Peptide Co. Inc. (Sunnyvale, CA). All other chemicals were of analytical grade (Sigma-Aldrich). Throughout the work, 1 mM CuCl₂ dissolved in 1 mM H₂SO₄ was used as the Cu(II) stock solution. A β (1–16) samples were prepared freshly by dissolving lyophilized powder samples in deionized water, while A β (1–42) samples were dissolved in dimethyl sulfoxide (DMSO) and subsequently diluted with phosphate buffer to desired concentrations. The AA stock solution (10 mM) was prepared before use.

Phosphate buffer (10 mM, pH 7.2) was employed for the kinetic experiments and preparations of A β (1–42) aggregates with or without Cu(II). The buffer was prepared with deionized water (18 M Ω ·cm⁻¹), followed by being passed through a Chelex 100 column (Bio-Rad Laboratories, Inc, Hercules, CA) to rid of the trace heavy metals. Using inductively coupled plasma mass spectrometry (ICP-MS), we found that concentrations of heavy metal ions after such a treatment were below 1 nM.

Kinetic Measurements

The AA oxidation was followed by UV-vis spectrophotometry (Cary 100 Bio, Varian Inc., Palo Alto, CA). The kinetic experiments were conducted at 25 °C in phosphate buffer (pH, 7.2). At this pH, AA exists predominately in the ascorbate monoanion form and throughout this work we use AA interchangeably to represent the ascorbate monoanion.⁴⁰ The initial oxidation rates of AA were measured from the decrease of absorbance at 265 nm, where AA shows a maximum absorption ($\epsilon = 1.5 \times 10^4 \text{ M}^{-1} \text{ cm}^{-1}$).⁴⁰ Within the initial 100 s, the absorbance decreases linearly with time. The initial reaction rate (V_0) is therefore taken as $m/\epsilon l$ (where m is the slope of the linear portion of the curve, ϵ the ascorbate absorption coefficient at 265 nm, and l the optical path length of the cuvette).

Under our experimental conditions, the autooxidation rate of AA without A β -Cu(II) was found to be always $< 0.1 \text{ nM s}^{-1}$. The oxygen contents in the sample were controlled by equilibrating the sample solution with N₂-O₂ gas mixtures of different compositions. Introducing the gas mixture into the solution was performed in a glove box (Plas Lab Inc., Lansing, MI) that was maintained under a N₂ atmosphere with a O₂ content less than 1 ppm. The N₂ and O₂ flows in the gas mixtures were regulated by mass flow controllers (Hastings Instruments, Hampton, VA) before being mixed in a stainless chamber and introduced into the solution. Prior to each experiment, the initial oxygen concentration was measured by a dissolved oxygen membrane electrode (080515MD, Thermo Fisher Scientific, Pittsburgh, PA). The A β -Cu(II) complex was freshly prepared by adding Cu(II) to an A β (i.e., either A β (1–16) or A β (1–42)) solution. In order to minimize the free Cu(II) concentration, unless otherwise stated, the [A β]/[Cu(II)] ratio was always kept at 20/1. At such a ratio, even if the binding constant between A β and Cu(II) were in the sub- μM range,^{10,26,29,31} copper would still exist predominantly in the form of a complex. We should emphasize that any uncomplexed Cu(II) can catalyze the reduction of

H₂O₂ to generate OH[•] (via the Harber–Weiss-like mechanism) as shown by the recent work of Guilloureau et al.¹⁴ Therefore, keeping the free Cu(II) concentration at the minimum by using a high [Aβ]/[Cu(II)] ratio is critical for avoiding such a complication. The Cu(II)-containing Aβ(1–42) aggregates were prepared by either preincubating Aβ(1–42) solution at 37 °C for a desired amount of time and then mixing with Cu(II) or incubating an Aβ(1–42)/Cu(II) mixture. In kinetic experiments, the abovementioned Aβ/Cu(II) mixture was spiked with AA and mixed with buffer equilibrated with a N₂-O₂ gas mixture.

Electrochemical Measurements

The electrochemical experiments were performed on a CHI 411 electrochemical workstation (CH Instruments, Austin, TX) using a homemade plastic electrochemical cell. A glassy carbon disk electrode and a platinum wire were used as the working and counter electrodes, respectively. The reference electrode was Ag/AgCl, and all of the potential values are reported with respect to this electrode unless otherwise stated. Prior to each experiment, the glassy carbon electrode was polished with diamond pastes of 15 and 3 μm and alumina pastes of 1 and 0.3 μm in diameter (Buehler, Lake Bluff, IL). The electrolyte solution was a phosphate buffer containing 0.1 M NaNO₃. The Aβ(1–16) stock was diluted with the same phosphate buffer to desired concentrations. In order to examine the reaction between AA and Aβ-Cu(II) without the complication from oxygen, the experiments were conducted in the same glove box under a N₂ atmosphere.

Atomic force microscopy

AFM images were obtained on an MFP-3D-SA microscope (Asylum Research, Santa Barbara, CA) equipped with a tapping mode. Lyophilized Aβ(1–42) was dissolved in 20 mM NaOH solution to prepare 1 mM stock solution. To generate Cu(II)-containing Aβ(1–42) aggregates, 200 μM Aβ(1–42) with or without 20 μM Cu(II) was incubated at 37 °C for a desired period of time. Aliquots of Aβ(1–42) or Aβ(1–42)/Cu(II) were taken out at predetermined incubation times, cast onto newly peeled mica, and left in contact with the mica substrate for 15 min. Thereafter, the slides were gently rinsed with water to remove any residual salt, and dried with nitrogen before imaging.

Results

Dependence of the AA Oxidation Rate on Its Concentration

Figure 1 shows that within 10–240 μM, the initial rate of AA or ascorbate oxidation varies with its concentration when 10 μM Aβ-Cu(II) and 240 μM O₂ are also present. The rate increases with [AA] almost linearly below 100 μM. As [AA] increases, the rate reaches a steady state. Although the rate-concentration dependence is characteristic of an enzymatic reaction, additional experimental results (*vide supra*) suggest a different reaction mechanism, which will be detailed in the Discussion Section.

Dependence of the AA Oxidation Rate on the Oxygen Concentration




Figure 2 shows the variation of the rate of AA oxidation with the oxygen concentration when [AA] and [Aβ-Cu(II)] were fixed at 240 and 10 μM respectively. Notice that we chose an initial AA concentration that far exceeds the upper limit of the linear range in Figure 1 so that AA is not the limiting reagent. Within 4.6–1170.1 μM [O₂], the AA oxidation rate was found to increase linearly with [O₂]. This contrasts the observation of a plateau at higher [O₂] when the histidine-Cu(II) complex was used as the catalyst,⁴⁰ suggesting that a different mechanism is at work.

Dependence of the initial AA oxidation rate on [Aβ-Cu(II)]

To further probe the AA oxidation or O₂ reduction mechanism, we investigated the effect of [Aβ-Cu(II)] on the AA oxidation rate. As shown in Figure 3, the initial AA oxidation rate also exhibits a linear relationship with the amount of [Aβ-Cu(II)] introduced to the sample solution. Notice that the solutions studied were saturated with air and contained a high [AA] so that these reactants are again not the limiting reagents.

The major product of oxygen reduction by AA that is facilitated by Aβ-Cu(II) is H₂O₂

The reduction of O₂ can form either H₂O or H₂O₂ or both. In most studies,^{40,43} even in the presence of a catalyst,^{41,43} H₂O₂ is assumed to be the product. However, usually, free Cu(II) catalyzes the decomposition of H₂O₂ as follows:^{12,46}



The above reactions have been shown to be relevant to the inhibition of OH• through the Cu(II) complexation by Aβ.¹⁴ Reactions (4)–(5) are essentially the Harber–Weiss reaction^{12,46} producing the hydroxyl radical OH• and reaction (6) is basically the Fenton reaction.⁴⁷ Thus, free Cu(II) decomposes H₂O₂ and produces the highly reactive hydroxyl radical.

For the reduction of oxygen by AA, it would be interesting to observe if the reaction is affected by the coordination of the metal ion. Along this line, we probed the reaction stoichiometry of the oxygen reduction catalyzed by free Cu(II) and compared it to that facilitated by the Aβ-Cu(II) complex. In a gastight cuvette that contained a solution with known amounts of O₂ and Aβ-Cu(II) and excess AA, the AA absorbance was monitored (Figure 4). As soon as oxygen was depleted, the reaction stopped. By calculating the amount of AA consumed, we obtained the stoichiometry of the oxidation reaction. In the presence of Aβ-Cu(II), the molar ratio between the consumed AA and O₂ is always close to 1:1. This suggests that H₂O₂ is the final product of the reaction $\text{AA} + \text{O}_2 + 2\text{H}^+ \rightarrow \text{dehydroascorbate} + \text{H}_2\text{O}_2$. In line with the results reported by us¹⁶ and others,³² Aβ-Cu(II), upon reduction by a reductant (e.g., AA), can accelerate the reduction of oxygen to H₂O₂ instead of H₂O. In sharp contrast, when only free Cu(II) was used, the ratio was found to vary between 1.5 and 2. Such a ratio is indicative of hydroxyl radical formation (or H₂O production from the four-electron reduction of O₂), an observation consistent with that reported by Faller's group.¹⁴

Aβ-Cu(II) can oxidize ascorbic acid

It has been reported that His-Cu(II) catalyzes the oxidation of AA by oxygen via the formation of a ternary intermediate among O₂, His-Cu(II) and AA.⁴⁰ This contrasts the alternating reduction and oxidation mechanism that involves Aβ-Cu(II) and Aβ-Cu(I). The absence of the His-Cu(II) reduction when the complex was mixed with AA was taken as the proof of the mechanism.⁴⁰ Interestingly, Aβ-Cu(II) has a higher oxidation potential than AA,^{14,16} and consequently should be able to convert AA to dehydroascorbate. Determining whether and how fast the reaction occurs will help pinpoint the exact reaction mechanism. To this end, we collected cyclic voltammograms of Aβ-Cu(II) in the absence and presence of AA.




The CV of A β -Cu(II) (red curve) exhibits a pair of oxidation ($E_{pa} = 0.20$ V in Figure 5) and reduction ($E_{pc} = -0.05$ V) peaks. This observation is consistent with our previously reported voltammograms.¹⁶ Specifically, A β -Cu(II) is reduced to A β -Cu(I) which can be reversibly oxidized back to A β -Cu(II). When an equivalent amount of AA was added, the characteristic irreversible AA oxidation peak was not observed (blue curve) and only the CV of A β -Cu(I) appeared. This indicates that within seconds the redox reaction between A β -Cu(II) and AA was completed. Further addition of AA generated a CV showing the AA oxidation peak at 0.37 V (black curve), which is slightly more positive than that of AA only (0.30 V; see also the inset of Figure 5). The oxidation peak is resulted mainly from the excessive and unoxidized AA. A careful examination of the initial scan of the CV shown as the black curve reveals a small shoulder peak at ~ 0.20 V, suggesting that A β -Cu(I) has been generated. The slight shift of the AA oxidation peak and the peak broadening may result from the contribution of the A β -Cu(I) oxidation.

O₂ reduction rates in the presence of Cu(II) complexes formed with A β peptides of different lengths and with different aggregates

We¹⁶ and others^{15,21} have shown that H₂O₂ can be measured as a stable product from the A β -Cu(II)-facilitated O₂ reduction by a biological reductant. By comparing A β (1–16)-Cu(II), A β (1–28)-Cu(II), and A β (1–42)-Cu(II), we have also shown that the amount of H₂O₂ produced decreases with the peptide length. The coordination environment of Cu(II) was attributed to the difference in the reactivity. In AD-afflicted brain, Cu(II) is bound to A β variants of 40–43 residues and accumulated in senile plaques.⁴⁸ Thus far, even though it is evident that A β aggregates can sequester Cu(II) essentially as strongly as the A β monomer,^{25,30} it is not clear if Cu(II) complexed by different A β aggregates can still facilitate the oxygen reduction reaction in the presence of AA. Gaining insight into this question should help unravel the role of metal-containing A β aggregates in inflicting oxidative stress. To this end, we compared the reaction rates of the AA oxidation reaction in the presence A β (1–16)-Cu(II), A β (1–42)-Cu(II) and Cu(II) bound by a mixture of A β (1–42) oligomers and fibrous aggregates. Shown in Figure 6 is the initial AA oxidation rates observed in the presence of these Cu(II)-containing species. Not surprisingly, free Cu(II) has the highest catalytic activity, and the coordination environment rendered by A β (1–16) and A β (1–42) (labeled as “A β (1–16)-Cu(II)” and “A β (1–42)-Cu(II)” in Figure 6, respectively) indeed has a significant influence on the activity of the respective complex. The decrease of reactivity with peptide length should be due to the aforementioned steric hindrance to the accesses of Cu(II) center by oxygen.¹⁶ Interestingly, Cu(II) bound by A β (1–42) aggregates (cf. bars labeled as “A β (1–42)-Cu(II), 48 h” and “A β (1–42) (48 h) + Cu(II)” in Figure 6, for example) can still facilitate the AA oxidation by O₂, though the reaction rate is much slower. The morphologies of incubated A β (1–42) are shown in Figure 7. At the beginning of the incubation, only a few globular aggregates of diameters around 5–8 nm (i.e., oligomers in images A and C) were present, whereas at 48 h of incubation protofibrils and fibrils coexist with some oligomers (images B and D). Adding Cu(II) into the A β (1–42) monomers followed by incubation (image B) or into a preincubated A β (1–42) solution (image D) produced similar aggregate mixtures. Incubation of A β (1–42) in the presence of Cu(II) results in the production of only a few amorphous aggregates (cf. the marked particle in image B whose height is larger than 5–8 nm). This is expected since the A β (1–42) concentration used for the experiments is much higher than the Cu(II) concentration. After 72 h incubation, the population of fibrous aggregates for both types of incubation becomes even greater with little oligomers and protofibrils discernable (data not shown). It is evident from Figure 6 that the fibrous aggregates, similar to the A β monomer, can also attenuate the ability of Cu(II) towards the AA oxidation. Given that free Cu(II) has a substantially higher catalytic activity, the lower activity in the incubated A β samples that was later spiked with Cu(II) suggests that the Cu(II) added must be sequestered (complexed) by the preformed aggregates. This is consistent with the finding reported by Karr and Szalai.²⁵

Discussion

As mentioned above, at least two mechanisms have been proposed for the oxidation of AA by molecular oxygen that is facilitated by Cu(II) complexed with different molecules.^{12,38–40} The formation of a ternary complex of O₂-Cu(II) complex-AA is one mechanism wherein Cu(II) is complexed with a ligand such as histidine. Since AA can readily reduce Aβ-Cu(II) to Aβ-Cu(I) (cf. Figure 5) and the linear dependence of reaction rate on oxygen concentration is not characteristic of such a mechanism, we propose the following mechanism:



where AA^{•−} and DA represent the ascobyl radical and dehydroascorbate, respectively. The overall reaction is determined by the turn-over rate of the Aβ-Cu(I), viz., reactions (7) and (8).

The rate of AA oxidation equals the rate of oxygen consumption once a steady state is established:

$$-\frac{d[AA]}{dt} = k_1[A\beta - Cu(II)][AA] = -\frac{d[O_2]}{dt} k_2[A\beta - Cu(I)][O_2] \quad (11)$$

When the reaction reaches a steady state, [Aβ-Cu(I)] becomes constant, i.e.,

$$\frac{d[A\beta - Cu(I)]}{dt} = k_1[AA][A\beta - Cu(II)] - k_2[A\beta - Cu(I)][O_2] = 0 \quad (12)$$

Based on the mass balance, we have

$$[A\beta - Cu(II)] + [A\beta - Cu(I)] = [Cu]_0 \quad (13)$$

where [Cu]₀ stands for the initial (formal) Cu(II) concentration.

Rearranging eq. 12 gives:

$$\frac{[A\beta - Cu(II)]}{[A\beta - Cu(I)]} = \frac{k_2[O_2]}{k_1[AA]} \quad (14)$$

At low [AA], little [Aβ-Cu(II)] is converted to [Aβ-Cu(I)] and therefore [Aβ-Cu(II)] approaches [Cu]₀. Consequently from eq. 12, the reaction rate is approximately equal to

$k_1[AA][Cu]_0$, and is independent of $[O_2]$. The rate constant k_1 for reaction (7) was deduced to be $117.2 \pm 15.4 \text{ M}^{-1}\text{s}^{-1}$ from the linear portion of curve in Figure 1. At high $[AA]$, all of the $[A\beta\text{-Cu(II)}]$ has been converted to $[A\beta\text{-Cu(I)}]$ which can be approximated to $[Cu]_0$, and the rate of reaction equals $k_2[Cu]_0[O_2]$. The rate constant k_2 was therefore estimated to be $15.8 \pm 2.8 \text{ M}^{-1}\text{s}^{-1}$ from Figure 2. The rate constant indeed shows that the turn-over of $A\beta\text{-Cu(I)}$ is the limiting step, and k_2 is about 7.4 times slower than k_1 . However, the comparability in kinetics of the two steps leads to the plateau in the rate- $[AA]$ relationship, reflecting the transition from the kinetics governed by step 1 to that controlled by step 2. The proposed mechanism explains all of the kinetic data extremely well. We should note that the rate of oxidation of ascorbate catalyzed by ascorbic acid oxidase was measured to be $1.7 \times 10^4 \text{ M}^{-1}\text{s}^{-1}$ previously.⁴⁹ Thus, compared to that catalyzed by the enzyme molecule of ascorbic acid oxidase, the reaction rate of AA oxidation by O_2 that is facilitated by $A\beta\text{-Cu(II)}$ is moderately high.

Our study demonstrates that the catalytic activity of $Cu(II)$ towards the reduction of O_2 is diminished upon its binding to $A\beta$ and is further decreased once $Cu(II)$ is incorporated in $A\beta$ aggregates. That complexation of $Cu(II)$ attenuates the ROS production is consistent with findings from previous studies.¹⁴ Recently, Fischer and coworker reported that free hydroxyl radicals are produced when $Cu(I)$ is directly complexed with $A\beta(1-28)_{38}$ and $A\beta(1-40)_{12}$ (instead of being created by the $A\beta\text{-Cu(II)}$ reduction). They also showed that $A\beta(1-40)$ is more efficient than $A\beta(1-28)$ in scavenging hydroxyl radicals through the oxidation of histidine residues that are bound to $Cu(I)$. Guilloreau et al. studied the reaction between O_2 and $A\beta\text{-Cu(II)}$ complexes in the presence of AA and found that hydroxyl radicals were produced but the amounts are less than the counterpart without $Cu(II)$ complexation.¹⁴ Murray et al. found that oxidative damage of lipid membranes is promoted by the $A\beta\text{-Cu(II)}$ complexes and AA, possibly through the hydroxyl radical generation.⁵⁰ We noticed that in these studies the $A\beta$ and $Cu(II)$ or $Cu(I)$ concentrations were typically at micromolar to sub-micromolar levels and maintained close to 1:1 stoichiometric ratio. As aforementioned, studies by us²⁶ and others^{10,29,31} have suggested that the binding affinity constant between $A\beta$ and $Cu(II)$ is between submicromolar to nanomolar. In fact, the mass spectra we collected using 1:1 stoichiometric ratio between $A\beta$ and $Cu(II)$ (both at micromolar concentrations) always showed the co-existence of free $A\beta$ and $A\beta\text{-Cu(II)}$.¹⁶ The observation of the free $A\beta$ peaks suggests that some $Cu(II)$ ions have remained free in the solution. Therefore, when $A\beta$ is not in substantial excess to $Cu(II)$, the free $Cu(II)$ in solution could continuously react with H_2O_2 produced by Reactions (1)–(2) to generate OH^\bullet via the Harber–Weiss reaction. In the case wherein $Cu(I)$ is directly complexed by $A\beta$, the excess $Cu(I)$ remaining free in solution could further react with H_2O_2 through the Fenton-like reaction to afford OH^\bullet . Therefore, these different results in the literature do not contradict our finding.

From our work it is also evident that the $Cu(II)$ center in $A\beta$ aggregates is still accessible by AA and O_2 . This is not entirely surprising since several reports have shown that the hydrophilic segment of $A\beta$, which is also the metal binding domain, populates the exterior of the aggregates.^{51–53} A study recently performed by Karr and Szalai has clearly demonstrated that aggregation of $A\beta$ does not affect its binding towards $Cu(II)$.²⁵ As for the decreased rate in O_2 reduction after the formation of higher-ordered $A\beta$ aggregates, we presume that the compact packing of the $A\beta$ aggregates must have created a $Cu(II)$ coordination environment that is different from the unstructured $A\beta$ monomer. A more compact structure should render greater steric hindrance to the access of O_2 and other solution species to the $Cu(II)$ center. Thus, it is conceivable that the activity of the $Cu(II)$ center will decrease as the aggregates become larger in size or higher in the packing order (e.g., monomer \rightarrow oligomers \rightarrow protofibrils \rightarrow fibrils⁴⁸). Nonetheless the reaction kinetics is still at least an order of magnitude higher than that of the autooxidation rate of AA ($< 1 \times 10^{-10} \text{ M s}^{-1}$ in the absence of redox metal ions⁴⁰).

Our finding has a significant biological relevance and may offer some insight into the role of A β in the AD etiology. A β appears to inhibit the production of the highly reactive hydroxyl radical from the O₂ reduction by complexing Cu(II) and producing H₂O₂ (a less reactive ROS). In this regard, our results argue for the protective role of A β . However, the in vivo situation is more complicated. First, in healthy brain there is little free cellular Cu(II)⁵⁴ and A β concentration is also very low (nM or less)⁵⁵. Several papers have suggested that only at low concentrations does A β become effective in alleviating oxidative stress.^{12,50,56} However, if the binding affinity between A β and Cu(II) were indeed at the sub-micromolar to nanomolar level,^{10,26,29,31} high concentrations of Cu(II) resulted from impaired metal homeostasis would not be adequately complexed by the low level of A β . Second, it has been proposed by Viles and coworkers that the elevated A β concentration in AD-afflicted brain might be a result of such a homeostasis.⁵⁶ The high propensity of A β to aggregate (especially at a high concentration) and precipitate will further facilitate the accumulation of Cu(II) near the cell membrane. The subsequent ROS generation in the proximity of the cell membrane will cause more severe membrane damages and a faster neuronal cell death. Based on such a model, perhaps a rather important and relevant finding of our work is that Cu(II) complexed by A β aggregates still possesses considerable activity of facilitating the H₂O₂ generation. The co-existence of copper and A β aggregates in senile plaques in AD brain and the availability of large amounts of AA further highlight the relevance of such H₂O₂ generation, as the presence of Cu(II)-containing A β aggregates would serve as a “catalyst” to continuously produce large quantities of H₂O₂. Since the rate constant is moderately high (only two orders of magnitude smaller than ascorbic acid oxidase⁴⁹), both the continuous consumption of AA or other biological redox species and the incessant production of H₂O₂ will wreak havoc to neuronal cells. It has also been suggested that A β may accumulate onto cell membrane prior to its oligomerization and aggregation.^{57,58} Thus Cu(II) sequestration by monomeric and oligomeric A β likely occurs in the early stage of AD development and continue on to the later stage wherein A β molecules have significantly aggregated. Finally, we should also point out that A β -Fe(III), with its metal center capable of cycling between Fe(III) and Fe(II),²² may behave in an analogous fashion as A β -Cu(II) to inflict oxidative stress and damage.

Conclusion

The mechanism of ascorbic acid (AA) oxidation by molecular oxygen facilitated by A β -Cu(II) complexes has been elucidated. The AA oxidation rate was found to be linearly dependent on the concentrations of A β -Cu(II) and dissolved oxygen, reaching a steady state at high [AA] (>100 μ M for [O₂] = 240 μ M). Voltammetric measurements indicate that A β -Cu(II) rapidly oxidizes AA. Based on these kinetic behaviors, we proposed a mechanism in which A β -Cu(II) behaves as a “catalyst” to facilitate the reduction of O₂ by AA and the production of H₂O₂ as the final product. The incessant production of H₂O₂ by a small amount of A β -Cu(II) complex will eventually deplete O₂, AA and/or other important redox species in highly localized regions, disrupting the cellular redox balance and expediting cell loss. The relatively high rate constant measured in this work substantiates such a possibility. These kinetic behaviors are different from those reported for the reaction between AA and O₂ that is catalyzed by the His-Cu(II) complex. When His-Cu(II) serves as the catalyst, the Cu(II) center is not directly reduced, instead a ternary complex among His-Cu(II), O₂, and AA is formed. Furthermore, in comparison with free Cu(II), Cu(II) bound by A β is less efficient in facilitating the O₂ reduction (or AA oxidation). When Cu(II) is complexed by A β aggregates, O₂ is still reduced, albeit at a slower rate. Based on the reaction stoichiometry, if [A β] is much greater than [Cu(II)], hydroxyl radicals are not produced. Overall, the rate of the O₂ reduction to H₂O₂ follows the order of free Cu(II) > Cu(II) bound by A β oligomers/fibrils > Cu(II) bound by A β fibrils. It is also evident that the coordination environment of and the accessibility to the Cu(II) center alter the reaction pathways and modulate the catalytic activity of the Cu(II) center. Our work

demonstrates that oxidative stress can be resulted from the interaction between redox metal ions and aggregates derived from A β

Acknowledgments

This work is supported by an NIH grant (Grant No: SC1NS070155) and the NIH-RIMI Program at California State University-Los Angeles (P20 MD001824-01). XL thanks the financial support from the State Key Laboratory of Environmental Chemistry and Ecotoxicology (No. KF2008-06) and the Graduate School of the Chinese Academy of Sciences. LL also thanks the China Scholarship funds for financial support.

References

- Hardy J, Selkoe DJ. Science 2002;297:353–356. [PubMed: 12130773]
- Selkoe DJ. Science 1997;275:630–631. [PubMed: 9019820]
- Varadarajan S, Yatin S, Aksenova M, Butterfield DA. J. Struct. Biol 2000;130:184–208. [PubMed: 10940225]
- Bush AI Trends Neurosci. Trends Neurosci 2003;26:207–214. [PubMed: 12689772]
- Liu G, Huang W, Moir RD, Vanderburg CR, Lai B, Peng Z, Tanzi RE, Rogers JT, Huang X. J. Struct. Biol 2006;155:45–51. [PubMed: 16503166]
- Lovell MA, Robertson JD, Teesdale WJ, Campbell JL, Mardesbery WR. J. Neurol. Sci 1998;158:47–52. [PubMed: 9667777]
- Jakob-Roetne R, Jacobsen H. Angew. Chem. Int. Ed 2009;48:3030–3059.
- Karr JW, Kaupp LJ, Szalai VA. J. Am. Chem. Soc 2004;126:13534–13538. [PubMed: 15479110]
- Shearer J, Szalai VA. J. Am. Chem. Soc 2008;130:17826–17835. [PubMed: 19035781]
- Syme CD, Nadal RC, Rigby SEJ, Viles JH. J. Biol. Chem 2004;279:18169–18177. [PubMed: 14978032]
- Dong J, Atwood CS, Anderson VE, Siedlak SL, Smith MA, Perry G, Carey PR. Biochemistry 2003;42:2768–2773. [PubMed: 12627941]
- Baruch-Suchodolsky R, Fischer B. Biochemistry 2009;48:4354–4370. [PubMed: 19320465]
- Faller P, Hureau C. Dalton Trans 2009:1080–1094. [PubMed: 19322475]
- Guilloureau L, Combalbert S, Sournia-Saquet A, Mazarguil H, Faller P. ChemBioChem 2007;8:1317–1325. [PubMed: 17577900]
- Huang X, Cuajungco MP, Atwood CS, Hartshorn MA, Tyntall JDA, Hanson GR, Stokes KC, Leopold M, Multhaup G, Goldstein LE, Scarpa RC, Saunders AJ, Lim J, Moir RD, Glabe C, Bowden EF, Masters CL, Fairlie DP, Tanzi RE, Bush AI. J. Biol. Chem 1999;274:37111–37116. [PubMed: 10601271]
- Jiang D, Men L, Wang J, Zhang Y, Chickenyen S, Wang Y, Zhou F. Biochemistry 2007;46:9270–9282. [PubMed: 17636872]
- Rottkamp CA, Raina AK, Zhu X, Bush AI, Atwood CS, Chevion M, Perry G, Smith MA. Free Rad. Biol. Med 2001;30:447–450. [PubMed: 11182300]
- Sayre LM, Perry G, Harris PLR, Liu Y, Schubert KA, Smith MA. J. Neurochem 2000;74:270–279. [PubMed: 10617129]
- Smith MA, Harris PLR, Sayre LM, Perry G. Proc. Natl. Acad. Sci. U. S. A 1997;94:9866–9868. [PubMed: 9275217]
- Sigel, A.; Sigel, H.; Sigel, RKO. Metal Ions in Life Sciences. West Sussex: John Wiley & Sons; 2006.
- Huang X, Atwood CS, Hartshorn MA, Multhaup G, Goldstein LE, Scarpa RC, Cuajungco MP, Gray DN, Lim J, Moir RD, Tanzi RE, Bush AI. Biochemistry 1999;38:7609–7614. [PubMed: 10386999]
- Jiang D, Li X, Williams R, Patel S, Men L, Wang Y, Zhou F. Biochemistry 2009;48:7939–7947. [PubMed: 19601593]
- Drew SC, Noble CJ, Masters CL, Hanson GR, Barnham KJ. J. Am. Chem. Soc 2009;131:1195–1207. [PubMed: 19119811]
- Karr JW, Akintoye H, Kaupp LJ, Szalai VA. Biochemistry 2005;44:5478–5487. [PubMed: 15807541]
- Karr JW, Szalai VA. Biochemistry 2008;47:5006–5016. [PubMed: 18393444]

26. Maiti NC, Jiang D, Wain AJ, Patel S, Dinh KL, Zhou F. *J. Phys. Chem. B* 2008;112:8406–8411. [PubMed: 18570397]
27. Atwood CS, Scarpa RC, Huang X, Moir RD, Jones WD, Fairlie DP, Tanzi RE, Bush AI. *J. Neurochem* 2000;75:1219–1233. [PubMed: 10936205]
28. Hatcher LQ, Hong L, Bush WD, Carducci T, Simon JD. *J. Phys. Chem. B* 2008;112:8160–8164. [PubMed: 18558757]
29. Ma Q-F, Hu J, Wu W-H, Liu H-D, Du J-T, Fu Y, Wu Y-W, Lei P, Zhao Y-F, Li Y-M. *Biopolymers* 2006;83:20–31. [PubMed: 16615111]
30. Sarell CJ, Syme CD, Rigby SEJ, Viles JH. *Biochemistry* 2009;48:4388–4402. [PubMed: 19338344]
31. Tougu V, Karafin A, Palumaa P. *J. Neurochem* 2008;104:1249–1259. [PubMed: 18289347]
32. Hewitt N, Rauk A. *J. Phys. Chem. B* 2009;113:1202–1209. [PubMed: 19123835]
33. Dryhurst, G.; Kadish, KM.; Scheller, F.; Renneberg, R. *Biological Electrochemistry*. New York, London: Academic Press; 1982.
34. da Silva GFZ, Tay WM, Ming L-J. *J. Biol. Chem* 2005;280:16601–16609. [PubMed: 15699049]
35. da Silva GFZ, Ming L-J. *Angew. Chem. Int. Ed* 2005;44:5501–5504.
36. Rice ME. *Trends Neurosci* 2000;23:209–216. [PubMed: 10782126]
37. Barabfis JB, Nagy E, Degrell I. *Arch. Gerontol. Geriat* 1995;21:43–48.
38. Baruch-Suchodolsky R, Fischer B. *Biochemistry* 2008;47:7796–7806. [PubMed: 18598056]
39. Shtamm EV, Purmal AP, Skurlatov YI. *Int. J. Chem. Kinet* 1979;XI:461–494.
40. Scarpa M, Vianello F, Signor L, Zennaro L, Rigo A. *Inorg. Chem* 1996;35:5201–5206.
41. Barron ESG, DeMeio RH, Klemperer F. *J. Biol. Chem* 1936;112:625–640.
42. Skurlatov YI, Kovner VY, Travin SO, Kirsh YE, Purmal AP, Kabanov VA. *Eur. Polym. J* 1979;15:811–815.
43. Gorbunova NV, Purmal AP, Skurlatov YI, Travin SO. *Int. J. Chem. Kinet* 1977:983–1005.
44. Lambert MP, Barlow AK, Chromy BA, Edwards C, Freed R, Liosatos M, Morgan TE, Pozovsky I, Trommer B, Viola KL, Wals P, Zhang C, Finch CC, Krafft GA, Klein WL. *Proc. Natl. Acad. Sci. U. S. A* 1998;95:6448–6453. [PubMed: 9600986]
45. Kaye R, Head E, Thompson JL, McIntire TM, Milton SC, Cotman CW, Glabe CG. *Science* 2003;300:486–489. [PubMed: 12702875]
46. Pecci L, Montefoschi G, Cavallini D. *Biochem. Biophys. Res. Commun* 1997;235:264–267. [PubMed: 9196074]
47. Barb WG, Baxendale JH, George P, Hargrave KR. *Trans. Faraday Soc* 1951;47:462–500.
48. Morgan C, Colombres M, Nuñez MT, Inestrosa NC. *Prog. Neurobiol* 2004;74:323–349. [PubMed: 15649580]
49. Kroneck, PMH.; Armstrong, FA.; Merkle, H.; Marchesini, A. *Advances in Chemistry, Ascorbic Acid: Chemistry, Metabolism, and Uses*. American Chemical Society: Washington, D. C.; 1982.
50. Murray IVJ, Sindoni ME, Axelsen PH. *Biochemistry* 2005;44:12606–12613. [PubMed: 16156673]
51. Luhrs T, Ritter C, Adrian M, Riek-Loher D, Bohrmann B, Dobeli H, Schubert D, Riek R. *Proc. Natl. Acad. Sci. U. S. A* 2005;102:17342–17347. [PubMed: 16293696]
52. Paul C, Axelsen PH. *J. Am. Chem. Soc* 2005;127:5754–5755. [PubMed: 15839650]
53. Kheterpal I, Williams A, Murphy C, Bledsoe B, Wetzel R. *Biochemistry* 2001;40:11757–11767. [PubMed: 11570876]
54. Rae TD, Schmidt PJ, Pufahl RA, Culotta VC, O'Halloran TV. *Science* 1999 284;:805–808. [PubMed: 10221913]
55. Seubert P, Vigo-Pelfrey C, Esch F, Lee M, Dovey H, Davis D, Sinha S, Schlossmacher M, Whaley J, Swindlehurst C, McCormack R, Wolfert R, Selkoe DJ, Lieberburg I, Schenk D. *Nature* 1992;359:325–327. [PubMed: 1406936]
56. Nadal RC, Rigby SEJ, Viles JH. *Biochemistry* 2008;47:11653–11664. [PubMed: 18847222]
57. Jiang D, Dinh KL, Ruthenburg TC, Zhang Y, Su L, Land DP, Zhou F. *J. Phys. Chem. B* 2009;113:3160–3168. [PubMed: 19260715]
58. Terzi E, Hölzemann G, Seelig J. *Biochemistry* 1997;36:14845–14852. [PubMed: 9398206]




Figure 1.

Dependence of the initial oxidation reaction rate of AA on its concentration. The rate was measured at room temperature with $[O_2] = 240 \mu\text{M}$ and $[A\beta\text{-Cu(II)}] = 10 \mu\text{M}$. Each data point is the average of at least three replicate measurements, and the RSD values, ranging from 1.9 to 13.2%, are represented by the error bars.




Figure 2.

The dependence of AA oxidation rate on the O_2 concentration. The rate was measured at room temperature with $[AA] = 240 \mu M$ and $[A\beta-Cu(II)] = 10 \mu M$. Each data point is the average of at least three replicate measurements, and the RSD values, ranging between 2.8 and 32.8%, are represented by the error bars




Figure 3.

The effect of $[\text{A}\beta\text{-Cu(II)}]$ on the initial AA oxidation rate. The rates were obtained with $[\text{AA}] = 240 \mu\text{M}$ in an air-saturated solution. Each data point is the average of at least three replicate measurements, and the RSD values, ranging between 3.5 and 9.5%, are represented by the error bars.




Figure 4. Change of AA absorbance as a function of reaction time in an air-saturated solution containing 10 μM A β -Cu(II) and 420 μM AA.




Figure 5.

The cyclic voltammograms of a solution containing 400 μM A β and 200 μM Cu(II) (red curve) and the same mixture spiked with 100 μM (blue curve) and 200 μM (black curve) AA. The inset shows the CV of AA only. The experiment was conducted at room temperature under a N_2 atmosphere in 10 mM phosphate buffer (pH 7.2) with 0.1 M NaNO_3 as the supporting electrolyte.




Figure 6.

Comparison of reaction rates of the AA oxidation in the presence of various Cu(II)-containing A β species. The rate was obtained under the conditions of 240 μ M AA, 10 μ M Cu(II), and 240 μ M A β in air-saturated phosphate buffer solutions (pH = 7.2 and [O₂] = 240 μ M). The Cu(II)-containing A β (1-42) aggregates were either produced with Cu(II) initially present in the A β (1-42) solution (bars labeled as “A β (1-42)-Cu(II), 48 h” and “A β (1-42)-Cu(II), 72 h” or with Cu(II) added into A β (1-42) solutions that had been preincubated for 48 h (bar labeled as “A β (1-42) (48 h) + Cu(II)”) or 72 h (bar labeled as “A β (1-42) (72 h) + Cu(II)”). The incubation temperature was 37 °C. Measurements were repeated at least three times for each solution, and the RSD values, ranging between 2.7 and 19.2%, are represented by the error bars.




Figure 7.

The morphologies of aggregates formed by incubating 200 μM A β in the presence and absence of 10 μM Cu(II) at 37°C for 48 h. Images A and C were taken right after the solution and surface preparations whereas images B and D were recorded at 48 h after the sample incubation. The area of each image is 5 $\mu\text{m} \times 5 \mu\text{m}$. The particle marked by the blue bar shown in image B is higher in height than the fibril marked in image D (cf. the cross-sectional contours at the bottoms of the respective images).