

Published in final edited form as:

*J Nat Prod.* 2011 May 27; 74(5): 1085–1092. doi:10.1021/np200029q.

## Gambogic Acid, a Natural Product Inhibitor of Hsp90

Jason Davenport<sup>†</sup>, Jacob R. Manjarrez<sup>†</sup>, Laura Peterson<sup>‡</sup>, Brian Krumm<sup>†</sup>, Brian S. J. Blagg<sup>†</sup>, and Robert L. Matts<sup>\*,†</sup>

<sup>†</sup>Department of Biochemistry and Molecular Biology, 246 Noble Research Center, Oklahoma State University, Stillwater, Oklahoma 74078

<sup>‡</sup>Department of Medicinal Chemistry, 1251 Wescoe Hall Drive, Malott 4070, The University of Kansas, Lawrence, Kansas 66045

### Abstract

A high-throughput screening of natural product libraries identified (–)-gambogic acid (**1**), a component of the exudate of *Garcinia harburyi*, as a potential Hsp90 inhibitor, in addition to the known Hsp90 inhibitor celastrol (**2**). Subsequent testing established that **1** inhibited cell proliferation, brought about the degradation of Hsp90 client proteins in cultured cells, and induced the expression of Hsp70 and Hsp90, which are hallmarks of Hsp90 inhibition. Gambogic acid also disrupted the interaction of Hsp90, Hsp70 and Cdc37 with the heme-regulated eIF2 $\alpha$  kinase (HRI—an Hsp90-dependent client) and blocked the maturation of HRI in vitro. Surface plasmon resonance spectroscopy indicated that **1** bound to the N-terminal domain of Hsp90 with a low micromolar  $K_d$ , in a manner that was not competitive with the Hsp90 inhibitor geldanamycin (**3**). Molecular docking experiments supported the posit that **1** binds Hsp90 at a site distinct from Hsp90's ATP binding pocket. The data obtained have firmly established **1** as a novel Hsp90 inhibitor and have provided evidence of a new site that can be targeted for the development of improved Hsp90 inhibitors.

The 90-kDa heat shock protein (Hsp90) is the core component of an oligomeric chaperone machine, the function of which is required for the viability of all eukaryotic cells. Hsp90 functions with a cohort of co-chaperones to facilitate the folding, activation, and stabilization of numerous client proteins, many of which have effects in regulating signal transduction pathways.<sup>1,2</sup> Among the plethora of Hsp90-dependent clients are proteins that function in pathways that represent all six hallmarks of cancer.<sup>3–6</sup> Thus, inhibition of Hsp90 function simultaneously incapacitates multiple client proteins, providing a combinatorial attack on cellular oncogenic processes. Consequently, Hsp90 has emerged as an exciting new target for the development of antitumor agents.

Natural product inhibitors of Hsp90 have been discovered that target binding sites in the N- and C-terminal domains of Hsp90. Geldanamycin (**3**) and radicicol, which are produced by the soil actinomycetes species *Streptomyces hygroscopicus* and the mycoparasitic fungus *Humicola fuscoatra*, respectively, bind to the ATP binding pocket in Hsp90's N-terminal domain,<sup>1,2</sup> while novobiocin (**4**), which is produced by *Streptomyces spheroides*, inhibits Hsp90 function by binding to Hsp90's C-terminal domain.<sup>7</sup> Other natural products with well-known antitumor and/ or chemopreventive properties, but poorly characterized mechanisms of action have been discovered to exhibit inhibitory activity toward Hsp90: epigallocatechin gallate (EGCG),<sup>8,9</sup> the well-known antioxidant found in green tea (*Camellia sinensis* (L.) Kuntze); gedunin, a tetranortriterpenoid isolated from the Indian neem tree (*Azadirachta indica* A. Juss.; Meliaceae),<sup>10,11</sup> celastrol (**2**), a quinone methide

\* Author to whom correspondence should be addressed. Tel.: (405)744-6200. Fax: (405)744-7799. robert.matts@okstate.edu.

triterpene, that is a pharmacologically active compound present in Thunder God Vine root extracts (*Tripterygium wilfordii* Hook.f.; Celastraceae);<sup>11,12</sup> and the rotenoid deguelin.<sup>13</sup>

Derivatives of geldanamycin (**3**) and other compounds that target the ATP-binding pocket in Hsp90's N-terminal domain have entered more than 20 clinical trials for the treatment of cancer.<sup>5,14,15</sup> Clinical complications have arisen in phase II trials of several Hsp90 inhibitors, with incidences of hepato-, cardio-, and ocular toxicity having dampened enthusiasm for the clinical use of Hsp90 inhibitors.<sup>5</sup> Consequently, there is an ongoing search for Hsp90 inhibitors with superior chemotherapeutic properties for the treatment of cancers.

To this end, we have screened natural product libraries for compounds that inhibit Hsp90-dependent refolding of thermally denatured firefly luciferase. It was presumed that natural products represent a fertile territory for the identification of new Hsp90-inhibitors, as it is reasonable to expect that evolutionary pressures give plants that producing secondary metabolites inhibitory to Hsp90 a competitive advantage, because such compounds might inhibit the growth and development of insect pests and other pathogens. Celastrol (**2**), a known Hsp90 inhibitor,<sup>11,12</sup> and (–)-gambogic acid (**1**), a component of *Garcinia hanburyi* Hook.f. (Clusiaceae), a species that has been used medicinally for centuries in southeast Asia, were identified as inhibitors of luciferase refolding in screens of two natural product libraries.

Gambogic acid (**1**), like Hsp90 inhibitors, has antitumor, antiangiogenic, and antimetastatic activities (reviewed in <sup>16–18</sup>), but a poorly characterized mechanism of action. In addition, like Hsp90 inhibitors <sup>19</sup>, **1** has been observed to be selectively cytotoxic to cancer versus normal cells <sup>20,21</sup>. While **1** has been reported to induce apoptosis in cancer cells by binding to the transferrin receptor,<sup>22</sup> the cytotoxic activity of this compound has also been found to have a transferrin receptor-independent component.<sup>23</sup> A recent publication on gambogic acid (**1**) indicates that **1** has recently been subjected to a phase I clinical trial in the People's Republic of China as an anti-cancer agent.<sup>24</sup> Herein, we present the characterization of the Hsp90 inhibitory activity of **1**, and compare its mechanism of action to those of other Hsp90 inhibitors.

## Results and Discussion

### Identification of Gambogic Acid (**1**) as a Putative Hsp90-inhibitor from a High-throughput Screen of Natural Product Libraries

Screening of natural product libraries purchased from Microsource and Biomol for compounds that inhibited Hsp90-dependent refolding of luciferase identified **1** as a potential Hsp90-inhibitor, along with the known Hsp90 inhibitor, celastrol (**2**), among other compounds. Neither celastrol nor **1** had any direct effect on the activity of native luciferase. Upon titration of various concentrations of the Two compounds into the refolding assay (Figure 1A), celastrol (**2**) and gambogic acid (**1**) were found to inhibit luciferase refolding by 50% (IC<sub>50</sub>) at a concentration of 20 and 2 μM, respectively.

Gambogic acid (**1**) has been demonstrated in numerous studies to inhibit the proliferation of a variety of cancer cell lines (reviewed in <sup>16–18</sup>). To determine whether antiproliferative activity of **1** could be correlated with its Hsp90-inhibitory activity, we examined the effect of varying concentrations of gambogic acid on the growth/ viability of HeLa cells, and MCF7 and SK-Br3 breast cancer cell lines. Gambogic acid (**1**) inhibited the proliferation of HeLa, MCF7, and SK-Br3 cells in a concentration dependent manner (Figure 1B). Growth of the HeLa, MCF7, and SK-Br3 cells was inhibited by 50% by treatment with 1.5, 2.0 and 0.8 μM **1**, respectively. The highest concentrations of **1** were cytotoxic as evidenced by

detachment of a significant number of cells from the surface of the culture flasks. Thus, the IC<sub>50</sub> of **1** for inhibition of cell proliferation correlated well with its IC<sub>50</sub> for the inhibition of luciferase refolding.

### Gambogic Acid (1)-induced Depletion of Hsp90-dependent Proteins

Treatment of cultured cells with known Hsp90 inhibitors depletes the cells of Hsp90-dependent proteins in a time- and concentration-dependent manner. To further characterize **1** as a potential Hsp90 inhibitor, MCF7 and Sk-Br3 cells were treated with varying concentration of **1** for 36 h, and equivalent amounts of protein from cell extracts were Western blotted for Hsp70 and Hsp90, and the Hsp90-dependent proteins Her2, Akt, and Raf-1, using actin as a loading control, geldanamycin (**3**) as a positive control for Hsp90-inhibition, and DMSO as a negative control. Gambogic acid was observed to deplete MCF7 and Sk-Br3 cells of Her2, Akt and Raf-1 in a concentration dependent fashion (Figure 2), which correlated well with the IC<sub>50</sub> value for inhibition of the proliferation of these cell lines induced by **1**. In addition, **1** induced Hsp90 and Hsp70 expression, another hallmark of Hsp90-inhibition. This compound had a similar effect on the levels of Her2, Raf-1, and Akt in HeLa cells (not shown). These results further support the hypothesis that the antiproliferative effect of **1** on cancer cell growth is mediated, at least in part, by its ability to inhibit Hsp90.

### Effect of Gambogic Acid (1) on the Association of Hsp90 Chaperone Components with HRI and HRI's Hsp90-dependent Maturation

Hsp90 inhibitors have distinct effects on the conformation and interactions of Hsp90 with co-chaperone partners and clients.<sup>25,26</sup> Geldanamycin (**3**), an Hsp90 inhibitor that binds to ATP pocket in the N-terminal domain of Hsp90 (N-terminal inhibitor), causes Hsp90 accumulate in complexes that contain intermediate co-chaperone components. Complexes that form between Hsp90 and kinase clients in the presence of **3** are salt-labile and lack Cdc37. Molybdate, on the other hand, inhibits Hsp90 by freezing it in complexes containing late co-chaperone components. Hsp90-kinase client complexes formed in the presence of molybdate are stable at high salt concentrations (e.g., 0.5 M NaCl) and contain Cdc37. Hsp90 inhibitors that bind to its C-terminal domain (C-terminal inhibitor), like novobiocin (**4**), appear to block the interaction of Hsp90 with most co-chaperones, as well as protein clients. Celastrol (**2**), which is thought to bind to the N-terminal domain of Hsp90 at a site distinct from the ATP binding pocket, inhibits Hsp90 in a manner distinct from geldanamycin (**3**) and novobiocin (**4**).<sup>25</sup>

HRI is a heme-regulated eIF2 $\alpha$  kinase that requires Hsp90, Hsp70 and Cdc37 for its maturation and activation under heme-deficient conditions.<sup>27–29</sup> To further characterize the mechanism of action of gambogic acid (**1**), the effects of **1** on the binding of Hsp90 chaperone components to HRI was compared to the effects of the N-terminal inhibitors geldanamycin (**3**) and celastrol (**2**), molybdate, and the C-terminal inhibitors novobiocin (**4**) and coumermycin A1 (**5**).<sup>25</sup> The non-activatable K199R mutant of HRI was used in these experiments as it is unable to mature and interacts constitutively with Hsp90. In these experiments, the immunoadsorbed samples were washed with low ionic strength buffer, as it distinguishes between the mechanism of action of geldanamycin (**3**) and celastrol (**2**), which bind to different sites in Hsp90's N-terminal domain. In agreement with results reported previously:<sup>30–32</sup> 1) geldanamycin (**3**) caused the accumulation of Hsp90 in complexes with HRI/K199R that contained increased amounts of the intermediate co-chaperone component Hsp70 compared to the DMSO control, but lacked Cdc37 (Figure 3A, lane 3); 2) molybdate stabilized the binding of Hsp90 and Cdc37 to HRI/K199R, while not affecting the interaction of Hsp70 with HRI/K199R (lane 4); and 3) novobiocin (**4**) and disrupted the interaction of Hsp90, Hsp70 and Cdc37 with HRI/K199R (lane 6), as did the

bivalent novobiocin related compound coumermycin A1 (**5**) (lane 7). Under similar conditions, celastrol (**2**) disrupted the binding of Hsp90, Hsp70, and Cdc37 to HRI (Figure 3A, lane 5), as did gambogic acid (**1**) (lane 8). These results indicate that **1** affected the interaction of Hsp90 chaperone components with HRI/K199R in a manner distinct from geldanamycin (**3**) and molybdate.

Subsequently, the effects of **1** on HRI activation were examined upon incubation in heme-deficient lysate. Hsp90-dependent maturation and activation of HRI is accompanied by a change in its electrophoretic mobility to a more slowly migrating species (Figure 3B). After 45 min of incubation of HRI in heme-deficient lysate, approximately half of the [<sup>35</sup>S]HRI had a slower electrophoretic mobility upon SDS-PAGE analysis compared to HRI incubated in heme-replete lysate (lane 2 vs. 1). Gambogic acid (**1**) inhibited the maturation/ activation of HRI in heme-deficient lysate to an extent similar to the known Hsp90 inhibitors geldanamycin (**3**), celastrol (**2**), and molybdate. The C-terminal Hsp90 inhibitors novobiocin (**4**) and coumermycin A1 (**5**) also inhibited the maturation/ activation of HRI, but it was also apparent that incubation with these compounds reduced the quantity of [<sup>35</sup>S]HRI that was present in the samples compared to the other Hsp90 inhibitors.

To determine whether the decreased recovery of [<sup>35</sup>S]HRI was a hallmark of C-terminal Hsp90 inhibitors inducing the breakdown of HRI, or whether it was simply due to less [<sup>35</sup>S]HRI being synthesized, aliquots were taken from each sample after protein synthesis had ceased, prior to the beginning of the maturational incubation in the presence of the compounds. Autoradiography of aliquots taken from samples prior to their maturational incubation indicated that an equivalent amount of [<sup>35</sup>S]HRI was present in each sample (Figure 3C, 0 min: upper panel). Incubation of [<sup>35</sup>S]HRI in the presence of novobiocin (**4**) or coumermycin A1 (**5**) resulted in a 90% and 80% loss of the [<sup>35</sup>S]HRI, respectively (Figure 3C, lanes 6 and 7). However, significantly less [<sup>35</sup>S]HRI was lost upon incubation of samples in the presence of geldanamycin (**3**) (0%), molybdate (10%), celastrol (**2**) (40%), or **1** (10%) compared to the DMSO control. A similar effect of novobiocin (**4**) and coumermycin A1 (**5**) has been observed on stability of Akt generated by TnT in reticulocyte lysate (not shown). Thus, the effect of Hsp90 inhibition on the stability of nascent kinases in reticulocyte lysate appears to be a property that further distinguishes inhibitors of Hsp90 that bind to its C-terminal domain from inhibitors that bind to its N-terminal domain.<sup>25,26</sup> The absence of an effect of **1** on HRI stability distinguishes its mechanism of action from that of C-terminal inhibitors. Previously, celastrol (**2**) was demonstrated to inhibit Hsp90 by a mechanism that was distinct from N-terminal inhibitors that bind to Hsp90's ATP binding pocket, such as geldanamycin (**3**).<sup>25</sup> The results presented here indicate that the mechanism of action of **1** is similar to that of celastrol (**2**), and distinct from that of **3**.

### Surface Plasmon Resonance (SPR) Analysis of the Binding of Gambogic Acid (**1**) to the N-terminal domain of Hsp90

To determine whether **1** is indeed interacting with Hsp90's N-terminal domain, we cloned the N-terminal domain of Hsp90 $\alpha$  (Hsp90NT; amino acids 1–241) with a 7-amino-acid extension at its C-terminus containing a C-terminal Cys residue. Recombinant Hsp90NT was biotinylated and immobilized onto a neutravidin sensor chip for analysis of the binding of **1** by SPR. Full-length Hsp90 and the C-terminal domain of Hsp90 (Hsp90CT, amino acids 531–732) were immobilized on sensor chips that were used as positive and negative controls, respectively. Gambogic acid (**1**) was observed to bind specifically to full-length Hsp90 (Figure 4A) and Hsp90NT (Figure 4B), but not to Hsp90CT (data not shown). Kinetic analysis of binding and dissociation kinetics indicated that **1** binds to both Hsp90 and Hsp90NT with similar association constants ( $k_a$ ), dissociation constants ( $k_d$ ) and calculated  $K_d$  values (9.8 versus 7.6  $\mu$ M, respectively, Table 1). Compound **1** was observed also to bind Hsp90NT in the presence of 20  $\mu$ M geldanamycin in the analyte buffer, with

geldanamycin (**3**) having no significant effect on the measured  $k_a$  and  $k_d$  or the calculated  $K_d$  (Table 1). Thus, the data indicate that gambogic acid binds to the N-terminal domain of Hsp90, and, like celastrol,<sup>11</sup> it binds to a site distinct from the ATP binding pocket. In addition, the  $K_d$  for the binding of **1** to Hsp90 correlated well with the  $IC_{50}$  values for gambogic acid (**1**)-induced inhibition of other Hsp90-dependent processes, further supporting Hsp90 as one of the physiological targets of this compound.

### Virtual Docking of Gambogic Acid (**1**) to Hsp90NT

We approached the problem of identifying the putative binding site of gambogic acid (**1**) in the N-terminal domain of Hsp90 using the “blind” docking method,<sup>33</sup> in which the entire Hsp90 N-terminal domain, not just the geldanamycin-binding site, was used to search for the lowest possible binding energy for **1**. The apo- and geldanamycin-bound Hsp90NT crystal structure, PDB ID 1YET, along with the open or unbound crystal structure of the Hsp90NT, PDB ID 1YES, were used for docking of **1** to determine if it would compete for the geldanamycin-binding site or bind elsewhere. These two structures share a 0.25 Å RMSD of the C $\alpha$  backbone spanning the entire 221 residues of the crystal structures with the majority of conformational changes occurring in residues 105–112.<sup>34</sup> In all three docking experiments, **1** preferentially bound to the cleft created by  $\alpha$ -helix 9 (H9) and  $\beta$ -sheet strand 8 (S8), residues 200–222 (Figure 5A) that are composed of both hydrophobic and positively charged elements (Figure 5B). The calculated average binding affinities for **1** for this region were –9.44, –9.50 and –9.37 kcal/mol for the apo, geldanamycin-bound, and open structures, respectively. Docking results predict that aliphatic hydrophobic interactions would occur between the gambogic acid substituent arms and the aliphatic side chains of H9 and Phe213 in the intervening loop, while the xanthene moiety would lie on the C $\alpha$  backbone of S8 making potential pi interaction with the carbonyl and amide bonds (Figure 5A). In addition, there is a potential salt bridge between Lys208 of H9 and both oxygens of the carboxylic acid moiety of **1**, and a hydrogen bond between Lys204 of H9 with the carbonyl moiety of the cyclohexene ring (Figure 5C). The regions of H9 and S8 undergo minor conformational changes with a 0.15 Å RMSD of the C $\alpha$  residues between residues 200–222.

The docking results are consistent with gambogic acid (**1**) being a non-competitive inhibitor of the Hsp90 N-terminal domain. The region of H9 and S8 of the HSP90 N-terminal domain undergoes little change between the geldanamycin-bound structure and the open conformation structure with a 0.15 Å RMSD of the C $\alpha$  backbone of residues 200–222. It should be noted that this site is distinct from that previously proposed for the binding of celastrol (**2**) to the Hsp90's N-terminal domain.<sup>12</sup>

Recently, the combination of gambogic acid (**1**) and celastrol (**2**) was reported to have a synergistic anti-tumor effect,<sup>35</sup> an observation that supports the notion that the two compounds interact with different sites within the N-terminal domain of Hsp90. The question of whether combinations of Hsp90 inhibitors that act by different mechanisms will demonstrate similar synergistic effects on cancer cell growth has yet to be explored, but the synergy exhibited by **1** and **2** suggests that such studies would be productive. For example, the ability of C-terminal inhibitors, such as novobiocin, to block the activation of HSF1<sup>36</sup> may potentiate the cytotoxicity of N-terminal inhibitors of Hsp90, which induce heat shock protein expression.

### Conclusion

The data presented above firmly establishes gambogic acid (**1**) as an inhibitor of Hsp90 and provides evidence of a new site that can be targeted for the development of improved Hsp90 inhibitors. Previous work with **1** has indicated that it reduces the activity or the levels of



expression of well known Hsp90-dependent clients (see: <http://www.picard.ch/downloads/Hsp90interactors.pdf> and <http://www.picard.ch/downloads/Hsp90facts.pdf>) that are involved in the regulation of cell immortality (telomerase),<sup>37,38</sup> cell growth (Akt, cSrc, Cdk2 and Cdk4),<sup>39–41</sup> apoptosis (Bcl-2, Bcl-xL and survivin),<sup>42–44</sup> angiogenesis (VEGFR),<sup>40,41</sup> and metastasis (MMP2 and MMP9).<sup>22,45</sup> Changes in levels of expression of proteins (Bcl-2, Bcl-xL, survivin, and telomerase) appeared to occur primary at the level of transcription rather than through destabilization of the protein, as mRNA levels coding for the protein were decreased. These changes, however, can still be accounted for by a mechanism invoking gambogic acid-induced inhibition of Hsp90, as NF- $\kappa$ B regulates the expression of Bcl-2, Bcl-xL and survivin, as well as VEGF,<sup>17,22</sup> and NF- $\kappa$ B activation is well known to be regulated by Hsp90.<sup>46</sup> This notion is supported by the observations that the Hsp90 inhibitor geldanamycin (**3**) inhibits the expression of Bcl-2 and Bcl-xL by suppressing the activation of NF- $\kappa$ B,<sup>47</sup> and has the same effects as gambogic acid on VEGFR signaling.<sup>48,49</sup> Furthermore, telomerase expression is regulated by c-Myc, and Hsp90 inhibition has recently been demonstrated to destabilize c-Myc,<sup>50</sup> in addition, to blocking the interaction of Hsp90 with the telomerase promoter which suppresses telomerase expression.<sup>51</sup> While this manuscript was in preparation, gambogic acid (**1**) was confirmed to block NF- $\kappa$ B activation through its ability to inhibit Hsp90.<sup>52</sup> Thus, many of the physiological effects of **1** on cultured cells reported previously in the literature, can be accounted for, in part, by its ability to inhibit Hsp90 function.

## Experimental Section

### Screen for Inhibitors of Hsp90-dependent Luciferase Refolding

Natural product libraries from Microsource and Biomol were screened as previously described for inhibitors of Hsp90-dependent refolding of thermally denatured firefly luciferase.<sup>53</sup> Positive hits were then screened against native luciferase to eliminate false-positives that were direct inhibitors of luciferase. Subsequently, celastrol (**2**, CalBiochem 219465,  $\geq 95\%$  by TLC) and gambogic acid (**1**, Biomol BML-AP305, 98% by TLC) were titrated into reticulocyte lysate containing thermally denatured luciferase to determine the concentration of test compound required to inhibit luciferase refolding by 50% (IC<sub>50</sub>) compared to the DMSO control, as previously described.<sup>54</sup> Reactions were carried out in triplicate at room temperature in 96-well microtiter plates, and experiments were repeated at least twice, with relative light unit (RLU) production measured using a L<sub>Max</sub> II (Molecular Devices) microplate reader and a 10 sec integration time.<sup>54</sup>

### Effect of Gambogic Acid (**1**) on Cell Proliferation

MCF-7 and HeLa cells were grown in Gibco Modified Essential Medium, supplemented with non-essential amino acids, 2 mM glutamine, and 10% fetal bovine serum. SkBr3 cells were maintained in DMEM/F12 (1:1) medium (GIBCO) with l-glutamine supplemented with streptomycin (500  $\mu$ g/mL), penicillin (100 units/mL), and 10% fetal bovine serum. Cells were grown to confluence in a humidified atmosphere (37 °C, 5% CO<sub>2</sub>). Cells were seeded at 2000 cells per well in a clear 96-well plate, the medium volume was brought to 100  $\mu$ L, and the cells were allowed to attach overnight. The next day, varying concentrations of compound or 1% DMSO vehicle control were added to the wells. Cells were then incubated at 37 °C for 72 h. Cell viability was determined using the Promega Cell Titer 96 Aqueous One Solution Cell Proliferation Assay, which makes use of a soluble tetrazolium compound that is converted into a chromophore by living cells. After incubation with compounds, 20  $\mu$ L of the assay substrate solution were added to the wells, and the plate was incubated at 37 °C for an additional 1 h. Absorbance at 490 nm was then read on a

Molecular Devices Versamax plate reader, and values were expressed as percent of absorbance from cells incubated in DMSO alone.

### **Effect of Gambogic Acid (1) on the interaction of Hsp90 and its Co-chaperones with Heme-regulated eIF2 $\alpha$ Kinase (HRI)**

His-tagged kinase-dead HRI/K199R was synthesized by coupled transcription-translation (TnT) in reticulocyte lysate in the presence of [<sup>35</sup>S]methionine.<sup>55,56</sup> After 20 min of synthesis, compounds or an equivalent volume of DMSO or water were added at the concentrations indicated in the figure legend and synthesis was continued for an additional 10 min. Subsequently, aliquots were taken for immunoadsorption of His-tagged HRI with anti-His-tag antibodies bound to anti-mouse-IgG agarose resin.<sup>55,56</sup> Agarose resins containing bound anti-His-tag antibody were washed four times with 10 mM PIPES (pH = 7.2) plus 100 mM NaCl and analyzed by SDS-PAGE and Western blotting for co-adsorbed Hsp90, Hsp70 and Cdc37, as previously described.<sup>55,56</sup> Lysate lacking plasmid encoding His-tagged HRI was used as the control for non-specific binding.

### **Effect of Gambogic Acid (1) on Hsp90-dependent HRI Maturation/ activation**

[<sup>35</sup>S]His-tagged wild type HRI or K199R mutant was generated by TnT in reticulocyte lysate, as described above. After 30 min, samples were diluted into 7 volumes of hemin-supplemented or heme-deficient lysate containing DMSO, water or test compounds at the same concentration present in the TnT lysate and incubated for an additional 45 min.<sup>55,56</sup> The His-tagged HRI was adsorbed from samples by the addition of NTA-Ni resin on ice for 1 h. Resins were washed, eluted by boiling in SDS-sample buffer, and samples were analyzed by SDS-PAGE and autoradiography after electrotransfer to PVDF membrane to detect a shift in the electrophoretic mobility of HRI, which is dependent upon its Hsp90-dependent maturation/activation.<sup>55,56</sup>

### **Gambogic Acid (1)-induced Depletion of Hsp90-dependent Proteins from Cultured MCF7 and SKBr3 Breast Cancer and HeLa cells**

MCF-7, SkBr3, and HeLa cells were grown to confluence as described above, seeded in culture dishes (1×10<sup>6</sup>/dish; brand) and allowed to attach overnight. Gambogic acid (1) was added at the concentrations indicated in figure 1, and the cells were incubated for an additional 36 h. Cells were harvested and analyzed for Hsp90 client protein degradation (Her2, Raf-1, and Akt) and heat shock protein induction (Hsp90 and Hsp70), as described previously.<sup>57</sup> For comparison, cells were incubated with DMSO (1%) or geldanamycin (3, 500 nM), and extracts (10  $\mu$ g of protein) were Western blotted for the indicated proteins with actin used as the loading control. One  $\mu$ g of SkBr3 extracts were used for the Western blot for Her2.

### **Surface Plasmon Resonance Spectroscopy of Gambogic Acid (1) Binding to Hsp90**

Insect Sf9 cells overexpressing human Hsp90 $\beta$  were cultured and harvested by the Baculovirus/Monoclonal Antibody Core facility at Baylor College of Medicine. Hsp90 $\beta$  was extracted and purified (>98% pure) as described previously,<sup>58,59</sup> but without the initial DEAE-cellulose chromatography step. Bacterial *E. coli* DE-3 Star cells carrying plasmids for the expression of either the N-terminal (Hsp90NT: amino acids 1–241 with a C-terminal -GELRSGC tail) or C-terminal (Hsp90CT: amino acids 531–732) domains of Hsp90 were cultured in LB medium and induced with IPTG. Recombinant Hsp90NT and Hsp90CT were purified using a NiNTA column. Following elution of the protein from the NiNTA column, the N-terminal His-tag was cleaved using TEV protease (Invitrogen), followed by size-exclusion chromatography on Superdex 200. The Hsp90-containing fractions were pooled, concentrated and dialyzed against 10 mM Hepes (pH 7.5) containing 150 mM NaCl and

10% glycerol, stored at  $-80^{\circ}\text{C}$  until use. Prior to the use of the Hsp90NT or Hsp90CT, each protein was reacted according to the manufacture's recommended protocol with the EZ-Link<sup>®</sup> Maleimide-PEG2-Biotin (Thermo Scientific) then buffer-exchanged into fresh 10 mM Hepes (pH 7.5) containing 150 mM NaCl to eliminate free biotin entities.

The surface of a SSO1 CO<sub>2</sub>H SPR sensor chip mounted in a SensiQ SPR instrument (ICX Nomadics) was activated by treatment with *N*'-3-dimethylaminopropyl-*N*'-ethylcarbodiimide hydrochloride and *N*-hydroxysuccinimide for preferential cross-linking of full length Hsp90's N-terminus to the surface. For immobilization of Hsp90, 250  $\mu\text{L}$  of Hsp90 (6.2 mg/mL) in 10 mM Hepes buffer (pH 7.4) containing 150 mM NaCl were injected at a flowrate of 10  $\mu\text{L}/\text{min}$ , resulting in 2000 response units of protein captured on the experimental surface of the chip. Then, 1 M ethanolamine (pH 8) was used to quench the remaining activated groups, and the surface washed with buffer containing 10 mM PIPES pH 7.4, 300 mM NaCl, and 2% DMSO. The surface of a SSO3 BioCap SPR sensor chip was mounted in a SensiQ SPR instrument (ICX Nomadics) and either the biotinylated Hsp90NT (7.0 mg/mL) or Hsp90CT (6.8 mg/mL) was discrete-injected over the experimental channel at a flow rate of 10  $\mu\text{L}/\text{min}$ , resulting in capture of Hsp90NT at 1400 response units and Hsp90CT at 1250 response units of protein on the experimental surface of the chip. The chips were then washed with assay buffer prior to experimental analysis.

Gambogic acid (**1**) was diluted in assay buffer containing 10 mM PIPES pH 7.4, 300 mM NaCl, and 2% DMSO and injected over the surface of each derivatized chip at a flow rate of 15  $\mu\text{L}/\text{min}$  at  $25^{\circ}\text{C}$  at the indicated concentrations. Additionally, for competition studies **1** was diluted in assay buffer containing 20  $\mu\text{M}$  geldanamycin (**3**). All measurements were done in triplicate. SPR binding curves were analyzed using QDAT software (ICX Nomadics) to calculate the  $k_a$ ,  $k_d$  and  $K_D$  data.

### Docking Studies of Geldanamycin (**3**) and Gambogic Acid (**1**) with the N-terminal Domain of Hsp90

*In silico* docking of geldanamycin and gambogic acid with the 3D coordinates of the X-ray crystal structures of the N-terminal domain of HSP90 with bound **3**, and in the open conformation, PDB IDs 1YET and 1YES respectively, was accomplished using the Autodock program<sup>60</sup> downloaded from the Molecular Graphic Laboratory of the Scripps Research Institute. The AutoDock program was chosen because it uses a genetic algorithm to generate the poses of the ligand inside a known or predicted binding site utilizing the Lamarckian version of genetic algorithm where the changes in conformations adopted by molecules after *in situ* optimization are used as subsequent poses for the offspring.

In the docking experiments carried out, water was removed from the 3D X-ray coordinate while Gasteiger charges were placed on the X-ray structures of the N-terminal domain of HSP90 along with geldanamycin (**3**) and **1** using tools from the Autodock suite. A grid box centered on the N-terminal HSP90 domain with definitions of  $126 \times 126 \times 126$  points and 0.4 Å spacing was chosen for ligand docking experiments. The docking parameters consisted of setting the population size to 300, the number of generations set to 27,000, and the number of evaluations set to 20,000,000, while the number of docking runs was set to 50 with a cutoff of 1 Å for the root mean square tolerance for the grouping of each docking run.

While the binding mode of geldanamycin with Hsp90 has been determined through X-ray crystallography,<sup>34</sup> the binding mode of **1** with Hsp90 has yet to be determined through either NMR spectroscopy or X-ray crystallography. The docking of geldanamycin to the apo X-ray structure of Hsp90NT, PDB ID 1YET, was used as a control to test and validate the docking parameters used. As expected, geldanamycin (**3**) docked to the binding site identified in the



crystal structure with an average binding energy of  $-9.65$  kcal/mol and a  $1 \text{ \AA}$  average root mean square deviation from the reference structure (Figure 6).

## Acknowledgments


The authors gratefully acknowledge support of this project by NIH (R01 CA125392), and the Oklahoma Agricultural Experiment Station (Project No. 1975).

## References

1. Pearl LH, Prodromou C, Workman P. *Biochem. J.* 2008; 410:439–453. [PubMed: 18290764]
2. Pratt WB, Toft DO. *Exp. Biol. Med.* (Maywood). 2003; 228:111–133. [PubMed: 12563018]
3. Neckers L. *J. Biosci.* 2007; 32:517–530. [PubMed: 17536171]
4. Hanahan D, Weinberg RA. *Cell.* 2000; 100:57–70. [PubMed: 10647931]
5. Holzbeierlein JM, Windsperger A, Vielhauer G. *Curr. Oncol. Rep.* 2010; 12:95–101. [PubMed: 20425593]
6. Bishop SC, Burlison JA, Blagg BS. *Curr. Cancer Drug Targets.* 2007; 7:369–388. [PubMed: 17979631]
7. Donnelly A, Blagg BS. *Curr. Med. Chem.* 2008; 15:2702–2717. [PubMed: 18991631]
8. Li Y, Zhang T, Jiang Y, Lee HF, Schwartz SJ, Sun D. *Mol. Pharm.* 2009; 6:1152–1159. [PubMed: 19438225]
9. Yin Z, Henry EC, Gasiewicz TA. *Biochemistry.* 2009; 48:336–345. [PubMed: 19113837]
10. Brandt GE, Schmidt MD, Prinszano TE, Blagg BS. *J. Med. Chem.* 2008; 51:6495–6502. [PubMed: 18816111]
11. Hieronymus H, Lamb J, Ross KN, Peng XP, Clement C, Rodina A, Nieto M, Du J, Stegmaier K, Raj SM, Maloney KN, Clardy J, Hahn WC, Chiosis G, Golub TR. *Cancer Cell.* 2006; 10:321–330. [PubMed: 17010675]
12. Zhang T, Hamza A, Cao X, Wang B, Yu S, Zhan CG, Sun D. *Mol. Cancer Ther.* 2008; 7:162–170. [PubMed: 18202019]
13. Oh SH, Woo JK, Yazici YD, Myers JN, Kim WY, Jin Q, Hong SS, Park HJ, Suh YG, Kim KW, Hong WK, Lee HY. *J. Natl. Cancer Inst.* 2007; 99:949–961. [PubMed: 17565155]
14. Porter JR, Fritz CC, Depew KM. *Curr. Opin. Chem. Biol.* 2010; 14:412–420. [PubMed: 20409745]
15. Kim YS, Alarcon SV, Lee S, Lee MJ, Giaccone G, Neckers L, Trepel JB. *Curr. Top. Med. Chem.* 2009; 9:1479–1492. [PubMed: 19860730]
16. Aggarwal BB, Kunnumakkara AB, Harikumar KB, Tharakan ST, Sung B, Anand P. *Planta Med.* 2008; 74:1560–1569. [PubMed: 18612945]
17. Gupta SC, Kim JH, Prasad S, Aggarwal BB. *Cancer Metastasis Rev.* 2010; 29:405–434. [PubMed: 20737283]
18. Han QB, Xu HX. *Curr. Med. Chem.* 2009; 16:3775–3796. [PubMed: 19747141]
19. Kamal A, Thao L, Sensintaffar J, Zhang L, Boehm MF, Fritz LC, Burrows FJ. *Nature.* 2003; 425:407–410. [PubMed: 14508491]
20. Yang Y, Yang L, You QD, Nie FF, Gu HY, Zhao L, Wang XT, Guo QL. *Cancer Lett.* 2007; 256:259–266. [PubMed: 17693016]
21. Zhao L, Guo QL, You QD, Wu ZQ, Gu HY. *Biol. Pharm. Bull.* 2004; 27:998–1003. [PubMed: 15256729]
22. Pandey MK, Sung B, Ahn KS, Kunnumakkara AB, Chaturvedi MM, Aggarwal BB. *Blood.* 2007; 110:3517–3525. [PubMed: 17673602]
23. Ortiz-Sanchez E, Daniels TR, Helguera G, Martinez-Maza O, Bonavida B, Penichet ML. *Leukemia.* 2009; 23:59–70. [PubMed: 18946492]
24. Ren Y, Yuan C, Chai HB, Ding Y, Li XC, Ferreira D, Kinghorn AD. *J. Nat. Prod.* 2011; 74 in press.
25. Matts RL, Brandt GE, Lu Y, Dixit A, Mollapour M, Wang S, Donnelly AC, Neckers L, Verkhivker G, Blagg BS. *Bioorg. Med. Chem.* 2011; 19:684–692. [PubMed: 21129982]


26. Matts RL, Manjarrez JR. *Curr. Top. Med. Chem.* 2009; 9:1462–1478. [PubMed: 19860729]
27. Shao J, Gramatikakis N, Scroggins B, Uma S, Huang W, Chen J-J, Hartson SD, Matts RL. *J. Biol. Chem.* 2001; 276:206–214. [PubMed: 11036079]
28. Uma S, Hartson SD, Chen J-J, Matts RL. *J. Biol. Chem.* 1997; 272:11648–11656. [PubMed: 9111082]
29. Uma S, Thulasiraman V, Matts RL. *Mol. Cell. Biol.* 1999; 19:5861–5871. [PubMed: 10454533]
30. Hartson SD, Thulasiraman V, Huang W, Whitesell L, Matts RL. *Biochemistry.* 1999; 38:3837–3849. [PubMed: 10090774]
31. Prince T, Matts RL. *J. Biol. Chem.* 2004; 279:39975–39981. [PubMed: 15258137]
32. Scroggins BT, Prince T, Shao J, Uma S, Huang W, Guo Y, Yun BG, Hedman K, Matts RL, Hartson SD. *Biochemistry.* 2003; 42:12550–12561. [PubMed: 14580201]
33. Hetenyi C, van der Spoel D. *Protein Sci.* 2002; 11:1729–1737. [PubMed: 12070326]
34. Stebbins CE, Russo AA, Schneider C, Rosen N, Hartl FU, Pavletich NP. *Cell.* 1997; 89:239–250. [PubMed: 9108479]
35. He D, Xu Q, Yan M, Zhang P, Zhou X, Zhang Z, Duan W, Zhong L, Ye D, Chen W. *BMC Cancer.* 2009; 9:343. [PubMed: 19778460]
36. Conde R, Belak ZR, Nair M, O'Carroll RF, Ovsenek N. *Biochem. Cell Biol.* 2009; 87:845–851. [PubMed: 19935870]
37. Guo QL, Lin SS, You QD, Gu HY, Yu J, Zhao L, Qi Q, Liang F, Tan Z, Wang X. *Life Sci.* 2006; 78:1238–1245. [PubMed: 16257012]
38. Wu ZQ, Guo QL, You QD, Zhao L, Gu HY. *Biol. Pharm. Bull.* 2004; 27:1769–1774. [PubMed: 15516720]
39. Liu Y, Li W, Ye C, Lin Y, Cheang TY, Wang M, Zhang H, Wang S, Zhang L. *J. Atheroscler. Thromb.* 2010; 17:901–913. [PubMed: 20543524]
40. Lu N, Yang Y, You QD, Ling Y, Gao Y, Gu HY, Zhao L, Wang XT, Guo QL. *Cancer Lett.* 2007; 258:80–89. [PubMed: 17920764]
41. Yi T, Yi Z, Cho SG, Luo J, Pandey MK, Aggarwal BB, Liu M. *Cancer Res.* 2008; 68:1843–1850. [PubMed: 18339865]
42. Shu W, Chen Y, Li R, Wu Q, Cui G, Ke W, Chen Z. *Basic Clin. Pharmacol. Toxicol.* 2008; 103:530–537. [PubMed: 19067680]
43. Xu X, Liu Y, Wang L, He J, Zhang H, Chen X, Li Y, Yang J, Tao J. *Int. J. Dermatol.* 2009; 48:186–192. [PubMed: 19200201]
44. Wang T, Wei J, Qian X, Ding Y, Yu L, Liu B. *Cancer Lett.* 2008; 262:214–222. [PubMed: 18248784]
45. Qi Q, Lu N, Wang XT, Gu HY, Yang Y, Liu W, Li C, You QD, Guo QL. *Biochem. Cell Biol.* 2008; 86:386–395. [PubMed: 18923540]
46. Salminen A, Paimela T, Suuronen T, Kaarniranta K. *Immunol. Lett.* 2008; 117:9–15. [PubMed: 18282612]
47. Bai J, Sui J, Demirjian A, Vollmer CM Jr, Marasco W, Callery MP. *Cancer Res.* 2005; 65:2344–2352. [PubMed: 15781649]
48. Masson-Gadais B, Houle F, Laferriere J, Huot J. *Cell Stress Chaperones.* 2003; 8:37–52. [PubMed: 12820653]
49. Dias S, Shmelkov SV, Lam G, Rafii S. *Blood.* 2002; 99:2532–2540. [PubMed: 11895790]
50. Regan PL, Jacobs J, Wang G, Torres J, Edo R, Friedmann J, Tang XX. *Int. J. Oncol.* 2011; 38:105–112. [PubMed: 21109931]
51. Kim RH, Kim R, Chen W, Hu S, Shin KH, Park NH, Kang MK. *Carcinogenesis.* 2008; 29:2425–2431. [PubMed: 18820283]
52. Zhang L, Yi Y, Chen J, Sun Y, Guo Q, Zheng Z, Song S. *Biochem. Biophys. Res. Commun.* 2010; 403:282–287. [PubMed: 21074517]
53. Galam L, Hadden MK, Ma Z, Ye QZ, Yun BG, Blagg BS, Matts RL. *Bioorg. Med. Chem.* 2007; 15:1939–1946. [PubMed: 17223347]
54. Thulasiraman V, Matts RL. *Biochemistry.* 1996; 35:13443–13450. [PubMed: 8873613]

55. Shao J, Irwin A, Hartson SD, Matts RL. *Biochemistry*. 2003; 42:12577–12588. [PubMed: 14580204]
56. Yun B-G, Huang W, Leach N, Hartson SD, Matts RL. *Biochemistry*. 2004; 43:8217–8229. [PubMed: 15209518]
57. Yu XM, Shen G, Neckers L, Blake H, Holzbeierlein J, Cronk B, Blagg BS. *J Am. Chem. Soc.* 2005; 127:12778–12779. [PubMed: 16159253]
58. Grenert JP, Sullivan WP, Fadden P, Haystead TA, Clark J, Mimnaugh E, Kruttsch H, Ochel HJ, Schulte TW, Sausville E, Neckers LM, Toft DO. *J. Biol. Chem.* 1997; 272:23843–23850. [PubMed: 9295332]
59. Owen BA, Sullivan WP, Felts SJ, Toft DO. *J. Biol. Chem.* 2002; 277:7086–7091. [PubMed: 11751892]
60. Morris GM, Goodsell DS, Halliday RS, Huey R, Hart WE, Belew RK, Olson AJ. *J. Comput. Chem.* 1998; 19:1639–1662.



**Figure 1.**

Effect of gambogic acid (1) and celastrol (2) on Hsp90-dependent luciferase refolding in reticulocyte lysate (A), and effect of 1 on cell proliferation of HeLa cells, and MCF7 and SkBr3 breast cancer cells. Experiments were carried out as described in the Experimental Section.




**Figure 2.**


Gambogic acid (**1**)-induced degradation of Hsp90 client proteins. Compound **1** was incubated with (A) MCF7 and (B) SkBr3 breast cancer cells at concentrations ( $\mu$ M) indicated in the figure. Gambogic acid (**1**) was evaluated for its ability to downregulate several client proteins as described in the Experimental Section. Geldanamycin (**3**) (500 nM) and DMSO were used as positive and negative controls, respectively. Cell extracts were prepared and equivalent amounts of protein were separated by SDS-PAGE and subsequently Western blotted for the indicated proteins as described in the Experimental Section.




## A. Co-immunoadsorption



## B. HRI Maturation




## C. HRI Stability



**Figure 3.**


Effect of Hsp90 inhibitors on the interaction of Hsp90 and its co-chaperones with HRI (A), on HRI's Hsp90-dependent maturation (B), and on HRI stability (C). A.  $^{35}\text{S}$ -His-tagged HRI/K199R was synthesized by TnT in reticulocyte lysate as described in the Experimental Section. After 10 min, DMSO (4% v/v, lanes 1 & 2), geldanamycin (3, 80  $\mu\text{M}$ , lane 3), sodium molybdate (20 mM, lane 4), celastrol (2, 100  $\mu\text{M}$ , lane 5), novobiocin (4, 4.0 mM, lane 6), coumermycin A1 (5, 400  $\mu\text{M}$ , lane 7), or gambogic acid (1, 50  $\mu\text{M}$ , lane 8) were added, followed by an additional 40 min of incubation.  $^{35}\text{S}$ -His-tagged HRI/K199R was then immunoadsorbed with anti-His antibodies and samples were analyzed for co-adsorbing Hsp90, Hsp70, and Cdc37 by SDS-PAGE and Western blotting. Lane 1: TnT lysate

containing no plasmid as the control for non-specific binding. Top panel: autoradiogram of immunoadsorbed [ $^{35}\text{S}$ ]HRI/K199R. B. [ $^{35}\text{S}$ ]His-tagged HRI was synthesized in reticulocyte lysate as described in the Experimental Section. After 20 min, DMSO (4% v/v, lanes 1, 2 and 3), geldanamycin (**3**, 80  $\mu\text{M}$ , lane 4), sodium molybdate (20 mM, lane 5), celastrol (**2**, 100  $\mu\text{M}$ , lane 6), novobiocin (**4**, 4.0 mM, lane 7), coumermycin A1 (**5**, 400  $\mu\text{M}$ , lane 8), or **1** (50  $\mu\text{M}$ , lane 9) were added, followed by an additional 10 min of incubation. An aliquot of the TnT lysate was then transferred to heme supplemented (20  $\mu\text{M}$ , lane 2) or heme-deficient (lanes 1 and 3–12) lysate containing and equivalent concentration of each addition, followed by an additional 45 min of incubation. The samples were then analyzed for HRI maturation by SDS-PAGE and autoradiography as described in the Experimental Section. Lane 1: TnT lysate containing no plasmid. [ $^{35}\text{S}$ ]HRI\*: mature, active HRI. C. [ $^{35}\text{S}$ ]His-tagged HRI was synthesized in reticulocyte lysate and treated with DMSO or Hsp90-inhibitors as described above. Aliquots of each reaction were taken prior to (upper panel, 0 min) and 45 min after (lower panel, 45 min) dilution into and incubation in heme-supplemented (lane 1) or heme-deficient (lanes 2–8) lysate. The band intensities in the lower panel were quantified by scanning densitometry and values below the lane numbers are given as %O.D./mm<sup>2</sup> of minus heme control.








**Figure 4.**

SPR analysis of the interaction of gambogic acid (**1**) with (A) full length Hsp90 and (B) the N-terminal domain of Hsp90. A. Injection of 1.0, 10, 25 and 50  $\mu\text{M}$  **1** over a SPR chip containing bound full length Hsp90. B. Injection of 0.5, 5, 15 and 25  $\mu\text{M}$  **1** over a SPR chip containing bound Hsp90NT. Black line: sensorgram of binding and dissociation; gray line with dots: curve fit.



**Figure 5.**

Models of gambogic acid (**1**) docked to geldanamycin-bound Hsp90NT. A. Ribbon diagram of Hsp90NT with the carbons of **1** and geldanamycin shown in yellow and green, respectively. B. Electrostatic surface potential of Hsp90NT shown in the same orientation as in A. C. Close-up showing the salt bridge and H-bond formed between K208 and K204 with **1**.





**Table 1**Constants for the Binding of Gambogic Acid (**1**) to Hsp90

protein	$k_a$ ( $M^{-1} S^{-1}$ )	$k_d$ ( $S^{-1}$ )	$K_D$ ( $\mu M$ )
full length Hsp90	$1.16(8)e^3$	0.0113(4)	9.8(2)
Hsp90NT	$1.47(6)e^3$	0.01122(7)	7.6(3)
Hsp90NT (+20 $\mu M$ GA)	$1.62(8)e^3$	0.0114(7)	7.0(4)
Hsp90CT	no binding		