

Published in final edited form as:

*J Am Chem Soc.* 2011 April 27; 133(16): 6158–6161. doi:10.1021/ja2013517.

## Conversion of Fatty Aldehydes to Alka(e)nes and Formate by a Cyanobacterial Aldehyde Decarbonylase: Cryptic Redox by an Unusual Di-metal Oxygenase

Ning Li<sup>1</sup>, Hanne Nørgaard<sup>2</sup>, Douglas M. Warui<sup>2</sup>, Squire J. Booker<sup>1,2,\*</sup>, Carsten Krebs<sup>1,2,\*</sup>, and J. Martin Bollinger Jr.<sup>1,2,\*</sup>

<sup>1</sup> Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University Park, Pennsylvania 16802, USA

<sup>2</sup> Department of Chemistry, The Pennsylvania State University, University Park, Pennsylvania 16802, USA

### Abstract

Cyanobacterial aldehyde decarbonylase (AD) catalyzes conversion of fatty aldehydes (R-CHO) to alka(e)nes (R-H) and formate. Curiously, although this reaction appears to be redox-neutral and formally hydrolytic, AD has a ferritin-like protein architecture and a carboxylate-bridged di-metal cofactor that are both structurally similar to those found in di-iron oxidases and oxygenases. In addition, the *in vitro* activity of the AD from *Nostoc punctiforme* (*Np*) was shown to require a reducing system similar to the systems employed by these O<sub>2</sub>-utilizing di-iron enzymes. Here, we resolve this conundrum by showing that aldehyde cleavage by the *Np* AD also requires dioxygen and results in incorporation of <sup>18</sup>O from <sup>18</sup>O<sub>2</sub> into the formate product. AD thus oxygenates, without oxidizing, its substrate. We posit that (i) O<sub>2</sub> adds to the reduced cofactor to generate a metal-bound peroxide nucleophile that attacks the substrate carbonyl and initiates a radical scission of the C1-C2 bond, and (ii) the reducing system delivers two electrons during aldehyde cleavage, ensuring a redox-neutral outcome, and two additional electrons to return an oxidized form of the cofactor back to the reduced, O<sub>2</sub>-reactive form.

Photosynthetic cyanobacteria use light to “fix” CO<sub>2</sub> into energy-rich biomolecules, including fatty acids.<sup>1,2</sup> A two-step pathway recently identified by Schirmer, et al. allows these organisms also to produce alkanes and alkenes from abundant saturated and unsaturated fatty acids.<sup>3</sup> Together, these two pathways could potentially be harnessed for a bioprocess that would effectively harvest solar energy, store it as a fungible fuel, and consume an important greenhouse gas.<sup>2</sup> The incalculable potential value of such a process has motivated a flurry of scientific,<sup>3,4</sup> intellectual-property,<sup>5,6</sup> and investment activity.<sup>7</sup> A deeper understanding of the cyanobacterial alkane-biosynthetic pathway might facilitate efforts to develop such a process.

The second step in the pathway is catalyzed by the enzyme aldehyde decarbonylase (AD<sup>8</sup>). Schirmer, et al. suggested that AD converts the C<sub>n</sub> fatty aldehyde product of the first enzyme into the corresponding C<sub>n-1</sub> alkane or alkene and carbon monoxide (CO),<sup>3</sup> but we recently showed that the C1-derived co-product from the *in vitro* conversion of octadecanal (R-<sup>13</sup>CHO, where R = *n*-C<sub>17</sub>H<sub>35</sub>) to heptadecane (R-H) by the *Nostoc punctiforme* (*Np*) AD

squire@psu.edu, ckrebs@psu.edu, jmb21@psu.edu.

Supporting Information Available: Figure illustrating determination of the ratio of NADPH oxidized to formate produced in the *Np* AD reaction. This material is available free of charge via the internet at <http://pubs.acs.org>

produced heterologously in *Escherichia coli* (*Ec*) is formate rather than CO.<sup>4</sup> On paper this reaction is formally *hydrolytic* (redox neutral). However, the structure of the *Prochlorococcus marinus* (*Pm*) MIT9313 ortholog showed that the ADs have a ferritin-like protein architecture and cofactor site that are similar to those found in non-heme di-iron oxygenases and oxidases,<sup>3</sup> including bacterial multicomponent monooxygenases (e.g., soluble methane monooxygenase)<sup>9,10</sup> and plant fatty acyl-ACP desaturases.<sup>11</sup> Moreover, the *in vitro* activity of the *Np* AD was observed (by both Schirmer, et al.<sup>3</sup> and us<sup>4</sup>) to require a reducing system (fulfilled by NADPH and spinach ferredoxin and ferredoxin reductase; hereafter N/F/FR), just as these di-iron oxidases and oxygenases require reducing systems to convert the Fe<sub>2</sub><sup>III/III</sup> “resting” forms of their cofactors to the O<sub>2</sub>-reactive Fe<sub>2</sub><sup>II/II</sup> states during each reaction cycle.<sup>9–11</sup> It appears that these analogies to known di-iron oxidases and oxygenases led to the incorrect depiction of the C1-derived co-product as CO<sub>2</sub> (which would be an oxidative outcome) and the designation of the cyanobacterial ADs as “decarboxylative monooxygenases” in a patent application from the company Joule Unlimited (Boston, MA).<sup>6</sup> The available data thus created a puzzle: how does an enzyme that looks and acts like an O<sub>2</sub>-utilizing oxidoreductase effect a formally hydrolytic reaction?

We suggested two possible resolutions to this conundrum.<sup>4,12</sup> First, the *Np* AD might emerge from expression and purification with its cofactor in an inactive oxidation state, requiring reduction of the cofactor and reaction with O<sub>2</sub> to regenerate the active state (Scheme 1A). This idea was founded on our understanding of class I ribonucleotide reductases, which (i) are structurally similar to the ADs,<sup>13,14</sup> (ii) contain stable, catalytically essential oxidants (a tyrosyl radical in class Ia<sup>15–17</sup> and Ib<sup>18,19</sup> and the Mn<sup>IV</sup> ion of a Mn/Fe cluster in class Ic<sup>20–22</sup>), (iii) can be inactivated by reduction of their essential oxidants,<sup>15,23,24</sup> and (iv) in the case of the class Ia enzymes, can then be reactivated by reduction of their Fe<sub>2</sub><sup>III/III</sup> clusters and subsequent reaction of the Fe<sub>2</sub><sup>II/II</sup> forms with O<sub>2</sub>.<sup>25</sup> Second, O<sub>2</sub> might be an obligatory co-substrate in some sort of (heretofore unprecedented) cryptically redox pathway for the formally hydrolytic aldehyde cleavage. This case would require that four electrons be delivered by the reducing system in each turnover for redox balance (Scheme 1B).

Both potential resolutions imply that O<sub>2</sub> should be required for AD activity, a prediction not tested to date. More diagnostically, if the N/F/FR reducing system and possibly O<sub>2</sub> should be required only for reactivation of the AD, but not for turnover, then it might be possible to observe activity after first exposing the enzyme to both in the absence of substrate and then removing one or both prior to initiating the reaction by addition of the substrate. By contrast, the second scenario would imply that omission of O<sub>2</sub> or any component of the reducing system should prevent turnover, irrespective of any pre-treatment. To clarify the nature and mechanism of the novel AD reaction, we experimentally evaluated these predictions.

We first assessed whether O<sub>2</sub> is required for activity. One set of reaction samples (see Figure 1 legend for their composition) was constituted with O<sub>2</sub>-free components inside an anoxic chamber (MBraun, Peabody, MA) by adding a pre-mixed solution of the N/F/FR reducing system and the *Np* AD (the affinity-tagged enzyme used in our previous study and purified from *Ec* as described therein<sup>4</sup>) to a solution containing the R-<sup>13</sup>CHO substrate and Triton x-100 detergent (0.2% in the complete sample). A matched set of samples was prepared by removing the pre-mixed solution of N/F/FR reducing system and *Np* AD from the anoxic chamber and adding it under ambient atmosphere to an air-saturated solution of the substrate and detergent. These two sets of samples were incubated at ambient temperature (23 ± 1 °C) for between 1.5 and 20 h and then analyzed for [<sup>13</sup>C]-formate by conversion to its 2-nitrophenylhydrazide (2NPH) derivative and quantification by liquid chromatography and mass spectrometry (LC-MS), as previously described.<sup>4</sup> Mass spectra of the samples incubated under air exhibit intense peaks at *m/z* = 181 (Figure 1A, black bars), indicating

that, as expected, formate was produced in these reactions (145  $\mu\text{M}$  in Figure 1A). Note that the peak at  $m/z = 180$  in the mass spectra of all samples arises from contaminating environmental formate, which also contributes weakly at  $m/z = 181$  due to the presence of natural-abundance M+1 isotopic species ( $\sim 8.9\%$  of the  $m/z = 180$  species) in its 2NPH derivative. By contrast, spectra of the samples prepared and incubated in the absence of  $\text{O}_2$  have much weaker peaks at  $m/z = 181$  (Figure 1A, gray bars), implying that much less formate (9  $\mu\text{M}$ ) was produced in these reactions. This result demonstrates that the *Np* AD requires  $\text{O}_2$  in some capacity. To assess whether  $\text{O}_2$  is required for activation (Scheme 1A) or catalysis (Scheme 1B), an additional matched sample was prepared by removing the solution of N/F/FR system and *Np* AD from the anoxic chamber and exposing it to air for 30 min, with periodic mixing to ensure permeation of  $\text{O}_2$  into the enzyme solution. The air-exposed (N/F/FR + *Np* AD) solution was then returned to the anoxic chamber and mixed therein with 9 equivalent volumes of an  $\text{O}_2$ -free solution of substrate and detergent. The sample was placed in a sealed vessel, incubated for 20 h, and analyzed as above. The weak peak at  $m/z = 181$  (Figure 1A, orange bars) implies that prior exposure of the *Np* AD to  $\text{O}_2$  in the presence of the reducing system does not activate it for subsequent  $\text{O}_2$ -independent turnover. These results suggest that  $\text{O}_2$  must be continuously present for activity.

In addition to scission of the C1-C2 bond and formation of a new C2-H bond, a second O-atom is added to C1 in conversion of R-CHO to R-H and  $\text{HCO}_2^-$ . The requirement for  $\text{O}_2$  raised the possibility that *it* could be the source of this O-atom, as occurs in other oxygenase reactions. Because substrate oxygenation *without net oxidation* is, to the best of our knowledge, unprecedented, we considered this possibility to be unlikely but tested for it nevertheless. Surprisingly, the mass spectrum of a sample in which the reaction was carried out under an atmosphere of  $^{18}\text{O}_2$  (Figure 1B, blue bars) exhibits a strong peak at  $m/z = 183$  (an increase of two mass units) that is absent in the spectrum of a control sample prepared identically but with natural-abundance  $\text{O}_2$  (red bars). It is important to note here that the analysis for formate, which involves its prior conversion to formyl-2NPH, leads to loss of one of two equivalent oxygen atoms, and so quantitative incorporation of a single  $^{18}\text{O}$  atom into the formate product should result in precisely 50% of the derivative having  $m/z = 183$  (the other 50% having  $m/z = 181$ ). Integration of the peaks in Figure 1B and analogous spectra from two other trials gave only  $(34 \pm 3)\%$  (mean and standard deviation) of the  $m/z = 183$  species. However, abortive coupling in the derivatization (i.e., carboxylate attack on the activating carbodiimide, EDC, followed by hydrolysis of the adduct) would lead to further loss of  $^{18}\text{O}$  and diminution of the  $m/z = 183$  peak relative to the  $m/z = 181$  peak. To assess whether the loss of  $\sim 16\%$  of the  $m/z = 183$  species in Figure 1B can be explained this way or reflects partial exchange of the  $\text{O}_2$ -derived atom with solvent during the AD reaction, control samples were constituted in  $\text{H}_2^{18}\text{O}$  (70% enrichment), subjected to the 2NPH coupling procedure, and analyzed by LC-MS (Figure 1C). They contained all reaction components except the substrate and were amended with either natural-abundance formate or  $[^{13}\text{C}]$ -formate. Prominent peaks at  $m/z = 182$  for the sample containing natural-abundance formate (green bars) and  $m/z = 183$  for the sample with  $\text{H}^{13}\text{CO}_2^-$  (purple bars) demonstrate that, as expected, the coupling reaction does result in partial exchange of the formate O-atoms with solvent. Comparison of the integrated intensities of the peaks corresponding to the  $^{16}\text{O}$ - and  $^{18}\text{O}$ -containing formyl-2NPH isotopologues gave 15% of the latter species, which corresponds to 20% when the  $^{18}\text{O}$  content of 70% is taken into account. This extent of exchange fully accounts for the diminution of the heavier isotope from the theoretical maximum of 50% in the  $^{18}\text{O}_2$  experiments in Figure 1B. These results show that (i)  $\text{O}_2$  is the source of the incorporated oxygen and (ii) to the limit of our analysis, there is no demonstrable exchange with solvent during the enzyme reaction. The AD reaction is thus the first example (of which we are aware) of substrate oxygenation without net oxidation.

The metal ions in the AD cofactor have not yet been identified. Nevertheless, the vast majority of proteins in this structural family that have been studied to date have di-iron clusters. In the reactions of several of these enzymes,  $(\mu\text{-peroxo})\text{-Fe}_2^{\text{III/III}}$  intermediates have been identified.<sup>26–31</sup> These intermediates either undergo O-O-bond cleavage to give high-valent ( $\text{Fe}_2^{\text{III/IV}}$  or  $\text{Fe}_2^{\text{IV/IV}}$ ) complexes that effect oxidation reactions or directly react as electrophiles (e.g., via attack of the  $\pi$ -electrons of an olefinic or aromatic substrate or the nitrogen of an amine or hydroxylamine), also leading to oxidative outcomes. However, studies of heme enzymes (cytochromes P-450) and inorganic complexes have shown that metal-bound peroxides can also act as *nucleophiles* to attack (among other electrophiles) aldehydes, leading to production of formate and oxidized co-products.<sup>32–38</sup> Moreover, recent computational studies by Hirao and Morokuma on *myo*-inositol oxygenase and hydroxyethylphosphonate dioxygenase suggest that, in each reaction, the nucleophilic attack of an Fe-coordinated peroxide on the carbonyl of a reaction intermediate may be a key step in effecting C-C-bond cleavage.<sup>39,40</sup> These reactions would provide precedent for the early steps of the hypothetical AD mechanism shown in Scheme 2. It involves addition of  $\text{O}_2$  to the reduced  $\text{Fe}_2^{\text{II/II}}$  cofactor (**A**) to form a peroxide intermediate (**B**) and attack of the peroxide on the C1 carbonyl of the substrate to form a peroxyhemiacetal complex (**C**). Breakdown of this intermediate leading to O-O and C1-C2 scission would explain production of formate with one O-atom from  $\text{O}_2$ . The intriguing and (to date) unprecedented aspect of this hypothetical mechanism would be breakdown of the peroxyhemiacetal intermediate to convert C2 of the substrate into a fully reduced (methyl) rather than partially oxidized (e.g. alcohol or olefinic methylene) center in the R-H product. One possibility, shown in Scheme 2, is that the reducing system would deliver an electron during cleavage of the O-O bond, forming a *gem*-diolyl radical and  $(\mu\text{-oxo})\text{-Fe}_2^{\text{III/III}}$  cluster (**D**). The *gem*-diolyl intermediate could undergo radical fragmentation of the C1-C2 bond, generating formate and the  $\text{R}^\bullet$  radical (**E**). Transfer of a hydrogen atom, either from the cofactor (as depicted) or from an amino acid in the active site, would produce R-H and either a  $(\mu\text{-oxo})\text{-Fe}_2^{\text{III/IV}}$  complex (analogous to the **X** intermediate that generates the tyrosyl radical in a class I RNR<sup>41,42</sup>) or an amino acid radical. The reducing system would then complete the reaction by quenching this remaining oxidizing equivalent, and subsequently deliver two more electrons to convert the  $\text{Fe}_2^{\text{III/III}}$  “product” form of the cofactor (**F**) back to the  $\text{O}_2$ -reactive  $\text{Fe}_2^{\text{II/II}}$  “reactant” state (**A**) in preparation for the next turnover.

The mechanism in Scheme 2 accounts for all the available data of which we are aware, including the requirement for a reducing system and the continuous presence of  $\text{O}_2$ , the identity of the C1-derived co-product (formate),<sup>4</sup> the origin of the new O-atom in the formate, and the origins of the hydrogen atoms in both formate and R-H products.<sup>4</sup> It also predicts an NADPH:formate (NADPH:R-H) reaction stoichiometry of 2:1 in order to account for the four electrons needed in each turnover for complete reduction of  $\text{O}_2$  (Scheme 1B). We tested this prediction by spectrophotometric determination of NADPH consumption and parallel quantification of formate at two reaction times (Figure S1). Experimental NADPH:formate ratios of 5.6 after 1 h and 6.7 after 2 h were calculated from the results. These ratios are sufficiently large to be consistent with the mechanism of Scheme 2 and also imply considerable uncoupling of NADPH consumption from R-CHO cleavage. This uncoupling is unsurprising, given that the reaction is so slow and that we are employing a heterologous reducing system (ferredoxin and ferredoxin reductase from spinach) to transfer the electrons from the reduced nicotinamide to the AD cofactor. It should be more informative to re-determine this ratio with the native reducing system, once it has been identified.

The mechanism of Scheme 2 implies that the cyanobacterial AD cofactor could indeed be a  $\text{Fe}_2$  cluster, as originally suggested by Schirmer, et al.<sup>3</sup> However, the unprecedented aspects of the mechanism leave room for the possibility that a different transition metal might be

present at one or both sites. Such a substitution might, for example, permit formation of a peroxyhemiacetal intermediate less disposed toward an oxidative outcome than the depicted Fe<sub>2</sub> complex is expected to be on the basis of the cited precedents. Similarly, mechanisms in which a metal-bound superoxide attacks C1 can also be formulated. In this case, a mixed-metal cluster might disfavor further reduction of the O<sub>2</sub> unit to the peroxide state and give the superoxo complex sufficient time to add to the carbonyl. Clearly, identification of the metals in the cofactor is urgently needed for a deeper understanding of this novel reaction.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments


This work was supported by the National Institutes of Health (GM-63847 to S.J.B.) and the Dreyfus Foundation (Teacher-Scholar Award to C.K.).

## References

1. Bryant, DA. The Molecular Biology of Cyanobacteria. Kluwer Academic Publishers; Dordrecht: 1994.
2. Ducat DC, Way JC, Silver PA. Trends in Biotechnology. 2011; 29:95–103. [PubMed: 21211860]
3. Schirmer A, Rude MA, Li X, Popova E, del Cardayre SB. Science. 2010; 329:559–562. [PubMed: 20671186]
4. Warui DM, Li N, Nørgaard H, Krebs C, Bollinger JM Jr, Booker SJ. J Am Chem Soc. 2011; 133:3316–3319. [PubMed: 21341652]
5. Schirmer, A.; Rude, M.; Brubaker, S. LS9. patent application. WO/2009/140696. 2009.
6. Reppas, NB.; Ridley, CP. patent. 7794969. 2010.
7. Service RF. Science. 2009; 325:379. [PubMed: 19628830]
8. Abbreviations: 2NPH, 2-nitrophenyl-hydrazide; AD, aldehyde decarbonylase; *Ec*, *Escherichia coli*; EDC, 1-ethyl-3-(3-dimethylaminopropyl)-carbodiimide; GC, gas chromatography; LC, liquid chromatography; MS, mass spectrometry; N/F/FR, NADPH, ferredoxin, ferredoxin reductase reducing system; *Np*, *Nostoc punctiforme*; *Pm*, *Prochlorococcus marinus* MIT9313; R, *n*-C<sub>17</sub>H<sub>35</sub>; SIM, single ion monitoring.
9. Merckx M, Kopp DA, Sazinsky MH, Blazyk JL, Müller J, Lippard SJ. Angew Chem Int Ed. 2001; 40:2782–2807.
10. Wallar BJ, Lipscomb JD. Chem Rev. 1996; 96:2625–2657. [PubMed: 11848839]
11. Fox BG, Lyle KS, Rogge CE. Acc Chem Res. 2004; 37:421–429. [PubMed: 15260504]
12. Krebs C, Bollinger JM Jr, Booker SJ. Curr Opin Chem Biol. 2011; 10.1016/j.cbpa.2011.02.019
13. Nordlund P, Sjöberg BM, Eklund H. Nature. 1990; 345:593–598. [PubMed: 2190093]
14. Nordlund P, Eklund H. Curr Opin Struct Biol. 1995; 5:758–66. [PubMed: 8749363]
15. Atkin CL, Thelander L, Reichard P, Lang G. J Biol Chem. 1973; 248:7464–7472. [PubMed: 4355582]
16. Sjöberg BM, Reichard P, Gräslund A, Ehrenberg A. J Biol Chem. 1977; 252:536–541. [PubMed: 188819]
17. Stubbe J. Curr Opin Chem Biol. 2003; 7:183–188. [PubMed: 12714050]
18. Cotruvo JA, Stubbe J. Biochemistry. 2010; 49:1297–1309. [PubMed: 20070127]
19. Cox N, Ogata H, Stolle P, Reijerse E, Auling G, Lubitz W. J Am Chem Soc. 2010; 132:11197–11213. [PubMed: 20698687]
20. Jiang W, Yun D, Saleh L, Barr EW, Xing G, Hoffart LM, Maslak MA, Krebs C, Bollinger JM Jr. Science. 2007; 316:1188–1191. [PubMed: 17525338]
21. Jiang W, Yun D, Saleh L, Bollinger JM Jr, Krebs C. Biochemistry. 2008; 47:13736–13744. [PubMed: 19061340]




22. Bollinger JM Jr, Jiang W, Green MT, Krebs C. *Curr Opin Struct Biol.* 2008; 18:650–657. [PubMed: 19046875]
23. Stubbe, J. *Advances in enzymology and related areas of molecular biology.* Vol. 63. Wiley; 1990. p. 349–419.
24. Jiang W, Xie J, Varano PT, Krebs C, Bollinger JM Jr. *Biochemistry.* 2010; 49:5340–5349. [PubMed: 20462199]
25. Wu CH, Jiang W, Krebs C, Stubbe J. *Biochemistry.* 2007; 46:11577–11588. [PubMed: 17880186]
26. Liu KE, Wang D, Huynh BH, Edmondson DE, Salifoglou A, Lippard SJ. *J Am Chem Soc.* 1994; 116:7465–7466.
27. Tong WH, Chen S, Lloyd SG, Edmondson DE, Huynh BH, Stubbe J. *J Am Chem Soc.* 1996; 118:2107–2108.
28. Yun D, García-Serres R, Chicales BM, An YH, Huynh BH, Bollinger JM Jr. *Biochemistry.* 2007; 46:1925–1932. [PubMed: 17256972]
29. Murray LJ, Naik SG, Ortillo DO, García-Serres R, Lee JK, Huynh BH, Lippard SJ. *J Am Chem Soc.* 2007; 129:14500–14510. [PubMed: 17967027]
30. Korboukh VK, Li N, Barr EW, Bollinger JM Jr, Krebs C. *J Am Chem Soc.* 2009; 131:13608–13609. [PubMed: 19731912]
31. Li N, Korboukh VK, Krebs C, Bollinger JM Jr. *Proc Natl Acad Sci USA.* 2010; 107:15722–15727. [PubMed: 20798054]
32. LeCloux DD, Barrios AM, Lippard SJ. *Bioorg Med Chem.* 1999; 7:763–772. [PubMed: 10400329]
33. Wertz DL, Sisemore MF, Selke M, Driscoll J, Valentine JS. *J Am Chem Soc.* 1998; 120:5331–5332.
34. Wertz, DL.; Valentine, JS. *Metal-Oxo and Metal-Peroxo Species in Catalytic Oxidations.* Vol. 97. Springer-Verlag Berlin; Berlin: 2000. p. 37–60.
35. Annaraj J, Suh Y, Seo MS, Kim SO, Nam W. *Chem Comm.* 2005:4529–4531. [PubMed: 16158103]
36. Cho J, Sarangi R, Annaraj J, Kim SY, Kubo M, Ogura T, Solomon EI, Nam W. *Nature Chemistry.* 2009; 1:568–572.
37. Annaraj J, Cho JH, Lee YM, Kim SY, Latifi R, de Visser SP, Nam W. *Angew Chem - Int Ed.* 2009; 48:4150–4153.
38. Cho J, Sarangi R, Kang HY, Lee JY, Kubo M, Ogura T, Solomon EI, Nam W. *J Am Chem Soc.* 2010; 132:16977–16986. [PubMed: 21062059]
39. Hirao H, Morokuma K. *J Am Chem Soc.* 2009; 131:17206–17214. [PubMed: 19929019]
40. Hirao H, Morokuma K. *J Am Chem Soc.* 2010; 132:17901–17909. [PubMed: 21121666]
41. Bollinger JM Jr, Edmondson DE, Huynh BH, Filley J, Norton JR, Stubbe J. *Science.* 1991; 253:292–298. [PubMed: 1650033]
42. Sturgeon BE, Burdi D, Chen S, Huynh BH, Edmondson DE, Stubbe J, Hoffman BM. *J Am Chem Soc.* 1996; 118:7551–7557.



**Figure 1.**


Reconstructed mass spectra illustrating the catalytic requirement for O<sub>2</sub> and the incorporation of <sup>18</sup>O from <sup>18</sup>O<sub>2</sub> into the formate product in the *Np* AD reaction. **(A)** Reactions in the continuous presence of O<sub>2</sub> (under air atmosphere, black bars), the absence of O<sub>2</sub> (gray), and the presence of O<sub>2</sub> during pre-incubation with the N/F/FR system (without substrate) but the absence of O<sub>2</sub> during the reaction (orange). **(B)** Reactions under an atmosphere of natural-abundance O<sub>2</sub> (red) or <sup>18</sup>O<sub>2</sub> (99% isotopic purity; blue). **(C)** Control reactions in which the 2NPH derivative of either natural-abundance formate (green) or [<sup>13</sup>C]-formate (purple) was generated in H<sub>2</sub><sup>18</sup>O (70% enrichment) to quantify the extent of exchange of the oxygen atoms during the coupling reaction. The reactions in **A** were carried out at 21 °C for 20 h and contained, in a final volume of 0.40 mL, 0.10 mM *Np* AD, 0.5 mM R-<sup>13</sup>CHO substrate, 2 mM NADPH, and 100 µg/mL each of spinach ferredoxin and ferredoxin reductase in air-saturated 100 mM HEPES buffer, pH 7.4, containing 0.2% triton X-100. The reactions in **B** had the same composition but were carried out for 2 h.



**Scheme 1. Two Alternative Explanations for the Similarity of *Np* AD to Di-iron Oxidases and Oxygenases and its Requirement for a Reducing System to Promote an Apparently Hydrolytic Reaction<sup>a</sup>**

<sup>a</sup>(A) The reducing system provides  $n$  electrons ( $\text{e}^-$ ), and the most reduced form of the cofactor reacts with  $\text{O}_2$  to generate the active state, which is more oxidized than the as-isolated form by  $4 - n$  units ( $x + y - i - j = 4 - n$ ); (B)  $\text{O}_2$  is reduced in every cycle by four electrons from the reducing system.





**Scheme 2.**  
Hypothetical Mechanism for the *Np* AD Reaction