

Published in final edited form as:

Clin Gastroenterol Hepatol. 2014 May ; 12(5): 738–748. doi:10.1016/j.cgh.2013.06.006.

Hepatitis B and C in African Americans: Current Status and Continued Challenges

Kimberly A. Forde, MD MHS^{1,2}, Orapin Tanapanpanit, MD¹, and K. Rajender Reddy, MD¹

¹Division of Gastroenterology and Hepatology, Department of Medicine, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA.

²Center for Clinical Epidemiology and Biostatistics, Department of Biostatistics and Epidemiology, Perelman School of Medicine, University of Pennsylvania, Philadelphia, PA.

Abstract

Viral hepatitis remains a public health concern in the US, resulting in excess morbidity and mortality for the individual and representing a burden to societies as evidenced by billions of dollars in healthcare expenditures. As with many chronic diseases, race and ethnicity influence various aspects of disease pathogenesis including mechanisms of persistence, disease progression, disease sequelae and response to therapy. For hepatitis B and C infections, African Americans disproportionately bear a large burden of disease in the US. The role and importance of African American race however has been less well characterized in the literature among the population of viral hepatitis-infected individuals. The differences in epidemiology, manifestations of liver disease, response to therapy and differential trends in liver transplantation in African Americans compared to other racial and ethnic groups deserve special attention.

This review will address the current status of hepatitis B and C infection in African Americans in the US and identify some of the remaining challenges in diagnosis, characterization of natural history, and treatment. For purposes of this review, the terms African American and black will be used interchangeably throughout the text.

Hepatitis B Infection

Epidemiology of Hepatitis B Infection

Globally, more than 2 billion persons have evidence of hepatitis B virus (HBV) infection and nearly 200 – 375 million of those have serologic evidence of chronic infection.¹ While prior estimates of HBV infection in the US have ranged from 800,000 – 1.4 million, emerging data suggest that this is an underestimate of disease burden, with infection in 2.2

© 2013 The American Gastroenterological Association. Published by Elsevier Inc. All rights reserved.

Corresponding Author: K. Rajender Reddy, MD Professor of Medicine Director of Hepatology Perelman School of Medicine at the University of Pennsylvania Division of Gastroenterology and Hepatology 3400 Spruce Street, 2 Dulles, Philadelphia, PA 19104 Telephone: (215) 662-4311, Fax: (215) 615-1601 rajender.reddy@uphs.upenn.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

The authors have no conflicts of interest to disclose.

million persons being proposed and of which immigrants bear the heaviest burden.¹⁻³ Furthermore, an estimated 1.4 million HBV infected individuals are unaware of their infectious status and important opportunities for treatment of chronic infection and surveillance for HCC are being missed.⁴

In the US, an estimated 35,000 new diagnoses of acute HBV infection were established in 2010.⁵ This represents a decline of nearly 80% when compared to the rate of new HBV infections in the early 1990s, prior to implementation of HBV vaccination programs. When examining incidence rates stratified by race and ethnicity, African Americans have the highest incidence of acute HBV infection, with 1.7 cases per 100,000 persons reported in 2010. This is in contrast to a rate of 0.6 per 100,000 persons in Asian Pacific Islanders and Hispanics, groups in the US with the lowest HBV incidence.⁵ Consistently, the incidence rate of acute HBV infection for blacks has been higher than that of other racial and ethnic groups since 2000.

Data from the National Health and Nutrition Examination Survey (NHANES) II and III suggest that the age-adjusted prevalence of chronic infection or serologic markers of past exposure to HBV in blacks was 5.5% and 4.9%, respectively; rates that failed to change over the years of study.⁶ In this group, rates of chronic infection ranged from 0.33-0.42%. In addition to identifying foreign-born status as a risk factor for increased HBV prevalence, as defined by hepatitis B surface antigen (HBsAg) or hepatitis B core (HBc) positivity, African American race was associated with an age-adjusted prevalence of 15.8% for years 1976-1980 and 11.9% for years 1988-1994. Furthermore, being black was associated with a 3.9 fold increase in odds of HBV positivity in comparison to whites.⁶ Additional analyses in NHANES data found that differences in HBV prevalence rates between African Americans and other groups persisted after adjustment for important risk factors including demographic, socioeconomic and behavioral characteristics.⁷ These data suggest that factors other than socioeconomics and other risk factors account for an increased prevalence of HBV infection in African Americans.

Natural History of Hepatitis B infection

The natural history of hepatitis B infection, including its incubation period, symptomatology and chronicity, has been well described.⁸ No studies to date have examined differences in the natural history of HBV in African Americans and made comparisons to other racial and ethnic groups.

Chronic HBV infection may be asymptomatic or result in a range of manifestations of liver disease including chronic hepatitis, cirrhosis or hepatocellular carcinoma (HCC). With regards to HCC, African Americans assume a greater burden of disease with a notable increase in both HCC incidence and HCC-related mortality when compared to whites.^{9, 10} Yu and colleagues compared HCC risk factors in African Americans and whites. They found that African Americans had higher rates of viral hepatitis, coinfection with HBV and HCV, and metabolic risk factors such as diabetes mellitus that increased their risk of HCC.¹¹ Additionally, given the high prevalence of genotype A HBV infection in blacks, a genotype associated with a 4-5 fold increase in risk of HCC in black Africans,¹² African Americans may have risk factors for HCC that result in a markedly elevated and additive risk of HCC.

Unfortunately, comparison data were not available for other racial groups such as Asians and Hispanics in the above study. Clearly, more research is required to further elucidate these aspects of HCC risk in African American patients with chronic HBV infection.

Because of the high risk of HCC in the context of viral hepatitis for many patient populations, surveillance for HCC is now recommended in select groups.¹³ One such group, Sub-Saharan Africans, not only tend to have a high incidence of HCC but also a high incidence of HCC at a younger age.¹⁴ In a cohort of Africans with HCC, HBV serologies including HBsAg and HBeAg were more prevalent in patients less than 30 years of age.¹⁵ Though there are several studies of Africans that have examined their risk of HCC and demonstrated the importance of HCC surveillance in this group, such data are not available for blacks from other regions including the US.¹⁵⁻¹⁷ However, given the increase in HBV infection in African Americans and their higher risk of HCC, it may be prudent to extend surveillance recommendations to this high-risk population.

Viral Genotype and Host Genetic Diversity in Hepatitis B Infection

HBV Genotypic Variability—HBV genotype is an important determinant of disease progression with a notable influence on disease severity and response to HBV-directed therapy.¹⁸⁻²³ A review of HBV genotypes suggested that there are differences in HBV genotype distribution by race and ethnicity and place of birth. With respect to African Americans, 84% were infected with genotype A followed by genotypes C and D at 6% and 4%, respectively.¹⁹ In whites, genotypes A and D were most prevalent, as seen in 71 and 21% of the cohort, respectively. In comparison, genotypes B and C were observed most frequently in Asians. In the study, genotypes A and C were associated with HBeAg and non-genotype B infection was an independent risk factor for abnormal alanine aminotransferase levels and decompensated cirrhosis. While data are limited, it can be gleaned that there may be differences in disease progression that may be mediated by differences in viral genotype distribution.

Host Genetic Diversity—Interleukin-10 (IL-10) promoter polymorphisms have been associated with HBV infectious outcomes, including spontaneous HBV clearance, HBeAg seroconversion, development of chronic liver disease and incidence of HCC.^{24, 25} Additionally, foci near the IL-19 and IL-20 genes have been associated with regulation of pro-inflammatory cytokine production early in viral infections. To further explore racial differences in response to acute HBV infection, Truelove and colleagues examined the effect of single nucleotide polymorphisms (SNPs) in the IL-10, IL-19 and IL-20 genes in African Americans and whites in a nested case-control study.²⁶ It was found that African Americans who were heterozygotes at the rs1518108 locus of IL-20 were more susceptible to chronic HBV infection. This locus is also of importance in the genetics of host response in hepatitis C virus (HCV) infection.²⁷ Furthermore, polymorphisms in the IL10 promoter region were observed to be an important determinant of HBV infection. In African Americans, SNP variants at the IL-10 and IL-20 loci determined the establishment of HBV infection whereas in those of European ancestry, IL-20 variants solely influenced recovery. Though more work needs to be done to clarify the genetic and immunologic basis of the development of HBV

infection stratified by race, differences in host genetics may contribute to the racial and ethnic differences observed in HBV prevalence.

Treatment and Outcomes in Chronic Hepatitis B Infection

There is a relative scarcity of data regarding treatment outcomes for chronic HBV infection in African Americans following interferon (IFN) and/or nucleoside/nucleotide based antiviral therapy. A long-term follow up study of patients who had undergone therapy with IFN observed that African American patients were much more likely to respond to therapy.²⁸ Of note, all African American responders not only cleared HBeAg and HBV DNA from the serum, but all cleared HBsAg, a relatively rare milestone with HBV therapy that occurs in up to 7.8% of patients on therapy.²⁹ Unfortunately, there were few African Americans who participated in the study (n=6) and these observations have yet to be further explored. With respect to use of nucleotide/nucleoside analogues, there is inadequate data about the efficacy and safety of these agents in African American patients with chronic HBV infection (Table 1).

Likewise, there are limited data on the number of treatment eligible African Americans with HBV infection that are initiated on therapy. A small study including subjects from an urban medical center found that only 7% of a predominantly African American and Hispanic population had been initiated on therapy.⁴⁵ The study also highlighted potential barriers to therapy including lack of insurance, history of non-adherence, and ongoing drug and alcohol abuse.

Liver Transplantation in Hepatitis B Infection

Sentinel work has demonstrated that disparities exist for those African American patients awaiting liver transplantation. In addition to being more likely to die and becoming too sick for liver transplantation, African Americans are less likely to be transplanted once listed.⁴⁶ Subsequent to this work in the pre-Model for End Stage Liver Disease (MELD) era, Moylan et al. demonstrated an improvement in racial disparities in liver transplantation that coincided with implementation of the MELD-based allocation system for liver transplantation that was adopted in 2002.⁴⁷ Though this data would seem to suggest a more equitable allocation of organ donation since implementation of the MELD score, the work also demonstrated that blacks are listed at higher MELD scores and their waiting times are shorter than whites, suggesting that access to care may still be a barrier for African Americans. Additionally, though there is a high prevalence of liver disease in African Americans, they represented a smaller proportion of those listed for liver transplantation than expected.

Shifting the focus to liver transplantation for HBV infection, the data on outcomes for African Americans are conflicting. In a review of outcomes after liver transplantation for HBV in the US, Kim et al. found that African Americans had a 35% increase in mortality compared to whites in the pre-MELD era.⁴⁸ However, Bzowej et al. failed to demonstrate race-based differences in HBV waitlist or post transplant outcomes.⁴⁹ This discrepancy in results, likely resulting from the limitations of observational studies and limited inclusion of

African Americans, speaks to the need for more research to address this and other questions regarding racial disparities in transplantation.

Challenges and Future Directions

Racial disparities exist in prevalence of chronic HBV infection in the US and there are unmet needs in those African Americans with chronic HBV (Table 2). Though those who are foreign-born bear the majority of the burden of HBV infection in the US, African Americans have a high prevalence of infection. Additionally, there remain differences in acquisition of infection, with African Americans having the highest rates of incident infection. Data presented also suggest that there are racial differences in genetic polymorphisms important in establishment of chronic infection and modulation of disease course. There are also vast differences in viral genotypic distribution by race and likely differences in response to therapy, a relationship that is not yet delineated. Continued research is needed to identify and address areas of disparity in chronic HBV infection and treatment and an improvement in enrollment of African Americans in clinical trials is essential for achievement of this goal.

Hepatitis C Infection

Epidemiology of Hepatitis C Infection

While African Americans remain a minority in the US, HCV infection is highly prevalent in this population compared to other racial groups,^{50, 51} and thus African Americans represent a group disproportionately affected by this disease. In NHANES III data, inclusive of years 1988 through 1994, it was estimated that approximately 3.9 million Americans, or 1.8% of the US population, had been exposed and 2.7 million, or 74%, chronically infected with HCV.⁵² Blacks had the highest prevalence of HCV exposure, equating to 3.2% of the population screened.

The NHANES III survey as well as a follow up study examining HCV prevalence in the US from 1999 through 2002, highlighted the tremendous HCV disease burden in those born between 1945 and 1965. Of note, the highest prevalence, 9.4%, was observed in black men 40-49 years.⁵³ Given the potential complications of chronic HCV infection, including cirrhosis, hepatic decompensation and HCC, it is this cohort of patients which is expected to have HCV-related complications. Mathematical modeling studies have suggested that the proportion of cases with advanced hepatic fibrosis secondary to HCV infection will continue to rise during the next decade with the number of cases of cirrhosis and hepatic decompensation peaking at years 2020 and 2022, respectively.⁵⁴ Unfortunately, given the burden of HCV infection in African Americans and other racial and ethnic minorities, these groups are likely to have increased rates of HCV-related death. Mortality data from 1999 – 2007 obtained from the National Center for Health Statistics suggest that the effect of HCV on mortality is already being seen. The mortality rate for HCV-related disease surpassed that of death among HIV-infected patients, a rate on the decline, and HBV-related death, a rate that has been relatively constant. This study also demonstrated racial and ethnic disparities in death rates for HCV infection, with African Americans having a 2 fold higher rate of death than expected.⁵⁵

Natural History of Hepatitis C Infection

Longitudinal studies addressing the natural history of HCV infection in African Americans are lacking. However cross-sectional and retrospective studies have observed a lower prevalence of cirrhosis among African Americans when compared with other groups.⁵⁶⁻⁵⁹ Additionally, most studies found a lower ALT, and less piecemeal necrosis and hepatic fibrosis in African Americans.^{56, 57} In contrast, a more recent study using mathematical modeling of disease progression suggested that there was little difference based on race in progression of hepatic fibrosis in HCV infection.⁶⁰ Hence, data on racial differences in hepatic fibrosis remain inconclusive. The rationale for differences in HCV natural history and disease progression based on racial and ethnicity are not completely understood and further research is necessary to determine mechanisms underlying these potential differences.

Given the differences suspected in the natural history of HCV infection in African Americans, it would be expected that they would have fewer disease related complications including hepatic decompensation and a lower incidence of HCC. No studies to date have explored differences in rates of hepatic decompensation stratified by race. However, there are race-based disparities noted in incidence of HCC. Recent data suggest that the incidence rate of HCC is 2 to 3 fold higher among blacks than whites.⁶¹ Moreover, African American men bear the highest burden of HCC, with an incidence rate of 5.4% per year noted, surpassing incidence rates for whites, Hispanics, and women.^{54, 62, 63}

Viral Genotype and Host Genetic Diversity in Hepatitis C Infection

HCV Genotypic Variability—HCV genotype is one of the major determinants of response to antiviral therapy. Genotype 1 infection has been recognized as being difficult to treat and is more frequent in the US, Europe and Japan.⁶⁴ Genotype 1 is the predominant genotype among African Americans, with 91% of this population infected with this genotype, in comparison to 70% of whites.^{52, 53, 65} Based on data from NHANES, African Americans are twice as likely as whites to be infected with genotype 1b.^{53, 66} Moreover, among genotype 1 infected patients, African Americans clear the virus at approximately half the rate of their white counterparts.⁶⁷⁻⁶⁹ The Study of Viral Resistance to Antiviral Therapy of Chronic Hepatitis C (Virahep-C) examined factors associated with non-response to peginterferon alfa (PEG-IFN α) and RBV therapy in patients infected with HCV genotype 1a or 1b. In the study, African Americans were found to have a poor response to treatment, a finding not explained on the basis of clinical factors including viral load, disease severity, or cumulative dose of therapy received.⁶⁹ As part of the Virahep-C study, an ancillary study on genetic variation in HCV genotype 1 strains found that higher inter-patient HCV genetic diversity was associated with a robust treatment response at day 28 of therapy. This increase in genotypic diversity was limited to a few genes including those in the NS3 and NS5A regions in genotype 1a, and the core and NS3 genes in genotype 1b.⁷⁰ Additionally, those subjects achieving a sustained virologic response (SVR) had more diverse HCV sequences than non-responders. This finding may suggest that African Americans are infected with more homogenous and perhaps more difficult to treat strains of HCV. Additionally, the relevance of subgenotypic variability of which was less important in an era of IFN-based dual therapy may generate more interest going forward given its importance in the context

of protease inhibitor based therapy. Genotype 1b is associated with a modestly improved SVR with protease inhibitor based therapy when compared with genotype 1a. This phenomenon is thought to be explained by genotype 1b's lower propensity for the development of viral resistance as compared with genotype 1a.⁷¹

Host Genetic Diversity—Genome-wide association studies (GWAS) were the first to identify single nucleotide polymorphisms (SNPs) near the IL-28B host gene that dictate the host's response to HCV infection. These SNPs not only predicted treatment-induced clearance and spontaneous recovery from HCV infection, but also explained some of the differences in response rates between whites and African Americans to HCV therapy with PEG-IFN α and RBV.⁷²⁻⁷⁴ The CC genotype at the rs12979860 locus of the IL-28B gene was shown to be associated with improved early viral kinetics and an increased likelihood of a rapid and early viral response and conferred approximately a 6 fold increased odds of achieving an SVR.⁷⁵⁻⁷⁷ In addition, those with the CC genotype were noted to have a higher probability of clearing HCV infection spontaneously than those with either the TC or TT genotypes. Two of these studies included African Americans and, though the sample size was small, found that the rs12979860 locus was associated with a favorable treatment outcome in these individuals.^{72, 75} Ge et al. showed that there was variation in the odds of SVR among different racial groups with the CC genotype, with a 7.3 fold increase noted in Caucasians, 5.6 fold increase in Hispanics, and 6.1 fold increase in African Americans.⁷² Thomson et al. demonstrated in multivariable analysis that African Americans with the CC genotype responded more favorably than whites with the non-CC IL-28B genotype but that the CC genotype was far less prevalent in African Americans.⁷⁵ Though IL-28B polymorphisms were the strongest pre-treatment predictor of SVR to PEG-IFN and RBV therapy, these SNPs failed to explain all of the treatment related variability observed in African Americans, with African American race remaining an independent negative predictor of outcome on multivariable regression analysis.⁷⁵

Treatment and Outcomes in Chronic Hepatitis C Infection

Treatment of HCV genotype 1 infection has changed over time, with each incremental modification in therapy being associated with consistent improvements in SVR rates (Table 3). The rates of SVR were approximately 6–12% with IFN monotherapy, 38–42% with conventional IFN and RBV, and as high as 55% with PEG-IFN and RBV.^{57, 78-81} In African-Americans, the response rates have been considerably lower and have ranged from 0-11% with IFN monotherapy^{57, 80, 82} and 8-28 % with PEG-IFN and RBV.^{67-69, 83} The introduction of two new direct acting antiviral (DAA) agents, telaprevir and boceprevir, approved by the FDA in 2011 for the treatment of chronic HCV genotype 1 infection, has brought with it the prospect of a shorter duration of therapy and an improvement in SVR rates. While efficacy of therapy has increased over the last 20 years, there remain challenges including management of complexity of treatment, adverse events, tolerability and cost. This is further exaggerated among African Americans because of the relatively lower SVR rates achieved and their limited representation in clinical trials, resulting in a dearth of meaningful observations until the post-marketing phase of drug development.

African Americans represented 10.6% of total enrollment in the most recent phase three clinical trials of telaprevir or boceprevir.⁹² Hence, data generated from these late phase clinical studies are limited for this underrepresented population. In the case of boceprevir however, given lower treatment response rates in African Americans, a separate treatment group for African American subjects was created, resulting in the enrollment of 159 subjects in this group, representing 15% of total study enrollment. The success of this targeted enrollment strategy may be a useful approach to increase African American enrollment going forward and it was invaluable for providing data on differences in response rates and adverse events in this population. Regardless of the strategy employed for enrollment, data suggest that many African Americans are more likely to be ineligible for trial enrollment. Melia and colleagues noted that African Americans were 65% less likely to be eligible for treatment.⁹³ Reasons cited for ineligibility included leukopenia, anemia and concomitant comorbid conditions like diabetes mellitus and chronic kidney disease. The authors therefore suggest that further consideration should be given to eligibility criteria, including a decrease in neutrophil requirement, to allow for improved study inclusion and moreover treatment rates in African Americans.

The introduction of triple therapy with boceprevir and telaprevir has demonstrated a substantial improvement in SVR rates for treatment naïve and treatment-experienced HCV patients. When comparing African American with non-African American patients, SVR rates in the ILLUMINATE study, inclusive of a telaprevir-based regimen, were 60% vs. 74%, respectively and the SPRINT-2 study, inclusive a boceprevir-based regimen, resulted in SVRs of 53% vs. 68%, respectively.^{90, 91} However, differences were still noted in overall treatment response rates in African Americans, a facet of this disease process that is not well understood. Our hope however is that the promise of new DAAs will bring with them the ability to eradicate infection in all difficult to treat populations.

Liver Transplantation in Hepatitis C Infection

In addition to disparities noted in HCV prevalence, natural history, and treatment response, African Americans also have a higher disease burden and higher prevalence of HCC than other racial and ethnic groups.^{53, 94, 95} Though the same eligibility criteria for liver transplantation apply, African Americans are more likely to have an advanced stage of HCC at presentation, and thus be considered high risk or unsuitable for liver transplantation.⁹⁶ Regardless of the indication for listing, African Americans were found to be under-represented on the liver transplantation waiting list even when matched to those with similar wait times. African Americans were also noted to be more likely to die while waiting for liver transplantation.⁴⁶

A sentinel study compared pre-MELD and MELD era liver transplantation survival in HCC patients with and without HCV. These investigators demonstrated that in the MELD era, HCV patients without HCC experienced a decrease in 5-year graft and patient survival. Those with HCV and HCC had improved graft and patient outcomes at 1 and 3 years; however this advantage nearly disappeared at 5 years. Not surprisingly, in both the pre-MELD and MELD eras, there were more white recipients and donors, regardless of HCV status. Another notable observation from this study was that African Americans with HCC

had a 30% lower survival rate than other races after adjusting for other factors.⁹⁷ Nair et al. also showed, in a study of clinical outcomes of liver transplantation recipients, that African Americans with HCV had higher rates of primary graft non-function, an almost twofold increase in graft failure due to chronic rejection, a higher 30 day mortality, and worsened 2 year survival than other races.⁹⁸ Furthermore, recipient race has been noted to be an independent predictor of survival in recipients with HCV.⁹⁹ African American race was an independent predictor of patient survival after liver transplantation compared with whites.⁹⁶ Though the data speaking to racial disparities in liver transplantation are relatively robust, the possible reasons for this inequality remain underexplored.

Challenges and Future Directions

Many barriers remain before chronic HCV infection can be eradicated in African Americans and, more globally, all infected patients. Because of the silent nature of disease prior to the onset of late stage sequelae, many patients harbor infection without knowledge of their infectious status. This results in missed opportunities for treatment and preventative care. Additionally, those at risk for disease-related complications have failed to be entered into surveillance programs, such as those that exist for HCC. In order to facilitate improvements in outcomes in African Americans and more broadly all patients with chronic HCV infection, more effective screening must be provided and more efficacious therapies must become available. Such strides undoubtedly can be facilitated through a promotion of awareness of HCV infection and its sequelae and additional research in difficult to treat populations, such as African Americans, to determine the basis of viral persistence, host viral responses and differences in response to therapy (Tables 4 and 5). Our ability to overcome these HCV specific factors will result in better outcomes for all patients with chronic HCV infection.

Conclusion

Viral hepatitis represent a public health concern whose impact on morbidity and mortality are only presently being recognized. As outlined in this review, there are many racial and ethnic disparities in the context of hepatitis B and C infection that have yet to be addressed in the literature. Attention to these disparities will not only increase our understanding of various aspects of disease but may also improve clinical outcomes for patients. Further understanding of these conditions, commitment to all encompassing research efforts and development of new, efficacious and cost effective treatment strategies will hopefully lead to eradication of these viral diseases in all, irrespective of race and ethnicity.

References

1. Ioannou GN. Hepatitis B virus in the United States: infection, exposure, and immunity rates in a nationally representative survey. *Ann Intern Med.* 2011; 154(5):319–28. Epub 2011/03/02. [PubMed: 21357909]
2. Weinbaum CM, Williams I, Mast EE, et al. Recommendations for identification and public health management of persons with chronic hepatitis B virus infection. *MMWR Recomm Rep.* 2008; 57(RR-8):1–20. Epub 2008/09/20. [PubMed: 18802412]

3. Kowdley KV, Wang CC, Welch S, et al. Prevalence of chronic hepatitis B among foreign-born persons living in the United States by country of origin. *Hepatology*. 2012; 56(2):422–33. Epub 2011/11/23. [PubMed: 22105832]
4. Cohen C, Holmberg SD, McMahon BJ, et al. Is chronic hepatitis B being undertreated in the United States? *J Viral Hepat*. 2011; 18(6):377–83. Epub 2010/12/15. [PubMed: 21143343]
5. Centers for Disease Control and Prevention. [January 31, 2012] Hepatitis B Information for Health Professionals. 2012. Available from: <http://www.cdc.gov/hepatitis/HBV/HBVfaq.htm>.
6. McQuillan GM, Coleman PJ, Kruszon-Moran D, et al. Prevalence of hepatitis B virus infection in the United States: the National Health and Nutrition Examination Surveys, 1976 through 1994. *Am J Public Health*. 1999; 89(1):14–8. Epub 1999/02/13. [PubMed: 9987458]
7. McQuillan GM, Kruszon-Moran D, Kottiri BJ, et al. Racial and ethnic differences in the seroprevalence of 6 infectious diseases in the United States: data from NHANES III, 1988-1994. *Am J Public Health*. 2004; 94(11):1952–8. Epub 2004/10/30. [PubMed: 15514236]
8. McMahon BJ. Natural history of chronic hepatitis B. *Clinics in liver disease*. 2010; 14(3):381–96. Epub 2010/07/20. [PubMed: 20638020]
9. El-Serag HB, Mason AC. Rising incidence of hepatocellular carcinoma in the United States. *N Engl J Med*. 1999; 340(10):745–50. Epub 1999/03/11. [PubMed: 10072408]
10. El-Serag HB. Hepatocellular carcinoma: recent trends in the United States. *Gastroenterology*. 2004; 127(5 Suppl 1):S27–34. Epub 2004/10/28. [PubMed: 15508094]
11. Yu L, Sloane DA, Guo C, et al. Risk factors for primary hepatocellular carcinoma in black and white Americans in 2000. *Clin Gastroenterol Hepatol*. 2006; 4(3):355–60. Epub 2006/03/11. [PubMed: 16527700]
12. Kew MC, Kramvis A, Yu MC, et al. Increased hepatocarcinogenic potential of hepatitis B virus genotype A in Bantu-speaking sub-saharan Africans. *J Med Virol*. 2005; 75(4):513–21. Epub 2005/02/17. [PubMed: 15714494]
13. Bruix J, Sherman M. Management of hepatocellular carcinoma: an update. *Hepatology*. 2011; 53(3):1020–2. Epub 2011/03/05. [PubMed: 21374666]
14. Kew MC. Hepatocellular carcinoma in African Blacks: Recent progress in etiology and pathogenesis. *World J Hepatol*. 2010; 2(2):65–73. Epub 2010/12/17. [PubMed: 21160975]
15. Kew MC, Macerollo P. Effect of age on the etiologic role of the hepatitis B virus in hepatocellular carcinoma in blacks. *Gastroenterology*. 1988; 94(2):439–42. Epub 1988/02/01. [PubMed: 2446950]
16. Kew MC, Marcus R, Geddes EW. Some characteristics of Mozambican Shangaans with primary hepatocellular cancer. *S Afr Med J*. 1977; 51(10):306–9. Epub 1977/03/05. [PubMed: 66757]
17. Umoh NJ, Lesi OA, Mendy M, et al. Aetiological differences in demographical, clinical and pathological characteristics of hepatocellular carcinoma in The Gambia. *Liver Int*. 2011; 31(2):215–21. Epub 2010/12/15. [PubMed: 21143369]
18. Chen CH, Eng HL, Lee CM, et al. Correlations between hepatitis B virus genotype and cirrhotic or non-cirrhotic hepatoma. *Hepatogastroenterology*. 2004; 51(56):552–5. Epub 2004/04/17. [PubMed: 15086200]
19. Chu CJ, Keeffe EB, Han SH, et al. Hepatitis B virus genotypes in the United States: results of a nationwide study. *Gastroenterology*. 2003; 125(2):444–51. Epub 2003/08/02. [PubMed: 12891547]
20. Kao JH, Chen PJ, Lai MY, et al. Hepatitis B genotypes correlate with clinical outcomes in patients with chronic hepatitis B. *Gastroenterology*. 2000; 118(3):554–9. Epub 2000/03/04. [PubMed: 10702206]
21. Kao JH, Wu NH, Chen PJ, et al. Hepatitis B genotypes and the response to interferon therapy. *J Hepatol*. 2000; 33(6):998–1002. Epub 2000/12/29. [PubMed: 11131465]
22. Lau GK, Piratvisuth T, Luo KX, et al. Peginterferon Alfa-2a, lamivudine, and the combination for HBeAg-positive chronic hepatitis B. *N Engl J Med*. 2005; 352(26):2682–95. Epub 2005/07/01. [PubMed: 15987917]
23. Wai CT, Chu CJ, Hussain M, et al. HBV genotype B is associated with better response to interferon therapy in HBeAg(+) chronic hepatitis than genotype C. *Hepatology*. 2002; 36(6):1425–30. Epub 2002/11/26. [PubMed: 12447868]

24. Cheong JY, Cho SW, Hwang IL, et al. Association between chronic hepatitis B virus infection and interleukin-10, tumor necrosis factor- α gene promoter polymorphisms. *J Gastroenterol Hepatol*. 2006; 21(7):1163–9. Epub 2006/07/11. [PubMed: 16824070]
25. Peng XM, Huang YS, Ma HH, et al. Interleukin-10 promoter polymorphisms are associated with the mode and sequel of HBeAg seroconversion in patients with chronic hepatitis B virus infection. *Liver Int*. 2006; 26(3):326–33. Epub 2006/04/06. [PubMed: 16584395]
26. Truelove AL, Oleksyk TK, Shrestha S, et al. Evaluation of IL10, IL19 and IL20 gene polymorphisms and chronic hepatitis B infection outcome. *Int J Immunogenet*. 2008; 35(3):255–64. Epub 2008/05/16. [PubMed: 18479293]
27. Oleksyk TK, Thio CL, Truelove AL, et al. Single nucleotide polymorphisms and haplotypes in the IL10 region associated with HCV clearance. *Genes Immun*. 2005; 6(4):347–57. Epub 2005/04/09. [PubMed: 15815689]
28. Lau DT, Everhart J, Kleiner DE, et al. Long-term follow-up of patients with chronic hepatitis B treated with interferon alfa. *Gastroenterology*. 1997; 113(5):1660–7. Epub 1997/11/14. [PubMed: 9352870]
29. Lok AS, McMahon BJ. Chronic hepatitis B: update 2009. *Hepatology*. 2009; 50(3):661–2. Epub 2009/08/29. [PubMed: 19714720]
30. Dienstag JL, Schiff ER, Wright TL, et al. Lamivudine as initial treatment for chronic hepatitis B in the United States. *N Engl J Med*. 1999; 341(17):1256–63. Epub 1999/10/21. [PubMed: 10528035]
31. Perrillo RP, Lai CL, Liaw YF, et al. Predictors of HBeAg loss after lamivudine treatment for chronic hepatitis B. *Hepatology*. 2002; 36(1):186–94. Epub 2002/06/27. [PubMed: 12085364]
32. Marcellin P, Chang TT, Lim SG, et al. Adefovir dipivoxil for the treatment of hepatitis B e antigen-positive chronic hepatitis B. *N Engl J Med*. 2003; 348(9):808–16. Epub 2003/02/28. [PubMed: 12606735]
33. Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, et al. Adefovir dipivoxil for the treatment of hepatitis B e antigen-negative chronic hepatitis B. *N Engl J Med*. 2003; 348(9):800–7. Epub 2003/02/28. [PubMed: 12606734]
34. Dienstag JL, Cianciara J, Karayalcin S, et al. Durability of serologic response after lamivudine treatment of chronic hepatitis B. *Hepatology*. 2003; 37(4):748–55. Epub 2003/04/02. [PubMed: 12668966]
35. Marcellin P, Lau GK, Bonino F, et al. Peginterferon alfa-2a alone, lamivudine alone, and the two in combination in patients with HBeAg-negative chronic hepatitis B. *N Engl J Med*. 2004; 351(12):1206–17. Epub 2004/09/17. [PubMed: 15371578]
36. Kuo A, Dienstag JL, Chung RT. Tenofovir disoproxil fumarate for the treatment of lamivudine-resistant hepatitis B. *Clin Gastroenterol Hepatol*. 2004; 2(3):266–72. Epub 2004/03/16. [PubMed: 15017612]
37. Chang TT, Gish RG, Hadziyannis SJ, et al. A dose-ranging study of the efficacy and tolerability of entecavir in Lamivudine-refractory chronic hepatitis B patients. *Gastroenterology*. 2005; 129(4):1198–209. Epub 2005/10/19. [PubMed: 16230074]
38. Hadziyannis SJ, Tassopoulos NC, Heathcote EJ, et al. Long-term therapy with adefovir dipivoxil for HBeAg-negative chronic hepatitis B for up to 5 years. *Gastroenterology*. 2006; 131(6):1743–51. Epub 2006/11/08. [PubMed: 17087951]
39. Chang TT, Gish RG, de Man R, et al. A comparison of entecavir and lamivudine for HBeAg-positive chronic hepatitis B. *N Engl J Med*. 2006; 354(10):1001–10. Epub 2006/03/10. [PubMed: 16525137]
40. Lai CL, Shouval D, Lok AS, et al. Entecavir versus lamivudine for patients with HBeAg-negative chronic hepatitis B. *N Engl J Med*. 2006; 354(10):1011–20. Epub 2006/03/10. [PubMed: 16525138]
41. Marcellin P, Chang TT, Lim SG, et al. Long-term efficacy and safety of adefovir dipivoxil for the treatment of hepatitis B e antigen-positive chronic hepatitis B. *Hepatology*. 2008; 48(3):750–8. Epub 2008/08/30. [PubMed: 18752330]
42. Marcellin P, Heathcote EJ, Buti M, et al. Tenofovir disoproxil fumarate versus adefovir dipivoxil for chronic hepatitis B. *N Engl J Med*. 2008; 359(23):2442–55. Epub 2008/12/05. [PubMed: 19052126]

43. Leung N, Peng CY, Hann HW, et al. Early hepatitis B virus DNA reduction in hepatitis B e antigen-positive patients with chronic hepatitis B: A randomized international study of entecavir versus adefovir. *Hepatology*. 2009; 49(1):72–9. Epub 2008/12/10. [PubMed: 19065670]
44. Liaw YF, Gane E, Leung N, et al. 2-Year GLOBE trial results: telbivudine is superior to lamivudine in patients with chronic hepatitis B. *Gastroenterology*. 2009; 136(2):486–95. Epub 2008/11/26. [PubMed: 19027013]
45. Widjaja D, Yarlagadda S, Singu BS, et al. Characteristics of patients with chronic hepatitis-B virus infection in an urban hospital. *J Natl Med Assoc*. 2007; 99(4):384–8. Epub 2007/04/21. [PubMed: 17444427]
46. Reid AE, Resnick M, Chang Y, et al. Disparity in use of orthotopic liver transplantation among blacks and whites. *Liver Transpl*. 2004; 10(7):834–41. Epub 2004/07/09. [PubMed: 15237365]
47. Moylan CA, Brady CW, Johnson JL, et al. Disparities in liver transplantation before and after introduction of the MELD score. *Jama*. 2008; 300(20):2371–8. Epub 2008/11/27. [PubMed: 19033587]
48. Kim WR, Poterucha JJ, Kremers WK, et al. Outcome of liver transplantation for hepatitis B in the United States. *Liver Transpl*. 2004; 10(8):968–74. Epub 2004/09/25. [PubMed: 15390321]
49. Bzowej N, Han S, Degertekin B, et al. Liver transplantation outcomes among Caucasians, Asian Americans, and African Americans with hepatitis B. *Liver Transpl*. 2009; 15(9):1010–20. Epub 2009/09/01. [PubMed: 19718627]
50. Kelen GD, Green GB, Purcell RH, et al. Hepatitis B and hepatitis C in emergency department patients. *N Engl J Med*. 1992; 326(21):1399–404. Epub 1992/05/21. [PubMed: 1373867]
51. Seeff LB, Miller RN, Rabkin CS, et al. 45-year follow-up of hepatitis C virus infection in healthy young adults. *Ann Intern Med*. 2000; 132(2):105–11. Epub 2000/01/22. [PubMed: 10644270]
52. Prysopoulou N, Jeffers L. Hepatitis C in African Americans. *Journal of clinical gastroenterology*. 2007; 41(2):185–93. Epub 2007/01/25. [PubMed: 17245218]
53. Armstrong GL, Wasley A, Simard EP, et al. The prevalence of hepatitis C virus infection in the United States, 1999 through 2002. *Ann Intern Med*. 2006; 144(10):705–14. Epub 2006/05/17. [PubMed: 16702586]
54. Davis GL, Alter MJ, El-Serag H, et al. Aging of hepatitis C virus (HCV)-infected persons in the United States: a multiple cohort model of HCV prevalence and disease progression. *Gastroenterology*. 2010; 138(2):513–21. Epub 2009/10/29. [PubMed: 19861128]
55. Ly KN, Jian X, Kleven RM, et al. The Increasing Burden of Mortality From Viral Hepatitis in the United States Between 1999 and 2007. *Ann Intern Med*. 2012; 156(4):271–W-52. [PubMed: 22351712]
56. Wiley TE, Brown J, Chan J. Hepatitis C infection in African Americans: its natural history and histological progression. *The American journal of gastroenterology*. 2002; 97(3):700–6. Epub 2002/04/02. [PubMed: 11922566]
57. Reddy KR, Hoofnagle JH, Tong MJ, et al. Racial differences in responses to therapy with interferon in chronic hepatitis C. Consensus Interferon Study Group. *Hepatology*. 1999; 30(3):787–93. Epub 1999/08/26. [PubMed: 10462387]
58. Kohla M, Iwata S, Ea R, et al. Histological Versus Clinical Cirrhosis in Chronic Hepatitis C: Does Race/Ethnicity Really Matter? *Dig Dis Sci*. 2012; 57(3):771–6. [PubMed: 21948357]
59. Crosse K, Umeadi OG, Anania FA, et al. Racial differences in liver inflammation and fibrosis related to chronic hepatitis C. *Clin Gastroenterol Hepatol*. 2004; 2(6):463–8. Epub 2004/06/08. [PubMed: 15181613]
60. Terrault NA, Im K, Boylan R, et al. Fibrosis Progression in African Americans and Caucasian Americans With Chronic Hepatitis C. *Clinical Gastroenterology and Hepatology*. 2008; 6(12):1403–11. [PubMed: 19081528]
61. El-Serag HB. Epidemiology of viral hepatitis and hepatocellular carcinoma. *Gastroenterology*. 2012; 142(6):1264–73. Epub 2012/04/28. [PubMed: 22537432]
62. Simard EP, Ward EM, Siegel R, et al. Cancers with increasing incidence trends in the United States: 1999 through 2008. *CA Cancer J Clin*. 2012 Epub 2012/01/28.

63. Alberti A, Vario A, Ferrari A, et al. Review article: chronic hepatitis C--natural history and cofactors. *Alimentary pharmacology & therapeutics*. 2005; 22(Suppl 2):74–8. Epub 2005/10/18. [PubMed: 16225479]
64. Mizokami M, Orito E. Molecular evolution of hepatitis viruses. *Intervirology*. 1999; 42(2-3):159–65. Epub 1999/10/12. [PubMed: 10516470]
65. Pyrsopoulos N, Jeffers L. Chronic hepatitis C in African Americans. *Clinics in liver disease*. 2005; 9(3):427–38. vii. Epub 2005/07/19. [PubMed: 16023975]
66. NIH Consensus Statement on Management of Hepatitis C: 2002. NIH consensus and state-of-the-science statements. 2002; 19(3):1–46. Epub 2004/02/11.
67. Jeffers LJ, Cassidy W, Howell CD, et al. Peginterferon alfa-2a (40 kd) and ribavirin for black American patients with chronic HCV genotype 1. *Hepatology*. 2004; 39(6):1702–8. [PubMed: 15185312]
68. Muir AJ, Bornstein JD, Killenberg PG, et al. Peginterferon alfa-2b and ribavirin for the treatment of chronic hepatitis C in blacks and non-Hispanic whites. *N Engl J Med*. 2004; 350(22):2265–71. Epub 2004/05/28. [PubMed: 15163776]
69. Conjeevaram HS, Fried MW, Jeffers LJ, et al. Peginterferon and Ribavirin Treatment in African American and Caucasian American Patients With Hepatitis C Genotype 1. *Gastroenterology*. 2006; 131(2):470–7. [PubMed: 16890601]
70. Donlin MJ, Cannon NA, Yao E, et al. Pretreatment sequence diversity differences in the full-length hepatitis C virus open reading frame correlate with early response to therapy. *Journal of virology*. 2007; 81(15):8211–24. Epub 2007/05/25. [PubMed: 17522222]
71. Lok AS, Gardiner DF, Lawitz E, et al. Preliminary study of two antiviral agents for hepatitis C genotype 1. *N Engl J Med*. 2012; 366(3):216–24. Epub 2012/01/20. [PubMed: 22256805]
72. Ge D, Fellay J, Thompson AJ, et al. Genetic variation in IL28B predicts hepatitis C treatment-induced viral clearance. *Nature*. 2009; 461(7262):399–401. Epub 2009/08/18. [PubMed: 19684573]
73. Suppiah V, Moldovan M, Ahlenstiel G, et al. IL28B is associated with response to chronic hepatitis C interferon-alpha and ribavirin therapy. *Nature genetics*. 2009; 41(10):1100–4. Epub 2009/09/15. [PubMed: 19749758]
74. Tanaka Y, Nishida N, Sugiyama M, et al. Genome-wide association of IL28B with response to pegylated interferon-alpha and ribavirin therapy for chronic hepatitis C. *Nature genetics*. 2009; 41(10):1105–9. Epub 2009/09/15. [PubMed: 19749757]
75. Thompson AJ, Muir AJ, Sulkowski MS, et al. Interleukin-28B polymorphism improves viral kinetics and is the strongest pretreatment predictor of sustained virologic response in genotype 1 hepatitis C virus. *Gastroenterology*. 2010; 139(1):120–9. e18. Epub 2010/04/20. [PubMed: 20399780]
76. McCarthy JJ, Li JH, Thompson A, et al. Replicated association between an IL28B gene variant and a sustained response to pegylated interferon and ribavirin. *Gastroenterology*. 2010; 138(7):2307–14. Epub 2010/02/24. [PubMed: 20176026]
77. Rauch A, Kutalik Z, Descombes P, et al. Genetic variation in IL28B is associated with chronic hepatitis C and treatment failure: a genome-wide association study. *Gastroenterology*. 2010; 138(4):1338–45. 45, e1–7. Epub 2010/01/12. [PubMed: 20060832]
78. Fried MW, Shiffman ML, Reddy KR, et al. Peginterferon alfa-2a plus ribavirin for chronic hepatitis C virus infection. *N Engl J Med*. 2002; 347(13):975–82. Epub 2002/09/27. [PubMed: 12324553]
79. Manns MP, McHutchison JG, Gordon SC, et al. Peginterferon alfa-2b plus ribavirin compared with interferon alfa-2b plus ribavirin for initial treatment of chronic hepatitis C: a randomised trial. *Lancet*. 2001; 358(9286):958–65. Epub 2001/10/05. [PubMed: 11583749]
80. McHutchison JG, Gordon SC, Schiff ER, et al. Interferon alfa-2b alone or in combination with ribavirin as initial treatment for chronic hepatitis C. Hepatitis Interventional Therapy Group. *N Engl J Med*. 1998; 339(21):1485–92. Epub 1998/11/20. [PubMed: 9819446]
81. Poynard T, Bedossa P, Chevallier M, et al. A comparison of three interferon alfa-2b regimens for the long-term treatment of chronic non-A, non-B hepatitis. Multicenter Study Group. *N Engl J Med*. 1995; 332(22):1457–62. Epub 1995/06/01. [PubMed: 7739681]

82. Theodore D, Shiffman ML, Sterling RK, et al. Intensive interferon therapy does not increase virological response rates in African Americans with chronic hepatitis C. *Digestive Diseases and Sciences*. 2003; 48(1):140–5. Epub 2003/03/21. [PubMed: 12645801]
83. Bacon BR, Gordon SC, Lawitz E, et al. Boceprevir for previously treated chronic HCV genotype 1 infection. *N Engl J Med*. 2011; 364(13):1207–17. Epub 2011/04/01. [PubMed: 21449784]
84. McHutchison JG, Poynard T, Pianko S, et al. The impact of interferon plus ribavirin on response to therapy in black patients with chronic hepatitis C. The International Hepatitis Interventional Therapy Group. *Gastroenterology*. 2000; 119(5):1317–23. Epub 2000/10/31. [PubMed: 11054390]
85. Hezode C, Forestier N, Dusheiko G, et al. Telaprevir and peginterferon with or without ribavirin for chronic HCV infection. *N Engl J Med*. 2009; 360(18):1839–50. Epub 2009/05/01. [PubMed: 19403903]
86. McHutchison JG, Everson GT, Gordon SC, et al. Telaprevir with peginterferon and ribavirin for chronic HCV genotype 1 infection. *The New England journal of medicine*. 2009; 360(18):1827–38. Epub 2009/05/01. [PubMed: 19403902]
87. McHutchison JG, Manns MP, Muir AJ, et al. Telaprevir for previously treated chronic HCV infection. *The New England journal of medicine*. 2010; 362(14):1292–303. Epub 2010/04/09. [PubMed: 20375406]
88. Kwo PY, Lawitz EJ, McCone J, et al. Efficacy of boceprevir, an NS3 protease inhibitor, in combination with peginterferon alfa-2b and ribavirin in treatment-naïve patients with genotype 1 hepatitis C infection (SPRINT-1): an open-label, randomised, multicentre phase 2 trial. *Lancet*. 2010; 376(9742):705–16. Epub 2010/08/10. [PubMed: 20692693]
89. Jacobson IM, McHutchison JG, Dusheiko G, et al. Telaprevir for previously untreated chronic hepatitis C virus infection. *N Engl J Med*. 2011; 364(25):2405–16. Epub 2011/06/24. [PubMed: 21696307]
90. Sherman KE, Flamm SL, Afdhal NH, et al. Response-guided telaprevir combination treatment for hepatitis C virus infection. *N Engl J Med*. 2011; 365(11):1014–24. Epub 2011/09/16. [PubMed: 21916639]
91. Poordad F, McCone J Jr, Bacon BR, et al. Boceprevir for untreated chronic HCV genotype 1 infection. *N Engl J Med*. 2011; 364(13):1195–206. Epub 2011/04/01. [PubMed: 21449783]
92. Burton MJ, Passarella MJ, McGuire BM. Telaprevir and boceprevir in african americans with genotype 1 chronic hepatitis C: implications for patients and providers. *Southern medical journal*. 2012; 105(8):431–6. Epub 2012/08/07. [PubMed: 22864102]
93. Melia MT, Muir AJ, McCone J, et al. Racial differences in hepatitis C treatment eligibility. *Hepatology (Baltimore, Md)*. 2011; 54(1):70–8. Epub 2011/04/14.
94. El-Serag HB, Davila JA, Petersen NJ, et al. The continuing increase in the incidence of hepatocellular carcinoma in the United States: an update. *Ann Intern Med*. 2003; 139(10):817–23. Epub 2003/11/19. [PubMed: 14623619]
95. Nguyen MH, Whittemore AS, Garcia RT, et al. Role of ethnicity in risk for hepatocellular carcinoma in patients with chronic hepatitis C and cirrhosis. *Clin Gastroenterol Hepatol*. 2004; 2(9):820–4. Epub 2004/09/09. [PubMed: 15354283]
96. Kemmer N, Neff G, Secic M, et al. Ethnic differences in hepatocellular carcinoma: implications for liver transplantation. *Dig Dis Sci*. 2008; 53(2):551–5. Epub 2007/06/15. [PubMed: 17562173]
97. Thuluvath PJ, Maheshwari A, Thuluvath NP, et al. Survival after liver transplantation for hepatocellular carcinoma in the model for end-stage liver disease and pre-model for end-stage liver disease eras and the independent impact of hepatitis C virus. *Liver Transpl*. 2009; 15(7):754–62. Epub 2009/06/30. [PubMed: 19562709]
98. Nair S, Eustace J, Thuluvath PJ. Effect of race on outcome of orthotopic liver transplantation: a cohort study. *Lancet*. 2002; 359(9303):287–93. Epub 2002/02/07. [PubMed: 11830194]
99. Charlton M, Seaberg E, Wiesner R, et al. Predictors of patient and graft survival following liver transplantation for hepatitis C. *Hepatology*. 1998; 28(3):823–30. Epub 1998/09/10. [PubMed: 9731579]

Table 1

African American Inclusion in Nucleotide/ Nucleoside Analogue Clinical Trials

Author	Treatment	Treatment Length Weeks	Total Patients N	African Americans N	African Americans %
Dienstag et al. 1999 ³⁰	Placebo	52	71	13	18
	Lamivudine	52	66	10	15
Perrillo et al. 2002 ³¹	Placebo	52	196	13	7
	Lamivudine		406	16	4
	IFN		68	3	4
	IFN+Lamivudine		135	5	4
Marcellin et al. 2003 ³²	Placebo	48	167	3	2
	Adefovir (10 mg)	48	171	8	5
	Adefovir (30 mg)	48	173	5	3
Hadziyannis et al. 2003 ³³	Placebo	48	61	1	2
	Adefovir	48	123	5	4
Dienstag et al. 2003 ³⁴	Lamivudine	144	40	1	3
Marcellin et al. 2004 ³⁵	PEG-IFN α -2a+Placebo	48	177	3	2
	PEG-IFN α -2a+Lamivudine	48	179	2	1
Kuo et al. 2004 ³⁶	Lamivudine	48	181	0	0
	Tenofovir	48	9	1	1
Chang et al. 2005 ³⁷	Lamivudine	76	45	0	0
	Entecavir (1.0 mg)	76	42	3	7
	Entecavir (0.5 mg)	76	47	1	2
	Entecavir (0.1 mg)	76	47	0	0
Hadziyannis et al. 2006 ³⁸	Placebo/Adefovir	97-240	55	1	2
	Adefovir/Adefovir	97-240	70	3	4
Chang et al. 2006 ³⁹	Lamivudine	52	355	8	2
	Entecavir	52	354	8	2

Author	Treatment	Treatment Length Weeks	Total Patients N	African Americans N	African Americans %
Lai et al. 2006 ⁴⁰	Lamivudine	52	313	7	2
	Entecavir	52	325	8	2
Marcellin et al. 2008 ⁴¹	Adefovir	48	171	8	5
	Adefovir	240	65	3	3
Marcellin et al. 2008 ⁴²	Adefovir	48	215	9	4
	Tenofovir	48	426	21	5
Leung et al. 2009 ⁴³	Adefovir	52	32	2	6
	Entecavir	52	33	2	6
Liaw et al. 2009 ⁴⁴	Lamivudine	104	687	8	1
	Telbivudine	104	680	4	1

IFN= Interferon.

Table 2

Unmet needs in African Americans with hepatitis B Infection

Understanding of high prevalence of HBV infection
Understanding of differences in natural history of HBV infection in African Americans compared to other racial/ ethnic groups
Understanding of host genetics and how they affect establishment of chronic HBV infection
Comparison of disease progression in African Americans and other racial/ ethnic groups
Focused studies of safety and efficacy of HBV anti-viral therapies in African Americans
More data on post-transplant patient and allograft outcomes in African Americans requiring liver transplantation for chronic HBV infection

Table 3

Review of HCV Treatment Studies

Author year (Study)	Treatment	Treatment Length Weeks	Total Patients N	African Americans N (%)	Non-AA SVR (%)	AA SVR (%)
Reddy et al. 1999 ⁵⁷	IFNα-2b or Consensus IFN	24	472	40 (8)	12	2
McHutchison et al. 2000 ⁸⁴	IFNα-2b	24	230	12 (5)	6	0
	IFNα-2b	48	489	13 (3)	16	0
	IFNα-2b+RBV	24	497	15 (3)	34	20
	IFNα-2b+RBV	48	496	13 (3)	41	23
Theodore et al. 2003 ⁸²	IFNα-2b (3 MU thrice weekly)	24	42	12 (29)	4	0
	IFNα-2b (5 MU daily)	24	32	8 (25)	10	11
Muir et al. 2004 ⁶⁸	PEG-IFNα-2b+RBV	48	200	100 (50)	52	19
Jeffers et al. 2004 ⁶⁷	PEG-IFNα-2a+RBV	48	106	78 (74)	39	26
Conjeevaram et al. 2006 ⁶⁹	PEG-IFNα-2a+RBV	48	401	196 (49)	52	28
Hezode et al. 2009 ⁸⁵	PEG-IFNα-2a+RBV	48	82	2 (2)	46	
	PEG-IFNα-2a+RBV+TVR	12	82	2 (2)	60	
	PEG-IFNα-2a+RBV+TVR / PEG-IFNα-2a+RBV	12/12	81	1 (1)	69	
	PEG-IFNα-2a+TVR	12	78	1 (1)	36	
McHutchison et al. 2009 ⁸⁶	PEG-IFNα-2a+RBV	48	75	9 (12)	41 [^]	11
	PEG-IFNα-2a+RBV+TVR	12	17	3 (18)	35 [^]	
	PEG-IFNα-2a+RBV+TVR / PEG-IFNα-2a+RBV	12/12	79	7 (9)	61 [^]	* 44
	PEG-IFNα-2a+RBV+TVR / PEG-IFNα-2a+RBV	12/36	79	8 (10)	67 [^]	
McHutchison et al. 2010 ⁸⁷	PEG-IFNα-2a+RBV	48	114	10(9)	14	10
	PEG-IFNα-2a+RBV+TVR / PEG-IFNα-2a+RBV	12/12	115	9 (8)	54	22
	PEG-IFNα-2a+RBV+TVR	24	111	10(9)	27	0
	PEG-IFNα-2a+RBV+TVR / PEG-IFNα-2a+RBV	24/24	113	11(10)	53	55

Author year (Study)	Treatment	Treatment Length Weeks	Total Patients N	African Americans N (%)	Non-AA SVR (%)	AA SVR (%)
Kwo et al. 2010 ⁸⁸	<u>Part 1</u>					
	PEG-IFN α -2b+RBV	48	104	16 (15)	42	13
	PEG-IFN α -2b+RBV/PEG-IFN α -2b+RBV+BOC	4/24	103	15 (15)	59	40
	PEG-IFN α -2b+RBV/PEG-IFN α -2b+RBV+BOC	4/44	103	15 (15)	78	53
	PEG-IFN α -2b+RBV+BOC	28	107	18 (17)	57	39
	PEG-IFN α -2b+RBV+BOC	48	103	14 (14)	73	29
Bacon et. al. 2011 ⁸³	<u>Part 2</u>					
	PEG-IFN α -2b+RBV+BOC	48	16	4 (25)	50	
	PEG-IFN α -2b+RBV (Low Dose)+BOC	48	59	16 (27)	36	
	PEG-IFN α -2b+RBV	48	80	12 (15)	24	8
Jacobson et. al. 2011 ⁸⁹	PEG-IFN α -2b+RBV/PEG-IFN α -2b+RBV +BOC (PEG-IFN α -2b+RBV)	4/32 (12)	162	18 (11)	58	61
	PEG-IFN α -2b+RBV/PEG-IFN α -2b+RBV+BOC	4/44	161	19 (12)	68	53
	PEG-IFN α -2a+RBV	48	361	28 (8)	46	25
Sherman et al. 2011 ⁹⁰	PEG-IFN α -2a+RBV+TVR/PEG-IFN α -2a+RBV [§]	12/12	162	17 (10)	92	88
	PEG-IFN α -2a+RBV+TVR/PEG-IFN α -2a+RBV [§]	12/36	160	17 (11)	87	88
	PEG-IFN α -2a+RBV+TVR/PEG-IFN α -2a+RBV	12/36	118	20 (17)	64	65
Poordad et.al. 2011 ⁹¹	PEG-IFN α -2b+RBV	48	363	52 (14)	40	23
	PEG-IFN α -2b+RBV/PEG-IFN α -2b+RBV+BOC (PEG-IFN α -2b+RBV)	4/20 (24)	368	52 (14)	67	42
	PEG-IFN α -2b+RBV/PEG-IFN α -2b+RBV+BOC	4/44	366	55 (15)	68	53

IFN= Interferon, PEG-IFN= Peginterferon, RBV= Ribavirin, TVR=Telaprevir, BOC= Boceprevir

* all Telaprevir groups combined

[^] represent overall SVR

[§] random drug regimen assignment in those subjects with an extended rapid virologic response.

Table 4
Differences in HCV Epidemiology, Natural History, Disease Progression and Outcomes

Factor	African Americans	White Americans
Prevalence	High prevalence of HCV Increased rate of HCV-related death	High prevalence of HCV
Genotype	Virus: Higher rate of genotype 1b More difficult to treat genotype of 1 Host: lower prevalence of CC IL-28B genotype	Virus: Higher rate of genotype 1a Host: Higher prevalence of CC IL-28B genotype (Non-CC genotype in whites responded to treatment less than CC genotype in Blacks) ⁷⁷
Disease progression	More frequent evolution to chronic state from acute infection Less severity and slower progression Lower ALT than other ethnic groups Lower prevalence of cirrhosis Hepatic fibrosis progression (inconclusive) High risk to disease sequelae	Less likely to evolve to chronic infection Higher prevalence of cirrhosis
Treatment response	Less responsive to interferon based therapy, including with the use of a protease inhibitor Fewer patients represented in clinical trials	Better response rates than African-Americans Better representation in clinical trials
HCC risk	Higher incidence rate of HCC HCC incidence rate of 5.4%	Incidence rates of HCC 1-4% per year
Liver transplantation	Underrepresented on liver transplant list Lower survival rate both on waiting list and post transplantation	More recipients and donors both pre-MELD and MELD era Higher survival rate post transplantation

Table 5

Unmet Needs in African Americans with Chronic HCV Infection

Better understanding of natural history of HCV infection in African Americans compared to other groups
More understanding of viral and host genetic diversity to generate better tailored treatments
Better accessibility, affordability and quality in both clinical trials and standard treatment
Determine mechanisms of differences in disease progression
Increase access to liver transplantation for racial and ethnic minorities
Improvement in allograft and patient survival in the post liver transplantation setting