

Published in final edited form as:

Front Biol (Beijing). 2013 August ; 8(4): 395–407. doi:10.1007/s11515-013-1258-y.

The Sac domain-containing phosphoinositide phosphatases: structure, function, and disease

FoSheng HSU and Yuxin MAO

Weill Institute for Cell and Molecular Biology and Department of Molecular Biology and Genetics, Cornell University, Ithaca, NY 14853, USA

Abstract

Phosphoinositides (PIs) have long been known to have an essential role in cell physiology. Their intracellular localization and concentration must be tightly regulated for their proper function. This spatial and temporal regulation is achieved by a large number of PI kinases and phosphatases that are present throughout eukaryotic species. One family of these enzymes contains a conserved PI phosphatase domain termed Sac. Although the Sac domain is homologous among different Sac domain-containing proteins, all appear to exhibit varied substrate specificity and subcellular localization. Dysfunctions in several members of this family are implicated in a range of human diseases such as cardiac hypertrophy, bipolar disorder, Down's syndrome, Charcot-Marie-Tooth disease (CMT) and Amyotrophic Lateral Sclerosis (ALS). In plant, several Sac domain-containing proteins have been implicated in the stress response, chloroplast function and polarized secretion. In this review, we focus on recent findings in the family of Sac domain-containing PI phosphatases in yeast, mammal and plant, including the structural analysis into the mechanism of enzymatic activity, cellular functions, and their roles in disease pathophysiology.

Keywords

lipid metabolism; membrane trafficking

1 Introduction

Phosphoinositides (PIs), discovered in the 1950s by Marble and Lowell Hokin (Hokin LE and Hokin MR, 1958; Hokin MR and Hokin LE, 1953), are phosphorylated forms of a precursor molecule called phosphatidylinositol (PtdIns). PtdIns is composed of diacylglycerol (DAG) connected to a phosphate followed by a cytoplasmic exposed headgroup *myo*-inositol ring. The five hydroxyl groups attached to the inositol headgroup can be reversibly phosphorylated at the 3', 4' and 5' positions to generate seven distinct PI isoforms, whereas positions 2' and 6' are obstructed presumably due to steric hindrance. In the past 30 years, the roles of PIs have been characterized extensively in both uni- and multi-cellular organisms. PtdIns comprise ~10% of total cellular lipids and PI constitute less than

Correspondence: Yuxin Mao, ym253@cornell.edu.

Compliance with ethics guidelines

FoSheng Hsu and Yuxin Mao declare that they have no conflict of interest.

This article does not contain any studies with human and animal subjects performed by any of the authors.

1%, yet they control vital cellular processes such as cell signaling, proliferation, organization of cytoskeleton, and membrane trafficking (Odorizzi et al., 2000; De Matteis and Godi, 2004; Di Paolo and De Camilli, 2006). It has also been shown that PIs play a role in regulating ion channel activity (Wang et al., 2012), transcription, mRNA trafficking, RNA splicing (Osborne et al., 2001), chromatin structure (Viiri et al., 2012), nuclear export (Dieck et al., 2012), and bacterial infection (Pizarro-Cerdá and Cossart, 2004; Ham et al., 2011). The mechanism for controlling such divergent processes is achieved mainly by three non-mutually exclusive schemes. First, each PI has its own unique subcellular localization to allow them to selectively recruit proteins containing PI recognition modules (e.g. ANTH, ENTH, FYVE, PX, PH, PDZ, PHD, PTB, C2, GRAM, PROPPINs, Tubby, and FERM) to specific organelle membranes (Cullen et al., 2001; Lemmon, 2003; Takenawa and Itoh, 2006; Krauss and Haucke, 2007). Second, the resulting heterogeneous anionic charge (produced by the number of phosphate groups) at each organelle membranes can differentially recruit soluble polycationic proteins (Hammond et al., 2012). Third, PI in combination with specific Rab GTPase can constitute coincidence detection via recruitment of effectors with dual recognition domains (Di Paolo and De Camilli, 2006; Jean and Kiger, 2012). Coincidence detection increases the affinity and therefore the specificity of effector localization.

The heterogeneous distribution of PIs imparts an “organelle identity” to the cell. The maintenance of the selective concentration of specific PI species, as well as the dynamic control of PI composition in response to acute signaling inputs is achieved by a large number of PI kinases and phosphatases that are present throughout eukaryotic species. One family of these enzymes contains a conserved PI phosphatase domain termed Sac (Suppressor of yeast *act*in mutations).

2 The Sac phosphatase domain family in yeast and mammal

The first Sac domain-containing phosphatase, Sac1, was discovered in two independent genetic screens searching for modifiers of actin cytoskeleton defects (Novick et al., 1989) and of trans-Golgi network exocytic failure caused by inactivation of Sec14p (Bankaitis et al., 1990). Subsequently, the Sac domain was found to share considerable homology with mammalian nerve terminal protein, synaptojanin (McPherson et al., 1996) and was demonstrated to possess phosphoinositide phosphatase activity (Guo et al., 1999). With the advancement in genome sequencing, all genes containing the Sac domain have been annotated in yeast, and mammal. There are five Sac domain-containing proteins in both human and yeast. Members include the transmembrane protein Sac1, cytosolic proteins Sac2/Inpp5f, Sac3/Fig4, and synaptojanin 1 and 2. These Sac domain-containing proteins can be divided into three subfamilies: 1) Sac domain with two transmembrane motifs (Sac1); 2) Sac domain followed by along C-terminal sequences with or without recognizable structural domains (Sac2 and Sac3); 3) The N-terminal Sac domain immediately followed by a central PI 5-phosphatase domain and a C-terminal proline rich region (synaptojanin 1 and 2) (Fig. 1A).

3 Sac domain structure and catalytic mechanism

The Sac domain of yeast Sac1 is the only crystal structure solved in the family to this day (Manford et al., 2010). The structure revealed that the Sac homology domain is comprised of two closely packed sub-domains: a novel N-terminal sub-domain and the catalytic phosphoinositide phosphatase sub-domain (Fig. 2A). The N-domain is comprised of a unique fold with three layers of β -sheet and three short and one long α -helix. The catalytic domain consists of a nine-stranded β -sheet flanked by five α -helices with two and three on each side of the β -sheet. The catalytic site of Sac domain consists of a peptide motif with a sequence of CX₅R (a catalytic cysteine residue followed by 5 amino acids and a conserved arginine residue). Structural comparison of the catalytic domain of Sac1 with other CX₅R motif based proteins and lipid phosphatases revealed that the topology of the catalytic domain is conserved with other phosphatases. All the CX₅R motif based phosphatases share a common four parallel β -strands and one α -helix at the core. The loop connecting this α -helix- and one of the β -strands harbors the catalytic CX₅R motif (P-loop) (Fig. 2A). The conserved structural topology of all lipid and protein tyrosine phosphatases suggests a conserved catalytic mechanism (Barford et al., 1998). First, the conserved arginine residue, together with the backbone amide groups of the P-loop will bind and position the phosphate group at the catalytic site. Second, the catalytic cysteine will attack the phosphate group to generate a cysteinyl-phosphate intermediate and the dephosphorylated product. Third, an aspartic acid residue either located in a so-called “WPD” loop (Barford et al., 1998) or within the catalytic P-loop motif will act as a general acid, donating a proton to the releasing product. Finally, the same aspartic acid residue will function as a general base to activate a water molecule, which will then hydrolyze the phosphate from the catalytic cysteine through a nucleophilic attack.

Since the lipid substrates of Sac1 are embedded in negatively charged membrane bilayer, it is not surprising that the overall surface charge of Sac1 is distributed in a way that is negatively charged on one surface and positively charged on the opposite surface. The catalytic CX₅R motif is localized on the positively charged surface, which will interface with the anionic membrane bilayer (Manford et al., 2010). Interestingly, this membrane interfacing surface is enriched with highly conserved residues as revealed by evolutionary trace method (Lichtarge et al., 1996) (Fig. 2B). This conservation suggests a general catalytic mechanism, including membrane interaction, and substrate recognition among all Sac domains.

Unlike other CX₅R-based PI phosphatases, such as PTEN (Lee et al., 1999) or MTM (Begley et al., 2003), the catalytic P-loop of Sac1 has a striking conformation in that the catalytic cysteine is far away from the conserved arginine in the CX₅R motif. These structural findings suggest that conformational change of the catalytic P-loop may be required to achieve functional arrangement of the catalytic residues. In agreement with the structural observation, Sac1 is found to be an allosteric enzyme and its activity can be stimulated by anionic phospholipids, PtdIns and phosphatidyl serine (PS) (Zhong et al., 2012). However, a detail picture of the catalytic mechanism would require co-crystal structure of Sac1 with its substrate or activator.

4 Sac1

4.1 Yeast Sac1

Yeast Sac1 is a 67kDa type II transmembrane protein that localizes to both the endoplasmic reticulum (ER) and Golgi apparatus (Whitters et al., 1993). The retention of yeast Sac1 in the ER is dependent on the interaction with the integral membrane protein Dpm1 during times of rapid cell growth (Faulhammer et al., 2005). Whereas the localization of Golgi is likely via interactions with other proteins such as Vps74 (the ortholog of human GOLPH3) (Wood et al., 2012). *In vitro* assays have shown that Sac1 dephosphorylates PI(3)P, PI(4)P, and PI(3,5)P₂ (Guo et al., 1999). In yeast, genetic ablation of Sac1 activity results in a nearly 10-fold increase in the steady-state levels of PI(4)P and a modest increase in the levels of PI(3)P (Rivas et al., 1999; Nemoto et al., 2000), suggesting that yeast Sac1 is the major enzyme in PI(4)P turnover *in vivo* (Table 1) (Guo et al., 1999; Strahl and Thorner, 2007). Loss of function of yeast Sac1 also affects a broad range of cellular defects such as disorganization of the actin cytoskeleton (Foti et al., 2001), secretory pathway (Schorr et al., 2001), inositol auxotrophy, cold sensitivity (Rivas et al., 1999), multiple drug sensitivity (Hughes et al., 1999), vacuolar function (Tahirovic et al., 2005), cell wall maintenance, ER protein quality control systems (Kochendörfer et al., 1999), and sphingolipid metabolism (Brice et al., 2009). Despite the broad impact of Sac1 on yeast cellular functions, Sac1 is not essential for cell viability.

Interestingly, deletion of Sac1 leads to a specific increase of PI(4)P at the plasma membrane (Foti et al., 2001; Stefan et al., 2011). Since Sac1 is localized on the ER and Golgi membrane, an intriguing question was raised as to whether Sac1 is able to hydrolyze PI(4)P in a *trans* mode. The crystal structure of Sac1 revealed a ~70-residues flexible linker between the catalytic domain and the C-terminal transmembrane domain that allows the catalytic domain of Sac1 to stretch more than 15 nm from its anchoring membrane (Manford et al., 2010). This idea is further supported by a recent finding that Sac1 can hydrolyze PI(4)P on the plasma membrane *in trans* at ER-plasma membrane contact sites in the presence of oxysterol binding homology (Osh) proteins (Stefan et al., 2011).

4.2 Mammalian Sac1

In mammals, Sac1 is ubiquitously expressed in most mouse tissues with a particularly high level in cerebellum, hippocampus, and heart (Nemoto et al., 2000). Similar to yeast Sac1, mammalian Sac1 shuttles between ER and Golgi apparatus. In quiescent cells, mammalian Sac1 accumulates at the Golgi to downregulate anterograde trafficking by depleting PI(4)P, thus slow secretion from the Golgi apparatus (Blagoveshchenskaya et al., 2008; Blagoveshchenskaya and Mayinger, 2009; Piao and Mayinger, 2012). The Golgi localization has been shown to be dependent on the leucine zipper region of mammalian Sac1 to form oligomer and the recruitment of the coat protein COPII complex. When quiescent cells are stimulated by mitogens, Sac1 dissociates from its oligomeric state and rapidly shuttles back to the endoplasmic reticulum (ER). The retrograde trafficking of Sac1 is mediated by the coatomer COPI complex through the interaction with a canonical KXXXX motif at C-terminus of mammalian Sac1 (Rohde et al., 2003) (Fig. 1A).

Unlike the case in yeast, genetic ablation of the single murine Sac1 gene results in progeny failing to progress past E3.5 (Liu et al., 2008) (Table 1). In mammalian cells, knockdown of Sac1 expression results in disorganization of Golgi membranes, and mitotic spindle, which leads to decrease in cell viability in cells due to a blockade in progression through the G2-M cell cycle (Liu et al., 2008; Cheong et al., 2010).

4.3 Sac1 in other species

In *Drosophila*, Sac1 has been shown to regulate axon guidance in the embryonic CNS midline and Sac1 mutants die as embryos with defects in dorsal closure (Wei et al., 2003; Lee et al., 2011). Loss of *Drosophila* Sac1 leads to improper activation of several key events during development: cell shape change in the amnioserosa, increase in JNK signaling (Wei et al., 2003) and activation of Hedgehog signaling (Yavari et al., 2010). These collective data suggest an evolutionarily conserved function and point to an essential housekeeping role of Sac1 in multicellular organisms.

5 Sac2/INPP5f

Sac2, a 128 kDa protein encoded by the gene INPP5f, is an evolutionarily conserved protein in multi-cellular organisms from nematode to human. It is the only Sac phosphatase that lacks an ortholog in *S. cerevisiae*, but is found in other Fungi genus such as *Aspergillus* spp. and *Yarrowia lipolytica*. PFAM domain searches of Sac2 reveal that in addition to the N-terminal Sac phosphatase domain, it possesses a homology Sac2 (hSac2) domain of approximately 120 amino acids of unknown function and structure (Fig. 1A). Interestingly, this unique domain is also present in the mouse tumor protein p63-regulated gene 1-like protein (TPRGL), which is a vertebrate-specific presynaptic protein in CNS (Kremer et al., 2007). Unlike Sac1, Sac2 does not hydrolyze mono-phosphorylated lipids (Minagawa et al., 2001). Initial cloning and characterization show that this enzyme exhibits 5-phosphatase activity specific for PI(4,5)P₂ and PI(3,4,5)P₃, generating PI(4)P and PI(3,4)P₂, respectively (Minagawa et al., 2001). However, given the fact that Sac1 and Sac2 both contain identical catalytic P-loop sequence (CMDCLDRT), the molecular details regarding how Sac2 can recognize di- and tri-phosphorylated lipids (as opposed to mono-phosphorylated lipids in Sac1) remain unclear.

In human, northern blot analysis shows that Sac2 has the highest level of expression in the brain, heart, skeletal muscle, kidney and placenta (Minagawa et al., 2001). In adult mouse heart, the regulation of Sac2 expression level is controlled by histone deacetylase-2 (Hdac2) (Trivedi et al., 2007). In Hdac2 deficient mice, Sac2 expression is increased, which results in attenuated cardiac hypertrophy due to constitutive activation of glycogen synthase kinase 3 beta (Gsk3β) via the activation of Akt. However, in Hdac2 transgenic mice, Sac2 expression is transcriptionally repressed, which results in augmented cardiac hypertrophy associated with inactivated Gsk3β (Trivedi et al., 2007). The role of Sac2 in regulating cardiac hypertrophy is further demonstrated in Sac2 knockout mice. In response to stress, the Sac2 knockout mice have abnormal fetal gene reactivation and heart hypertrophy compared to wild type littermates (Zhu et al., 2009) (Table 1). These data suggest that Sac2 is an important regulator in cardiac hypertrophy response and the regulation may be through its

enzymatic activity on PI(3,4,5)P₃, which in turn downregulates PI3K/AKT signaling pathway.

6 Sac3/Fig4p

6.1 Yeast Sac3/Fig4p

Sac3/Fig4p, a 97 kDa protein encoded by the gene *Fig4* (Factor Induced Gene), was originally identified in a large-scale transposon tagging screen for genes induced by mating pheromone in *S. cerevisiae* (Erdman et al., 1998). Sac3/Fig4p is a phosphoinositide 5-phosphatase that specifically hydrolyzes PI(3,5)P₂ to generate PI(3)P both *in vitro* and *in vivo* (Duex et al., 2006a, 2006b) (Table 1). Unlike other Sac domain proteins, the dephosphorylation reaction is Mg²⁺ dependent (Rudge et al., 2004). Mutations in *Fig4* result in abnormal actin distribution at the shmoo tip, a failure to establish mating cell polarity leading to enlarged cells, and reduced mating efficiency (Erdman et al., 1998). Fig4p-GFP is localized to the limiting membrane of yeast vacuole, and this localization is dependent on a scaffold protein Vac14 (Rudge et al., 2004). Interestingly, Vac14 also positively regulates the Fab1 kinase that is responsible for generating PI(3,5)P₂ from PI(3)P, suggesting the formation of a multi-protein complex in regulating PI(3,5)P₂ levels (Rudge et al., 2004). In this Vac14-Fig4-Fab1 complex, Fig4 is unexpectedly required for the activation of Fab1 to synthesize PI(3,5)P₂ (Gary et al., 2002). For this reason, deletion of Fig4 elicits reduced, rather than elevated levels of PI(3,5)P₂. In agreement with this notion, Fig4 is found to be responsible for the hyperosmotic shock-induced increase as well as the turnover in PI(3,5)P₂ levels (Duex et al., 2006a).

6.2 Mammalian Sac3/Fig4p function and disease

In mouse, Sac3/Fig4 RNA transcript is ubiquitously detected in all tissues, with highest level in the testes, spleen and heart (Chow et al., 2007), whereas Sac3/Fig4 protein level is highest in the brain (Guo et al., 2012; Sbrissa et al., 2007). The Vac14-Fig4-Fab1 ternary protein complex is also conserved in mammals and is responsible for the acute regulation of subcellular levels of PI(3,5)P₂ (Jin et al., 2008; Sbrissa et al., 2008).

Genetic mutations of Sac3/Fig4 lead to a number of diseases in human and mouse, including an autosomal recessive Charcot-Marie-Tooth disorder (CMT4J) and a subset of ALS in human (Chow et al., 2007; Zhang et al., 2008; Chow et al., 2009) as well as neurodegeneration in the “pale tremor” mouse that exhibits severe tremor, abnormal gait and diluted pigmentation (Chow et al., 2007; Lenk et al., 2011) (Table 1). *Fig4* null mice reveal a dramatic reduction in myelin in the brain, spongiform degeneration, gliosis, and juvenile lethality (Nicholson et al., 2011; Winters et al., 2011; Ferguson et al., 2012). These neurological defects have been proposed to be caused by enlarged vacuoles in neurons, which may physically interfere with normal vesicular trafficking. In support of this hypothesis, time-lapse imaging of fibroblasts from CMT4J patients displays impaired trafficking of intracellular organelles due to enlarged vacuoles (Zhang et al., 2008). Furthermore, *Fig4*^{-/-} derived fibroblasts and neurons exhibit enlarged endosomal and lysosomal compartments, and impaired organelle trafficking (Ferguson et al., 2009).

6.3 Sac3/Fig4p disease mutations

Sac3 is a large protein with an N-terminal Sac homology domain (Fig. 1A). Several missense mutations in the gene are found to be responsible for the CMT4J (I41T and E302K) (Chow et al., 2007; Lenk et al., 2011) and ALS disorder (D48G, D53Y, R388G, and I411V) (Chow et al., 2009). Based on homology structure modeling of the Sac phosphatase domain of human Sac3 to the crystal structure of yeast Sac1, I41T, which is a recessive mutation found in patients with CMT4J, is mapped to the Sac N-terminal sub-domain and 40 Å away from the catalytic site (Fig. 3). The I41T substitution may not affect the catalytic activity of Fig4, but the protein stability and/or protein binding ability. Consistent with this, I41T mutant shows no significant difference in its PI(3,5)P₂ phosphatase activity (Chow et al., 2007). In contrast to WT Fig4, I41T mutation leads to a rapid decline in protein level even in the presence of ArPIKfyve (human homolog of Vac14) (Ikononov et al., 2010) and in yeast two-hybrid system, interaction between Fig4 I41T and Vac14 is significantly impaired (Lenk et al., 2011). Another CMT4J mutation identified at high frequency is E302K. This residue is located on a β-strand of the catalytic domain and is at the interface with the N-domain where it makes several hydrogen bonds with the surrounding residues (Fig. 3). Mutation of E302K is predicted to destabilize the protein and render the protein inactive. This is supported by the fact that Fig4 E302K construct fails to rescue the enlarged vacuole observed in *Fig4* mutant in yeast (Nicholson et al., 2011).

D48G and D53Y, which have been predicted to be responsible for a subset of ALS disease (Chow et al., 2009), is mapped to the exposed surface area on the N-domain. These two mutations may affect the surface property and consequently may interfere with protein–protein interactions mediated through the N-domain. The R388G mutation is located in a loop close to the catalytic P-loop (Fig. 3). This arginine residue likely contributes to positive electrostatic potentials at the catalytic site. R388G mutation may directly affect substrate binding and thus the catalytic activity of the enzyme. In fact, R388G mutant fails to rescue enlarged vacuole phenotype in *Fig4* yeast strain (Chow et al., 2009). The other mutation, I411V is found in a hydrophobic core on the opposite site of the P-loop (Fig. 3). The deleterious effect of this substitution by a valine residue with similar hydrophobic property as isoleucine is not clear and the I411V mutant behaves similar to wild type in yeast rescue assays (Chow et al., 2009).

7 Synaptojanins

7.1 Enzyme activity and domain organization

Synaptojanins contain an N-terminal Sac domain, a central 5-phosphatase domain, a RNA binding domain (RBD) that is conserved in multi-cellular organisms, but not in yeast, and a C-terminal PRD (Fig. 1A). The Sac phosphatase domain is ~30% sequence identical to yeast Sac1. Biochemical characterization shows that the N-terminal Sac domain of synaptojanins predominantly dephosphorylates PI(3)P and PI(4)P, whereas the 5-phosphatase domain dephosphorylates PI(4,5)P₂ and PI(3,4,5)P₃ at the 5' position of the inositol ring (Guo et al., 1999; Nemoto et al., 2001) (Table 1). It is speculated that the Sac domain and the 5-phosphatase domains may function in a concerted way. In the concerted model, the 5-

phosphatase domain converts PI(4,5)P₂ to PI(4)P, which is directly channeled to Sac domain for further dephosphorylation to PI(3)P without accumulation of the intermediate en route.

7.2 Yeast synaptojanins function and regulation

S. cerevisiae encodes in its genome three synaptojanin-like genes, *sjl1*, *sjl2*, and *sjl3*, which are involved in endocytic membrane trafficking pathway (Singer-Krüger et al., 1998; Stolz et al., 1998). It is interesting to note that the Sac domain in *sjl1* lacks the conserved CX₅R motif and consequently does not exhibit phosphatase activity (Guo et al., 1999). Although neither of these genes is essential for viability, triple mutant of *sjl1-3* is inviable under normal growth conditions (Srinivasan et al., 1997) and *sjl1 sjl2* double mutant displays a pronounced increase in PI(4,5)P₂ level (Stefan et al., 2002). The hydrolysis of PI(4,5)P₂ is critical for clathrin and actin-mediated endocytosis, including vesicle scission (Hauke, 2005). Consistent with this, *sjl2* is shown to localize to cortical actin patches via recruitment of actin patch component Abp1. The physical interaction is critical for the regulation of vesicle formation and fission during endocytosis (Stefan et al., 2005). In addition, *sjl1* and *sjl2* have been implicated in eisosome receptor-mediated endocytosis based on studies using eisosome marker Pil1 (Murphy et al., 2011). Besides regulating the level of PI(4,5)P₂, *sjl2* and *sjl3* are also involved in mediating PI(3)P level. In *ymr1 sjl3* double mutant (*ymr1* is the ortholog of MTM family of PI-3-phosphatase), cells display more than two fold increase in PI(3)P level compare to either single mutant (Parrish et al., 2004). The role of *sjl3* in PI(3)P level likely contributes to the AP-1/clathrin transport from TGN (trans Golgi network) to endosome (Ha et al., 2003).

7.3 Mammalian synaptojanins function and regulation

There are two members that are present in mammals: synaptojanin 1 and 2. The founding member of this family, the mammalian synaptojanin 1 was first cloned and characterized in the search for presynaptic factors involved in synaptic vesicle recycling (McPherson et al., 1996). Synaptojanin 1 can be alternatively spliced to form a 145 kDa and 170 kDa isoforms. The 145 kDa isoform is highly expressed in adult neurons and at nerve terminals where it is involved in clathrin-dependent synaptic vesicle recycling (Cremona et al., 1999). On the other hand, the 170 kDa isoform, containing an additional C-terminal fragment involved in protein-protein interactions, is ubiquitously expressed in a variety of tissues (Ramjaun and McPherson, 1996). These two splicing isoforms have been shown to be differentially targeted to and regulate the maturation of clathrin-coated pits through the interactions with an array of proteins at sites of clathrin-mediated endocytosis (Perera et al., 2006). It has been shown that both isoforms interact with the BAR domain protein endophilin to preferentially remove PI(4,5)P₂ from curved membranes as opposed to flat ones (Chang-Ileto et al., 2011). Other interacting partners include amphiphysin (Ramjaun and McPherson, 1998), intersectin (Koh et al., 2004; Irie et al., 2005), Grb2 (growth-factor-receptor-bound protein 2) (Ringstad et al., 1997), Snx9 (Yeow-Fong et al., 2005), and Myosin 1E (Krendel et al., 2007). All these interactions are critical for proper localization at clathrin-mediated endocytic sites and/or catalytic activity of the enzyme. However, only the 170 kDa isoform directly interacts with clathrin and its adaptor protein AP-2 (Haffner et al., 2000) via its unique C-terminal portion. These studies suggest a role of synaptojanin in endocytosis. The hydrolysis

of PI(4,5)P₂ by synaptojanins may facilitate the disassembly of clathrin from endocytic vesicles (Slepnev and De Camilli, 2000).

The *in vivo* functions of synaptojanin are also well studied in knock mice. Deletion of this gene in mice causes neurological defects and early postnatal lethality (Cremona et al., 1999). In neurons of synaptojanin mutant mouse, abnormal amount of clathrin-coated vesicles are observed to accumulate at the nerve termini and a delay in re-entry of recycling vesicles, which correlates with increased PI(4,5)P₂ levels (Cremona et al., 1999; Kim et al., 2002; Mani et al., 2007). These mutant mice are further shown to have impaired constitutive and induced endocytosis of post synaptic AMPA receptors (Gong and De Camilli, 2008). In contrast to synaptojanin 1 deletion, overexpression of synaptojanin 1 in transgenic mice or neuroblastoma cell line leads to enlarged endosomes (Cossec et al., 2012). Interestingly, synaptojanin 1 has been shown to be a trisomic gene located at chromosome 21 that is overexpressed in Down's syndrome patients. The increased levels of synaptojanin 1 may account for the observed enlarged endosomes in neurons and brain dysfunction in Down's syndrome patients (Arai et al., 2002; Voronov et al., 2008). Mutations of synaptojanin 1 have also been linked to bipolar disorder (Saito et al., 2001; Stopkova et al., 2004). However, the mechanism for the association of synaptojanin 1 with bipolar disorder is not clear.

Similar to synaptojanin 1, synaptojanin 2 also consists of several splice isoforms that are referred to as 2A and 2B, in which the 2B variant can undergo further alternative splicing to generate 2B1 and 2B2 (Khvotchev and Südhof, 1998). The 2A isoform is broadly expressed in many tissues, while the 2B isoform is predominantly expressed at nerve terminals in the brain as well as at spermatid manchette in adult testis (Nemoto and De Camilli, 1999; Nemoto et al., 2001). Unlike synaptojanin 1, the function of synaptojanin 2 is less well understood. It has been shown that synaptojanin 2 can be recruited to plasma membrane via the direct interaction with active Rac1 (Malecz et al., 2000). The interaction of synaptojanin 2 with Rac1 has been suggested to play a role in human glioma cell invasion and migration (Chuang et al., 2004). In mouse mutant strain called Mozart, a mutation in synaptojanin 2 that abolishes its enzymatic activity has been shown to be responsible for the progressive hearing loss (Manji et al., 2011), suggesting a link between phosphoinositide metabolism to hair cell survival and hearing.

7.4 Synaptojanins in other species

The phenotypes observed in both mice and human are further corroborated by studies in *Danio rerio*, *Drosophila melanogaster*, and *Caenorhabditis elegans*. In *D. rerio*, synaptojanin 1 mutant displays a defect in cone photoreceptor ribbon anchoring, abnormal synaptic transmission at synapses (Holzhausen et al., 2009) as well as a role in maintaining the quantity, fusion and release of synaptic vesicles at hair-cell ribbon synapses (Trapani et al., 2009). In *D. melanogaster*, synaptojanin is recruited by endophilin to promote synaptic vesicle uncoating (Verstreken et al., 2003). Overexpression of synaptojanin in *Drosophila* leads to abnormal synaptic morphology (Chang and Min, 2009). In *C. elegans*, mutations in synaptojanin (unc-26) not only disrupt synaptic vesicle recycling (Harris et al., 2000), but also enhance polyglutamine toxicity which is a phenotype observed in Huntington-

interacting protein 1 mutant implicated in neuronal function in Huntington's disease (Parker et al., 2007). These data point to a role of synaptojanins in clathrin-mediated endocytic processes particularly in synaptic vesicle recycling at the nerve terminus.

8 Plant Sac phosphatase family

Phosphoinositides in plants regulate many cellular activities such as vesicle trafficking (Kim et al., 2001), stomatal movement (Jung et al., 2002), polar tip growth of pollen tube and root hairs (Kost et al., 1999), and responses to stress and hormonal treatments (Pical et al., 1999; De Wald et al., 2001). In *Arabidopsis thaliana*, a genome wide analysis reveals the presence of nine Sac domain-containing phosphatases with high sequence similarity (55%–69%) to Sac domain of yeast Sac1, which are subsequently named AtSac1 to AtSac9 (Zhong and Ye, 2003). These nine genes can be sub-divided into three groups based on sequence similarity: 1) Group I contains C-terminal sequences with a range of 252–338 amino acids. 2) Group II contains a C-terminal transmembrane motif. 3) Group III consists of only one member, AtSac9, with a long stretch of C-terminal sequences of ~1100 amino acids (Zhong and Ye, 2003) (Fig. 1B). Although other Sac homologs are found in other plant species, the following section will mainly focus on the Sac domain phosphatases in the context of *Arabidopsis*.

The group I enzymes consist of AtSac1–AtSac5, which are considered as Fig4 homologs based on sequence identity. Their gene expression is widely distributed in most plant organs, with relatively lower level in mature leaves. Among this group, AtSac1 has been well characterized. AtSac1 is a PI-phosphatase specific for PI(3,5)P₂ and mutations in AtSac1 cause defects in cell morphogenesis, such as a decrease in cell wall synthesis, a reduction in cell elongation, defect in actin cytoskeleton and a global change in plant architecture (Zhong et al., 2005).

The group II, which consists of AtSac6–AtSac8, is most closely related to yeast Sac1, which contains two transmembrane motifs at its C-terminal end. AtSac7 and AtSac8 are widely expressed in most plant organs, whereas AtSac6 is only present in flowers (Zhong and Ye, 2003). Among the three enzymes, only AtSac7 has been characterized so far. AtSac7 has been shown to be enriched at the TGN-like compartment at the tips of growing root hairs (Thole et al., 2008). Mutation of this gene is responsible for the root hair defective 4-1 (rhd4-1) phenotypes that displayed aberrant root hair and polarized tip growth. This phenotype is associated with an accumulation of PI(4)P in the tip root hairs, consistent with the *in vitro* assay showing its preferred PI phosphatase activity toward PI(4)P (Thole and Nielsen, 2008; Thole et al., 2008).

AtSac9 is the only enzyme belongs in Group III due to its distinct domain features that include a long C-terminal domain and the presence of a WW domain within the Sac domain (Fig. 1B). Microarray analysis shows highest expression in the roots and lower expression in the leaves and flowers (Zhong and Ye, 2003). AtSac9 mutant displays elevated PI(4,5)P₂ and water soluble Ins(1,4,5)P₃ levels in the roots compared to wild type plants (Williams et al., 2005). The AtSac9 mutants are hypersensitive to external stress and constitutively

overexpress stress-induced genes and overaccumulate reactive-oxygen species (Williams et al., 2005).

9 Concluding remarks

It has only been a little more than 20 years since the discovery of the first Sac domain-containing protein, the yeast Sac1. With the advance of whole-genome sequencing, all the Sac homology domain-containing PI phosphatases have been annotated in quite a few species, including yeast, mammal, and plant. Extensive functional studies on this family of PI phosphatases have been documented and many genetic diseases have been linked to mutations in genes encoding PI phosphatases in this family. Meanwhile, crystal structure for the Sac phosphatase domain provides an excellent model for molecular understanding of genetic mutations occurred on Sac domain-containing phosphatases. Despite the current progress in this field, there are still several key questions remain to be addressed. Given that the catalytic pocket residues are highly conserved across species (Fig. 2B), the mechanistic detail for how substrate recognition and specificity is achieved would require detail structural information of the Sac domain with its cognate substrate. Moreover, how the activity of individual Sac domain containing proteins is precisely controlled by sophisticated signaling networks under physiological or disease state remain to be explored in the future. With the availability of disease models that have deletions or mutations of the Sac domain family members from a variety of organisms, future studies to screen for small-molecules that interfere with phosphoinositide metabolism may lead to potential therapeutic agents for the treatment of human diseases.

Acknowledgments

This work is supported by grants from NIH: 1R01GM094347 (to Y.M.) and is funded by the Cornell University Harry Samuel Mann Award (to F.H.)

References

- Arai Y, Ijuin T, Takenawa T, Becker LE, Takashima S. Excessive expression of synaptojanin in brains with Down syndrome. *Brain Dev.* 2002; 24(2):67–72.10.1016/S0387-7604(01)00405-3 [PubMed: 11891094]
- Bankaitis VA, Aitken JR, Cleves AE, Dowhan W. An essential role for a phospholipid transfer protein in yeast Golgi function. *Nature.* 1990; 347(6293):561–562.10.1038/347561a0 [PubMed: 2215682]
- Barford D, Das AK, Egloff MP. The structure and mechanism of protein phosphatases: insights into catalysis and regulation. *Annu Rev Biophys Biomol Struct.* 1998; 27(1):133–164.10.1146/annurev.biophys.27.1.133 [PubMed: 9646865]
- Begley MJ, Taylor GS, Kim SA, Veine DM, Dixon JE, Stuckey JA. Crystal structure of a phosphoinositide phosphatase, MTMR2: insights into myotubular myopathy and Charcot-Marie-Tooth syndrome. *Mol Cell.* 2003; 12(6):1391–1402.10.1016/S1097-2765(03)00486-6 [PubMed: 14690594]
- Blagoveshchenskaya A, Cheong FY, Rohde HM, Glover G, Knödler A, Nicolson T, Boehmelt G, Mayinger P. Integration of Golgi trafficking and growth factor signaling by the lipid phosphatase SAC1. *J Cell Biol.* 2008; 180(4):803–812.10.1083/jcb.200708109 [PubMed: 18299350]
- Blagoveshchenskaya A, Mayinger P. SAC1 lipid phosphatase and growth control of the secretory pathway. *Mol Biosyst.* 2009; 5(1):36–42.10.1039/b810979f [PubMed: 19081929]
- Brice SE, Alford CW, Cowart LA. Modulation of sphingolipid metabolism by the phosphatidylinositol-4-phosphate phosphatase Sac1p through regulation of phosphatidylinositol in

- Saccharomyces cerevisiae*. J Biol Chem. 2009; 284(12):7588–7596.10.1074/jbc.M808325200 [PubMed: 19139096]
- Chang KT, Min KT. Upregulation of three *Drosophila* homologs of human chromosome 21 genes alters synaptic function: implications for Down syndrome. Proc Natl Acad Sci USA. 2009; 106(40): 17117–17122.10.1073/pnas.0904397106 [PubMed: 19805187]
- Chang-Ileto B, Frere SG, Chan RB, Voronov SV, Roux A, Di Paolo G. Synaptojanin 1-mediated PI(4,5)P2 hydrolysis is modulated by membrane curvature and facilitates membrane fission. Dev Cell. 2011; 20(2):206–218.10.1016/j.devcel.2010.12.008 [PubMed: 21316588]
- Cheong FY, Sharma V, Blagoveshchenskaya A, Oorschot VM, Brankatschk B, Klumperman J, Freeze HH, Mayinger P. Spatial regulation of Golgi phosphatidylinositol-4-phosphate is required for enzyme localization and glycosylation fidelity. Traffic. 2010; 11(9):1180–1190.10.1111/j.1600-0854.2010.01092.x [PubMed: 20573065]
- Chow CY, Landers JE, Bergren SK, Sapp PC, Grant AE, Jones JM, Everett L, Lenk GM, McKenna-Yasek DM, Weisman LS, Figlewicz D, Brown RH, Meisler MH. Deleterious variants of FIG4, a phosphoinositide phosphatase, in patients with ALS. Am J Hum Genet. 2009; 84(1):85–88.10.1016/j.ajhg.2008.12.010 [PubMed: 19118816]
- Chow CY, Zhang Y, Dowling JJ, Jin N, Adamska M, Shiga K, Szigeti K, Shy ME, Li J, Zhang X, Lupski JR, Weisman LS, Meisler MH. Mutation of FIG4 causes neurodegeneration in the pale tremor mouse and patients with CMT4J. Nature. 2007; 448(7149):68–72.10.1038/nature05876 [PubMed: 17572665]
- Chuang YY, Tran NL, Rusk N, Nakada M, Berens ME, Symons M. Role of synaptojanin 2 in glioma cell migration and invasion. Cancer Res. 2004; 64(22):8271–8275.10.1158/0008-5472.CAN-04-2097 [PubMed: 15548694]
- Cossec JC, Lavaur J, Berman DE, Rivals I, Hoischen A, Stora S, Ripoll C, Mircher C, Grattau Y, Olivomarin JC, de Chaumont F, Lecourtois M, Antonarakis SE, Veltman JA, Delabar JM, Duyckaerts C, Di Paolo G, Potier MC. Trisomy for synaptojanin1 in Down syndrome is functionally linked to the enlargement of early endosomes. Hum Mol Genet. 2012; 21(14):3156–3172.10.1093/hmg/dds142 [PubMed: 22511594]
- Cremona O, Di Paolo G, Wenk MR, Lüthi A, Kim WT, Takei K, Daniell L, Nemoto Y, Shears SB, Flavell RA, McCormick DA, De Camilli P. Essential role of phosphoinositide metabolism in synaptic vesicle recycling. Cell. 1999; 99(2):179–188.10.1016/S0092-8674(00)81649-9 [PubMed: 10535736]
- Cullen PJ, Cozier GE, Banting G, Mellor H. Modular phosphoinositide-binding domains—their role in signalling and membrane trafficking. Current Biol, CB. 2001; 11:R882–893.
- De Matteis MA, Godi A. PI-loting membrane traffic. Nat Cell Biol. 2004; 6(6):487–492.10.1038/ncb0604-487 [PubMed: 15170460]
- DeWald DB, Torabinejad J, Jones CA, Shope JC, Cangelosi AR, Thompson JE, Prestwich GD, Hama H. Rapid accumulation of phosphatidylinositol 4,5-bisphosphate and inositol 1,4,5-trisphosphate correlates with calcium mobilization in salt-stressed *Arabidopsis*. Plant Physiol. 2001; 126(2):759–769.10.1104/pp.126.2.759 [PubMed: 11402204]
- Di Paolo G, De Camilli P. Phosphoinositides in cell regulation and membrane dynamics. Nature. 2006; 443(7112):651–657.10.1038/nature05185 [PubMed: 17035995]
- Dieck CB, Boss WF, Perera IY. A role for phosphoinositides in regulating plant nuclear functions. Front Plant Sci. 2012; 3:50. [PubMed: 22645589]
- Duex JE, Nau JJ, Kauffman EJ, Weisman LS. Phosphoinositide 5-phosphatase Fig 4p is required for both acute rise and subsequent fall in stress-induced phosphatidylinositol 3,5-bisphosphate levels. Eukaryot Cell. 2006a; 5(4):723–731.10.1128/EC.5.4.723-731.2006 [PubMed: 16607019]
- Duex JE, Tang F, Weisman LS. The Vac14p-Fig4p complex acts independently of Vac7p and couples PI3,5P2 synthesis and turnover. J Cell Biol. 2006b; 172(5):693–704.10.1083/jcb.200512105 [PubMed: 16492811]
- Erdman S, Lin L, Malczynski M, Snyder M. Pheromone-regulated genes required for yeast mating differentiation. J Cell Biol. 1998; 140(3):461–483.10.1083/jcb.140.3.461 [PubMed: 9456310]
- Faulhammer F, Konrad G, Brankatschk B, Tahirovic S, Knödler A, Mayinger P. Cell growth-dependent coordination of lipid signaling and glycosylation is mediated by interactions between

- Sac1p and Dpm1p. *J Cell Biol.* 2005; 168(2):185–191.10.1083/jcb.200407118 [PubMed: 15657391]
- Ferguson CJ, Lenk GM, Jones JM, Grant AE, Winters JJ, Dowling JJ, Giger RJ, Meisler MH. Neuronal expression of Fig4 is both necessary and sufficient to prevent spongiform neurodegeneration. *Hum Mol Genet.* 2012; 21(16):3525–3534.10.1093/hmg/dds179 [PubMed: 22581779]
- Ferguson CJ, Lenk GM, Meisler MH. Defective autophagy in neurons and astrocytes from mice deficient in PI(3,5)P2. *Hum Mol Genet.* 2009; 18(24):4868–4878.10.1093/hmg/ddp460 [PubMed: 19793721]
- Foti M, Audhya A, Emr SD. Sac1 lipid phosphatase and Stt4 phosphatidylinositol 4-kinase regulate a pool of phosphatidylinositol 4-phosphate that functions in the control of the actin cytoskeleton and vacuole morphology. *Mol Biol Cell.* 2001; 12(8):2396–2411. [PubMed: 11514624]
- Gary JD, Sato TK, Stefan CJ, Bonangelino CJ, Weisman LS, Emr SD. Regulation of Fab1 phosphatidylinositol 3-phosphate 5-kinase pathway by Vac7 protein and Fig4, a polyphosphoinositide phosphatase family member. *Mol Biol Cell.* 2002; 13(4):1238–1251.10.1091/mbc.01-10-0498 [PubMed: 11950935]
- Gong LW, De Camilli P. Regulation of postsynaptic AMPA responses by synaptojanin 1. *Proc Natl Acad Sci USA.* 2008; 105(45):17561–17566.10.1073/pnas.0809221105 [PubMed: 18987319]
- Guo J, Ma YH, Yan Q, Wang L, Zeng YS, Wu JL, Li J. Fig4 expression in the rodent nervous system and its potential role in preventing abnormal lysosomal accumulation. *J Neuropathol Exp Neurol.* 2012; 71(1):28–39.10.1097/NEN.0b013e31823deda8 [PubMed: 22157617]
- Guo S, Stolz LE, Lemrow SM, York JD. SAC1-like domains of yeast SAC1, INP52, and INP53 and of human synaptojanin encode polyphosphoinositide phosphatases. *J Biol Chem.* 1999; 274(19):12990–12995.10.1074/jbc.274.19.12990 [PubMed: 10224048]
- Ha SA, Torabinejad J, DeWald DB, Wenk MR, Lucast L, De Camilli P, Newitt RA, Aebersold R, Nothwehr SF. The synaptojanin-like protein Inp53/Sjl3 functions with clathrin in a yeast TGN-to-endosome pathway distinct from the GGA protein-dependent pathway. *Mol Biol Cell.* 2003; 14(4):1319–1333.10.1091/mbc.E02-10-0686 [PubMed: 12686590]
- Haffner C, Di Paolo G, Rosenthal JA, de Camilli P. Direct interaction of the 170 kDa isoform of synaptojanin 1 with clathrin and with the clathrin adaptor AP-2. *Current Biol, CB.* 2000; 10:471–474.
- Ham H, Sreelatha A, Orth K. Manipulation of host membranes by bacterial effectors. *Nat Rev Microbiol.* 2011; 9(9):635–646.10.1038/nrmicro2602 [PubMed: 21765451]
- Hammond GR, Fischer MJ, Anderson KE, Holdich J, Koteci A, Balla T, Irvine RF. PI4P and PI(4,5)P2 are essential but independent lipid determinants of membrane identity. *Science.* 2012; 337(6095):727–730.10.1126/science.1222483 [PubMed: 22722250]
- Harris TW, Hartwig E, Horvitz HR, Jorgensen EM. Mutations in synaptojanin disrupt synaptic vesicle recycling. *J Cell Biol.* 2000; 150(3):589–600.10.1083/jcb.150.3.589 [PubMed: 10931870]
- Hauke V. Phosphoinositide regulation of clathrin-mediated endocytosis. *Biochem Soc Trans.* 2005; 33(Pt 6):1285–1289.10.1042/BST20051285 [PubMed: 16246100]
- Hokin LE, Hokin MR. Phosphoinositides and protein secretion in pancreas slices. *J Biol Chem.* 1958; 233(4):805–810. [PubMed: 13587496]
- Hokin MR, Hokin LE. Enzyme secretion and the incorporation of P32 into phospholipides of pancreas slices. *J Biol Chem.* 1953; 203(2):967–977. [PubMed: 13084667]
- Holzhausen LC, Lewis AA, Cheong KK, Brockerhoff SE. Differential role for synaptojanin 1 in rod and cone photoreceptors. *J Comp Neurol.* 2009; 517(5):633–644.10.1002/cne.22176 [PubMed: 19827152]
- Hughes WE, Pocklington MJ, Orr E, Paddon CJ. Mutations in the *Saccharomyces cerevisiae* gene *SAC1* cause multiple drug sensitivity. *Yeast.* 1999; 15(11):1111–1124.10.1002/(SICI)1097-0061(199908)15:11<1111::AID-YEA440>3.0.CO;2-H [PubMed: 10455234]
- Ikonomov OC, Sbrissa D, Fligger J, Delvecchio K, Shisheva A. ArPIKfyve regulates Sac3 protein abundance and turnover: disruption of the mechanism by Sac3I41T mutation causing Charcot-Marie-Tooth 4J disorder. *J Biol Chem.* 2010; 285(35):26760–26764.10.1074/jbc.C110.154658 [PubMed: 20630877]

- Irie F, Okuno M, Pasquale EB, Yamaguchi Y. EphrinB-EphB signalling regulates clathrin-mediated endocytosis through tyrosine phosphorylation of synaptojanin 1. *Nat Cell Biol.* 2005; 7(5):501–509.10.1038/ncb1252 [PubMed: 15821731]
- Jean S, Kiger AA. Coordination between RAB GTPase and phosphoinositide regulation and functions. *Nat Rev Mol Cell Biol.* 2012; 13(7):463–470.10.1038/nrm3379 [PubMed: 22722608]
- Jin N, Chow CY, Liu L, Zolov SN, Bronson R, Davisson M, Petersen JL, Zhang Y, Park S, Duex JE, Goldowitz D, Meisler MH, Weisman LS. VAC14 nucleates a protein complex essential for the acute interconversion of PI3P and PI(3,5)P(2) in yeast and mouse. *EMBO J.* 2008; 27(24):3221–3234.10.1038/emboj.2008.248 [PubMed: 19037259]
- Jung JY, Kim YW, Kwak JM, Hwang JU, Young J, Schroeder JJ, Hwang I, Lee Y. Phosphatidylinositol 3- and 4-phosphate are required for normal stomatal movements. *Plant Cell.* 2002; 14(10):2399–2412.10.1105/tpc.004143 [PubMed: 12368494]
- Khvotchev M, Südhof TC. Developmentally regulated alternative splicing in a novel synaptojanin. *J Biol Chem.* 1998; 273(4):2306–2311.10.1074/jbc.273.4.2306 [PubMed: 9442075]
- Kim DH, Eu YJ, Yoo CM, Kim YW, Pih KT, Jin JB, Kim SJ, Stenmark H, Hwang I. Trafficking of phosphatidylinositol 3-phosphate from the trans-Golgi network to the lumen of the central vacuole in plant cells. *Plant Cell.* 2001; 13(2):287–301. [PubMed: 11226186]
- Kim WT, Chang S, Daniell L, Cremona O, Di Paolo G, De Camilli P. Delayed reentry of recycling vesicles into the fusion-competent synaptic vesicle pool in synaptojanin 1 knockout mice. *Proc Natl Acad Sci USA.* 2002; 99(26):17143–17148.10.1073/pnas.222657399 [PubMed: 12481038]
- Kochendörfer KU, Then AR, Kearns BG, Bankaitis VA, Mayinger P. Sac1p plays a crucial role in microsomal ATP transport, which is distinct from its function in Golgi phospholipid metabolism. *EMBO J.* 1999; 18(6):1506–1515.10.1093/emboj/18.6.1506 [PubMed: 10075922]
- Koh TW, Verstreken P, Bellen HJ. Dap160/intersectin acts as a stabilizing scaffold required for synaptic development and vesicle endocytosis. *Neuron.* 2004; 43(2):193–205.10.1016/j.neuron.2004.06.029 [PubMed: 15260956]
- Kost B, Lemichez E, Spielhofer P, Hong Y, Tolias K, Carpenter C, Chua NH. Rac homologues and compartmentalized phosphatidylinositol 4, 5-bisphosphate act in a common pathway to regulate polar pollen tube growth. *J Cell Biol.* 1999; 145(2):317–330.10.1083/jcb.145.2.317 [PubMed: 10209027]
- Krauss M, Haucke V. Phosphoinositides: regulators of membrane traffic and protein function. *FEBS Lett.* 2007; 581(11):2105–2111.10.1016/j.febslet.2007.01.089 [PubMed: 17316616]
- Kremer T, Kempf C, Wittenmayer N, Nawrotzki R, Kuner T, Kirsch J, Dresbach T. Mover is a novel vertebrate-specific presynaptic protein with differential distribution at subsets of CNS synapses. *FEBS Lett.* 2007; 581(24):4727–4733.10.1016/j.febslet.2007.08.070 [PubMed: 17869247]
- Krendel M, Osterweil EK, Mooseker MS. Myosin 1E interacts with synaptojanin-1 and dynamin and is involved in endocytosis. *FEBS Lett.* 2007; 581(4):644–650.10.1016/j.febslet.2007.01.021 [PubMed: 17257598]
- Lee JO, Yang H, Georgescu MM, Di Cristofano A, Maehama T, Shi Y, Dixon JE, Pandolfi P, Pavletich NP. Crystal structure of the PTEN tumor suppressor: implications for its phosphoinositide phosphatase activity and membrane association. *Cell.* 1999; 99(3):323–334.10.1016/S0092-8674(00)81663-3 [PubMed: 10555148]
- Lee S, Kim S, Nahm M, Kim E, Kim TI, Yoon JH, Lee S. The phosphoinositide phosphatase Sac1 is required for midline axon guidance. *Mol Cells.* 2011; 32(5):477–482.10.1007/s10059-011-0168-6 [PubMed: 22042447]
- Lemmon MA. Phosphoinositide recognition domains. *Traffic.* 2003; 4(4):201–213.10.1034/j.1600-0854.2004.00071.x [PubMed: 12694559]
- Lenk GM, Ferguson CJ, Chow CY, Jin N, Jones JM, Grant AE, Zolov SN, Winters JJ, Giger RJ, Dowling JJ, Weisman LS, Meisler MH. Pathogenic mechanism of the FIG4 mutation responsible for Charcot-Marie-Tooth disease CMT4J. *PLoS Genet.* 2011; 7(6):e1002104.10.1371/journal.pgen.1002104 [PubMed: 21655088]
- Lichtarge O, Bourne HR, Cohen FE. An evolutionary trace method defines binding surfaces common to protein families. *J Mol Biol.* 1996; 257(2):342–358.10.1006/jmbi.1996.0167 [PubMed: 8609628]


- Liu Y, Boukhelifa M, Tribble E, Morin-Kensicki E, Uetrecht A, Bear JE, Bankaitis VA. The Sac1 phosphoinositide phosphatase regulates Golgi membrane morphology and mitotic spindle organization in mammals. *Mol Biol Cell*. 2008; 19(7):3080–3096.10.1091/mbc.E07-12-1290 [PubMed: 18480408]
- Malecz N, McCabe PC, Spaargaren C, Qiu R, Chuang Y, Symons M. Synaptojanin 2, a novel Rac1 effector that regulates clathrin-mediated endocytosis. *Current Biol*, CB. 2000; 10:1383–1386.
- Manford A, Xia T, Saxena AK, Stefan C, Hu F, Emr SD, Mao Y. Crystal structure of the yeast Sac1: implications for its phosphoinositide phosphatase function. *EMBO J*. 2010; 29(9):1489–1498.10.1038/emboj.2010.57 [PubMed: 20389282]
- Mani M, Lee SY, Lucast L, Cremona O, Di Paolo G, De Camilli P, Ryan TA. The dual phosphatase activity of synaptojanin1 is required for both efficient synaptic vesicle endocytosis and reavailability at nerve terminals. *Neuron*. 2007; 56(6):1004–1018.10.1016/j.neuron.2007.10.032 [PubMed: 18093523]
- Manji SS, Williams LH, Miller KA, Ooms LM, Bahlo M, Mitchell CA, Dahl HH. A mutation in synaptojanin 2 causes progressive hearing loss in the ENU-mutagenised mouse strain Mozart. *PLoS ONE*. 2011; 6(3):e17607.10.1371/journal.pone.0017607 [PubMed: 21423608]
- Martí-Renom MA, Stuart AC, Fiser A, Sánchez R, Melo F, Sali A. Comparative protein structure modeling of genes and genomes. *Annu Rev Biophys Biomol Struct*. 2000; 29(1):291–325.10.1146/annurev.biophys.29.1.291 [PubMed: 10940251]
- McPherson PS, Garcia EP, Slepnev VI, David C, Zhang X, Grabs D, Sossin WS, Bauerfeind R, Nemoto Y, De Camilli P. A presynaptic inositol-5-phosphatase. *Nature*. 1996; 379(6563):353–357.10.1038/379353a0 [PubMed: 8552192]
- Minagawa T, Ijuin T, Mochizuki Y, Takenawa T. Identification and characterization of a sac domain-containing phosphoinositide 5-phosphatase. *J Biol Chem*. 2001; 276(25):22011–22015.10.1074/jbc.M101579200 [PubMed: 11274189]
- Murphy ER, Boxberger J, Colvin R, Lee SJ, Zahn G, Loor F, Kim K. Pil1, an eisosome organizer, plays an important role in the recruitment of synaptojanins and amphiphysins to facilitate receptor-mediated endocytosis in yeast. *Eur J Cell Biol*. 2011; 90(10):825–833.10.1016/j.ejcb.2011.06.006 [PubMed: 21872358]
- Nemoto Y, De Camilli P. Recruitment of an alternatively spliced form of synaptojanin 2 to mitochondria by the interaction with the PDZ domain of a mitochondrial outer membrane protein. *EMBO J*. 1999; 18(11):2991–3006.10.1093/emboj/18.11.2991 [PubMed: 10357812]
- Nemoto Y, Kearns BG, Wenk MR, Chen H, Mori K, Alb JG Jr, De Camilli P, Bankaitis VA. Functional characterization of a mammalian Sac1 and mutants exhibiting substrate-specific defects in phosphoinositide phosphatase activity. *J Biol Chem*. 2000; 275(44):34293–34305.10.1074/jbc.M003923200 [PubMed: 10887188]
- Nemoto Y, Wenk MR, Watanabe M, Daniell L, Murakami T, Ringstad N, Yamada H, Takei K, De Camilli P. Identification and characterization of a synaptojanin 2 splice isoform predominantly expressed in nerve terminals. *J Biol Chem*. 2001; 276(44):41133–41142.10.1074/jbc.M106404200 [PubMed: 11498538]
- Nicholson G, Lenk GM, Reddel SW, Grant AE, Towne CF, Ferguson CJ, Simpson E, Scheuerle A, Yasick M, Hoffman S, Blouin R, Brandt C, Coppola G, Biesecker LG, Batish SD, Meisler MH. Distinctive genetic and clinical features of CMT4J: a severe neuropathy caused by mutations in the PI(3,5)P(2) phosphatase *FIG4*. *Brain*. 2011; 134:1959–1971. [PubMed: 21705420]
- Novick P, Osmond BC, Botstein D. Suppressors of yeast actin mutations. *Genetics*. 1989; 121(4):659–674. [PubMed: 2656401]
- Odorizzi G, Babst M, Emr SD. Phosphoinositide signaling and the regulation of membrane trafficking in yeast. *Trends Biochem Sci*. 2000; 25(5):229–235.10.1016/S0968-0004(00)01543-7 [PubMed: 10782093]
- Osborne SL, Thomas CL, Gschmeissner S, Schiavo G. Nuclear PtdIns(4,5)P2 assembles in a mitotically regulated particle involved in pre-mRNA splicing. *J Cell Sci*. 2001; 114(Pt 13):2501–2511. [PubMed: 11559758]
- Parker JA, Metzler M, Georgiou J, Mage M, Roder JC, Rose AM, Hayden MR, Neri C. Huntingtin-interacting protein 1 influences worm and mouse presynaptic function and protects *Caenorhabditis*

C. elegans neurons against mutant polyglutamine toxicity. *J Neurosci.* 2007; 27:11056–11064. [PubMed: 17928447]

- Parrish WR, Stefan CJ, Emr SD. Essential role for the myotubularin-related phosphatase Ymr1p and the synaptojanin-like phosphatases Sjl2p and Sjl3p in regulation of phosphatidylinositol 3-phosphate in yeast. *Mol Biol Cell.* 2004; 15(8):3567–3579.10.1091/mbc.E04-03-0209 [PubMed: 15169871]
- Perera RM, Zoncu R, Lucast L, De Camilli P, Toomre D. Two synaptojanin 1 isoforms are recruited to clathrin-coated pits at different stages. *Proc Natl Acad Sci USA.* 2006; 103(51):19332–19337.10.1073/pnas.0609795104 [PubMed: 17158794]
- Piao H, Mayinger P. Growth and metabolic control of lipid signalling at the Golgi. *Biochem Soc Trans.* 2012; 40(1):205–209.10.1042/BST20110637 [PubMed: 22260691]
- Pical C, Westergren T, Dove SK, Larsson C, Sommarin M. Salinity and hyperosmotic stress induce rapid increases in phosphatidylinositol 4,5-bisphosphate, diacylglycerol pyrophosphate, and phosphatidylcholine in *Arabidopsis thaliana* cells. *J Biol Chem.* 1999; 274(53):38232–38240.10.1074/jbc.274.53.38232 [PubMed: 10608898]
- Pizarro-Cerdá J, Cossart P. Subversion of phosphoinositide metabolism by intracellular bacterial pathogens. *Nat Cell Biol.* 2004; 6(11):1026–1033.10.1038/ncb1104-1026 [PubMed: 15516995]
- Ramjaun AR, McPherson PS. Tissue-specific alternative splicing generates two synaptojanin isoforms with differential membrane binding properties. *J Biol Chem.* 1996; 271(40):24856–24861.10.1074/jbc.271.40.24856 [PubMed: 8798761]
- Ramjaun AR, McPherson PS. Multiple amphiphysin II splice variants display differential clathrin binding: identification of two distinct clathrin-binding sites. *J Neurochem.* 1998; 70(6):2369–2376.10.1046/j.1471-4159.1998.70062369.x [PubMed: 9603201]
- Ringstad N, Nemoto Y, De Camilli P. The SH3p4/Sh3p8/SH3p13 protein family: binding partners for synaptojanin and dynamin via a Grb2-like Src homology 3 domain. *Proc Natl Acad Sci USA.* 1997; 94(16):8569–8574.10.1073/pnas.94.16.8569 [PubMed: 9238017]
- Rivas MP, Kearns BG, Xie Z, Guo S, Sekar MC, Hosaka K, Kagiwada S, York JD, Bankaitis VA. Pleiotropic alterations in lipid metabolism in yeast *sac1* mutants: relationship to “bypass *Sec14p*” and inositol auxotrophy. *Mol Biol Cell.* 1999; 10(7):2235–2250. [PubMed: 10397762]
- Rohde HM, Cheong FY, Konrad G, Paiha K, Mayinger P, Boehmelt G. The human phosphatidylinositol phosphatase SAC1 interacts with the coatamer I complex. *J Biol Chem.* 2003; 278(52):52689–52699.10.1074/jbc.M307983200 [PubMed: 14527956]
- Rudge SA, Anderson DM, Emr SD. Vacuole size control: regulation of PtdIns(3,5)P₂ levels by the vacuole-associated Vac14-Fig4 complex, a PtdIns(3,5)P₂-specific phosphatase. *Mol Biol Cell.* 2004; 15(1):24–36.10.1091/mbc.E03-05-0297 [PubMed: 14528018]
- Saito T, Guan F, Papolos DF, Lau S, Klein M, Fann CS, Lachman HM. Mutation analysis of SYNJ1: a possible candidate gene for chromosome 21q22-linked bipolar disorder. *Mol Psychiatry.* 2001; 6(4):387–395.10.1038/sj.mp.4000871 [PubMed: 11443522]
- Sbrissa D, Ikononov OC, Fenner H, Shisheva A. ArPIKfyve homomeric and heteromeric interactions scaffold PIKfyve and Sac3 in a complex to promote PIKfyve activity and functionality. *J Mol Biol.* 2008; 384(4):766–779.10.1016/j.jmb.2008.10.009 [PubMed: 18950639]
- Sbrissa D, Ikononov OC, Fu Z, Ijuin T, Gruenberg J, Takenawa T, Shisheva A. Core protein machinery for mammalian phosphatidylinositol 3,5-bisphosphate synthesis and turnover that regulates the progression of endosomal transport. Novel Sac phosphatase joins the ArPIKfyve-PIKfyve complex. *J Biol Chem.* 2007; 282(33):23878–23891.10.1074/jbc.M611678200 [PubMed: 17556371]
- Schor M, Then A, Tahirovic S, Hug N, Mayinger P. The phosphoinositide phosphatase Sac1p controls trafficking of the yeast Chs3p chitin synthase. *Current Biol, CB.* 2001; 11:1421–1426.
- Singer-Krüger B, Nemoto Y, Daniell L, Ferro-Novick S, De Camilli P. Synaptojanin family members are implicated in endocytic membrane traffic in yeast. *J Cell Sci.* 1998; 111(Pt 22):3347–3356. [PubMed: 9788876]
- Slepnev VI, De Camilli P. Accessory factors in clathrin-dependent synaptic vesicle endocytosis. *Nat Rev Neurosci.* 2000; 1(3):161–172.10.1038/35044540 [PubMed: 11257904]

- Srinivasan S, Seaman M, Nemoto Y, Daniell L, Suchy SF, Emr S, De Camilli P, Nussbaum R. Disruption of three phosphatidylinositol-polyphosphate 5-phosphatase genes from *Saccharomyces cerevisiae* results in pleiotropic abnormalities of vacuole morphology, cell shape, and osmohomeostasis. *Eur J Cell Biol.* 1997; 74(4):350–360. [PubMed: 9438131]
- Stefan CJ, Audhya A, Emr SD. The yeast synaptojanin-like proteins control the cellular distribution of phosphatidylinositol (4,5)-bisphosphate. *Mol Biol Cell.* 2002; 13(2):542–557.10.1091/mbc.01-10-0476 [PubMed: 11854411]
- Stefan CJ, Manford AG, Baird D, Yamada-Hanff J, Mao Y, Emr SD. Osh proteins regulate phosphoinositide metabolism at ER-plasma membrane contact sites. *Cell.* 2011; 144(3):389–401.10.1016/j.cell.2010.12.034 [PubMed: 21295699]
- Stefan CJ, Padilla SM, Audhya A, Emr SD. The phosphoinositide phosphatase Sjl2 is recruited to cortical actin patches in the control of vesicle formation and fission during endocytosis. *Mol Cell Biol.* 2005; 25(8):2910–2923.10.1128/MCB.25.8.2910-2923.2005 [PubMed: 15798181]
- Stolz LE, Huynh CV, Thorner J, York JD. Identification and characterization of an essential family of inositol polyphosphate 5-phosphatases (INP51, INP52 and INP53 gene products) in the yeast *Saccharomyces cerevisiae*. *Genetics.* 1998; 148(4):1715–1729. [PubMed: 9560389]
- Stopkova P, Vevera J, Paclt I, Zukov I, Lachman HM. Analysis of SYNJ1, a candidate gene for 21q22 linked bipolar disorder: a replication study. *Psychiatry Res.* 2004; 127(1–2):157–161.10.1016/j.psychres.2004.03.003 [PubMed: 15261714]
- Strahl T, Thorner J. Synthesis and function of membrane phosphoinositides in budding yeast, *Saccharomyces cerevisiae*. *Biochim Biophys Acta.* 2007; 1771(3):353–404.10.1016/j.bbalip.2007.01.015 [PubMed: 17382260]
- Tahirovic S, Schorr M, Mayinger P. Regulation of intracellular phosphatidylinositol-4-phosphate by the Sac1 lipid phosphatase. *Traffic.* 2005; 6(2):116–130.10.1111/j.1600-0854.2004.00255.x [PubMed: 15634212]
- Takenawa T, Itoh T. Membrane targeting and remodeling through phosphoinositide-binding domains. *IUBMB Life.* 2006; 58(5–6):296–303.10.1080/15216540600732039 [PubMed: 16754321]
- Thole JM, Nielsen E. Phosphoinositides in plants: novel functions in membrane trafficking. *Curr Opin Plant Biol.* 2008; 11(6):620–631.10.1016/j.pbi.2008.10.010 [PubMed: 19028349]
- Thole JM, Vermeer JE, Zhang Y, Gadella TW Jr, Nielsen E. Root hair defective4 encodes a phosphatidylinositol-4-phosphate phosphatase required for proper root hair development in *Arabidopsis thaliana*. *Plant Cell.* 2008; 20(2):381–395.10.1105/tpc.107.054304 [PubMed: 18281508]
- Trapani JG, Obholzer N, Mo W, Brockerhoff SE, Nicolson T. Synaptojanin1 is required for temporal fidelity of synaptic transmission in hair cells. *PLoS Genet.* 2009; 5(5):e1000480.10.1371/journal.pgen.1000480 [PubMed: 19424431]
- Trivedi CM, Luo Y, Yin Z, Zhang M, Zhu W, Wang T, Floss T, Goettlicher M, Noppinger PR, Wurst W, Ferrari VA, Abrams CS, Gruber PJ, Epstein JA. Hdac2 regulates the cardiac hypertrophic response by modulating Gsk3 beta activity. *Nat Med.* 2007; 13(3):324–331.10.1038/nm1552 [PubMed: 17322895]
- Verstreken P, Koh TW, Schulze KL, Zhai RG, Hiesinger PR, Zhou Y, Mehta SQ, Cao Y, Roos J, Bellen HJ. Synaptojanin is recruited by endophilin to promote synaptic vesicle uncoating. *Neuron.* 2003; 40(4):733–748.10.1016/S0896-6273(03)00644-5 [PubMed: 14622578]
- Viiri K, Mäki M, Lohi O. Phosphoinositides as regulators of protein-chromatin interactions. *Sci Signal.* 2012; 5(222):pe19.10.1126/scisignal.2002917 [PubMed: 22550339]
- Voronov SV, Frere SG, Giovedi S, Pollina EA, Borel C, Zhang H, Schmidt C, Akeson EC, Wenk MR, Cimasoni L, Arancio O, Davisson MT, Antonarakis SE, Gardiner K, De Camilli P, Di Paolo G. Synaptojanin 1-linked phosphoinositide dyshomeostasis and cognitive deficits in mouse models of Down's syndrome. *Proc Natl Acad Sci USA.* 2008; 105(27):9415–9420.10.1073/pnas.0803756105 [PubMed: 18591654]
- Wang X, Zhang X, Dong XP, Samie M, Li X, Cheng X, Goschka A, Shen D, Zhou Y, Harlow J, Zhu MX, Clapham DE, Ren D, Xu H. TPC proteins are phosphoinositide-activated sodium-selective ion channels in endosomes and lysosomes. *Cell.* 2012; 151(2):372–383.10.1016/j.cell.2012.08.036 [PubMed: 23063126]

- Wei HC, Sanny J, Shu H, Baillie DL, Brill JA, Price JV, Harden N. The Sac1 lipid phosphatase regulates cell shape change and the JNK cascade during dorsal closure in *Drosophila*. *Current Biol*. 2003; 13:1882–1887.
- Whitters EA, Cleves AE, McGee TP, Skinner HB, Bankaitis VA. SAC1p is an integral membrane protein that influences the cellular requirement for phospholipid transfer protein function and inositol in yeast. *J Cell Biol*. 1993; 122(1):79–94.10.1083/jcb.122.1.79 [PubMed: 8314848]
- Williams ME, Torabinejad J, Cohick E, Parker K, Drake EJ, Thompson JE, Hortter M, Dewald DB. Mutations in the Arabidopsis phosphoinositide phosphatase gene SAC9 lead to overaccumulation of PtdIns(4,5)P2 and constitutive expression of the stress-response pathway. *Plant Physiol*. 2005; 138(2):686–700.10.1104/pp.105.061317 [PubMed: 15923324]
- Winters JJ, Ferguson CJ, Lenk GM, Giger-Mateeva VI, Shrager P, Meisler MH, Giger RJ. Congenital CNS hypomyelination in the Fig4 null mouse is rescued by neuronal expression of the PI(3,5)P(2) phosphatase Fig4. *J Neurosci*. 2011; 31:17736–17751. [PubMed: 22131434]
- Wood CS, Hung CS, Huoh YS, Mousley CJ, Stefan CJ, Bankaitis V, Ferguson KM, Burd CG. Local control of phosphatidylinositol 4-phosphate signaling in the Golgi apparatus by Vps74 and Sac1 phosphoinositide phosphatase. *Mol Biol Cell*. 2012; 23(13):2527–2536.10.1091/mbc.E12-01-0077 [PubMed: 22553352]
- Yavari A, Nagaraj R, Owusu-Ansah E, Folick A, Ngo K, Hillman T, Call G, Rohatgi R, Scott MP, Banerjee U. Role of lipid metabolism in smoothened derepression in hedgehog signaling. *Dev Cell*. 2010; 19(1):54–65.10.1016/j.devcel.2010.06.007 [PubMed: 20643350]
- Yeow-Fong L, Lim L, Manser E. SNX9 as an adaptor for linking synaptojanin-1 to the Cdc42 effector ACK1. *FEBS Lett*. 2005; 579(22):5040–5048.10.1016/j.febslet.2005.07.093 [PubMed: 16137687]
- Zhang X, Chow CY, Sahenk Z, Shy ME, Meisler MH, Li J. Mutation of FIG4 causes a rapidly progressive, asymmetric neuronal degeneration. *Brain*. 2008; 131:1990–2001. [PubMed: 18556664]
- Zhong R, Burk DH, Nairn CJ, Wood-Jones A, Morrison WH 3rd, Ye ZH. Mutation of SAC1, an *Arabidopsis* SAC domain phosphoinositide phosphatase, causes alterations in cell morphogenesis, cell wall synthesis, and actin organization. *Plant Cell*. 2005; 17(5):1449–1466.10.1105/tpc.105.031377 [PubMed: 15805481]
- Zhong R, Ye ZH. The SAC domain-containing protein gene family in *Arabidopsis*. *Plant Physiol*. 2003; 132(2):544–555.10.1104/pp.103.021444 [PubMed: 12805586]
- Zhong S, Hsu F, Stefan CJ, Wu X, Patel A, Cosgrove MS, Mao Y. Allosteric activation of the phosphoinositide phosphatase Sac1 by anionic phospholipids. *Biochemistry*. 2012; 51(15):3170–3177.10.1021/bi300086c [PubMed: 22452743]
- Zhu W, Trivedi CM, Zhou D, Yuan L, Lu MM, Epstein JA. Inpp5f is a polyphosphoinositide phosphatase that regulates cardiac hypertrophic responsiveness. *Circ Res*. 2009; 105(12):1240–1247.10.1161/CIRCRESAHA.109.208785 [PubMed: 19875726]

**Figure 1.**

Sac domain-containing phosphatase family. (A) The yeast and mammalian Sac phosphatases can be divided into three subfamilies. TM: transmembrane motif; HSac2: homology of Sac2; KKKXX: COPI binding motif; RBD: RNA binding domain; PRD: proline rich domain. (B) The plant Sac domain-containing phosphatases.




Figure 2.

Sac domain structure of yeast Sac1. (A) The Sac phosphatase domain is comprised of two sub-domains: a novel N-terminal domain (blue) and the catalytic domain (yellow). The loop colored red represents the CX₅R motif. (B) Space filling model of the Sac domain with residues colored based on evolutionary conservation, which is calculated by an evolution trace method (Lichtarge et al., 1996). The least and highest conserved residues are colored from white to yellow respectively.




Figure 3. Disease mutations mapping on the modeled structure of the Sac domain of human Sac3/ Fig4. The Sac domain of human Sac3 is modeled based on the crystal structure of yeast Sac1 with the MODELER program (Martí-Renom et al., 2000). The local structural environment of six known CMT4J and ALS associated mutations are shown in zoomed-in panels.

Table 1

Sac domain enzymatic activities and associated phenotypes.

Protein name	<i>in vitro</i> substrate	Major <i>in vivo</i> substrate	Phenotype
Yeast Sac1	PI(3)P, PI(4)P, PI(5)P, PI(3,5)P ₂	PI(4)P	Cold-sensitive growth, actin reorganization, inositol auxotrophy, etc
Mammalian Sac1	PI(3)P, PI(4)P, PI(5)P, PI(3,5)P ₂	PI(4)P	Preimplantation lethality in mouse
<i>Aspergillus</i> spp.	N/A	N/A	N/A
Mammalian Sac2	PI(4,5)P ₂ , PI(3,4,5)P ₃	PI(3,4,5)P ₃ ?	Stress-induced cardiac hypertrophy
Yeast Fig4	PI(3,5)P ₂	PI(3,5)P ₂	Vacuole morphology defects
Mammalian Sac3	PI(3)P, PI(4)P, PI(3,5)P ₂	PI(3,5)P ₂	CMT4I, ALS, neurodegeneration in the pale tremor mouse and juvenile lethality
Synaptojanin 1	PI(4)P, PI(4,5)P ₂ , PI(3,4,5)P ₃	PI(4,5)P ₂ , PI(3,4,5)P ₃	Bipolar disorder, Down's syndrome, early postnatal lethality in mouse.
Synaptojanin 2	PI(4)P, PI(4,5)P ₂ , PI(3,4,5)P ₃	PI(4,5)P ₂	Hearing loss associated with cochlea hair loss
Sji1	PI(4,5)P ₂	PI(4,5)P ₂	Endocytic defects, increase sensitivity to temperature and neomycin
Sji2	PI(3)P, PI(4)P, PI(3,5)P ₂ , PI(4,5)P ₂	PI(4,5)P ₂	Endocytic defects, increase sensitivity to temperature and neomycin
Sji3	PI(3)P, PI(4)P, PI(3,5)P ₂ , PI(4,5)P ₂	PI(4,5)P ₂	Endocytic defects
AtSac1	PI(3,5)P ₂	PI(3,5)P ₂	Cell morphology defects
AtSac2	N/A	N/A	N/A
AtSac3	N/A	N/A	N/A
AtSac4	N/A	N/A	N/A
AtSac5	N/A	N/A	N/A
AtSac6	N/A	N/A	N/A

Protein name	<i>in vitro</i> substrate	Major <i>in vivo</i> substrate	Phenotype
AtSac7	PI(4)P, PI(3,5)P ₂	PI(4)P	Defective root hair and polarized growth
AtSac8	N/A	N/A	N/A
AtSac9	N/A	PI(4,5)P ₂ ?	Hypersensitive to stress