

Published in final edited form as:

*J Genet Couns.* 2014 December ; 23(6): 1034–1044. doi:10.1007/s10897-014-9732-5.

# An Internal Performance Assessment of CancerGene Connect: An Electronic Tool to Streamline, Measure and Improve the Genetic Counseling Process

Mary Pritzlaff<sup>1,2</sup>, Arielle Yorczyk<sup>1,2</sup>, Linda S Robinson<sup>1</sup>, Sara Pirzadeh-Miller<sup>1</sup>, Tirun Lin<sup>1</sup>,  
David Euhus<sup>3</sup>, and Theodora S Ross<sup>1,\*</sup>

<sup>1</sup>Department of Cancer Genetics, the University of Texas Southwestern Medical Center's Simmons Comprehensive Cancer Center, Dallas and Moncrief Cancer Institute, Fort Worth, Texas <sup>3</sup>Department of Surgery, Johns Hopkins Hospital, Baltimore, Maryland

## Abstract

CancerGene Connect (*CGC*) is a web-based program that combines the collection of family and medical history, cancer risk assessment, psychosocial assessment, report templates, a result tracking system, and a patient follow up system. The performance of *CGC* was assessed in several ways: pre-appointment completion data analyzed for demographic and health variables; a time study to assess overall time per case and to compare the data entry by the genetic counselor compared to the patient, and a measured quality assessment of the program via observation and interview of patients. Prior to their appointment, 52.3% of 2,414 patients completed the online patient questionnaire section of *CGC*. There were significant differences in completion rates among racial and ethnic groups. County hospital patients were less likely to complete the questionnaire than insured patients ( $p < 0.0001$ ); and likewise uninsured patients and patients with Medicare/Medicaid were less likely to complete the questionnaire than private patients ( $p < 0.0001$ ). The average genetic counseling time per case was 82 minutes, with no significant differences whether the counselor or the patient completed *CGC*. *CGC* reduces genetic counselor time by approximately 14%-46% compared to average time per case using traditional risk assessment and documentation methods previously reported. All surveyed users felt the questionnaire was easy to understand. *CGC* is an effective tool that streamlines workflow, and provides a standardized data collection tool that can be used to evaluate and improve the genetic counseling process.

## Keywords

CancerGene Connect; health care information technology; genetic counseling; hereditary cancer risk assessment; web-based program

\*Corresponding Author: Theodora Ross, UT Southwestern Medical Center, 5323 Harry Hines Blvd MC 8852, Dallas, Texas 75390-8852, P: 214 648-7026 F: 214 645-7027, theo.ross@utsouthwestern.edu.

<sup>2</sup>The authors contributed equally to this work.

Mary Pritzlaff, Arielle Yorczyk, Linda S Robinson, Sara Pirzadeh-Miller, Tirun Lin, David Euhus and Theodora Ross, declare that they have no conflict of interest.

## Introduction

Due to recent government initiatives and the increasing demands on healthcare practitioners, the field of healthcare information technology (IT) is rapidly expanding. The Health IT for Economic and Clinical Health (HITECH) Act was enacted in 2009 to promote the adoption and meaningful use of health IT and to incentivize the use of IT in medical practice. While health IT is rapidly expanding as an industry, its application to cancer genetics risk assessment is more limited. In 1998, we developed one of the first electronic cancer risk assessment programs; CancerGene.

Across health care professions, electronic programs have been used to improve efficiency (Chaudhry et al., 2006; Euhus, 2001; Lepanto et al., 2006) and have been shown to decrease face-to-face time spent with patients in a genetic counseling setting (S. A. Cohen & McIlvried, 2011). The incorporation of a computer program to assist with the genetic counseling process would likely allow more time for patient focused care during the genetic counseling process (Euhus, 2001).

One of the most time-consuming components of the genetic counseling process is family history collection. It is also one of the most crucial components of a genetic counseling session, with specific guidelines published by the American Society of Clinical Oncology (Lu et al., 2014), the Commission on Cancer (COC, 2012), and the National Accreditation Program for Breast Cancers (NAPBC, 2012). Additionally, even though eliciting a family history is crucial, its accuracy cannot be assured. Multiple studies have shown that patient recall of their family history is often inadequate (Dominguez et al., 2007; Gaff et al., 2004; Glanz et al., 1999; King et al., 2002; Mai et al., 2011; Murff et al., 2004; Ozanne et al., 2012; Pinsky et al., 2003; Quillin et al., 2006; Weinrich et al., 2002; Wideroff et al., 2010; Ziogas & Anton-Culver, 2003) and can vary by education level (Wideroff et al., 2010), income (Weinrich et al., 2002), race (Dominguez et al., 2007; Pinsky et al., 2003; Wideroff et al., 2010), gender (Glanz et al., 1999; Pinsky et al., 2003; Wideroff et al., 2010) and cancer site of origin (King et al., 2002; Mai et al., 2011; Murff et al., 2004; Ziogas & Anton-Culver, 2003). Asking patients to report their family history before the genetic counseling session provides an opportunity for the patient to confer with relatives and gather information, allowing for more comprehensive and accurate family history collection.

In addition to potentially improving the accuracy of the family history and decreasing the time it takes to collect it during the genetic counseling appointment, integration of an electronic family history data collection tool into genetic counseling clinics has the potential to decrease errors due to illegibility and incomplete data collection. There is a paucity of studies that explore this potential benefit within the genetic counseling setting. However, this benefit is documented in other areas of health care, with significant reductions in prescription errors (Shamliyan et al., 2008), legibility issues (Ali et al., 2010; Jani et al., 2008; Smith et al., 2009), and incomplete prescriptions (Ali et al., 2010; Jani et al., 2008). Similar benefits would be expected when electronic systems are applied to family history collection.

The majority of the electronic tools created to assist in the genetic counseling process center around the collection of family history. Electronic family history collection, versus handwritten notes or lists, is beneficial because an electronic tool can automatically organize this data for patients and clinicians, often before the appointment. Patient acceptance of electronic family history collection has been previously reported (Westman et al., 2000). Researchers have also compared family histories collected by electronic tool versus those collected by a genetic counselor and found the majority of affected family members were reported on both pedigrees (Acheson et al., 2006). And when there were discrepancies, it was more common for cancer to be recorded by their electronic tool only, versus reported by the genetic counselor only (Acheson et al., 2006). Several family history collection tools that can be used to assist the genetic counseling process are displayed in Table I.

Despite the number of family history tools available, there are barriers to their use. Widmer et al. (2013) found that only 16% of the genetic counselors they surveyed had ever had access to an electronic version of their patients' family histories, and a smaller percentage had patient populations who consistently had access. Barriers to use include concern over ease of use and the inability to modify family history with additional information collected during the session (Widmer et al., 2013). Also, many family history collection tools do not allow both patient and provider to input and modify family history information within the electronic system, or collect medical history, which is an important component of the cancer risk assessment.

Furthermore, the majority of these tools are unable to automatically generate mutation risk estimates. In addition to entering the patient information into an electronic system to document the encounter, the provider would need to re-enter the patient's medical and family history information into a computer risk program, such as CancerGene to generate a risk estimate.

The impetus for developing *CGC* was to expand the use of technology within the genetic counseling process beyond the collection of family history to include streamlining cancer risk assessment and clinic documentation. The limitations and barriers to the use of other electronic tools for genetic counseling were addressed by the model for *CGC*. The goal was to create an all-encompassing program where the patient's targeted medical and family history could be entered one time and then used to generate risk assessment reports, track results, and manage patient follow-up. As a result, patient care would be improved by obtaining more complete and more accurate intake information, as well as facilitating clinical research. Genetic counseling throughput could be maximized by allowing the patient to enter their own medical and family history prior to the appointment, improving the accuracy and completeness of information collected from the patient and enhancing efficacy and efficiency of genetic counseling sessions. As a result, patient care would be improved by allowing more time for addressing, rather than assessing, patient's risk, and future research would be improved by standardized data collection.

The purpose of this research was to determine whether *CGC* helps achieve these goals and streamline the cancer genetic counseling session. Here, three aspects of *CGC* were evaluated (Figure 1). First, completion rates were assessed to determine whether demographic and

health status variables were associated with completion of the patient-accessible portion of *CGC*. Second, we asked whether *CGC* was able to improve genetic counseling workflow by comparing time spent per patient when *CGC* was used versus traditional genetic counseling processes without *CGC*. Finally, we discuss measures taken to determine whether the questions within the patient-accessible portion of *CGC* were easy to understand, and whether patient responses to the questions in *CGC* were valid.

## Methods

### CGC Completion Rates and Demographics

**Participants**—The analyses included all patients entered in the *CGC* database between January 1, 2010 and June 30, 2013, ( $N=5,157$ ) seen at our four primary sites: Simmons Cancer Center (SCC); Moncrief Cancer Institute (MCI); John Peter Smith Hospital (JPS); and Parkland Memorial Hospital (PMH). Data regarding patient completion were collected beginning May 1, 2011. However, the Spanish version of the online patient questionnaire portion of *CGC* was not available until March 2013, and patients seen at our county safety net clinics (JPS and PMH) (30% of our patients) were not asked to enter their own data until that time. Therefore, patient completion of the online *CGC* questionnaire was evaluated from May 2011 to June 2013 at our two primary clinics (SCC and MCI). This included a total of 2,414 patients. For comparison between clinic populations and testing funding source, completion rates were separately analyzed from April 2013 to June 2013 for all four clinics, after the Spanish version of the online questionnaire became available, and when the online questionnaire was routinely offered in all clinics. This analysis included 499 patients. Of these patients, 153 patients did not have genetic testing, and were not included in the analysis of completion by testing funding source. IRB approval was obtained (IRB #STU 072013-051- Evaluation of Utilization of CancerGene Connect and Genetics Navigation in a Clinical Cancer Risk Assessment Program), and an informed consent waiver for this retrospective data review was granted.

**Data Analysis**—The pre-appointment patient completion rates for *CGC* were analyzed by several demographic and health status categories (gender, age, race, ethnicity, cancer status, and genetic test status) to assess if these categories affect the likelihood that the patient would complete *CGC*. The association between patient completion and demographic and health status variables were analyzed using the Fisher's Exact Test or 2-tailed Chi-square tests for independence using GraphPad InStat software. In this study, we did not adjust for multiple comparisons.

### Effects of CGC use on Genetic Counseling Workflow

**Procedures**—A time study was conducted from May 2011 to August 2011 to analyze the genetic counselor's time in two scenarios: when the patient completed the CancerGene Connect program prior to their appointment, and when the genetic counselor had to do the data entry themselves. A time tracking form was developed to capture time spent on various genetic counseling tasks, including pre-appointment chart review, time in clinic, documentation of clinic visit, follow up activities, calling results, and documenting and sending results. The genetic counselor used this form to self-report the time they spent on

each of these tasks as they worked through each case. The genetic counselor also documented the reasons why patients did not complete the online patient questionnaire portion of *CGC* on this tracking form.

**Participants**—Five genetic counselors were each asked to prospectively record the time spent on each task for the first 10 patients that completed the online *CGC* questionnaire prior to their appointment and for the first 10 patients that did not from the study initiation. Data were collected on a total of 89 patients; 39 patients in this cohort completed the online *CGC* questionnaire prior to their appointment, and 50 did not.

**Data Analysis**—Time spent on various counseling tasks was compared based on patient completion of the online questionnaire portion of *CGC*. The data were analyzed using the 2-tailed unpaired t-test with Welsch correction.

### Quality Assurance Testing of Patient Comprehension of *CGC*

**Procedures**—In 2012, the online questionnaire section of *CGC* was translated to Spanish. UT Southwestern's Clinical Sciences team then performed cognitive interviews on over 50 patients (both English and Spanish speaking) to ensure that questions were understandable and responses are valid in both languages. Cognitive interviewing is a type of semi-structured interview intended to refine survey instruments and behavioral interventions (Weidmar et al., 1999; Hurtado et al., 2005; Willis, 2004 ). The cognitive interviews consisted of various questions about items included in the computer program as well as the formatting and usability of the program. We assessed whether participants understood what the questions in the computer program were asking and what the various response options meant. Prior to this evaluation, we determined the completion rate for each question to determine questions that were not being answered by the participants. Though some sections of the interface had 100% completion rates for those accessing the system, other sections, such as the physical activity section, had completion rates of only 35%. Qualitative data from the cognitive interviews were analyzed through techniques outlined by Willis, 2004, Weidmar 1999, and Hurtado 2005. Interviews were transcribed and a code book was developed. For each survey item, a list of codes relevant to the cognitive processing of the question was developed. Codes included: respondent has difficulty understanding the question (comprehension), respondent has difficulty remembering the question (recall), and respondent has difficulty understanding the meaning of particular words or concepts. Coders applied the codebook to each survey question using study notes, audiotapes and transcripts. Coding discrepancies were identified and brought back to the language validation team for resolution. After all discrepancies were worked out, appropriate changes were incorporated to the final versions which were then used for pilot testing. These participants were also administered a separate questionnaire with Likert-style and open ended questions to assess their reaction to the program (IRB #STU 102011-028 CancerGene Connect Study funded by CPRIT, Cancer Prevention Research Institute of Texas).

**Participants**—Altogether, 50 users underwent cognitive interviews; 32 users underwent interviews to assess the initial format, and 18 users (11 English, 7 Spanish) were interviewed to assess the revised format. These participants were recruited from our PMH general cancer

genetics clinic. All English and Spanish speaking patients being scheduled in this clinic between the ages of 18 and 64 were invited to participate in this study, and were enrolled until the target enrollment was reached. Informed consent was obtained from study participants after the nature of the procedure was fully explained to them.

### Description of CGC

*CGC* was developed by the UT Southwestern Cancer Genetics team under the direction of Dr. David Euhus and Linda Robinson, MS, CGC. It was first piloted in late 2009 at Moncrief Cancer Institute (UT Southwestern-affiliated) and UT Southwestern Simmons Cancer Center.

*CGC* was designed as a secure, password-protected web-based application. When patients are scheduled for a genetics appointment, they are provided with a secure ID and password and a link to the *CGC* website that allows them to complete the online questionnaire from their home prior to appointment. The online, patient-accessible questionnaire consists of over 130 targeted questions that cover the topics of: patient demographics; referring physician information; obstetric/gynecologic (Ob/Gyn) history; hormonal and birth control history; surgical history (targeted for breast and gynecological surgery); colon history (including surveillance, surgery, and risk factors); general health; social history (including alcohol, tobacco use and exercise habits); psychological screens (discussed below); and family history.

The online questionnaire is designed with branching logic, such that patients are only prompted to complete applicable information. For example, males are not asked about gynecological, hormonal, or birth control use history; patients who have not had a breast biopsy are not asked for biopsy results, etc. By patient report and observation, it typically takes the patient 20 minutes to complete this online questionnaire, but length of time may vary. The website is hosted by UT Southwestern server using HTTPS secure encryption. Data are encrypted and transferred to a secure UT Southwestern SQL server where it is stored.

All data input by the patient is immediately accessible by the genetic counselor, and does not need to be printed by the patient. A pedigree is automatically generated. This, and the patient's targeted medical history, can be viewed by patient and provider. At this time, the program is not integrated directly into the medical record. The data within the program can be printed and scanned into the patient's electronic medical record or copied and pasted.

*CGC* incorporates a number of previously validated cancer risk assessment models, including the Gail, Claus, BRCAPRO, MMRpro, PancPro, and MelaPro models (Chen et al., 2006). Links to the PENN II, LAMBDA, Tyrer-Cuzick, BODICEA, and PTEN models, as well as the Myriad tables, MMR predict and NCI colon cancer risk model are also included on the provider interface of *CGC*. Once the patient has completed the online questionnaire, the provider can review patient responses, run the risk assessment models, print the pedigree and patient summary, and document patient encounters. The provider has the ability to update and edit any of the information that the patient has entered. The provider also can complete additional fields that are only accessible to the provider, such as head



circumference, maternal and paternal ancestry, consanguinity, and genetic testing status (e.g. pending, negative, variant of uncertain significance, etc). All fields can be updated by the provider (e.g. when test result status changes from “pending” to “negative”).

*CGC* also includes previously validated psychological scales that assess depression, social support and post-traumatic stress (CES-D, ISEL, IES-R, respectively) (S. Cohen & Hoberman, 1983; Horowitz et al., 1979; Radloff, 1977). These scales use Likert-style questions to provide a raw score for each of these categories. *CGC* reports both the raw score and a qualitative “yes/no” indicator that the patient’s responses suggest that they have one or more of these psychosocial conditions. Psychological scales allow for standardization of the psychosocial assessment process. Standardization of psychosocial assessment is consistent with the 2012 American College of Surgeons Commission on Cancer Program Standard 2.3, mandating psychosocial assessment as part of the hereditary cancer risk assessment process (Greene, 2013). The provider can view a summary of the patient’s scores and can incorporate recommendations for psychological support and/or document referrals for psychological services into the risk assessment report using templates. *CGC* also includes a follow-up screen in which the provider can document prophylactic surgeries, surveillance, chemoprevention, and psychosocial support interventions for the patient. This feature allows the provider to create reminders for patient surveillance, or otherwise follow patients to ensure appropriate management.

Several report templates are built into the provider interface of *CGC*, including risk assessment reports, family letters, and positive, negative, and variant of uncertain significance result reports. The reports include drop-down menus with editable text. The positive results templates incorporate management recommendations based on literature review and/or National Comprehensive Cancer Network (NCCN) guidelines, which are cited within the templates’ text. The templates are revised as new versions of NCCN guidelines and/or new literature becomes available.

After piloting the program in late 2009, the online patient questionnaire portion of *CGC* was updated based on patient and genetic counselor feedback, and the UT Southwestern Cancer Genetics group began entering all patients into *CGC* in January 2010. The program has been regularly updated since its creation. Updates include the creation of additional fields to capture method of test payment and other clinical/administrative data, the incorporation of updates to national management guidelines, and the addition of test menus and patient fact sheets.

Beyond its clinical application, *CGC* is a powerful administrative, patient care, and research tool. Every data point collected within the patient and provider interface is stored within a secure database. This information can be used to analyze patient volumes, referral patterns, and result statistics. The information can also be used for patient care, to generate lists of patients with pending test results or lists of patients positive for a given gene mutation. Finally, the database can facilitate research, by providing a method to identify health disparities or identify patients that might meet criteria for a given protocol. At UT Southwestern, the *CGC* database has been linked with the Informatics for Integrating Biology and the Bedside program, a tool that allows database queries for researchers.

## RESULTS

### CGC Completion Rates and Demographics

Data collection of completion rates began May 1, 2011 and ended June 30, 2013. During that time, completion rates based on demographic variables were assessed at our main clinic sites (Simmons Cancer Center and Moncrief Cancer Institute), where patients were routinely offered the online *CGC* questionnaire prior to their appointment. A total of 52.3% of patients in this cohort completed the questionnaire. Thus 47.7% either did not start the questionnaire or did not complete enough of it such that the genetic counselor needed to go back into *CGC* and complete it themselves. For this cohort, there were statistically significant differences in completion rates based on age ( $p=0.026$ ); 30.4% of individuals younger than 18 years completed the questionnaire, versus 55.1% of individuals aged 18–29 years, 54.2% aged 30–49 years, 52.7% aged 50–64 years, and 46.6% aged 65 and older. Post-hoc comparison indicates that patients ages 18–64 had a higher completion rate compared to patients under 18 and above 65, although patients under 18 had a much lower completion rate ( $p=0.0052$ ). The differences in completion rates among racial groups were statistically significant ( $p<0.0001$ ), and are as follows; American Indian/Native Alaskans 53.3%, Asians 52.6%, African Americans 37.0%, Native Hawaiian/Pacific Islanders 8.3%, and White/Caucasians 55.2%. Post-hoc comparison indicates that completion rates were significantly less among African Americans and Hawaiian/Pacific Islanders compared with the other racial groups ( $p<0.0001$ ). Individuals of reported Hispanic ethnicity (35.5%) were less likely to complete the questionnaire than individuals who did not report Hispanic ethnicity (55.9%) ( $p<0.0001$ ). The differences in completion rates by gender and cancer status were not statistically significant.

For comparison, completion rates for our main clinics and county hospital clinics were assessed from April 1, 2013 to June 30, 2013, when the online questionnaire was routinely offered at all sites. Patients seen at our county hospital clinics (21.5%) were less likely to complete the questionnaire than patients seen in our main clinics (SCC 47.3%; MCI 56.7%) ( $p<0.0001$ ). Significant differences were observed in completion rates among patients without insurance, patients with Medicare/Medicaid, and patients with private insurance ( $p<0.0001$ ). Patients without insurance (27.0%) were significantly less likely to complete *CGC* compared to patients with private insurance (50.0%) ( $p<0.0001$ ); as were patients with Medicare/Medicaid (29.6%) ( $p=0.015$ ). See Table II for more detail and information on non-statistically significant associations.

Reasons for non-completion were not collected during the entire time frame of the study. However, reasons for non-completion were documented by the genetic counselors from 43 patients seen between May 2011 and August 2011 (prior to the implementation of the Spanish version of the program). The reasons given for non-completion were either because the patient: was not offered the questionnaire at their clinic site ( $n=11$ ); was seen as a walk-in ( $n=10$ ); did not have internet access ( $n=8$ ); did not speak English ( $n=7$ ); had technical difficulties using the program ( $n=2$ ); forgot their password ( $n=2$ ); forgot to complete the questionnaire ( $n=2$ ); and declined to complete the questionnaire ( $n=1$ ). If patients had not



done so before arriving to clinic, they no longer had the opportunity to complete the online questionnaire.

### Effects of CGC use on Genetic Counseling Workflow

The average genetic counseling time per case using *CGC* was 82 minutes. Completion of the online questionnaire portion of *CGC* prior to the appointment did not result in statistically significant reductions (or increases) in time spent on any of the genetic counseling tasks that were measured. Anecdotally, genetic counselors reported that while the same amount of time was spent with completers vs. non-completers, they were able to spend more time in the session *addressing* risk vs. *assessing* risk with the completers, giving the genetic counselors a greater feeling of efficacy with these patients.

We also used the time study to compare the total time a genetic counselor spends on a cancer genetics case from start to finish using *CGC* with the historical time commitment for a case. Prior to *CGC*, the genetic counselor would enter the pedigree obtained in the session into CancerGene, draw the pedigree in Progeny and write the follow up documentation in another system. At our institution, the estimated average time spent per case was approximately 2.5-3 hours prior to the implementation of *CGC*, which is similar to genetic counseling times previously reported in the literature (reports vary from 85 to 150 min/case) (McPherson et al., 2008; Uhlman et al., 2011; Wham et al., 2010; Cohen, McIvried. 2011). Compared to genetic counseling times reported in the literature, use of *CGC* reduced the time spent per case by approximately 14% to 46%. At our institution, the implementation of *CGC* reduced the overall time spent on cancer genetics cases by half compared with estimated time spent per case prior to implementation of the program.

### Quality Assurance Testing of Patient Comprehension of CGC

Seventy two percent of users strongly agreed that the program was easy to use and 100% agreed or strongly agreed that the questions were easy to understand. In response to open ended questions asked about their satisfaction with the program, users commented that they enjoyed creating their family history, they liked being asked about how they feel (referring to the psychological assessment), and that the program was easy/simple/fast to use. Patient assessment of the program is summarized in Table III. Based on this pilot study, an updated “2.0” version of *CGC* was implemented in March 2013 where over 1,500 patients have been entered.

## DISCUSSION

### Major Findings and Practice Implications

*CGC* is a web-based application that saves time, provides patients and providers the ability to continually update family history information, performs hereditary risk and cancer risk calculations, and includes modifiable macro-generated reports. *CGC* is a tool that has the potential to provide benefit to all components of the genetic counseling session including medical and family history collection, psychosocial assessment, reporting of test results to patient and provider, and documentation of follow-up procedures.

The workflow analysis showed that this program reduces the time spent per case from 14% to 46% compared to previously reported genetic counseling times. In short, *CGC* saves genetic counselors time regardless of whether or not the patient completes the online questionnaire section prior to the session. With implementation of *CGC*, there is greater opportunity to reduce redundancy in clinic workflows because medical and family history information only needs to be entered once in order to be documented and to generate risk estimates and pedigrees.

*CGC* also enhances the quality of the genetic counseling encounter when patients complete the questionnaire before the appointment. By having the information before the appointment, the provider is given a better idea of what needs to be addressed in the session before they see the patient. Thus, when the provider is aware of patients' individualized mutation risks, cancer risks, and psychosocial profile before the session, they can spend less time collecting information during the session, and more time establishing rapport and addressing individualized concerns. This finding is consistent with a previously predicted benefit following integration of a computer program to assist the genetic counseling process (Euhus, 2001).

*CGC* also includes a validated psychosocial assessment tool to accommodate NCCN guidelines (not evaluated here). This is used to standardize the psychosocial assessment of cancer genetics patients. This standardization should help eliminate counselor variability in identifying psychosocial issues. Incorporating a psychosocial assessment into a pre-visit questionnaire provides the genetic counselor with the opportunity to easily identify which patients may require additional resources, support, or referrals to psychological services prior to the session. Based on psychological assessment scores, UT Southwestern genetic counselors have identified and documented the need for additional support and resources for over 600 patients, and have identified psychosocial disparities within our clinic population. Patients are routinely referred to psychological services at our institution based upon this needs assessment.

In addition to the benefits listed above, *CGC* has transformed the administrative tasks of compiling clinic volumes and tracking test results. This information is regularly requested by hospital administration to evaluate productivity, as well as for department administrative purposes such as evaluating clinic workload, referral patterns and need for additional resources; utilization for clinical purposes includes following up with positive patients. Previously, these data were captured by an administrative assistant entering every patient encounter into a database, recording the clinic site, visit type, referring physician, patient demographic information, genetic counselor, and test results. *CGC* has eliminated this process at UT Southwestern, as all of this information is gathered in the *CGC* database, and this information can be quickly retrieved via database query.

The modifiable macros and templates make it possible to incorporate ever-evolving updates of the national management guidelines, thereby preventing outdated patient information from being a barrier to use. *CGC* capitalizes on its ability to use health IT to improve accuracy, efficiency, and standardization of the genetic counseling process, and provides a

standardized data collection platform from which evidence based outcomes of genetic counseling can be measured.

We found that patients from our safety net clinics were less likely to complete the online questionnaire section of *CGC* than our private patients, and patients with private insurance were more likely to complete the online questionnaire than patients with Medicare, Medicaid, or no insurance. There may be several reasons for this, including differences in scheduling personnel for different clinics, patient access to the internet, computer literacy, and psychosocial factors. Computer illiteracy (Hilton et al., 2012; Simon et al., 2008) and the inability to access the internet at home (Simon et al., 2008) are associated with inability to engage in web-based health information. However, other research has shown that patients living in either suburban or urban areas were equally likely to access the internet, even though suburban residents were more likely to own a computer and urban residents had a higher odds of being uninsured (Bond et al., 2012). Household income cannot always predict internet use (Appleby-Tagoe et al., 2012; Bond et al., 2012; Simon et al., 2008). Research has shown that mental health issues and substance use have not been barriers to online health information either (Hilton et al., 2012).

Due to the limited, but undeniable, number of individuals who are unable to access the internet, a non-electronic option should always be available (Simon et al., 2008), even though, research has shown that race, age, education, personal history of cancer, or family history of cancer does not affect patients' *desire* to use the internet (Simon et al., 2008). A paper version of the *CGC* questionnaire is available to our patients. Completion rates and demographic information of those who filled out the paper questionnaire, versus the online version, were not collected, however.

There are several topics that should be discussed with the patient when she/he is asked to complete the online family history data collection tool. It is important to explain to patients that they will have the opportunity to clarify any questions regarding their family history when they meet with the genetic counselor. The belief that patients will not have the opportunity to discuss their family history directly with health care providers is a reason some patients would not fill out an electronic family history tool (Simon et al., 2008). Privacy concerns should also be discussed, as the potential for additional loss of confidentiality has been established as a barrier to electronic family history intake (Simon et al., 2008). Furthermore, patients should be made aware of the importance of genetic counseling and that gathering a complete family history will allow them to maximize their benefit from the genetic counseling process (Appleby-Tagoe et al., 2012).

*CGC* does not obviate the need for review of family history during the genetic counseling session. For example, genetic counselors can clarify whether cancer represents a new primary versus a metastasis site (Acheson et al., 2006). Additional questions used to clarify family history can help to improve accuracy (Kadan-Lottick et al., 2003), including information about third degree relatives, which is not collected by *CGC*. Pathology should be used to verify cancer reports whenever possible, particularly when there is significant overrepresentation of malignancy within a family history (Parent et al., 1997).

## Study Limitations

*CGC* is specific to the hereditary cancer genetic counseling process, and is not applicable for use in prenatal, pediatric, or other genetic counseling subspecialties. This study is limited by the fact that it was conducted within one cancer genetics program, and results may vary at other institutions. Assessment of patient completion rates is limited by the relatively short time that the Spanish translation of the questionnaire was available, and the relatively short time that the questionnaire was routinely offered to at all clinic sites. The statistical analysis is limited in that corrections for multiple comparisons were not performed. The qualitative assessment of the program is limited by the small number of participants that were included.

## Research Recommendations

*CGC* can be further enhanced to increase the efficiency of the genetic counselor as well as provide patients with self-care tools. For example, we plan to target specific cancer risk factors, such as alcohol use, smoking and obesity, and offer the patient resources as they complete the online questionnaire portion of *CGC* based on their responses, such as links to alcohol and smoking cessation tools or weight management programs. We will continue to evaluate patient completion rates and analyze the outcomes of interventions to improve these, such as adding tablets in our waiting rooms for patients to complete the questionnaire prior to their appointment. We will continue to evaluate disparities in psychological assessment scores and develop interventions. We are currently developing a follow-up tool to generate automated reminders to our BRCA and Lynch positive patients for recommended screening tests. To measure the effect of cancer genetic testing and the role of genetic counseling in the reduction of cancer incidence, we are using the data obtained from our *CGC* program to model the uptake of prophylactic surgeries and compliance with NCCN management guidelines to determine the impact on the long term cancer incidence. A systematic data collection system is needed to measure genetic counseling outcomes. As the demand for services continues to increase, a technological tool will be needed to address inefficiencies. CancerGene Connect is a tool that can meet this need.

## Acknowledgments

This work was supported by the Cancer Prevention and Research Institute of Texas (grant #PP110220). The authors thank the support of the entire staff of the UT Southwestern Cancer Genetics group, Parkland Hospital Oncology and John Peter Smith Hospital, as well as the UT Southwestern Clinical Sciences Department, including Dr. Celette Skinner, Saddyna Belmashkan, and Maria Funes.

All procedures followed were in accordance with the ethical standards of the responsible committee on human experimentation (institutional and national) and with the Helsinki Declaration of 1975, as revised in 2000. An informed consent waiver was obtained from the Institutional Review Board for all patients that were included in the study.

## References


- Acheson LS, Zyzanski SJ, Stange KC, Deptowicz A, Wiesner GL. Validation of a self-administered, computerized tool for collecting and displaying the family history of cancer. *J Clin Oncol*. 2006; 24(34):5395–5402. [PubMed: 17088568]
- Ali J, Barrow L, Vuylsteke A. The impact of computerised physician order entry on prescribing practices in a cardiothoracic intensive care unit. *Anaesthesia*. 2010; 65(2):119–123. [PubMed: 19889112]

- Appleby-Tagoe JH, Foulkes WD, Palma L. Reading between the lines: a comparison of responders and non-responders to a family history questionnaire and implications for cancer genetic counselling. *J Genet Couns.* 2012; 21(2):273–291. [PubMed: 21845493]
- Bond MC, Klemm R, Merlis J, Kopinski JE, Hirshon JM. Computer access and Internet use by urban and suburban emergency department customers. *J Emerg Med.* 2012; 43(1):159–165. [PubMed: 22142670]
- Britten N. Qualitative Research: Qualitative interviews in medical research. *BMJ.* 1995; 12(6999): 251–253. doi: <http://dx.doi.org/10.1136/bmj.311.6999.251>. [PubMed: 7627048]
- Chaudhry B, Wang J, Wu S, Maglione M, Mojica W, Roth E, et al. Systematic review: impact of health information technology on quality, efficiency, and costs of medical care. *Ann Intern Med.* 2006; 144(10):742–752. [PubMed: 16702590]
- Chen S, Wang W, Lee S, et al. PRediction of germline mutations and cancer risk in the lynch syndrome. *JAMA.* 2006; 296(12):1479–1487. [PubMed: 17003396]
- Cohen SA, McIlvried DE. Impact of computer-assisted data collection, evaluation and management on the cancer genetic counselor's time providing patient care. *Fam Cancer.* 2011; 10(2):381–389. [PubMed: 21240560]
- Cohen, Sheldon; Hoberman, Harry M. Positive Events and Social Supports as Buffers of Life Change Stress1. *Journal of applied social psychology.* 1983; 13(2):99–125.
- Commission on Cancer (COC). Cancer Program Standards 2012: Ensuing Patient-Centered Care. 2012. p. 140 <http://www.facs.org/cancer/coc/programstandards2012.pdf>
- Dominguez FJ, Lawrence C, Halpern EF, Drohan B, Grinstein G, Black DM, et al. Accuracy of self-reported personal history of cancer in an outpatient breast center. *J Genet Couns.* 2007; 16(3):341–345. [PubMed: 17508275]
- Euhus DM. Understanding mathematical models for breast cancer risk assessment and counseling. *Breast Journal.* 2001; 7(4):224–232. [PubMed: 11678799]
- Gaff CL, Aragona C, MacInnis RJ, Cowan R, Payne C, Giles GG, et al. Accuracy and completeness in reporting family history of prostate cancer by unaffected men. *Urology.* 2004; 63(6):1111–1116. [PubMed: 15183962]
- Glanz K, Grove J, Le Marchand L, Gotay C. Underreporting of family history of colon cancer: correlates and implications. *Cancer Epidemiol Biomarkers Prev.* 1999; 8(7):635–639. [PubMed: 10428202]
- Greene, FL. Cancer Program Standards 2012: Ensuring Patient-Centered Care, Version 1/2. Commission on Cancer. 2013. Retrieved from Cancer Program Standards 2012: Ensuring Patient-Centered Care, Version 1.2 website: <http://www.facs.org/cancer/coc/programstandards2012.pdf>
- Hilton JF, Barkoff L, Chang O, Halperin L, Ratanawongsa N, Sarkar U, et al. A cross-sectional study of barriers to personal health record use among patients attending a safety-net clinic. *PLoS One.* 2012; 7(2):e31888. [PubMed: 22363761]
- Horowitz M, Wilner N, Alvarez W. Impact of Event Scale: a measure of subjective stress. *Psychosomatic Medicine.* 1979; 41(3):209–218. [PubMed: 472086]
- Hurtado MP, Angeles J, Blahut SA, Hays RD. Assessment of the equivalence of the Spanish and English versions of the CAHPS Hospital Survey on the quality of inpatient care. *Health Serv Res.* 2005; 40:2140–2161. [PubMed: 16316442]
- Jani YH, Ghaleb MA, Marks SD, Cope J, Barber N, Wong IC. Electronic prescribing reduced prescribing errors in a pediatric renal outpatient clinic. *J Pediatr.* 2008; 152(2):214–218. [PubMed: 18206691]
- Kadan-Lottick NS, Friedman DL, Mertens AC, Whitton JA, Yasui Y, Strong LC, et al. Self-reported family history of cancer: the utility of probing questions. *Epidemiology.* 2003; 14(6):737–740. [PubMed: 14569191]
- King TM, Tong L, Pack RJ, Spencer C, Amos CI. Accuracy of family history of cancer as reported by men with prostate cancer. *Urology.* 2002; 59(4):546–550. [PubMed: 11927311]
- Lepanto L, Pare G, Gauvin A. Impact of PACS deployment strategy on dictation turnaround time of chest radiographs. *Acad Radiol.* 2006; 13(4):447–452. [PubMed: 16554224]


- Lu KH, Wood ME, Daniels M, Burke C, Ford J, Kauff ND, et al. American Society of Clinical Oncology expert statement: collection and use of a cancer family history for oncology providers). *J Clin Oncol*. 2014; 32(8):833–40. [PubMed: 24493721]
- Mai PL, Garceau AO, Graubard BI, Dunn M, McNeel TS, Gonsalves L, et al. Confirmation of family cancer history reported in a population-based survey. *J Natl Cancer Inst*. 2011; 103(10):788–797. [PubMed: 21562245]
- McPherson E, Zaleski C, Benishek K, McCarty CA, Giampietro PF, Reynolds K, Rasmussen K. Clinical genetics provider real-time workflow study. *Genetics in Medicine*. 2008; 10(9):699–706. [PubMed: 18978682]
- Murff HJ, Spigel DR, Syngal S. Does this patient have a family history of cancer? An evidence-based analysis of the accuracy of family cancer history. *JAMA*. 2004; 292(12):1480–1489. [PubMed: 15383520]
- National Accreditation Program for Breast Centers (NAPBC). 2012 Breast Cancer Standards Manual. 2012. p. 72 <http://napbc-breast.org/standards/2012standardsmanual.pdf>
- Ozanne EM, O'Connell A, Bouzan C, Bosinoff P, Rourke T, Dowd D, et al. Bias in the reporting of family history: implications for clinical care. *J Genet Couns*. 2012; 21(4):547–556. [PubMed: 22237666]
- Parent ME, Ghadirian P, Lacroix A, Perret C. The reliability of recollections of family history: implications for the medical provider. *J Cancer Educ*. 1997; 12(2):114–120. [PubMed: 9229275]
- Pinsky PF, Kramer BS, Reding D, Buys S. Reported family history of cancer in the prostate, lung, colorectal, and ovarian cancer screening trial. *Am J Epidemiol*. 2003; 157(9):792–799. [PubMed: 12727673]
- Quillin JM, Ramakrishnan V, Borzelleca J, Bodurtha J, Bowen D, Baer Wilson D. Paternal relatives and family history of breast cancer. *Am J Prev Med*. 2006; 31(3):265–268. [PubMed: 16905040]
- Radloff, Lenore Sawyer. The CES-D scale a self-report depression scale for research in the general population. *Applied psychological measurement*. 1977; 1(3):385–401.
- Resource document for curriculum development in cancer genetics education. American Society of Clinical Oncology. *J Clin Oncol*. 1997; 15(5):2157–2169. [PubMed: 9164231]
- Shamliyan TA, Duval S, Du J, Kane RL. Just what the doctor ordered. Review of the evidence of the impact of computerized physician order entry system on medication errors. *Health Serv Res*. 2008; 43(1 Pt 1):32–53. [PubMed: 18211517]
- Simon C, Acheson L, Burant C, Gerson N, Schramm S, Lewis S, et al. Patient interest in recording family histories of cancer via the Internet. *Genet Med*. 2008; 10(12):895–902. [PubMed: 19092442]
- Smith M, Dang D, Lee J. E-prescribing: clinical implications for patients with diabetes. *J Diabetes Sci Technol*. 2009; 3(5):1215–1218. [PubMed: 20144439]
- Uhlmann, WR.; Schuette, JL.; Yashar, B. A guide to genetic counseling. 2011. Wiley.com
- Weinrich SP, Faison-Smith L, Hudson-Priest J, Royal C, Powell I. Stability of self-reported family history of prostate cancer among African American men. *J Nurs Meas*. 2002; 10(1):39–46. [PubMed: 12048968]
- Westman J, Hampel H, Bradley T. Efficacy of a touchscreen computer based family cancer history questionnaire and subsequent cancer risk assessment. *J Med Genet*. 2000; 37(5):354–360. [PubMed: 10807694]
- Wham D, Vu T, Chan-Smutko G, Kobelka C, Urbauer D, Heald B. Assessment of clinical practices among cancer genetic counselors. *Familial Cancer*. 2010; 9(3):459–468. [PubMed: 20140525]
- Wideroff L, Garceau AO, Greene MH, Dunn M, McNeel T, Mai P, et al. Coherence and completeness of population-based family cancer reports. *Cancer Epidemiol Biomarkers Prev*. 2010; 19(3):799–810. [PubMed: 20160272]
- Weidmar B, Brown J, Garcia L. Translating the CAHPS 1.0 Survey Instrument into Spanish. *Med Care*. 1999; 37(3 Suppl):MS89–MS96. [PubMed: 10098563]
- Widmer C, Deshazo JP, Bodurtha J, Quillin J, Creswick H. Genetic counselors' current use of personal health records-based family histories in genetic clinics and considerations for their future adoption. *J Genet Couns*. 2013; 22(3):384–392. [PubMed: 23242928]



- Willis, GB. Cognitive interviewing: A tool for improving questionnaire design. Thousand Oaks, CA: Sage Publications; 2004.
- Ziogas A, Anton-Culver H. Validation of family history data in cancer family registries. *Am J Prev Med.* 2003; 24(2):190–198. [PubMed: 12568826]



**Figure 1.**  
Overview of the Assessment of the CancerGene



**Figure 2.**  
Comparison of Time Spent on Genetic Counseling Tasks

**Table I**

## Electronic Family History Collection Tools

Name	Creator(s)	Website
<b>CancerGene</b> *	UT Southwestern Medical Center and The BayesMendel Group	<a href="http://www4.utsouthwestern.edu/breasthealth/cagene/cagene.asp">http://www4.utsouthwestern.edu/breasthealth/cagene/cagene.asp</a>
<b>CancerGene Connect</b>	UT Southwestern Cancer Genetics Department	<a href="https://www4.utsouthwestern.edu/breasthealth/cgc/default.asp">https://www4.utsouthwestern.edu/breasthealth/cgc/default.asp</a>
<b>Cyrillic</b> *	Cyrillic Software	<a href="http://www.cyrillicsoftware.com/">http://www.cyrillicsoftware.com/</a>
<b>Family Health Link</b>	Ohio State University	<a href="https://familyhealthlink.osumc.edu/Notice.aspx">https://familyhealthlink.osumc.edu/Notice.aspx</a>
<b>Family Healthware</b>	Center for Disease Control and Prevention (currently undergoing clinical trials)	<a href="http://www.cdc.gov/genomics/famhistory/famhx.htm">http://www.cdc.gov/genomics/famhistory/famhx.htm</a>
<b>Health Heritage</b>	University of Virginia Health System-Department of Public Health	<a href="http://www.healthheritage.net">www.healthheritage.net</a>
<b>Hughes Risk Apps</b> *	Consortium comprised of Massachusetts General Hospital, University of Massachusetts at Lowell, and the Newton-Wellesley Hospital and originally headed by Kevin S. Hughes, MD and Sherwood S. Hughes, MD	<a href="http://www.hughesriskapps.net">http://www.hughesriskapps.net</a>
<b>Medical Family Tree</b>	Monroe-Meyer Institute for Genetics and the Eppley Cancer Center of the University of Nebraska	<a href="http://app1.unmc.edu/gencancer">http://app1.unmc.edu/gencancer</a>
<b>My Family Health Portrait</b>	U.S. Surgeon General	<a href="https://familyhistory.hhs.gov/fhh-web/home.action">https://familyhistory.hhs.gov/fhh-web/home.action</a>
<b>My Generations</b>	NorthShore University Medical Center	<a href="http://www.northshore.org/genetics/mygenerations">http://www.northshore.org/genetics/mygenerations</a>
<b>Progeny</b> *	Progeny software, LLC	<a href="http://www.progenygenetics.com/clinical">http://www.progenygenetics.com/clinical</a>

\* indicates provider-only accessibility

**Table II**

Patient Completion Rates by Gender, Age, Race, Ethnicity, Cancer Status, Clinic Site, and Insurance Status

Total patients		Patients who completed <i>CGC</i> online prior to appointment	% who completed <i>CGC</i> online prior to appointment
<b>Gender: (<math>p = 0.21</math>)</b>			
Male	282	158	56.0%
Female	2132	1106	51.9%
<b>Age: (<math>p = 0.026</math>)</b>			
Under 18	23	7	30.4%
18-29	158	87	55.1%
30-49	913	495	54.2%
50-64	922	486	52.7%
65 and Older	367	171	46.6%
<b>Race: (<math>p &lt; 0.0001</math>)</b>			
American Indian/ Native Alaskan	15	8	53.3%
Asian	78	41	52.6%
Black/ African American	332	123	37.0%
Native Hawaiian/ Pacific Islander	12	1	8.3%
White/ Caucasian	1977	1091	55.2%
<b>Ethnicity: (data taken prior to implementation of Spanish version) (<math>p &lt; 0.0001</math>)</b>			
Hispanic	417	148	35.5%
Non-Hispanic	1996	1115	55.9%
<b>Cancer Status: (<math>p = 0.21</math>)</b>			
Patient has Current or Previous History of Cancer	1495	767	51.3%
Patient has Not had Cancer	918	496	54.0%
<b>Clinic (4/1/13-6/30/13): (<math>p &lt; 0.0001</math>)</b>			
Simmons Comprehensive Cancer Center (Dallas)	186	88	47.3%
Moncrief Cancer Institute (Fort Worth)	127	72	56.7%
County Hospitals (John Peter Smith and Parkland)	186	40	21.5%
<b>Source of Funding for Genetic Testing (4/1/13-6/30/13): (<math>p &lt; 0.0001</math>)</b>			
Testing covered 100% by financial assistance (uninsured)	152	41	27.0%
Medicare or Medicaid	54	16	29.6%
Private Insurance	140	70	50.0%

**Table III**

## User Assessment of CancerGene Connect

Question	Strongly agree n (%)	Agree n (%)	Disagree n (%)	Strongly disagree n (%)	Don't Know n (%)
1. The computer was easy to use. Do you...	13 (72%)	4 (22%)	0	0	1 (6%)
2. You felt rushed using the computer. Do you...	0	1 (6%)	9 (50%)	8 (44%)	0
3. Answering the questions was frustrating for you. Do you...	0	2 (11%)	10 (56%)	6 (33%)	0
4. The questions were easy to understand. Do you...	8 (44%)	10 (56%)	0	0	0
5. You liked answering the questions about you and your family. Do you...	6 (33%)	12 (67%)	0	0	0
6. Words on the computer screen were easy to see. Do you...	7 (39%)	8 (44%)	3 (17%)	0	0
7. You enjoyed using the computer program. Do you...	5 (28%)	13 (72%)	0	0	0
8. How much help did you need to use the computer program?	A lot of help 2 (11%)	Some help 6 (33%)	A little help 4 (22%)	No help 6 (33%)	0
9. In the future, do you do think you would choose to complete the questionnaire...	In-clinic before appointment 7 (39%)	Internet before appointment 11 (61%)			