

Published in final edited form as:

Curr Opin Pharmacol. 2014 December ; 0: 31–37. doi:10.1016/j.coph.2014.06.006.

Brain peptides and the modulation of postoperative gastric ileus

Andreas Stengel^a and Yvette Taché^b

^aCharité Center for Internal Medicine and Dermatology, Division of General Internal and Psychosomatic Medicine; Charité-Universitätsmedizin Berlin, Campus Benjamin Franklin, Berlin, Germany

^bCURE: Digestive Diseases Research Center, Center for Neurobiology of Stress and Women's Health, Department of Medicine, Digestive Diseases Division at the University of California Los Angeles, and VA Greater Los Angeles Health Care System, CA 90073, USA

Abstract

Postoperative ileus (POI) develops after abdominal surgery irrespective of the site of surgery. When prolonged, POI can lead to longer hospitalization times and higher healthcare costs. Moreover, it is associated with complaints for the patient. In order to develop new strategies to treat this condition, a deeper understanding of the pathophysiology of the POI is necessary. This review will focus on brain peptides (ghrelin, nesfatin-1, somatostatin, corticotropin-releasing factor, thyrotropin-releasing hormone and calcitonin gene-related peptide) involved in the mediation of POI and the possible modulation of these pathways to shorten the time of POI. Lastly, the role of vagal signaling and the possibility of reducing ambient temperature or chewing gum as potential treatment strategies of alleviating symptoms of POI is discussed.

List of compounds discussed in the article

α-CGRP; astressin; calcitonin gene-related peptide; corticotropin-releasing factor; ghrelin; NUCB2/nesfatin-1; RX-77368; S-406-028; somatostatin; thyrotropin-releasing hormone; ulimorelin

INTRODUCTION

Postoperative ileus regularly develops after abdominal surgery and is characterized by a delay in gastric emptying and a prolongation of intestinal transit [1]. Two decades ago

© 2014 Elsevier Ltd. All rights reserved.

Correspondence to Yvette Taché, PhD, Professor of Medicine, Center for Neurobiology of Stress, CURE Building 115, Room 117, VA GLA Healthcare System, 11301 Wilshire Blvd, Los Angeles, CA, 90073, USA Tel: +1 310 312 9275, Fax: +1 310 268 4963, ytache@mednet.ucla.edu.

Conflicts of interest

There are no conflicts of interest.

The authors have nothing to disclose.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

postoperative ileus was defined as uncomplicated delayed transit that occurred after surgery and resolved spontaneously within two to three days [2]. It is clinically characterized by symptoms such as absence of bowel movements, flatus and defecation, abdominal bloating and pain. When this conditions persists over a longer period of time (> three days after surgery), it is often referred to as paralytic postoperative ileus (or prolonged or pathological ileus) [2]. Another – more recently suggested – definition of postoperative ileus describes the ileus as time from the surgery until passage of flatus or stool together with the time to adequate oral nutrient intake [3]. This type of postoperative ileus has to be clearly distinguished from the secondary postoperative ileus aggravated by complications such as leakage, peritonitis or abscess [3, 4]. In the recovery period, the functions of stomach and small intestine are restored more quickly (within 24 to 48 h) than those of the large bowel function (up to 72 h) [5]. Besides suffering for the patient, prolonged postoperative ileus also leads to higher healthcare costs mainly due to longer hospitalization times [6, 7].

In light of these data a deeper understanding of the pathophysiological mechanisms leading to postoperative ileus is necessary in order to develop effective pharmacological treatment strategies to shorten the time of postoperative ileus. At the cellular level, the pathophysiology of postoperative ileus can be separated in an early neural and humoral [8] as well as a delayed immunological phase [9]. The present review will focus on the early phase with brain peptides involved in the mediation of postoperative ileus and also on the role of vagal signaling in the modulation of postoperative ileus and associated treatment strategies such as chewing gum or reducing ambient temperature, whereas the inflammatory mediators involved and the possible mediation of this phase to alleviate symptoms of postoperative ileus are reviewed elsewhere [10].

MODULATION OF POSTOPERATIVE ILEUS BY BRAIN PEPTIDES AND NEURONAL ACTIVATION

Ghrelin – expression and regulation

Ghrelin was discovered in 1999 as the endogenous ligand of the growth hormone secretagogue receptor 1a (GHS-R1a) [11] which was later renamed ghrelin (GRLN) receptor due to its main ligand [12]. Ghrelin is a 28 amino acid peptide that bears a unique fatty acid residue on the third amino acid essential for the binding to its receptor [11]. The enzyme catalyzing this acylation was long unknown and identified few years ago as a member of the superfamily of membrane-bound O-acyltransferases (MBOATs) and termed ghrelin-O-acyltransferase (GOAT) [13, 14]. The by far major source of ghrelin is the stomach [15] with smaller quantities produced in the intestinal tract [16], pancreas [17] and other peripheral organs including liver, kidney, adipose tissue, heart and skin [18]. Lastly, ghrelin is also produced – although in smaller quantities – directly in the brain in neurons adjacent to the third ventricle [19] and in the arcuate nucleus of the hypothalamus (Arc) [20].

Early on, ghrelin plasma levels were shown to vary in relation to feeding status with an increase in the fasting state and before a meal and a decline thereafter [21] giving rise to a physiological role of ghrelin in the modulation of feeding. Indeed, the food-intake stimulating action of ghrelin has been well documented in humans [22]. In addition, ghrelin

is also regulated by long term changes in body weight with a decrease of ghrelin levels under conditions of obesity [23]. Interestingly, the variation of circulating ghrelin levels also shows a close association with the appearance of gastric migrating motor complexes [24]. Subsequent studies established a physiological role of endogenous ghrelin in the stimulation of gastric motility in rodents. This assumption is supported by the finding that peripheral administration of ghrelin stimulates antral motility in rodents [25] and humans [26] contributing to the acceleration of gastric emptying [27]. However, it is important to note that only pharmacological/supraphysiological doses exert gastroprokinetic effects under conditions of gastroparesis [28].

Ghrelin – involvement in postoperative ileus

Animal studies showed that ghrelin levels are reduced following abdominal surgery with a rapid decline of acyl ghrelin occurring within the first 30 min post surgery followed by a sustained inhibition of these levels over a period of 5 h with a restoration within 24 h [29]. The decline in plasma ghrelin levels was associated with a reduction of gastric and plasma GOAT protein concentrations [29] likely contributing to the decrease in plasma ghrelin due to reduced acylation. Moreover, a stimulated de-acylation of ghrelin may contribute to the lower levels observed as recently reported following the injection of lipopolysaccharide [30]. In line with the assumption of a causal role of ghrelin for the observed changes after surgery, the intravenous injection of ghrelin or GRLN agonists reversed the surgery-induced delay of gastric emptying in rodents [31, 32] and humans [33] and accelerated the time to the first bowel movement in humans [33] resulting in a shorter hospitalization period [34] (Table 1). However, a recent randomized, placebo-controlled phase 3 trial investigating the effect of intravenously injected ulimorelin, a GRLN agonist, did not detect differences in the time to the first bowel movement after surgery compared to placebo [35]. Whether this is related to the dose or pharmacokinetic of the compound will have to be further established. Taken together, the stimulation of ghrelin signaling by GRLN mimetics or activation of GOAT may be a useful strategy to promote gastrointestinal motility after abdominal surgery.

Nesfatin-1 – expression and regulation

Nesfatin-1 was discovered in 2006 in the rat hypothalamus and introduced as a potential cleavage product of nucleobindin2 (NUCB2) [36]. Following the description of nesfatin-1 in food intake-regulatory nuclei in the hypothalamus and brainstem [36], several studies showed the expression of NUCB2/nesfatin-1 in other distinct nuclei throughout the rodent brain [37, 38] pointing towards a pleiotropic action of this peptide. The expression of NUCB2/nesfatin-1 has also been detected in the human Edinger-Westphal nucleus [39] likely contributing to the mature nesfatin-1 detected in the cerebrospinal fluid [40]. As observed before for several other peptides once thought to be restricted to the brain, NUCB2/nesfatin-1 was also detected in peripheral organs such as stomach [41], adipose tissue [42], and pancreas [43] of humans likely contributing to the circulating peptide levels of nesfatin-1 [44]. Interestingly, ghrelin and NUCB2/nesfatin-1 were detected in the same gastric cell, the X/Alike endocrine cell, in stomachs of obese human subjects [41] highlighting this cell type as a possible dual regulator of food intake with a stimulation or inhibition of feeding depending on the peptide product released.

The brain expression of NUCB2/nesfatin-1 was shown to be decreased under conditions of fasting [36] which was in line with a reduction of plasma NUCB2/nesfatin-1 levels that were restored after re-feeding in rats [45] indicative of a role in the regulation of food intake. Nesfatin-1 is likely also involved in the regulation of long term body weight as body weight modulation affects circulating NUCB2/nesfatin-1 levels with higher levels under conditions of obesity reflected in a positive correlation with body mass index [46].

Nesfatin-1 – involvement in postoperative ileus

Recent studies indicate that intracerebroventricular injection of nesfatin-1 delays gastric emptying in rats [45] and reduces gastroduodenal motility in mice [47]. In addition, there is evidence that abdominal surgery activates several NUCB2/nesfatin-1 positive brain nuclei in rats [48] suggesting that activated NUCB2/nesfatin-1 signaling under conditions of abdominal surgery contributes to the observed decrease of food intake and gastrointestinal transit (Table 1). This hypothesis warrants further investigation and the modulation of NUCB2/nesfatin-1 signaling by nesfatin-1 antagonists should be tested after abdominal surgery in humans. However, these studies are hampered so far by the lack of identification of the NUCB2/nesfatin-1 receptor which will be a great leap forward and allow for the generation of specific NUCB2/nesfatin-1 antagonists.

Somatostatin – expression and regulation

Somatostatin was discovered in 1973 as a growth hormone-inhibitory factor isolated from ovine hypothalamus [49] and by now known to exert its effects *via* the interaction with five somatostatin (sst) receptor subtypes (sst₁₋₅) [50]. Subsequent studies showed the widespread distribution of somatostatin in the whole rodent brain except the cerebellum [51] and the human brain [52] in line with the multiple central actions described including the stimulation of food intake, involvement in thermoregulation and a modulation of behavior and autonomic nervous system activity in addition of the well-established endocrine growth hormone suppression [53]. Support for a physiological role of brain somatostatin in the regulation of food intake comes from the observation that hypothalamic somatostatin expression shows a circadian variation with a peak at the beginning of the dark phase when rats as nocturnal eaters show their maximal food consumption and lowest levels in the early light phase [54]. Besides its expression in the brain, somatostatin is also widely expressed in the gastrointestinal tract [55] where it is known to act as an inhibitory modulator of several gastrointestinal peptides and functions [56].

Somatostatin – involvement in postoperative ileus

Since the stomach is a major expression site of somatostatin [57] as well as the somatostatin receptor 2 (sst₂) [58], several studies also investigated the role of somatostatin-sst₂ signaling under conditions of abdominal surgery. The intravenous injection of the selective peptide sst₂ antagonist, S-406-028 prevented the surgery-induced reduction of circulating ghrelin [29]. This somatostatin-sst₂-ghrelin signaling pathway may represent a paracrine mode of action as somatostatin-producing D cells are located in close proximity to the sst₂-bearing ghrelin-producing X/A-like cells of the stomach [29]. However, the blockade of sst₂ signaling under conditions of abdominal surgery did not modulate the surgery-induced

gastric ileus [29]. This observation is in line with a study showing that brain activation of sst₂ signaling does not alter the surgery-induced delay of gastric emptying, whereas the inhibition of food intake was restored [59]. This was associated with a restoration of circulating ghrelin levels to physiological concentrations observed under conditions of fasting [59] giving rise to a dissociation of ghrelin's orexigenic and prokinetic actions under conditions of abdominal surgery when modulated by sst₂ signaling (Table 1). In summary, the blockade of sst₂ signaling may be a useful strategy to promote appetite following abdominal surgery.

Corticotropin-releasing factor – expression and regulation

The stimulatory action of corticotropin-releasing factor (CRF) to release pituitary adrenocorticotrophic hormone (ACTH) in response to various stressors was described in 1950 [60]. This factor was purified five years later [61], however, it took until 1981 until CRF was identified and characterized as a 41 amino acid peptide [62]. Besides the key involvement in the endocrine response to stress, CRF showed to have pleiotropic actions in the modulation of stress-related alterations of autonomic, visceral, immune and behavioral functions [63]. The peptide is expressed in specific brain nuclei with major localization sites in the paraventricular nucleus (PVN) of the hypothalamus, cerebral cortex, amygdala-hippocampal complex and the pontine Barrington's nucleus. CRF is also peripherally expressed in the gastrointestinal tract, enteric nervous system, adipose tissue, heart, lung and testis [64]. It is well established that acute stressors and brain or peripheral injection of CRF reduces upper gastrointestinal motility, while colonic motility and secretion are enhanced [65].

Corticotropin-releasing factor – involvement in postoperative gastric ileus

Key nuclei that mediate autonomic outflow to the stomach such as the PVN and the dorsal vagal complex in the brainstem are activated under conditions of abdominal surgery [66] responsible for the mediation of the CRF-induced inhibition of gastric motility [65]. This action can be blocked by peripheral (intravenous) as well as brain injection of a non-selective CRF antagonist, astressin [67]. It is important to note that the central gastroparetic action of CRF is independent of a stimulation of the hypothalamic-pituitary-adrenal axis based on the observation that the gastric-inhibitory response to CRF is still visible in hypophysectomized or adrenalectomized rats [68]. Since CRF is known to reduce vagal efferent activity [69] and acetylcholine stimulates ghrelin release from the stomach [70], the postoperative CRF-induced delay of gastric emptying may be mediated by a reduction of vagal efferent activity. These actions are likely to be CRF₁ mediated as in mice lacking the CRF₁ no postoperative gastric ileus following midline celiotomy, cecal exteriorization and palpation was observed [71] (Table 1).

Calcitonin gene-related peptide - expression and regulation

Alpha-calcitonin gene-related peptide (α -CGRP) is a 37 amino acid peptide and part of the calcitonin family of peptides that acts in the brain to exert its biological actions [72]. CGRP is – besides its expression in the periphery including the gastrointestinal tract – expressed in the nucleus of the solitary tract of the brainstem and immunoreactive fibers make close contact to gastric efferent motor neurons [73]. This finding is in line with the identification

of the CGRP receptor on neurons of the dorsal motor nucleus of the vagus nerve [74], providing the neuroanatomical basis for the decrease of vagal efferent activity observed after intracisternal injection of α -CGRP [75].

Calcitonin gene-related peptide - involvement in postoperative ileus

The α -CGRP-induced decrease of efferent vagal activity is likely to be involved in the mediation of postoperative gastric ileus as shown by experimental evidence that peripheral injection of the CGRP receptor antagonist, CGRP₈₋₃₇ reversed the abdominal surgery-induced delay of gastric emptying in dogs [76]. Similarly, immunoneutralization of CGRP using an anti-CGRP antibody reversed the inhibition of gastric emptying [77] and also of colonic transit, the reduction of fecal pellet output and also the anorexigenic response following abdominal surgery in rats [78] indicating a major involvement of CGRP signaling in the mediation of alterations of gastrointestinal functions under conditions of abdominal surgery (Table 1).

Vagal signaling - involvement in postoperative ileus

Since the vagus nerve is a major driving source of gastrointestinal motility [79], the modulation of vagal activity is an interesting target in order to influence symptoms of abdominal surgery-induced postoperative ileus. It is well known that a reduction in ambient temperature activates the thyrotropin-releasing hormone (TRH) signaling system [80] expressed in the raphe pallidus, raphe obscurus and parapyramidal regions projecting to the dorsal vagal complex of the brainstem [81] resulting in a vagal dependent cholinergic stimulation of gastric motility and secretion through activation of gastric myenteric cholinergic neurons [80]. In line with these findings, brainstem (intracisternal) injection of the stable TRH agonist, RX-77368 or endogenous TRH induced by cold ambient temperature (4–6 °C) prevents the abdominal surgery-induced delay of gastric emptying [82] (Table 1). Interestingly, this was associated with a complete restoration of circulating ghrelin levels reduced by abdominal surgery [82] indicating the advantage of stimulating vagal activity following abdominal surgery.

Recent studies point towards the benefit of chewing sugarless gum as an applicable mean to stimulate vagal activity which has proven effective to shorten the time to the first flatus and bowel movement and to reduce the period of hospitalization as indicated in a meta-analysis reviewing the data of nine studies with a combined population of more than 400 patients that underwent abdominal surgery [83]. The treatment appears to be safe as no difference in complication rates was observed for chewing gum *versus* placebo [84] (Table 1). These data are corroborated by a recent randomized clinical trial investigating women after cesarean section that showed a shortened time to the first bowel movement, first passage of flatus, first defecation and first hunger sensation in the gum *versus* control group [85].

SUMMARY

Postoperative ileus is a frequent condition occurring after abdominal surgery that may lead to suffering for the individual patient and increased health care costs for the society. Although our understanding of the pathophysiology underlying postoperative ileus increased

during the past years, this did not lead to widely accepted and generally recommended treatment strategies for this medical condition. Promising pharmacological targets encompass the promotion of prokinetic pathways inhibited by surgery namely vagal activation, ghrelin, and to block receptors activated by abdominal surgery namely CRF, CGRP and sst₂ and inhibit gastric motor function. Clinical studies providing evidence that chewing gum reduces the duration of postoperative ileus corroborate the role of activating central vagal mechanisms as an important approach consistent with experimental studies.

Acknowledgments

This work was supported by NIH R01 DK-33061, NIH Center Grant DK-41301 (Animal Core), VA Research Career Scientist (Y.T.), VA Merit Award (Y.T.) German Research Foundation STE 1765/3-1 (A.S.) and Charité University Funding UFF 89-441-176 (A.S.).

REFERENCES AND RECOMMENDED READING

Papers of particular interest have been highlighted as:

* of special interest

** of outstanding interest

1. Luckey A, Livingston E, Taché Y. Mechanisms and treatment of postoperative ileus. *Arch Surg*. 2003; 138:206–14. [PubMed: 12578422]
2. Livingston EH, Passaro EP Jr. Postoperative ileus. *Dig Dis Sci*. 1990; 35:121–32. [PubMed: 2403907]
3. Kehlet H. Postoperative ileus--an update on preventive techniques. *Nat Clin Pract Gastroenterol Hepatol*. 2008; 5:552–8. [PubMed: 18695704]
4. Delaney CP, Senagore AJ, Viscusi ER, Wolff BG, Fort J, Du W, et al. Postoperative upper and lower gastrointestinal recovery and gastrointestinal morbidity in patients undergoing bowel resection: pooled analysis of placebo data from 3 randomized controlled trials. *Am J Surg*. 2006; 191:315–9. [PubMed: 16490538]
5. Benson, MJ.; Wingate, DL. Ileus and mechanical obstruction. In: Kumar, D.; Wingate, DL., editors. *An illustrated guide to gastrointestinal motility*. London: Churchill Livingstone; 1993. p. 547–66.
- 6*. Asgeirsson T, El-Badawi KI, Mahmood A, Barletta J, Luchtefeld M, Senagore AJ. Postoperative ileus: it costs more than you expect. *J Am Coll Surg*. 2010; 210:228–31. This paper describes postoperative ileus as a frequent condition and gives an overview over the costs of longer hospitalization periods and the treatment of associated complications. [PubMed: 20113944]
7. Zeinali F, Stulberg JJ, Delaney CP. Pharmacological management of postoperative ileus. *Can J Surg*. 2009; 52:153–7. [PubMed: 19399212]
8. Bueno L, Fioramonti J, Ruckebusch Y. Postoperative intestinal motility in dogs and sheep. *Am J Dig Dis*. 1978; 23:682–9. [PubMed: 685935]
9. Kalff JC, Carlos TM, Schraut WH, Billiar TR, Simmons RL, Bauer AJ. Surgically induced leukocytic infiltrates within the rat intestinal muscularis mediate postoperative ileus. *Gastroenterology*. 1999; 117:378–87. [PubMed: 10419919]
- 10*. van Bree SH, Nemethova A, Cailotto C, Gomez-Pinilla PJ, Matteoli G, Boeckxstaens GE. New therapeutic strategies for postoperative ileus. *Nat Rev Gastroenterol Hepatol*. 2012; 9:675–83. This review focuses on the inflammatory pathways activated under conditions of abdominal surgery that contribute to the delayed phase of postoperative ileus. Possible novel approaches to tackle these inflammatory pathways as treatment options of postoperative ileus are also discussed. [PubMed: 22801725]
11. Kojima M, Hosoda H, Date Y, Nakazato M, Matsuo H, Kangawa K. Ghrelin is a growth-hormone-releasing acylated peptide from stomach. *Nature*. 1999; 402:656–60. [PubMed: 10604470]

12. Davenport AP, Bonner TI, Foord SM, Harmar AJ, Neubig RR, Pin JP, et al. International Union of Pharmacology. LVI. Ghrelin receptor nomenclature, distribution, and function. *Pharmacol Rev.* 2005; 57:541–6. [PubMed: 16382107]
13. Yang J, Brown MS, Liang G, Grishin NV, Goldstein JL. Identification of the acyltransferase that octanoylates ghrelin, an appetite-stimulating peptide hormone. *Cell.* 2008; 132:387–96. [PubMed: 18267071]
14. Gutierrez JA, Solenberg PJ, Perkins DR, Willency JA, Knierman MD, Jin Z, et al. Ghrelin octanoylation mediated by an orphan lipid transferase. *PNAS U S A.* 2008; 105:6320–5.
15. Jeon TY, Lee S, Kim HH, Kim YJ, Son HC, Kim DH, et al. Changes in plasma ghrelin concentration immediately after gastrectomy in patients with early gastric cancer. *J Clin Endocrinol Metab.* 2004; 89:5392–6. [PubMed: 15531487]
16. Date Y, Kojima M, Hosoda H, Sawaguchi A, Mondal MS, Suganuma T, et al. Ghrelin, a novel growth hormone-releasing acylated peptide, is synthesized in a distinct endocrine cell type in the gastrointestinal tracts of rats and humans. *Endocrinology.* 2000; 141:4255–61. [PubMed: 11089560]
17. Date Y, Nakazato M, Hashiguchi S, Dezaki K, Mondal MS, Hosoda H, et al. Ghrelin is present in pancreatic alpha-cells of humans and rats and stimulates insulin secretion. *Diabetes.* 2002; 51:124–9. [PubMed: 11756331]
18. Gnanapavan S, Kola B, Bustin SA, Morris DG, McGee P, Fairclough P, et al. The tissue distribution of the mRNA of ghrelin and subtypes of its receptor, GHS-R, in humans. *J Clin Endocrinol Metab.* 2002; 87:2988. [PubMed: 12050285]
19. Cowley MA, Smith RG, Diano S, Tschop M, Pronchuk N, Grove KL, et al. The distribution and mechanism of action of ghrelin in the CNS demonstrates a novel hypothalamic circuit regulating energy homeostasis. *Neuron.* 2003; 37:649–61. [PubMed: 12597862]
20. Lu S, Guan JL, Wang QP, Uehara K, Yamada S, Goto N, et al. Immunocytochemical observation of ghrelin-containing neurons in the rat arcuate nucleus. *Neurosci Lett.* 2002; 321:157–60. [PubMed: 11880196]
21. Cummings DE, Purnell JQ, Frayo RS, Schmidova K, Wisse BE, Weigle DS. A preprandial rise in plasma ghrelin levels suggests a role in meal initiation in humans. *Diabetes.* 2001; 50:1714–9. [PubMed: 11473029]
22. Druce MR, Wren AM, Park AJ, Milton JE, Patterson M, Frost G, et al. Ghrelin increases food intake in obese as well as lean subjects. *Int J Obes.* 2005; 29:1130–6.
23. Cummings DE, Weigle DS, Frayo RS, Breen PA, Ma MK, Dellinger EP, et al. Plasma ghrelin levels after diet-induced weight loss or gastric bypass surgery. *N Eng J Med.* 2002; 346:1623–30.
24. Ariga H, Tsukamoto K, Chen C, Mantyh C, Pappas TN, Takahashi T. Endogenous acyl ghrelin is involved in mediating spontaneous phase III-like contractions of the rat stomach. *Neurogastroenterol Motil.* 2007; 19:675–80. [PubMed: 17640183]
25. Fujino K, Inui A, Asakawa A, Kihara N, Fujimura M, Fujimiya M. Ghrelin induces fasted motor activity of the gastrointestinal tract in conscious fed rats. *J Physiol.* 2003; 550:227–40. [PubMed: 12837928]
26. Tack J, Depoortere I, Bisschops R, Delpoort C, Coulie B, Meulemans A, et al. Influence of ghrelin on interdigestive gastrointestinal motility in humans. *Gut.* 2006; 55:327–33. [PubMed: 16216827]
27. Levin F, Edholm T, Schmidt PT, Gryback P, Jacobsson H, Degerblad M, et al. Ghrelin stimulates gastric emptying and hunger in normal-weight humans. *J Clin Endocrinol Metab.* 2006; 91:3296–302. [PubMed: 16772353]
28. Murray CD, Martin NM, Patterson M, Taylor SA, Ghatei MA, Kamm MA, et al. Ghrelin enhances gastric emptying in diabetic gastroparesis: a double blind, placebo controlled, crossover study. *Gut.* 2005; 54:1693–8. [PubMed: 16085693]
- 29*. Stengel A, Goebel-Stengel M, Wang L, Shaikh A, Lambrecht NW, Rivier JE, et al. Abdominal surgery inhibits circulating acyl ghrelin and ghrelin-O-acyltransferase levels in rats: role of the somatostatin receptor subtype 2. *Am J Physiol Gastrointest Liver Physiol.* 2011; 301:G239–48. This paper demonstrates that abdominal surgery decreases circulating ghrelin levels *via* a gastric somatostatin-somatostatin receptor 2 signaling pathway. [PubMed: 21636529]

30. Satou M, Nishi Y, Yoh J, Hattori Y, Sugimoto H. Identification and characterization of acyl-protein thioesterase 1/lysophospholipase I as a ghrelin deacylation/lysophospholipid hydrolyzing enzyme in fetal bovine serum and conditioned medium. *Endocrinology*. 2010; 151:4765–75. [PubMed: 20685872]
31. Trudel L, Tomasetto C, Rio MC, Bouin M, Plourde V, Eberling P, et al. Ghrelin/motilin-related peptide is a potent prokinetic to reverse gastric postoperative ileus in rat. *Am J Physiol Gastroint Liver Physiol*. 2002; 282:G948–52.
32. Poitras P, Polvino WJ, Rocheleau B. Gastrokinetic effect of ghrelin analog RC-1139 in the rat. Effect on post-operative and on morphine induced ileus. *Peptides*. 2005; 26:1598–601. [PubMed: 16112398]
- 33*. Falken Y, Webb DL, Abraham-Nordling M, Kressner U, Hellstrom PM, Naslund E. Intravenous ghrelin accelerates postoperative gastric emptying and time to first bowel movement in humans. *Neurogastroenterol Motil*. 2013; 25:474–80. In this randomized trial the authors show that ghrelin infusion accelerates gastric emptying and shortens the time to the first bowel movement under inhibited conditions following abdominal surgery giving rise to the use of ghrelin/ghrelin mimetics under these conditions. [PubMed: 23527561]
34. Popescu I, Fleshner PR, Pezzullo JC, Charlton PA, Kosutic G, Senagore AJ. The Ghrelin agonist TZP-101 for management of postoperative ileus after partial colectomy: a randomized, dose-ranging, placebo-controlled clinical trial. *Dis Colon Rectum*. 2010; 53:126–34. [PubMed: 20087086]
35. Shaw M, Pediconi C, McVey D, Mondou E, Quinn J, Chamblin B, et al. Safety and efficacy of ulimorelin administered postoperatively to accelerate recovery of gastrointestinal motility following partial bowel resection: results of two randomized, placebo-controlled phase 3 trials. *Dis Colon Rectum*. 2013; 56:888–97. [PubMed: 23739196]
36. Oh-I S, Shimizu H, Satoh T, Okada S, Adachi S, Inoue K, et al. Identification of nesfatin-1 as a satiety molecule in the hypothalamus. *Nature*. 2006; 443:709–12. [PubMed: 17036007]
37. Goebel M, Stengel A, Wang L, Lambrecht NW, Taché Y. Nesfatin-1 immunoreactivity in rat brain and spinal cord autonomic nuclei. *Neurosci Lett*. 2009; 452:241–6. [PubMed: 19348732]
38. Goebel-Stengel M, Wang L, Stengel A, Taché Y. Localization of nesfatin-1 neurons in the mouse brain and functional implication. *Brain Res*. 2011; 1396:20–34. [PubMed: 21555116]
39. Bloem B, Xu L, Morava E, Faludi G, Palkovits M, Roubos EW, et al. Sex-specific differences in the dynamics of cocaine- and amphetamine-regulated transcript and nesfatin-1 expressions in the midbrain of depressed suicide victims vs. controls. *Neuropharmacology*. 2012; 62:297–303. [PubMed: 21803054]
40. Tan BK, Hallschmid M, Kern W, Lehnert H, Randeve HS. Decreased cerebrospinal fluid/plasma ratio of the novel satiety molecule, nesfatin-1/NUCB-2, in obese humans: evidence of nesfatin-1/NUCB-2 resistance and implications for obesity treatment. *J Clin Endocrinol Metab*. 2011; 96:E669–73. [PubMed: 21252251]
41. Stengel A, Hofmann T, Goebel-Stengel M, Lembke V, Ahnis A, Elbelt U, et al. Ghrelin and NUCB2/nesfatin-1 are expressed in the same gastric cell and differentially correlated with body mass index in obese subjects. *Histochem Cell Biol*. 2013; 139:909–18. [PubMed: 23515787]
42. Ramanjaneya M, Addison M, Randeve HS. Possible role of NUCB2/nesfatin-1 in adipogenesis. *Curr Pharm Des*. 2013; 19:6976–80. [PubMed: 23537078]
43. Riva M, Nitert MD, Voss U, Sathanoori R, Lindqvist A, Ling C, et al. Nesfatin-1 stimulates glucagon and insulin secretion and beta cell NUCB2 is reduced in human type 2 diabetic subjects. *Cell Tissue Res*. 2011; 346:393–405. [PubMed: 22108805]
44. Tsuchiya T, Shimizu H, Yamada M, Osaki A, Oh IS, Ariyama Y, et al. Fasting Concentrations of Nesfatin-1 Are Negatively Correlated with Body Mass Index in Non-Obese Males. *Clin Endocrinol (Oxf)*. 2010; 73:484–90. [PubMed: 20550530]
45. Stengel A, Goebel M, Wang L, Rivier J, Kobelt P, Mönnikes H, et al. Central nesfatin-1 reduces dark-phase food intake and gastric emptying in rats: differential role of corticotropin-releasing factor2 receptor. *Endocrinology*. 2009; 150:4911–9. [PubMed: 19797401]

46. Ramanjaneya M, Chen J, Brown JE, Tripathi G, Hallschmid M, Patel S, et al. Identification of nesfatin-1 in human and murine adipose tissue: a novel depot-specific adipokine with increased levels in obesity. *Endocrinology*. 2010; 151:3169–80. [PubMed: 20427481]
47. Atsuchi K, Asakawa A, Ushikai M, Ataka K, Tsai M, Koyama K, et al. Centrally administered nesfatin-1 inhibits feeding behaviour and gastroduodenal motility in mice. *Neuroreport*. 2010; 21:1008–11. [PubMed: 20827224]
48. Stengel A, Goebel M, Wang L, Taché Y. Abdominal surgery activates nesfatin-1 immunoreactive brain nuclei in rats. *Peptides*. 2010; 31:263–70. [PubMed: 19944727]
49. Brazeau P, Vale W, Burgus R, Ling N, Butcher M, Rivier J, et al. Hypothalamic polypeptide that inhibits the secretion of immunoreactive pituitary growth hormone. *Science*. 1973; 179:77–9. [PubMed: 4682131]
50. Olias G, Viollet C, Kusserow H, Epelbaum J, Meyerhof W. Regulation and function of somatostatin receptors. *J Neurochem*. 2004; 89:1057–91. [PubMed: 15147500]
51. Viollet C, Lepousez G, Loudes C, Videau C, Simon A, Epelbaum J. Somatostatinergic systems in brain: networks and functions. *Mol Cell Endocrinol*. 2008; 286:75–87. [PubMed: 17997029]
52. Mengod G, Rigo M, Savasta M, Probst A, Palacios JM. Regional distribution of neuropeptide somatostatin gene expression in the human brain. *Synapse*. 1992; 12:62–74. [PubMed: 1357764]
53. Stengel A, Goebel M, Wang L, Rivier J, Kobelt P, Mönnikes H, Taché Y. Selective central activation of somatostatin2 receptor increases food intake, grooming behavior and rectal temperature in rats. *J Physiol Pharmacol*. 2010; 61:399–407. [PubMed: 20814067]
54. Gardi J, Obal F Jr, Fang J, Zhang J, Krueger JM. Diurnal variations and sleep deprivation-induced changes in rat hypothalamic GHRH and somatostatin contents. *Am J Physiol*. 1999; 277:R1339–44. [PubMed: 10564205]
55. Shulkes A. Somatostatin: physiology and clinical applications. *Baillieres Clin Endocrinol Metab*. 1994; 8:215–36. [PubMed: 7907862]
56. Low MJ. Clinical endocrinology and metabolism. The somatostatin neuroendocrine system: physiology and clinical relevance in gastrointestinal and pancreatic disorders. *Best Pract Res Clin Endocrinol Metab*. 2004; 18:607–22. [PubMed: 15533778]
57. Sachs G, Zeng N, Prinz C. Physiology of isolated gastric endocrine cells. *Annu Rev Physiol*. 1997; 59:243–56. [PubMed: 9074763]
58. Schindler M, Humphrey PP. Differential distribution of somatostatin sst2 receptor splice variants in rat gastric mucosa. *Cell Tissue Res*. 1999; 297:163–8. [PubMed: 10398894]
59. Stengel A, Goebel-Stengel M, Wang L, Luckey A, Hu E, Rivier J, et al. Central administration of pan-somatostatin agonist ODT8-SST prevents abdominal surgery-induced inhibition of circulating ghrelin, food intake and gastric emptying in rats. *Neurogastroenterol Motil*. 2011; 23:e294–308. [PubMed: 21569179]
60. De Groot J, Harris GW. Hypothalamic control of the anterior pituitary gland and blood lymphocytes. *J Physiol*. 1950; 111:335–46. [PubMed: 14795443]
61. Guillemin R, Rosenberg B. Humoral hypothalamic control of anterior pituitary: a study with combined tissue cultures. *Endocrinology*. 1955; 57:599–607. [PubMed: 13270723]
62. Vale W, Spiess J, Rivier C, Rivier J. Characterization of a 41-residue ovine hypothalamic peptide that stimulates secretion of corticotropin and beta-endorphin. *Science*. 1981; 213:1394–7. [PubMed: 6267699]
63. Stengel A, Taché Y. Corticotropin-releasing factor signaling and visceral response to stress. *Exp Biol Med* (Maywood). 2010; 235:1168–78. [PubMed: 20881321]
64. Taché Y, Bonaz B. Corticotropin-releasing factor receptors and stress-related alterations of gut motor function. *J Clin Invest*. 2007; 117:33–40. [PubMed: 17200704]
65. Stengel A, Taché Y. Neuroendocrine Control of the Gut During Stress: Corticotropin-Releasing Factor Signaling Pathways in the Spotlight. *Annu Rev Physiol*. 2009; 71:219–39. [PubMed: 18928406]
66. Wang L, Goebel-Stengel M, Stengel A, Wu SV, Ohning G, Taché Y. Comparison of CRF-immunoreactive neurons distribution in mouse and rat brains and selective induction of Fos in rat hypothalamic CRF neurons by abdominal surgery. *Brain Res*. 2011; 1415:34–46. [PubMed: 21872218]

67. Martinez V, Rivier J, Taché Y. Peripheral injection of a new corticotropin-releasing factor (CRF) antagonist, astressin, blocks peripheral CRF- and abdominal surgery-induced delayed gastric emptying in rats. *J Pharmacol Exp Ther.* 1999; 290:629–34. [PubMed: 10411571]
68. Taché Y, Maeda-Hagiwara M, Turkelson CM. Central nervous system action of corticotropin-releasing factor to inhibit gastric emptying in rats. *Am J Physiol.* 1987; 253:G241–5. [PubMed: 3497585]
69. Adelson DW, Kosoyan HP, Wang Y, Steinberg JZ, Taché Y. Gastric vagal efferent inhibition evoked by intravenous CRF is unrelated to simultaneously recorded vagal afferent activity in urethane-anesthetized rats. *J Neurophysiol.* 2007; 97:3004–14. [PubMed: 17314242]
70. Shrestha YB, Wickwire K, Giraudo SQ. Direct effects of nutrients, acetylcholine, CCK, and insulin on ghrelin release from the isolated stomachs of rats. *Peptides.* 2009; 30:1187–91. [PubMed: 19463754]
71. Luckey A, Wang L, Jamieson PM, Basa NR, Million M, Czimmer J, et al. Corticotropin-releasing factor receptor 1-deficient mice do not develop postoperative gastric ileus. *Gastroenterology.* 2003; 125:654–9. [PubMed: 12949710]
72. Amara SG, Arriza JL, Leff SE, Swanson LW, Evans RM, Rosenfeld MG. Expression in brain of a messenger RNA encoding a novel neuropeptide homologous to calcitonin gene-related peptide. *Science.* 1985; 229:1094–7. [PubMed: 2994212]
73. Ladic LA, Buchan AM. Three-dimensional spatial relationship of neuropeptides and receptors in the rat dorsal vagal complex. *Brain Res.* 1998; 795:312–24. [PubMed: 9622662]
74. Ma W, Chabot JG, Powell KJ, Jhamandas K, Dickerson IM, Quirion R. Localization and modulation of calcitonin gene-related peptide-receptor component protein-immunoreactive cells in the rat central and peripheral nervous systems. *Neuroscience.* 2003; 120:677–94. [PubMed: 12895509]
75. Taché Y. Intracisternal injection of calcitonin-gene related peptide inhibits experimental gastric ulcers. In: Garner, A.; O'Brien, PE., editors. *Mechanisms of injury, protection and repair of the upper gastrointestinal tract.* Chichester: John Wiley & Sons; 1991. p. 383-93.
76. Trudel L, Bouin M, Tomasetto C, Eberling P, St-Pierre S, Bannon P, et al. Two new peptides to improve post-operative gastric ileus in dog. *Peptides.* 2003; 24:531–4. [PubMed: 12860196]
77. Plourde V, Wong HC, Walsh JH, Raybould HE, Taché Y. CGRP antagonists and capsaicin on celiac ganglia partly prevent postoperative gastric ileus. *Peptides.* 1993; 14:1225–9. [PubMed: 7510881]
78. Zittel TT, Lloyd KC, Rothenhofer I, Wong H, Walsh JH, Raybould HE. Calcitonin gene-related peptide and spinal afferents partly mediate postoperative colonic ileus in the rat. *Surgery.* 1998; 123:518–27. [PubMed: 9591004]
79. Stengel, A.; Taché, Y. The physiological relationships between the brainstem, vagal stimulation and feeding. In: Preedy, VR., editor. *The International Handbook of Behavior, Diet and Nutrition.* London: Springer; 2011.
80. Taché Y, Yang H, Miampamba M, Martinez V, Yuan PQ. Role of brainstem TRH/TRH-R1 receptors in the vagal gastric cholinergic response to various stimuli including sham-feeding. *Auton Neurosci.* 2006; 125:42–52. [PubMed: 16520096]
81. Lynn RB, Kreider MS, Miselis RR. Thyrotropin-releasing hormone-immunoreactive projections to the dorsal motor nucleus and the nucleus of the solitary tract of the rat. *J Comp Neurol.* 1991; 311:271–88. [PubMed: 1753019]
82. Stengel A, Goebel M, Luckey A, Yuan PQ, Wang L, Taché Y. Cold ambient temperature reverses abdominal surgery-induced delayed gastric emptying and decreased plasma ghrelin levels in rats. *Peptides.* 2010; 31:2229–35. [PubMed: 20817059]
- 83**. Noble EJ, Harris R, Hosie KB, Thomas S, Lewis SJ. Gum chewing reduces postoperative ileus? A systematic review and meta-analysis. *Int J Surg.* 2009; 7:100–5. This meta-analysis shows that chewing gum post surgically reduces the time to flatus, time to first bowel movement and the duration of hospitalization. Despite the fact that rigorous randomized controlled trials are lacking so far the authors conclude that chewing gum might be a useful and applicable approach to reduce symptoms of postoperative ileus. [PubMed: 19261555]

84. Fitzgerald JE, Ahmed I. Systematic review and meta-analysis of chewing-gum therapy in the reduction of postoperative paralytic ileus following gastrointestinal surgery. *World J Surg.* 2009; 33:2557–66. [PubMed: 19763686]
85. Mohsenzadeh Ledari F, Barat S, Delavar MA, Banihosini SZ, Khafri S. Chewing sugar-free gum reduces ileus after cesarean section in nulliparous women: a randomized clinical trial. *Iran Red Crescent Med J.* 2013; 15:330–4. [PubMed: 24083008]

KEY POINTS

- Postoperative ileus is a frequent condition after abdominal surgery causing suffering for the patient and significant costs for the healthcare system.
- Several brain peptides are involved in the development of postoperative ileus.
- Chewing gum as a mean to stimulate vagal signaling may be an applicable method to alleviate symptoms of postoperative ileus.

Highlights

- Postoperative ileus (POI) is a frequent condition after abdominal surgery
- POI causes suffering for patients and significant costs for the healthcare system
- Several brain peptides are involved in the development of POI
- Chewing gum may be an applicable method to alleviate symptoms of POI

Table 1

Modulation of postoperative ileus by centrally acting peptides.

Peptide	Proposed physiological function	Modulation of postoperative ileus	Reference
Ghrelin	stimulation of food intake and gastrointestinal motility	reversal of surgery-induced delay of gastric emptying; acceleration of time to first bowel movement; earlier hospital discharge	[33, 34]
Nesfatin-1	inhibition of food intake and gastrointestinal motility	no data available so far under conditions of postoperative ileus; NUCB2/nesfatin-1 brain nuclei are activated by abdominal surgery	[48]
Somatostatin	peripheral: inhibition of food intake and gastrointestinal motility; central: stimulation of food intake	peripheral: sst ₂ blockade restores ghrelin levels but does not alter the surgery-induced gastric ileus; central: sst ₂ activation does not alter the surgery-induced delay of gastric emptying but restores circulating ghrelin levels and stimulates food intake	[29, 59]
CRF	key hormone in the response to stress, reduction of upper gastrointestinal and stimulation of colonic motility and secretion	CRF ₁ mediates delay of gastric emptying in mice	[71]
CGRP	decrease of vagal efferent activity	immunoneutralization of CGRP reverses postoperative inhibition of gastric emptying, delay of colonic transit, reduction of fecal pellet output and the anorexigenic response	[77, 78]
TRH	stimulation of gastrointestinal motility and secretion	cold-induced TRH-mediated vagal activation restores gastric emptying and ghrelin levels post surgery; chewing gum shortens time to first flatus and bowel movement after surgery and reduces time of hospitalization	[82, 83]

Abbreviations: CGRP, calcitonin gene-related peptide; CRF, corticotropin-releasing factor; CRF₁, corticotropin-releasing factor receptor subtype 1; sst₂, somatostatin receptor subtype 2; TRH, thyrotropin-releasing hormone