

## ERRATUM

# Erratum for Mandelbrot et al., Placental Transfer of Rilpivirine in an *Ex Vivo* Human Cotyledon Perfusion Model

Laurent Mandelbrot,<sup>a,b,c</sup> Dominique Duro,<sup>a</sup> Emilie Belissa,<sup>d</sup> Gilles Peytavin<sup>d,e</sup>

Assistance Publique-Hôpitaux de Paris, Hôpital Louis Mourier, Service de Gynécologie-Obstétrique, Hôpitaux Universitaires Paris-Nord Val de Seine, Colombes, France<sup>a</sup>; Université Paris-Diderot, Université Sorbonne Paris-Cité, DHU Risks in Pregnancy, Paris, France<sup>b</sup>; INSERM U1018, Kremlin-Bicêtre, France<sup>c</sup>; Assistance Publique-Hôpitaux de Paris, Hôpital Bichat-Claude Bernard, Clinical Pharmacology-Toxicology Department, Hôpitaux Universitaires Paris-Nord Val de Seine, Paris, France<sup>d</sup>; IAME, UMR 1137, INSERM, Paris, France<sup>e</sup>

Volume 59, no. 5, p. 2901–2903, 2015. Page 2902: In Table 1, the placenta 6 CM and CF values were inadvertently repeated for placenta 7. The table should appear as shown below.

**TABLE 1** Placental transfer of rilpivirine in *ex vivo*-perfused human cotyledons

Perfusion time (Min)	Placenta 1 <sup>a</sup>				Placenta 2				Placenta 3				Placenta 4			
	CM (ng/ml)	CF (ng/ml)	FTR (%)	CLI (%)	CM (ng/ml)	CF (ng/ml)	FTR (%)	CLI (%)	CM (ng/ml)	CF (ng/ml)	FTR (%)	CLI (%)	CM (ng/ml)	CF (ng/ml)	FTR (%)	CLI (%)
15	388	40	10	52	544	37	7	13	262	67	26	38	423	44	10	44
30	217	45	21	64	399	62	16	26	249	82	33	47	400	92	23	47
45	622	76	12	31	463	86	19	29	375	78	21	26	400	162	41	71
60	598	50	8	23	410	105	26	45	243	79	33	42	420	96	23	42
75	210	65	31	139	404	86	21	33	314	77	25	29	387	90	23	101
90	341	75	22	61	387	118	30	59	304	106	35	58	340	110	32	225
Mean	396	58.5	17	62	434.50	82.33	20	34	291.17	81.50	29	40	395	99	25	88
SD	179.78	15.60	9	41	59.75	29.26	8	16	50.32	13.03	6	12	30.16	38.09	10	71

  

Perfusion time (Min)	Placenta 5				Placenta 6				Placenta 7				Placenta 8			
	CM (ng/ml)	CF (ng/ml)	FTR (%)	CLI (%)	CM (ng/ml)	CF (ng/ml)	FTR (%)	CLI (%)	CM (ng/ml)	CF (ng/ml)	FTR (%)	CLI (%)	CM (ng/ml)	CF (ng/ml)	FTR (%)	CLI (%)
15	387	42	11	41	438	53	12	27	499	30	6	18	420	36	9	31
30	406	250	62	134	399	131	33	50	403	58	14	33	408	66	16	40
45	418	101	24	52	451	129	29	48	393	62	16	34	401	159	40	71
60	350	291	83	223	430	168	39	106	360	80	22	58	403	189	47	72
75	398	103	26	70	413	153	37	60	405	85	21	58	418	196	47	93
90	799	113	14	33	440	147	33	63	379	107	28	98	358	191	53	84
Mean	459.67	150	37	92	428.5	130.17	31	59	406.5	70.33	18	50	401.33	139.5	35	65
SD	167.85	97.49	29	74	19.19	40.48	10	26	48.31	26.46	8	28	22.59	70.41	18	24

<sup>a</sup> CM, rilpivirine concentration in the maternal compartment; CF, rilpivirine concentration in the fetal compartment; FTR, fetal transfer rate; CLI, clearance index.

**Citation** Mandelbrot L, Duro D, Belissa E, Peytavin G. 2015. Erratum for Mandelbrot et al., Placental transfer of rilpivirine in an *ex vivo* human cotyledon perfusion model. *Antimicrob Agents Chemother* 59:5869. doi:10.1128/AAC.01495-15.

Copyright © 2015, American Society for Microbiology. All Rights Reserved.

doi:10.1128/AAC.01495-15