

Outcomes of minimally invasive double valve surgery

Orlando Santana¹, Steve Xydas¹, Roy F. Williams¹, Angelo LaPietra¹, Maurice Mawad¹, Frederick Hasty², Esteban Escolar¹, Christos G. Mihos³

¹Division of Cardiology, Mount Sinai Heart Institute, the Columbia University, Miami Beach, FL, USA; ²Department of Anesthesia, Mount Sinai Medical Center, Miami Beach, FL, USA; ³Cardiac Ultrasound Laboratory, Massachusetts General Hospital, Harvard Medical School, Boston, MA, USA

Contributions: (I) Conception and design: O Santana, CG Mihos; (II) Administrative support: None; (III) Provision of study materials or patients: None; (IV) Collection and assembly of data: O Santana, CG Mihos; (V) Data analysis and interpretation: All authors; (VI) Manuscript writing: All authors; (VII) Final approval of manuscript: All authors.

Correspondence to: Orlando Santana, MD. Echocardiography Laboratory, Columbia University Division of Cardiology, Mount Sinai Heart Institute, 4300 Alton Road, Miami Beach, Florida 33140, USA. Email: osantana@msmc.com.

Background: Double valve surgery is associated with an increased peri-operative morbidity and mortality. A less invasive right thoracotomy approach may be a viable alternative to median sternotomy surgery in these higher-risk patients.

Methods: We retrospectively analyzed the baseline demographics, operative characteristics, and post-operative outcomes of patients who underwent minimally invasive double valve surgery between January 2009 and December 2011 at our institution.

Results: The cohort consisted of 117 patients, of which 68 (58.1%) were female. The mean age was 73±11 years, and the mean left ventricular ejection fraction was 52±11%. There were 43 (36.8%) patients with a history of congestive heart failure, 45 (38.5%) with chronic obstructive pulmonary disease, and 5 (4.3%) had a history of chronic kidney disease. The patients underwent primary (90.6%) or re-operative (9.4%) double valve surgery, which consisted of 50 (42.7%) aortic valve replacement and mitral valve repair, 31 (26.5%) mitral and tricuspid valve repair, 18 (15.4%) aortic and mitral valve replacement, 17 (14.5%) mitral valve replacement with tricuspid valve repair, and 1 (0.9%) aortic valve replacement with tricuspid valve repair. Post-operatively, there were 40 (34.2%) cases of prolonged ventilation, 9 (7.7%) acute kidney injury, 6 (5.1%) re-operations for bleeding, 1 (0.9%) cerebrovascular accident, and 15 (12.8%) cases of atrial fibrillation. The mean total hospital length of stay was 12±12 days, with an in-hospital mortality of 2 (1.7%).

Conclusions: A minimally invasive right thoracotomy approach to primary or re-operative double valve surgery is feasible, may be utilized with acceptable peri-operative morbidity and mortality.

Keywords: Aortic valve replacement; double valve surgery; minimally invasive valve surgery; mitral valve surgery; tricuspid valve repair

Submitted Apr 18, 2017. Accepted for publication May 08, 2017.

doi: 10.21037/jtd.2017.05.62

View this article at: <http://dx.doi.org/10.21037/jtd.2017.05.62>

Introduction

The benefits of minimally invasive valve surgery are well recognized when compared with a standard median sternotomy, in terms of reduced blood loss and less post-operative pain, and reductions in morbidity, hospital lengths of stay, and costs (1-4). Of all the potential benefits, an enhanced recovery with a faster return to normal activity is

the most consistent finding (5). These advantages are seen particularly in high-risk populations, such as elderly and obese patients, and in those who undergoing re-operative valve surgery (6-12). However, data evaluating the feasibility and benefits of minimally invasive double valve surgery are limited (13-15). Our study aim was to analyze the outcomes of minimally invasive double valve surgery, performed via a right thoracotomy approach.

Methods

With Institutional Review Board approval, the institutional Society of Thoracic Surgeons database was retrospectively reviewed from January 2009 to December 2011, to identify patients who underwent minimally invasive repair or replacement of two simultaneous valves (double valve surgery). Excluded were patients in cardiogenic shock, those requiring emergency surgery, patients with prior valve surgery, or those who underwent concomitant cardiac surgical procedures such as coronary artery bypass grafting, surgery of the aorta, or triple valve surgery. We have previously published our results of patients undergoing combined mitral and tricuspid valve surgery, and some of those patients are included in this study (13).

All collected data were reviewed, including pertinent diagnostic testing prior to admission, operative reports, post-operative outcomes and discharge documentation. The total operative times, as well as the intensive care unit and postoperative length of stays were reviewed. The outcome variables evaluated were: prolonged mechanical ventilation (>24 hours), re-intubation, development of acute kidney injury, bleeding requiring re-operation, cerebrovascular accident, sepsis, post-operative atrial fibrillation, and in-hospital mortality.

Our approach to minimally invasive double valve surgery has been previously described in detail, and involves the utilization of a right thoracotomy (14). In summary, cardiopulmonary bypass was established utilizing a femoral platform. For combined aortic and mitral valve surgery, and for combined aortic and tricuspid procedures, a 6–7 cm incision was performed over the 4th intercostal space starting at the mid-clavicular line and extended laterally. For mitral and tricuspid valve surgery, the incision was made at the right 4th to 5th intercostal space, at the anterior axillary line. In the patients with prior coronary artery bypass grafting undergoing mitral and tricuspid valve surgery, moderate-to-deep hypothermia (24–26 °C) was utilized with fibrillatory arrest.

Statistical analysis

All variables are reported as the mean \pm 1 standard deviation (SD), median and interquartile range (IQR, 25–75%), or number (N) and percentage. The statistical software utilized for the analyses was the Statistical Package for Social Sciences, version 21 (SPSS Inc., Chicago, IL, USA).

Results

A total of 117 patients underwent double valve surgery during the study timeframe, consisting of 68 (58.1%) females and 49 (41.9%) males. The mean age and left ventricular ejection fraction were 73 ± 11 years and $52\pm 11\%$, respectively. Forty-three (36.8%) patients had congestive heart failure, 45 (38.5%) had chronic obstructive pulmonary disease, and 5 (4.3%) had a history of chronic kidney disease. There were 11 (9.4%) patients who had undergone prior coronary artery bypass graft surgery (*Table 1*). The types of surgery performed consisted of 50 (42.7%) aortic valve replacement and mitral valve repair, 31 (26.5%) mitral and tricuspid valve repair, 18 (15.4%) aortic and mitral valve replacement, 17 (14.5%) mitral valve replacement and tricuspid valve repair, and 1 (0.9%) aortic valve replacement and tricuspid valve repair. The median aortic cross-clamp and cardiopulmonary bypass times were 110 minutes (IQR 94–136), and 141 minutes (IQR 121–180) (*Table 2*).

No patient required conversion to median sternotomy. Post-operatively, the median intensive care unit length of stay was 70 hours (IQR 46–135). There were 40 (34.2%) cases of prolonged mechanical ventilation, 9 (7.7%) cases of acute kidney injury, 6 (5.1%) re-operations for bleeding, 1 (0.9%) cerebrovascular accident, and 15 (12.8%) cases of atrial fibrillation. The mean total inpatient length of stay was 12 ± 12 days, with an in-hospital mortality of 2 (1.7%) (*Table 3*).

Discussion

Double valve surgery accounts for 10% of valve procedures performed in the United States (16). Patients requiring a multiple valve operation have an operative mortality that is more than twice that of single valve operations, which is cited at 9.7% for patients undergoing combined mitral and tricuspid valve surgery, 10.7% for those undergoing aortic and mitral valve surgery, and 13.2% for those undergoing aortic and tricuspid valve surgery (17). Since a minimally invasive approach has been shown to be associated with a decreased morbidity, shorter hospital lengths of stay, and an enhanced recovery with a faster return to normal activity when compared with median sternotomy valve operations, we hypothesized that higher-risk patients such as those undergoing double valve surgery may benefit from this approach (5,18).

The present study demonstrated an excellent in-hospital

Table 1 Patient baseline characteristics

Variables	n=117
Age (mean \pm SD)	73 \pm 11
Males	49 (41.9%)
Ejection fraction (mean \pm SD)	52 \pm 11%
Hypertension	108 (92.3%)
Diabetes mellitus	38 (32.5%)
Renal failure	5 (4.3%)
Chronic obstructive pulmonary disease	45 (38.5%)
Coronary artery disease	39 (33.3%)
Previous coronary artery bypass surgery	11 (9.4%)
Previous percutaneous coronary intervention	22 (18.8%)
Congestive heart failure	43 (36.8%)
History of endocarditis	8 (7%)
Pulmonary artery systolic pressure (mean \pm SD)	37 \pm 14
Pre-operative creatinine (mg/dL, mean \pm SD)	1.17 \pm 0.9
Pre-operative Hematocrit (mean \pm SD)	38 \pm 5

SD, standard deviation.

Table 2 Operative characteristics

Variables	n=117
Aortic valve replacement with mitral valve repair	50 (42.7%)
Aortic and mitral valve replacement	18 (15.4%)
Mitral valve replacement with tricuspid valve repair	17 (14.5%)
Mitral and tricuspid valve repair	31 (26.5%)
Aortic valve replacement with tricuspid valve repair	1 (0.9%)
Aortic cross-clamp time (minutes, median, IQR)	110 [94–136]
Cardiopulmonary bypass time (minutes, median, IQR)	141 [121–180]

IQR, interquartile range.

mortality of 1.7% utilizing a minimally invasive right thoracotomy in patients requiring double valve surgery. It also demonstrated acceptable outcomes in the incidence of post-operative cerebrovascular accidents, renal failure, and re-operation for bleeding. The most common post-operative complication noted was the incidence of prolonged mechanical ventilation, which occurred in 34% of the patients. This may be explained by the relatively high prevalence of chronic obstructive pulmonary disease in the

Table 3 Post-operative outcomes

Variables	n=117
Ventilation time (hours, median, IQR)	21 [18–43]
Intensive care unit length of stay (hours, median, IQR)	70 [46–135]
Prolonged mechanical ventilation	40 (34.2%)
Re-intubation	12 (10.3%)
Acute kidney injury	9 (7.7%)
Re-operation for bleeding	6 (5.1%)
Cerebrovascular accident	1 (0.9%)
Sepsis	3 (2.6%)
Atrial fibrillation	15 (12.8%)
Post-operative hospital length of stay (days, mean \pm SD)	12 \pm 12
In-hospital mortality	2 (1.7%)

SD, standard deviation; IQR, interquartile range.

cohort, as well as the timeframe in which the study took place. Following this institutional era, we implemented several changes in our post-cardiac surgery extubation practices, including standardized multidisciplinary ventilation weaning protocols. As a result, we observed a significant reduction in the length of mechanical ventilation and intensive care unit stay in subsequent analyses, in which a minimally invasive approach often proved superior to median sternotomy (6-9).

Some authors have expressed concern regarding the possibility of having lower repair rates of the mitral valve with minimally invasive valve surgery (19). This was not found to be an issue in the present study, as 80 (69%) of the 116 patients who underwent a mitral valve operation were able to have their mitral valve repaired. This repair rate of 69% is higher than the 57.4% reported by the Society of Thoracic Surgeons (20). This issue is important because mitral valve repair does not only demonstrate a lower mortality than replacement for isolated mitral valve surgery, but it is also associated with a lower mortality when performed in those undergoing multiple valve surgery, such as the cohort of the present study (21-23).

Another concern with minimally invasive valve surgery is obtaining adequate exposure, and this is especially true in patients undergoing double valve surgery and re-operative valve surgery. In circumstances where adequate exposure cannot be obtained, the minimally invasive surgery may need

to be converted to a standard median sternotomy. Although data for double valve surgery are scarce, the conversion rate of minimally invasive surgery to median sternotomy is 1.0–4.0% during single valve operations (24,25). In the present study, even though 9.4% underwent re-operative surgery, adequate exposure was obtained in all patients, and none required conversion to a median sternotomy.

Data involving minimally invasive double valve surgery are somewhat limited. A propensity-matched analysis by Atik *et al.*, compared the outcomes of 162 patients undergoing aortic and mitral valve surgery via a right thoracotomy (N=81) or conventional median sternotomy (N=81) (13). No difference was found in the outcomes of operative mortality (6.2% versus 2.5%, $P=0.4$), acute kidney injury (4.9% versus 1.2%, $P=0.4$), cerebrovascular accidents (2.5% versus 2.5%, $P=1.0$), or re-operation for bleeding (8.6% versus 4.9%, $P=0.5$). Survival at 10 years was also similar, being 82% and 76% for the minimally invasive and median sternotomy group, respectively ($P=0.7$). The authors concluded that minimally invasive surgery had cosmetic and blood product use advantages over conventional surgery, with no apparent detriments. Pfannmüller *et al.* reported their experience with 441 consecutive patients who underwent mitral and tricuspid valve surgery over a 10-year period, with the utilization of a right thoracotomy approach (14). Post-operative outcomes included 37 (8%) re-operations for bleeding, 10 (2%) cerebrovascular accidents, and 19 (4%) deaths at 30 days. The actuarial survival at 5 years was $77.2 \pm 2.5\%$. Finally, Mihos *et al.* evaluated the outcomes of 132 patients who had combined mitral and tricuspid valve surgery via a right thoracotomy, of which 12% underwent re-operative double valve surgery (15). Post-operative outcomes included 6 (5%) cases of acute kidney injury, 6 (5%) re-operations for bleeding, 4 (3%) cerebrovascular accidents, and 5 (4%) in-hospital deaths. The 1- and 5-year survival rates were 93% and 88%, respectively.

The current study is subject to the limitations inherent in a single center, retrospective design. Firstly, the study cohort is made up of a heterogeneous group of patients, with the results being limited to in-hospital outcomes. Secondly, the types of valve operations performed varied, and the etiologies of the valve dysfunction were not available. These are important factors that typically present in distinct demographic groups, and are associated with differing clinical and surgical risk profiles. What impact these variables may have had on the observed outcomes is unknown. Thirdly, there is no median sternotomy

control group, which limits the conclusions that can be drawn from the current data. Finally, all of the operations were performed via a right thoracotomy approach, and cannot be extrapolated to represent outcomes that may be obtained from other minimally invasive approaches. The results should be viewed as an observational study of a consecutive series of minimally invasive right thoracotomy double valve surgery, and as supportive of the hypothesis that this approach may be an alternative option to median sternotomy.

In summary, in patients requiring primary, or re-operative, double valve surgery, a minimally invasive right thoracotomy approach may be utilized with an acceptable peri-operative morbidity and mortality. It may be considered as a feasible alternative to the standard median sternotomy approach.

Acknowledgements

None.

Footnote

Conflicts of Interest: The authors have no conflicts of interest to declare.

Ethical Statement: This retrospective study was approved by the Institutional Review Board at the Mount Sinai Medical Center, Miami Beach, Florida, USA.

References

1. Cohn LH, Adams DH, Couper GS, et al. Minimally invasive cardiac valve surgery improves patient satisfaction while reducing costs of cardiac valve replacement and repair. *Ann Surg* 1997;226:421-6; discussion 427-8.
2. Phan K, Xie A, Di Eusano M, et al. A meta-analysis of minimally invasive versus conventional sternotomy for aortic valve replacement. *Ann Thorac Surg* 2014;98:1499-511.
3. Cheng DC, Martin J, Lal A, et al. Minimally invasive versus conventional open mitral valve surgery: a meta-analysis and systematic review. *Innovations (Phila)* 2011;6:84-103.
4. Santana O, Larrauri-Reyes M, Zamora C, et al. Is a minimally invasive approach for mitral valve surgery more cost-effective than median sternotomy? *Interact Cardiovasc Thorac Surg* 2016;22:97-100.
5. Schmitto JD, Mokashi SA, Cohn LH. Minimally-invasive valve surgery. *J Am Coll Cardiol* 2010;56:455-62.

6. Lamelas J, Sarria A, Santana O, et al. Outcomes of minimally invasive valve surgery versus median sternotomy in patients age 75 years or greater. *Ann Thorac Surg* 2011;91:79-84.
7. Santana O, Reyna J, Grana R, et al. Outcomes of minimally invasive valve surgery versus standard sternotomy in obese patients undergoing isolated valve surgery. *Ann Thorac Surg* 2011;91:406-10.
8. Santana O, Reyna J, Benjo AM, et al. Outcomes of minimally invasive valve surgery in patients with chronic obstructive pulmonary disease. *Eur J Cardiothorac Surg* 2012;42:648-52.
9. Mihos CG, Santana O, Lamas GA, et al. Outcomes of right minithoracotomy mitral valve surgery in patients with previous sternotomy. *Ann Thorac Surg* 2011;91:1824-7.
10. Moscarelli M, Fattouch K, Casula R, et al. What Is the Role of Minimally Invasive Mitral Valve Surgery in High-Risk Patients? A Meta-Analysis of Observational Studies. *Ann Thorac Surg* 2016;101:981-9.
11. Moscarelli M, Margaryan R, Cerillo A, et al. Minimally Invasive Mitral Valve Surgery in Truly High-Risk Patients: Are We Pushing the Boundaries?: An Observational Study. *Innovations (Phila)* 2015;10:328-33.
12. Pineda AM, Santana O, Lamas GA, et al. Is a minimally invasive approach for re-operative aortic valve replacement superior to standard full resternotomy? *Interact Cardiovasc Thorac Surg* 2012;15:248-52.
13. Atik FA, Svensson LG, Blackstone EH, et al. Less invasive versus conventional double-valve surgery: a propensity-matched comparison. *J Thorac Cardiovasc Surg* 2011;141:1461-8.e4.
14. Pfannmüller B, Davierwala P, Hirnle G, et al. Concomitant tricuspid valve repair in patients with minimally invasive mitral valve surgery. *Ann Cardiothorac Surg* 2013;2:758-64.
15. Mihos CG, Pineda AM, Davila H, et al. Combined Mitral and Tricuspid Valve Surgery Performed via a Right Minithoracotomy Approach. *Innovations (Phila)* 2015;10:304-8.
16. Lee R, Li S, Rankin JS, et al. Fifteen-year outcome trends for valve surgery in North America. *Ann Thorac Surg* 2011;91:677-84; discussion p 684.
17. Vassileva CM, Li S, Thourani VH, et al. Outcome characteristics of multiple-valve surgery: comparison with single-valve procedures. *Innovations (Phila)* 2014;9:27-32.
18. Santana O, Xydias S, Williams RF, et al. Minimally invasive valve surgery in high-risk patients. *J Thorac Dis* 2017. [Epub ahead of print].
19. Anyanwu AC, Adams DH. Should complex mitral valve repair be routinely performed using a minimally invasive approach? *Curr Opin Cardiol* 2012;27:118-24.
20. Badhwar V, Rankin JS, He X, et al. The Society of Thoracic Surgeons Mitral Repair/Replacement Composite Score: A Report of The Society of Thoracic Surgeons Quality Measurement Task Force. *Ann Thorac Surg* 2016;101:2265-71.
21. D'Agostino RS, Jacobs JP, Badhwar V, et al. The Society of Thoracic Surgeons Adult Cardiac Surgery Database: 2017 Update on Outcomes and Quality. *Ann Thorac Surg* 2017;103:18-24.
22. Rankin JS, He X, O'Brien SM, et al. The Society of Thoracic Surgeons risk model for operative mortality after multiple valve surgery. *Ann Thorac Surg* 2013;95:1484-90.
23. Gillinov AM, Blackstone EH, Cosgrove DM 3rd, et al. Mitral valve repair with aortic valve replacement is superior to double valve replacement. *J Thorac Cardiovasc Surg* 2003;125:1372-87.
24. Vollroth M, Seeburger J, Garbade J, et al. Minimally invasive mitral valve surgery is a very safe procedure with very low rates of conversion to full sternotomy. *Eur J Cardiothorac Surg* 2012;42:e13-5; discussion e16.
25. Tabata M, Umakanthan R, Khalpey Z, et al. Conversion to full sternotomy during minimal-access cardiac surgery: reasons and results during a 9.5-year experience. *J Thorac Cardiovasc Surg* 2007;134:165-9.

Cite this article as: Santana O, Xydias S, Williams RF, LaPietra A, Mawad M, Hasty F, Escolar E, Mihos CG. Outcomes of minimally invasive double valve surgery. *J Thorac Dis* 2017;9(Suppl 7):S602-S606. doi: 10.21037/jtd.2017.05.62