



Published in final edited form as:

*Eur Respir J*. 2017 May ; 49(5): . doi:10.1183/13993003.01505-2016.

## A meta-analysis of genome-wide association studies of asthma in Puerto Ricans

Qi Yan, PhD<sup>1</sup>, John Brehm, MD, MPH<sup>1</sup>, Maria Pino-Yanes, PhD<sup>2,3</sup>, Erick Forno, MD, MPH<sup>1</sup>, Jerome Lin<sup>4</sup>, Sam S. Oh, PhD<sup>5</sup>, Edna Acosta-Perez, PhD<sup>6</sup>, Cathy C. Laurie, PhD<sup>7</sup>, Michelle M. Cloutier, MD<sup>8</sup>, Benjamin A. Raby, MD, MPH<sup>9</sup>, Adrienne M. Stilp, PhD<sup>7</sup>, Tamar Sofer, PhD<sup>7</sup>, Donglei Hu, PhD<sup>5</sup>, Scott Huntsman<sup>5</sup>, Celeste S. Eng<sup>5</sup>, Matthew P. Conomos, PhD<sup>7</sup>, Deepa Rastogi, MD<sup>10</sup>, Kenneth Rice, PhD<sup>7</sup>, Glorisa Canino, PhD<sup>6</sup>, Wei Chen, PhD<sup>1</sup>, R. Graham Barr, MD, DrPH<sup>11,13</sup>, Esteban G. Burchard, MD, MPH<sup>5,12,13</sup>, and Juan C. Celedón, MD, DrPH<sup>1,13,\*</sup>

<sup>1</sup>Division of Pediatric Pulmonary Medicine, Allergy, and Immunology, Children's Hospital of Pittsburgh of UPMC, University of Pittsburgh, Pittsburgh, PA 15224, USA

<sup>2</sup>CIBER de Enfermedades Respiratorias, Instituto de Salud Carlos III, Madrid, Spain

<sup>3</sup>Research Unit, Hospital Universitario N.S. de Candelaria, Universidad de La Laguna, Santa Cruz de Tenerife, Spain

<sup>4</sup>Department of Human Genetics, Graduate School of Public Health, University of Pittsburgh, Pittsburgh, PA 15261, USA

<sup>5</sup>Department of Medicine, University of California San Francisco, San Francisco, CA 94143, USA

<sup>6</sup>Behavioral Sciences Research Institute, University of Puerto Rico, San Juan, Puerto Rico

<sup>7</sup>Department of Biostatistics, University of Washington, Seattle, WA 98195, USA

<sup>8</sup>Department of Pediatrics, University of Connecticut, Farmington, CT 06030, USA

<sup>9</sup>Channing Division of Network Medicine, Department of Medicine, Brigham and Women's Hospital, Harvard Medical School, MA 02115, USA

<sup>10</sup>Department of Pediatrics, Albert Einstein College of Medicine, Bronx, NY 10461, USA

<sup>11</sup>Department of Epidemiology, Columbia University, New York, NY 10027, USA

<sup>12</sup>Department of Bioengineering and Therapeutic Sciences, University of California San Francisco, San Francisco, CA 94143, USA

### Abstract

**Corresponding author:** Juan C. Celedón, MD, DrPH, Division of Pulmonary Medicine, Allergy and Immunology, Children's Hospital of Pittsburgh of UPMC, 4401 Penn Avenue, Pittsburgh, PA 15224, Phone: 412.692.8429; Fax 412.692.7636, [juan.celedon@chp.edu](mailto:juan.celedon@chp.edu).

<sup>13</sup>These authors contributed equally to this work

**Author Contributions:** Conception and study design: G.R.B., E.G.B., and J.C.C.; Data acquisition, analysis and interpretation: Q.Y., M.P.Y., J.B., E.F., J.L., S.O., E.A-P., C.L., M.M.C., B.A.R., A.S., T.S., D.H., S.H., C.E., M.P.C., D.R., K.R., G.C., and W.C.; and drafting of the manuscript for intellectual content: Q.Y., R.G.B., E.G.B., and J.C.C. All authors approved the final version of the manuscript prior to submission

**Rationale**—No genome-wide association study (GWAS) of asthma has been conducted in Puerto Ricans.

**Objective**—To identify susceptibility genetic variants for asthma in Puerto Ricans.

**Methods**—We conducted a meta-analysis of GWAS of asthma, including Puerto Rican participants from: GALA I-II, the Hartford-Puerto Rico Study, and the Hispanic Community Health Study. Moreover, we examined whether susceptibility loci identified in previous meta-analyses of GWAS are associated with asthma in Puerto Ricans.

**Results**—The only locus to achieve a genome-wide significant association with asthma in an analysis of 2,144 cases and 2,893 controls was chromosome 17q21, as evidenced by our top SNP, rs907092 (OR = 0.71,  $P = 1.2 \times 10^{-12}$ ) on *IKZF3*. Similar to findings in non-Puerto Ricans, SNPs in genes in the same LD block as *IKZF3* (e.g. *ZBP2*, *ORMDL3* and *GSDMB*) were also significantly associated with asthma in Puerto Ricans. With regard to results from a meta-analysis in Europeans, we replicated findings for the SNP at *GSDMB*, but not for SNPs in any other genes. On the other hand, we replicated results from a meta-analysis of North American populations for SNPs in *IL1RL1*, *TSLP* and *GSDMB* but not for *IL33*.

**Conclusions**—Common variants on chromosome 17q21 have the greatest effects on asthma in Puerto Ricans, a high-risk ethnic group.

## Keywords

GWAS; asthma; Puerto Ricans; chromosome 17q21

## INTRODUCTION

Asthma is a disease with substantial heritability.<sup>(1, 2)</sup> Genome-wide association studies (GWAS) have identified several susceptibility loci for asthma, including the chromosome 17q21 region.<sup>(3, 4)</sup> This locus, which contains genes *IKZF3*, *ZBP2*, *GSDMB* and *ORMDL3*, has been consistently replicated across diverse ethnic groups.<sup>(3–13)</sup>

The burden of asthma varies among racial or ethnic groups. In the United States, current asthma is more common in Puerto Ricans (16.1%) than in non-Hispanic blacks (11.2%), non-Hispanic whites (7.7%) or Mexicans (5.4%).<sup>(14)</sup> Moreover, Puerto Ricans have greater morbidity from asthma than members of other racial or ethnic groups,<sup>(15)</sup> and thus studying asthma in this ethnic group is relevant to public health and understanding disease pathogenesis. Although two previous GWAS of asthma included Puerto Ricans who participated in the Genetics of Asthma in Latino Americans Study (GALA I)<sup>(3)</sup> and in the Genes-environments & Admixture in Latino Americans Study (GALA II)<sup>(13)</sup>, no separate analysis was presented for Puerto Ricans.

We conducted a meta-analysis of GWAS of asthma, including only Puerto Rican participants from: GALA I, GALA II, the Hartford-Puerto Rico Study (Hartford-PR)<sup>(16)</sup> and the Hispanic Community Health Study/Study of Latinos (HCHS/SOL).<sup>(17, 18)</sup> Moreover, we examined whether susceptibility loci identified in two previous meta-analyses of GWAS of asthma (conducted by the GABRIEL and EVE consortiums<sup>(3, 4)</sup>) are associated with asthma in Puerto Ricans.

## METHODS

### Study Populations

**Hartford-PR**—Children were recruited in Hartford, CT (n=449) and San Juan, PR (n=678), as reported elsewhere.<sup>(16)</sup> At both study sites, the main recruitment tool was a screening questionnaire given to parents of children ages 6 to 14 years. All participants had to have four Puerto Rican grandparents. Asthma was defined as physician-diagnosed asthma and at least one episode of wheeze in the prior year. Control subjects had neither physician-diagnosed asthma nor wheeze in the prior year. Genome-wide genotyping was conducted using the HumanOmni2.5 BeadChip platform (Illumina Inc., San Diego, CA), as previously described.<sup>(16)</sup> Imputation of non-genotyped single nucleotide polymorphisms (SNPs) was performed with IMPUTE2,<sup>(19)</sup> using data from the Phase I (November 2010 release) of the 1000 Human Genomes Project as the reference panel. After quality control measures, 948 children (523 with asthma) and ~7 million genotyped and imputed SNPs were included in the genome-wide association analysis. This analysis was conducted using logistic regression under an additive genetic model, adjusting for age, gender, study site and principal components (PCs) calculated using smartPCA.<sup>(20)</sup> Written parental consent was obtained for participating children, from whom written assent was also obtained. The study was approved by the Institutional Review Boards of the University of Puerto Rico (San Juan, PR), Brigham and Women's Hospital (Boston, MA) and the University of Pittsburgh (Pittsburgh, PA).

**GALA I**—This study comprised subjects with and without asthma, recruited mainly from Puerto Rico, but also from New York City, NY.<sup>(21, 22)</sup> Subjects were included in the study if they were aged 8 to 40 years and self-identified all four grandparents as Puerto Rican. Cases had physician-diagnosed asthma and had experienced two or more symptoms in the previous two years (wheezing, coughing, and/or shortness of breath). Control subjects had no symptoms of asthma or allergies, had no physician-diagnosis of asthma, and no history of other chronic respiratory illness or allergies (eczema, hives, or hay fever). All subjects were genotyped on the Affymetrix 6.0 GeneChip (Affymetrix, Santa Clara, CA), and quality control was performed as described elsewhere.<sup>(23)</sup> Genotyped data was phased with SHAPE-IT<sup>(24)</sup> and imputation was performed with IMPUTE2,<sup>(19)</sup> using all populations from 1000 Genomes Project Phase I v3<sup>(25)</sup> as reference. Association testing was conducted using logistic regression under an additive genetic model, adjusting for age, gender, and African and Native American ancestry estimates (calculated using ADMIXTURE<sup>(26)</sup>). A total of 437 unrelated subjects (251 with asthma) with complete data for all covariates were used in the current analysis. Written parental consent was obtained for participating children, from whom written assent was also obtained. The study was approved by the Institutional Review Boards of the University of California at San Francisco (UCSF) and at each participating center.

**GALA II**—This is a case-control study of asthma in Latino children.<sup>(27)</sup> Cases and control subjects were recruited using a combination of community and clinic-based approaches from centers throughout the U.S. (Chicago [IL], Bronx [NY], Houston [TX], San Francisco Bay Area [CA] and Puerto Rico). Subjects were eligible if they were aged 8 to 21 years, had <10 pack-years of smoking history and were not current smokers, and self-reported having four

grandparents of Latino ethnicity. Asthma was defined based on physician diagnosis and report of symptoms and medication use within the last two years. Control subjects had no history of asthma or allergies, and no wheeze or shortness of breath during their lifetime. The current analysis focused on 1,786 participants (892 with asthma) with self-reported Puerto Rican ethnicity. All subjects were genotyped on the Axiom<sup>®</sup> LAT1 array (World Array 4, Affymetrix, Santa Clara, CA), and quality control was performed as described elsewhere.<sup>(28)</sup> Imputation and association testing was performed as described above for GALA I. The study was approved by the Institutional Review Boards of UCSF and at each participating center. All subjects and their parents provided written informed assent and written informed consent, respectively.

**HCHS/SOL**—This is a community-based cohort study of self-identified Hispanic/Latino individuals aged 18–74 years, who were recruited at four centers (Chicago, IL; Miami, FL; Bronx, NY; San Diego, CA). Participants were recruited using a two-stage sampling scheme, in which census block units were sampled first, and then household and individuals from these units.<sup>(17, 18)</sup> The HCHS/SOL study was approved by institutional review boards at participating institutions, and written informed consent was obtained from all participants.

The HCHS/SOL includes individuals who self-identified as Mexican, Central American, South American, Puerto Rican, Dominican or Cuban. However, genetic analysis groups were constructed based on a combination of these self-identified Hispanic/Latino background and genetic similarity. These genetic analysis groups largely overlap with the self-identified groups, but using the genetic analysis groups in association testing and stratified analyses has advantages, as previously described.<sup>(29)</sup> Individuals from the Puerto Rican genetic analysis group were used in the current analysis. Cases (n=478) had self-reported current physician-diagnosed asthma. Control subjects (n=1,388) had never been diagnosed with asthma. Individuals were genotyped at Illumina on the HCHS/SOL custom 15041502 B3 array, and imputed to 1000 Human Genomes Phase I data. Details about genotyping, imputation, and quality control are provided elsewhere.<sup>(29)</sup>

Association analysis was performed using GMMAT,<sup>(30)</sup> a mixed-model logistic regression that accounts for correlations due to relatedness, shared household and block group. The analysis was adjusted for five genetic PCs, study center, sampling weights (to prevent potential selection bias resulting from the sampling scheme), age, gender, smoking status, and pack-years of cigarette smoking.

In our primary meta-analysis, we included three cohorts of children (Hartford-PR, GALA I, and GALA II) and one cohort of adults (HCHS/SOL) for maximum sample size. Since childhood asthma likely has genetic determinants that differ from those for adult-onset asthma, we repeated the meta-analysis after excluding HCHS/SOL.

## Statistical Methods

METAL<sup>(31)</sup> software was used to perform the meta-analysis of the four GWAS of asthma. METAL takes *P*-values across independent studies as input, with sample size and effect direction taken into account. First, for each SNP, the coded and alternative alleles are determined, and a Z-score is calculated based on the *P*-values and direction of effect in each

study. Specifically, large positive  $Z$ -scores indicate small  $P$ -values where the coded allele is the risk allele, and large negative  $Z$ -scores indicate small  $P$ -values where the coded allele is protective. Formally the  $Z$ -score is

$$Z_i = \Phi^{-1} \left( 1 - \frac{P_i}{2} \right) \times \text{sign}(\Delta_i),$$

where  $Z_i$  is the  $Z$ -score for study  $i$ ,  $P_i$  is the  $P$ -value for study  $i$ ,  $\Delta_i$  is the direction of effect for study  $i$ , and  $\Phi^{-1}$  gives the percentile of a standard normal distribution. Then, the overall  $Z$ -score and  $P$ -value are calculated from a weighted sum of the individual  $Z$ -scores,

$$Z = \frac{\sum_i Z_i w_i}{\sqrt{\sum_i w_i^2}},$$

$$P = 2\Phi(|Z|)$$

where  $Z$  is the overall  $Z$ -score,  $P$  is the overall  $P$ -value, and  $w_i$  is the weight for study  $i$ ,

$$w_i = \frac{\text{MAF}_i(1 - \text{MAF}_i)N_i^{\text{cas}}N_i^{\text{con}}}{(N_i^{\text{cas}} + N_i^{\text{con}})}$$

where  $\text{MAF}$  is the minor allele frequency for study  $i$ ,  $N_i^{\text{cas}}$  is the number of cases for study  $i$  and  $N_i^{\text{con}}$  is the number of controls for study  $i$ . This weighting is intended to closely approximate the results that would be obtained combining subject-level data across the studies, in an analysis that adjusts for study. Summary odds ratios (ORs) were calculated by averaging the study-specific log-odds ratios, with weights reflecting the standard errors from the study-specific ORs.

We used the varLD<sup>(32, 33)</sup> software to assess whether there were statistically significant differences in regional linkage disequilibrium (LD) structures between our study populations. VarLD uses a Monte Carlo approach to calculate  $P$ -values, which were based on 10,000 permutations. For a comparison of two populations, a  $P < 0.05$  indicates a significant difference in LD structures.

## RESULTS

The characteristics of participants in each of the four studies included in the meta-analysis are shown in Table 1. The results reported here are calculated based on 7,485,508 million imputed and genotyped SNPs in 2,144 subjects with asthma and 2,893 control subjects.

QQ plots for the meta-analysis (Figures S1 and S3) show that neither the results from each of the four component studies nor the combined results are inflated in their test statistics. Moreover, these plots reveal an abundance of small  $P$ -values in the combined study. We expected to see SNPs with  $P$ -value  $< 5 \times 10^{-8}$ , the standard significance threshold for GWAS.

In the combined study, we found that 89 SNPs on chromosome 17q21 are significant at this level (Figure 1 and Table S1). Of these 89 SNPs, rs907092 on *IKZF3* showed the most significant association with asthma ( $P = 1.2 \times 10^{-12}$ , Figures 2 and 3). This SNP is also associated with asthma at  $P < 0.023$  in all three studies of children and at  $P = 0.092$  in HCHS/SOL, a study of adults (Figures 2 and S6).

Given our results for chromosome 17q21 and asthma, particularly in studies including children, we examined the LD pattern of this region among Puerto Ricans in Hartford-PR, GALA I and GALA II. Figures S8–S10 show that subjects in these studies have similar LD patterns (varLD  $P$ -value = 0.08 for Hartford-PR vs. GALA I;  $P$ -value = 0.46 for Hartford-PR vs. GALA II [see Table S2]), and that nearly all significant SNPs gather in an LD bin spanning ~200 Kb (see Figures S8–S10). The *IKZF3-ZBP2-GSDMB-ORMDL3* locus is included in this bin. We then compared the LD pattern of chromosome 17q21 in Puerto Ricans to that of other ethnic groups (Figure S11). Puerto Ricans have an LD pattern that differs from that in Mexicans or Europeans (varLD  $P$ -value = 0.0007 for Hartford-PR vs. Mexicans (in GALA I or GALA II) or 1000G Europeans [see Table S2]). For this analysis, Mexican genotypes were extracted from GALA I and GALA II, and European genotypes were extracted from the 1000 Genomes Project.

Next, we examined whether SNPs previously associated with asthma in either a meta-analysis of GWAS of Europeans<sup>(4)</sup> or a meta-analysis of ethnically diverse North American populations<sup>(3)</sup> were also associated with asthma in Puerto Ricans (Table 2). For the meta-analysis in Europeans, we were able to replicate associations for the SNP at *GSDMB* at  $P < 0.005$  (Bonferroni correction for 10 tests). For the meta-analysis of ethnically diverse populations, we were able to replicate the association with SNPs at *IL1RL1*, *TSLP* and *GSDMB* at  $P < 0.01$  (Bonferroni correction for 5 tests). The two SNPs (rs2305480 and rs11078927) at *GSDMB* identified in the two cited studies consistently have small  $P$ -values in Hartford-PR, GALA I, GALA II, as well as in the meta-analysis. We did not replicate associations with SNPs in *IL33*, reported by both previous meta-analyses. Since this may be caused by differences in LD patterns in or near *IL33* between Puerto Ricans and ethnic groups included in the prior meta-analyses (e.g., Europeans and Mexicans, Figure S12), we also show the top *IL33* SNPs from our meta-analysis in Table S3. In this analysis, no *IL33* SNP was significantly associated with asthma ( $P > 0.00035$  in all instances, Bonferroni correction for 142 tests).

Since childhood asthma likely has genetic determinants that differ from those of adult-onset asthma, we repeated the meta-analysis of GWAS of asthma without HCHS/SOL, obtaining similar results (Figures S4, S5 and S7) to those from the meta-analysis including HCHS/SOL: only SNPs on chromosome 17q21 reached a significance level of  $P < 5 \times 10^{-8}$ . Moreover, we were able to replicate the same SNPs from the two previous meta-analyses of GWAS of asthma in the meta-analysis excluding HCHS/SOL (Table S4). In addition, SNPs at *IL18R1* and *GSDMA* were replicated at  $P < 0.005$  (Bonferroni correction for 10 tests).

In order to check if there were independent SNPs associated with asthma, we performed conditional analyses in chromosome 17q21 by adjusting for the top SNP in our analysis (rs907092). Since adults did not show a strong genetic association in this region, we



conducted the conditional analyses in a meta-analysis of the three cohorts of children. The results showed that no SNPs in chromosome 17q21 reached the Bonferroni-corrected significance level,  $\alpha=4.1\times 10^{-5}$  (1,215 overlapped SNPs with  $MAF > 0.05$  across the three cohorts) and no SNPs in the *IKZF3-ZPBP2-GSDMB-ORMDL3* locus reached the Bonferroni-corrected significance level,  $\alpha=2.5\times 10^{-4}$  (197 overlapped SNPs with  $MAF > 0.05$  across the three cohorts). Thus, in addition to SNPs in LD with rs907092, there seemed to be no other independent signals in chromosome 17q21.

## DISCUSSION

We report findings from the first GWAS of asthma conducted solely in Puerto Ricans. Although two of the cohorts included in the current meta-analysis (GALA I and II) were part of Hispanic/Latino cohorts included in previous studies, those studies did not separately analyze data from Puerto Ricans but instead combined data from Puerto Ricans and Mexicans. Such approach can reduce statistical power, since Puerto Ricans and Mexicans differ markedly with regard to racial ancestry (i.e. Puerto Ricans have, on average, a greater proportion of African ancestry but a lower proportion of Native American ancestry than Mexicans), asthma burden (as Mexicans have a much lower asthma burden than Puerto Ricans), and allelic frequencies.<sup>(15)</sup>

In our meta-analysis of four GWAS, the only locus to achieve a genome-wide significant association with asthma was chromosome 17q21, as evidenced by our top SNP, rs907092 ( $P = 1.2\times 10^{-12}$ ), located on *IKZF3*. Similar to previous findings in non-Puerto Ricans, SNPs in genes located in the same LD block as *IKZF3* on chromosome 17q21 (e.g. *ZPBP2*, *ORMDL3* and *GSDMB*) were also significantly associated with asthma in our meta-analysis. A previous candidate-gene study conducted in 399 Puerto Ricans and 301 Mexicans in GALA I, as well as 261 African Americans, found significant associations between two SNPs in *ORMDL3* (rs4378650 and rs12603332) and asthma in Mexicans and African Americans, but not in Puerto Ricans ( $P=0.08$  for both SNPs).<sup>(6)</sup> In our meta-analysis, these two SNPs ( $P = 4.0\times 10^{-6}$  for rs4378650 and  $P = 5.5\times 10^{-6}$  for rs12603332) were replicated for an association with asthma.

Consistent with prior findings suggesting stronger effects for SNPs in chromosome 17q21 on childhood asthma than in adult asthma among Europeans<sup>(4)</sup>, we found genome-wide significant results for this locus in a meta-analysis restricted to the three cohorts of children, but not in the GWAS of asthma among adults in HCHS/SOL.

We also attempted to replicate previous findings for potential asthma-susceptibility SNPs from meta-analyses conducted in Europeans (by the GABRIEL consortium)<sup>(4)</sup> and ethnically diverse North American populations (by the EVE consortium).<sup>(3)</sup> With regard to results from GABRIEL, we replicated findings for the SNP at *GSDMB* (on chromosome 17q21), but not for SNPs in any other gene. On the other hand, we report the first replication of results from the multi-ethnic EVE consortium for SNPs in *IL1RL1*, *TSLP* and *GSDMB* in a cohort composed exclusively of Puerto Ricans. Our lack of replication of most findings from GABRIEL in Puerto Ricans mimics negative findings from the EVE consortium, and

may be due to ethnic differences in risk variants for asthma. In fact, EVE only replicated findings for chromosome 17q21 and *IL33* (albeit for different SNPs) from GABRIEL.

We did not replicate results for *PYHIN1* or *IL33* from EVE. *PYHIN1* was only associated with asthma among African-Americans and Afro-Caribbeans, and ancestry differences may account for our negative results. For SNPs with MAF = 0.37 (the MAF of *IL33* SNP rs2381416 in our study cohorts), we had 70% statistical power to detect an OR = 1.10 at  $\alpha = 0.01$  in our meta-analysis.<sup>(34)</sup> Under the same assumptions, the statistical power to detect an OR = 1.20 at  $\alpha = 0.01$  was > 99% in our meta-analysis. Of the 142 SNPs in *IL33* that were tested for association with asthma in the current study, only 6 were associated at  $P < 0.01$ , and none remained significantly associated with asthma after a Bonferroni correction.

Expression of the 17q21 gene, *ORMDL3*, is regulated by asthma-associated SNPs,<sup>(35)</sup> as replicated in our previous eQTL study (rs8067378,  $P = 2.6 \times 10^{-10}$ ) using the Hartford-PR data.<sup>(36)</sup> Lately, *ORMDL3* has been implicated in various cellular processes that could be relevant to asthma.<sup>(35)</sup> In addition to its effects on asthma susceptibility, the study from Tavendale *et al.* also showed that *ORMDL3* is associated with asthma exacerbations<sup>(11)</sup> and that the *ORMDL3* SNP-mediated expression is affected by rhinovirus infection, a common trigger of such exacerbations.<sup>(37)</sup> The SNP in the *IKZF3-ZBP2-GSDMB-ORMDL3* region that was associated with asthma in the EVE meta-analysis was rs11078927 in *GSDMB*, which is replicated in our study ( $P = 8.1 \times 10^{-12}$  [see Table 2]). In EVE, the reported ORs for this SNP and asthma were 0.80 in European Americans and 0.78 in Latinos (non-significant in African Americans). In our study, we estimated ORs of 0.72 in all subjects and 0.66 in children only, suggesting a stronger effect of the *IKZF3-ZBP2-GSDMB-ORMDL3* locus on childhood asthma in Puerto Ricans than in other ethnic groups. Thus, the effects of the major allele for SNP rs11078927 (i.e. OR of 1.51 [1/0.66] in children) may contribute to asthma etiology in Puerto Ricans.

The *ZBP2-GSDMB-ORMDL3* locus was detected to have allelic differences in expression levels in lymphoblastoid cell lines (LCLs) and CD4+ cells.<sup>(38, 39)</sup> Early studies of gene expression<sup>(40–42)</sup> revealed that the asthma-associated SNPs regulate the expression of *ORMDL3* and *GSDMB*, and that these two genes might be coregulated.<sup>(38)</sup> Findings from a prior study suggest that an asthma-associated 17q21 regulatory haplotype affects transcriptional activity of *ZBP2*, *GSDMB*, and *ORMDL3*, with one non-coding variant in *ZBP2*, rs12936231, differentially influencing the binding of the insulator protein CTCF in an allele-specific manner;<sup>(38)</sup> subsequent eQTL studies showed that SNP rs12936231 increases *ORMDL3* and *GSDMB* expression in primary lymphocytes, whole blood, and lung tissue.<sup>(43, 44)</sup> However, SNP rs12936231 (which has a minor allele frequency=0.44–0.46 in our cohorts) was non-significantly associated with asthma in our meta-analysis ( $P = 2.0 \times 10^{-6}$ ) and is only in moderate LD ( $r^2 > 0.51$ ) with our top SNP (rs907092) in *IKZF3*. This suggests that other functional polymorphisms on chromosome 17q21 may affect asthma risk in Puerto Ricans, likely through changes in expression of *ORMDL3* (shown to cause experimental asthma in a transgenic murine model),<sup>(45)</sup> *GSDMB*, or *ZBP2*. Another SNP located within the promoter region of *ZBP2* (rs4795397,  $P = 3.7 \times 10^{-12}$  in our meta-analysis [Table S1]) is a putative functional polymorphism that shows asthma-associated allele specific nucleosome occupancy.<sup>(39)</sup> However, the SNP's strong influence on *ZBP2*



promoter activity is masked by DNA methylation of exon 1 of this gene. In contrast, the *ORMDL3* promoter is fully unmethylated. The *ZPBP2* and *ORMDL3* genes show allelic differences in expression.<sup>(38, 39)</sup> It has been shown that the *IKZF3-ZPBP2-GSDMB-ORMDL3* haplotype acts differently in regulation of transcripts between Europeans and Africans,<sup>(38)</sup> with a stronger association in Europeans than in Africans. In a study by Verlaan *et al.*,<sup>(38)</sup> SNP rs8067378 was the most significant *cis*-eQTL in this four-gene region in both Europeans ( $P = 1.1 \times 10^{-18}$ ) and Africans ( $P = 4.3 \times 10^{-9}$ ). In an eQTL study in Puerto Ricans using our Hartford-PR dataset,<sup>(36)</sup> the same SNP was also the most significant *cis*-eQTL ( $P = 2.6 \times 10^{-10}$ ) in the four-gene region. Although the two studies<sup>(36, 38)</sup> might not be comparable due to different statistical models and different tissues (i.e., LCL used in the Verlaan *et al.* study<sup>(38)</sup> and globin-cleaned whole blood used in the Chen *et al.* study<sup>(36)</sup>), we speculate that the significance of associated SNPs with expression in Puerto Ricans is between Europeans and Africans, since these are two ancestral populations in Puerto Ricans. We further performed eQTL analyses between our top SNP (rs907092) and expression of the *IKZF3-ZPBP2-GSDMB-ORMDL3* locus using whole-blood RNA microarray data (n=121) from the Hartford-PR study. Findings from this secondary analysis, conducted using linear regression and adjusting for age, gender, asthma status, study site and the first two PCs, suggest that our top SNP (rs907092) may regulate expression of the *IKZF3-ZPBP2-GSDMB-ORMDL3* locus ( $P = 0.07$  for *IKZF3*,  $P = 0.02$  for *ZPBP2*,  $P = 0.009$  for *GSDMB* and  $P = 0.0004$  for *ORMDL3*).

We recognize several study limitations. First, we had insufficient statistical power to detect weak genetic effects of common SNPs (e.g., ORs between 0.9 and 1.10 for *IL33*) or rare susceptibility variants. Second, subjects in our study lived in different sites in the U.S. mainland and Puerto Rico, and thus environmental differences may have confounded our results despite adjustment for study site in the four component studies. However, most minor allelic frequencies and effect estimates were similar across study cohorts. Third, misclassification of COPD as asthma is possible among adults in the HCHS/SOL cohort, given that 980 (52.5%) participants were former or current smokers (mean pack-years=18.6). Such misdiagnosis could account for the less significant associations reported in HCHS/SOL, even after adjustment for cigarette smoking. In summary, our findings suggest that common allelic variants in the chromosome 17q21 locus have the greatest effects on, and are thus particularly important in, the pathogenesis of asthma in Puerto Ricans. We confirmed susceptibility variants in two previously reported genes (*IL1RL1* and *TLSP*). Future studies should aim to characterize functional variants that cause asthma in Puerto Ricans, either through primary genetic effects or through interactions with relevant exposures (e.g., second-hand smoke and air pollution).

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

We thank the participants and staff of the Hartford-PR study, GALA I and GALA II, and the HCHS/SOL study for their contributions to this work.

**Funding:** This work was supported by grants HL079966 and HL117191 from the U.S. National Institutes of Health (NIH), and by The Heinz Endowments. Dr. Yan's contribution was supported by grant from Children's Hospital of Pittsburgh of UPMC. Dr. Forno's contribution was supported by grant HL125666 from the U.S. NIH. Dr. Rastogi's contribution is supported by grant HL118733 from the NIH. The GALA II study was supported by U.S. NIH grants to E.G.B: HL088133, HL004464, HL117004, ES015794, ES24844, TRDRP 24RT 0025, MD006902, and GM007546. EGB was also supported by the RWJF Amos Medical Faculty Development Award, the American Asthma Foundation and the Sandler Foundation. MPY was supported by grant AC15/00015 from the Instituto de Salud Carlos III within the ERACoSysMed 1<sup>st</sup> Joint Transnational Call (SysPharmPedia 99) from the European Union, under the Horizon 2020. The HCHS/SOL was supported by NIH contracts N01-HC65233, N01-HC65234, N01-HC65235, N01-HC65236, N01-HC65237, and HHSN268201300005C AM03. The baseline examination of HCHS/SOL was carried out as a collaborative study supported by contracts from the National Heart, Lung, and Blood Institute (NHLBI) to the University of North Carolina (N01-HC65233), University of Miami (N01-HC65234), Albert Einstein College of Medicine (N01-HC65235), Northwestern University (N01-HC65236), and San Diego State University (N01-HC65237). The following Institutes/Centers/Offices contributed to the first phase of HCHS/SOL through a transfer of funds to the NHLBI: National Institute on Minority Health and Health Disparities, National Institute on Deafness and Other Communication Disorders, National Institute of Dental and Craniofacial Research (NIDCR), National Institute of Diabetes and Digestive and Kidney Diseases, National Institute of Neurological Disorders and Stroke, NIH Institution-Office of Dietary Supplements. The Genetic Analysis Center at the University of Washington was supported by NHLBI and NIDCR contracts (HHSN268201300005C AM03 and MOD03).

## References


- Duffy DL, Martin NG, Battistutta D, Hopper JL, Mathews JD. Genetics of asthma and hay fever in australian twins. *The American review of respiratory disease*. 1990; 142:1351–1358. [PubMed: 2252253]
- Nieminen MM, Kaprio J, Koskenvuo M. A population-based study of bronchial asthma in adult twin pairs. *Chest*. 1991; 100:70–75. [PubMed: 2060393]
- Torgerson DG, Ampleford EJ, Chiu GY, Gauderman WJ, Gignoux CR, Graves PE, Himes BE, Levin AM, Mathias RA, Hancock DB, Baurley JW, Eng C, Stern DA, Celedon JC, Rafaels N, Capurso D, Conti DV, Roth LA, Soto-Quiros M, Togias A, Li X, Myers RA, Romieu I, Van Den Berg DJ, Hu D, Hansel NN, Hernandez RD, Israel E, Salam MT, Galanter J, Avila PC, Avila L, Rodriguez-Santana JR, Chapela R, Rodriguez-Cintron W, Diette GB, Adkinson NF, Abel RA, Ross KD, Shi M, Faruque MU, Dunston GM, Watson HR, Mantese VJ, Ezurum SC, Liang L, Ruczinski I, Ford JG, Huntsman S, Chung KF, Vora H, Li X, Calhoun WJ, Castro M, Sienra-Monge JJ, del Rio-Navarro B, Deichmann KA, Heinzmann A, Wenzel SE, Busse WW, Gern JE, Lemanske RF Jr, Beaty TH, Bleecker ER, Raby BA, Meyers DA, London SJ. Mexico City Childhood Asthma Study SoG-E, Admixture in Latino A, Study of African Americans AG, Environments, Martinez FD, Childhood Asthma R, Education N, Weiss ST, Childhood Asthma Management P, Williams LK, Study of Asthma P, Pharmacogenomic Interactions by R-E, Barnes KC, Genetic Research on Asthma in African Diaspora S, Ober C, Nicolae DL. Meta-analysis of genome-wide association studies of asthma in ethnically diverse north american populations. *Nat Genet*. 2011; 43:887–892. [PubMed: 21804549]
- Moffatt MF, Gut IG, Demenais F, Strachan DP, Bouzigon E, Heath S, von Mutius E, Farrall M, Lathrop M, Cookson WO, Consortium G. A large-scale, consortium-based genomewide association study of asthma. *N Engl J Med*. 2010; 363:1211–1221. [PubMed: 20860503]
- Bouzigon E, Corda E, Aschard H, Dizier MH, Boland A, Bousquet J, Chateigner N, Gormand F, Just J, Le Moual N, Scheinmann P, Siroux V, Vervloet D, Zelenika D, Pin I, Kauffmann F, Lathrop M, Demenais F. Effect of 17q21 variants and smoking exposure in early-onset asthma. *N Engl J Med*. 2008; 359:1985–1994. [PubMed: 18923164]
- Galanter J, Choudhry S, Eng C, Nazario S, Rodriguez-Santana JR, Casal J, Torres-Palacios A, Salas J, Chapela R, Watson HG, Meade K, LeNoir M, Rodriguez-Cintron W, Avila PC, Burchard EG. Ormdl3 gene is associated with asthma in three ethnically diverse populations. *American journal of respiratory and critical care medicine*. 2008; 177:1194–1200. [PubMed: 18310477]
- Halapi E, Gudbjartsson DF, Jonsdottir GM, Bjornsdottir US, Thorleifsson G, Helgadóttir H, Williams C, Koppelman GH, Heinzmann A, Boezen HM, Jonasdottir A, Blondal T, Gudjonsson SA, Jonasdottir A, Thorlacius T, Henry AP, Altmueller J, Krueger M, Shin HD, Uh ST, Cheong HS, Jonsdottir B, Ludviksson BR, Ludviksdottir D, Gislason D, Park CS, Deichmann K, Thompson PJ,

- Wjst M, Hall IP, Postma DS, Gislason T, Kong A, Jonsdottir I, Thorsteinsdottir U, Stefansson K. A sequence variant on 17q21 is associated with age at onset and severity of asthma. *European journal of human genetics* : EJHG. 2010; 18:902–908. [PubMed: 20372189]
8. Leung TF, Sy HY, Ng MC, Chan IH, Wong GW, Tang NL, Waye MM, Lam CW. Asthma and atopy are associated with chromosome 17q21 markers in chinese children. *Allergy*. 2009; 64:621–628. [PubMed: 19175592]
  9. Madore AM, Tremblay K, Hudson TJ, Laprise C. Replication of an association between 17q21 snps and asthma in a french-canadian familial collection. *Human genetics*. 2008; 123:93–95. [PubMed: 17992541]
  10. Sleiman PM, Annaiah K, Imielinski M, Bradfield JP, Kim CE, Frackelton EC, Glessner JT, Eckert AW, Otieno FG, Santa E, Thomas K, Smith RM, Glaberson W, Garriss M, Gunnlaugsson S, Chiavacci RM, Allen J, Spergel J, Grundmeier R, Grunstein MM, Magnusson M, Bisgaard H, Grant SF, Hakonarson H. Ormdl3 variants associated with asthma susceptibility in north americans of european ancestry. *The Journal of allergy and clinical immunology*. 2008; 122:1225–1227. [PubMed: 18760456]
  11. Tavendale R, Macgregor DF, Mukhopadhyay S, Palmer CN. A polymorphism controlling ormdl3 expression is associated with asthma that is poorly controlled by current medications. *The Journal of allergy and clinical immunology*. 2008; 121:860–863. [PubMed: 18395550]
  12. Bisgaard H, Bonnelykke K, Sleiman PM, Brasholt M, Chawes B, Kreiner-Moller E, Stage M, Kim C, Tavendale R, Baty F, Phipper CB, Palmer CN, Hakonarson H. Chromosome 17q21 gene variants are associated with asthma and exacerbations but not atopy in early childhood. *American journal of respiratory and critical care medicine*. 2009; 179:179–185. [PubMed: 19029000]
  13. Galanter JM, Gignoux CR, Torgerson DG, Roth LA, Eng C, Oh SS, Nguyen EA, Drake KA, Huntsman S, Hu D, Sen S, Davis A, Farber HJ, Avila PC, Brigino-Buenaventura E, LeNoir MA, Meade K, Serebrisky D, Borrell LN, Rodriguez-Cintron W, Estrada AM, Mendoza KS, Winkler CA, Klitz W, Romieu I, London SJ, Gilliland F, Martinez F, Bustamante C, Williams LK, Kumar R, Rodriguez-Santana JR, Burchard EG. Genome-wide association study and admixture mapping identify different asthma-associated loci in latinos: The genes-environments & admixture in latino americans study. *The Journal of allergy and clinical immunology*. 2014; 134:295–305. [PubMed: 24406073]
  14. Akinbami LJ, Moorman JE, Bailey C, Zahran HS, King M, Johnson CA, Liu X. Trends in asthma prevalence, health care use, and mortality in the united states, 2001–2010. *NCHS Data Brief*. 2012:1–8.
  15. Rosser FJ, Forno E, Cooper PJ, Celedon JC. Asthma in hispanics. An 8-year update. *Am J Respir Crit Care Med*. 2014; 189:1316–1327. [PubMed: 24881937]
  16. Brehm JM, Acosta-Perez E, Klei L, Roeder K, Barmada MM, Boutaoui N, Forno E, Cloutier MM, Datta S, Kelly R, Paul K, Sylvia J, Calvert D, Thornton-Thompson S, Wakefield D, Litonjua AA, Alvarez M, Colon-Semidey A, Canino G, Celedon JC. African ancestry and lung function in puerto rican children. *J Allergy Clin Immunol*. 2012; 129:1484–1490. e1486. [PubMed: 22560959]
  17. Lavange LM, Kalsbeek WD, Sorlie PD, Aviles-Santa LM, Kaplan RC, Barnhart J, Liu K, Giachello A, Lee DJ, Ryan J, Criqui MH, Elder JP. Sample design and cohort selection in the hispanic community health study/study of latinos. *Ann Epidemiol*. 2010; 20:642–649. [PubMed: 20609344]
  18. Sorlie PD, Aviles-Santa LM, Wassertheil-Smoller S, Kaplan RC, Daviglus ML, Giachello AL, Schneiderman N, Raj L, Talavera G, Allison M, Lavange L, Chambless LE, Heiss G. Design and implementation of the hispanic community health study/study of latinos. *Ann Epidemiol*. 2010; 20:629–641. [PubMed: 20609343]
  19. Howie BN, Donnelly P, Marchini J. A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS genetics*. 2009; 5:e1000529. [PubMed: 19543373]
  20. Price AL, Patterson NJ, Plenge RM, Weinblatt ME, Shadick NA, Reich D. Principal components analysis corrects for stratification in genome-wide association studies. *Nat Genet*. 2006; 38:904–909. [PubMed: 16862161]

21. Burchard EG, Avila PC, Nazario S, Casal J, Torres A, Rodriguez-Santana JR, Toscano M, Sylvia JS, Alioto M, Salazar M, Gomez I, Fagan JK, Salas J, Lilly C, Matallana H, Ziv E, Castro R, Selman M, Chapela R, Sheppard D, Weiss ST, Ford JG, Boushey HA, Rodriguez-Cintron W, Drazen JM, Silverman EK, Genetics of Asthma in Latino Americans S. Lower bronchodilator responsiveness in puerto rican than in mexican subjects with asthma. *American journal of respiratory and critical care medicine*. 2004; 169:386–392. [PubMed: 14617512]
22. Choudhry S, Burchard EG, Borrell LN, Tang H, Gomez I, Naqvi M, Nazario S, Torres A, Casal J, Martinez-Cruzado JC, Ziv E, Avila PC, Rodriguez-Cintron W, Risch NJ. Ancestry-environment interactions and asthma risk among puerto ricans. *Am J Respir Crit Care Med*. 2006; 174:1088–1093. [PubMed: 16973984]
23. Torgerson DG, Gignoux CR, Galanter JM, Drake KA, Roth LA, Eng C, Huntsman S, Torres R, Avila PC, Chapela R, Ford JG, Rodriguez-Santana JR, Rodriguez-Cintron W, Hernandez RD, Burchard EG. Case-control admixture mapping in latino populations enriches for known asthma-associated genes. *J Allergy Clin Immunol*. 2012; 130:76–82. e12. [PubMed: 22502797]
24. Delaneau O, Marchini J, Zagury JF. A linear complexity phasing method for thousands of genomes. *Nature methods*. 2012; 9:179–181.
25. Genomes Project C, Abecasis GR, Altshuler D, Auton A, Brooks LD, Durbin RM, Gibbs RA, Hurles ME, McVean GA. A map of human genome variation from population-scale sequencing. *Nature*. 2010; 467:1061–1073. [PubMed: 20981092]
26. Alexander DH, Novembre J, Lange K. Fast model-based estimation of ancestry in unrelated individuals. *Genome research*. 2009; 19:1655–1664. [PubMed: 19648217]
27. Nishimura KK, Galanter JM, Roth LA, Oh SS, Thakur N, Nguyen EA, Thyne S, Farber HJ, Serebrisky D, Kumar R, Brigino-Buenaventura E, Davis A, LeNoir MA, Meade K, Rodriguez-Cintron W, Avila PC, Borrell LN, Bibbins-Domingo K, Rodriguez-Santana JR, Sen S, Lurmann F, Balmes JR, Burchard EG. Early-life air pollution and asthma risk in minority children. The gala ii and sage ii studies. *American journal of respiratory and critical care medicine*. 2013; 188:309–318. [PubMed: 23750510]
28. Pino-Yanes M, Thakur N, Gignoux CR, Galanter JM, Roth LA, Eng C, Nishimura KK, Oh SS, Vora H, Huntsman S, Nguyen EA, Hu D, Drake KA, Conti DV, Moreno-Estrada A, Sandoval K, Winkler CA, Borrell LN, Lurmann F, Islam TS, Davis A, Farber HJ, Meade K, Avila PC, Serebrisky D, Bibbins-Domingo K, Lenoir MA, Ford JG, Brigino-Buenaventura E, Rodriguez-Cintron W, Thyne SM, Sen S, Rodriguez-Santana JR, Bustamante CD, Williams LK, Gilliland FD, Gauderman WJ, Kumar R, Torgerson DG, Burchard EG. Genetic ancestry influences asthma susceptibility and lung function among latinos. *J Allergy Clin Immunol*. 2015; 135:228–235. [PubMed: 25301036]
29. Conomos MP, Laurie CA, Stilp AM, Gogarten SM, McHugh CP, Nelson SC, Sofer T, Fernandez-Rhodes L, Justice AE, Graff M, Young KL, Seyerle AA, Avery CL, Taylor KD, Rotter JI, Talavera GA, Daviglus ML, Wassertheil-Smoller S, Schneiderman N, Heiss G, Kaplan RC, Franceschini N, Reiner AP, Shaffer JR, Barr RG, Kerr KF, Browning SR, Browning BL, Weir BS, Aviles-Santa ML, Papanicolaou GJ, Lumley T, Szpiro AA, North KE, Rice K, Thornton TA, Laurie CC. Genetic diversity and association studies in us hispanic/latino populations: Applications in the hispanic community health study/study of latinos. *American journal of human genetics*. 2016; 98:165–184. [PubMed: 26748518]
30. Chen H, Wang C, Conomos MP, Stilp AM, Li Z, Sofer T, Szpiro AA, Chen W, Brehm JM, Celedon JC, Redline S, Papanicolaou GJ, Thornton TA, Laurie CC, Rice K, Lin X. Control for population structure and relatedness for binary traits in genetic association studies via logistic mixed models. *American journal of human genetics*. 2016
31. Willer CJ, Li Y, Abecasis GR. Metal: Fast and efficient meta-analysis of genomewide association scans. *Bioinformatics*. 2010; 26:2190–2191. [PubMed: 20616382]
32. Teo YY, Fry AE, Bhattacharya K, Small KS, Kwiatkowski DP, Clark TG. Genome-wide comparisons of variation in linkage disequilibrium. *Genome research*. 2009; 19:1849–1860. [PubMed: 19541915]
33. Ong RT, Teo YY. Varld: A program for quantifying variation in linkage disequilibrium patterns between populations. *Bioinformatics*. 2010; 26:1269–1270. [PubMed: 20308177]

34. Lara M, Akinbami L, Flores G, Morgenstern H. Heterogeneity of childhood asthma among hispanic children: Puerto rican children bear a disproportionate burden. *Pediatrics*. 2006; 117:43–53. [PubMed: 16396859]
35. Ono JG, Worgall TS, Worgall S. 17q21 locus and ormdl3: An increased risk for childhood asthma. *Pediatric research*. 2014; 75:165–170. [PubMed: 24165737]
36. Chen W, Brehm JM, Lin J, Wang T, Forno E, Acosta-Perez E, Boutaoui N, Canino G, Celedon JC. Expression quantitative trait loci (eqtl) mapping in puerto rican children. *PLoS One*. 2015; 10:e0122464. [PubMed: 25816334]
37. Caliskan M, Bochkov YA, Kreiner-Moller E, Bonnelykke K, Stein MM, Du G, Bisgaard H, Jackson DJ, Gern JE, Lemanske RF Jr, Nicolae DL, Ober C. Rhinovirus wheezing illness and genetic risk of childhood-onset asthma. *The New England journal of medicine*. 2013; 368:1398–1407. [PubMed: 23534543]
38. Verlaan DJ, Berlivet S, Hunninghake GM, Madore AM, Lariviere M, Moussette S, Grundberg E, Kwan T, Ouimet M, Ge B, Hoberman R, Swiatek M, Dias J, Lam KC, Koka V, Harmsen E, Soto-Quiros M, Avila L, Celedon JC, Weiss ST, Dewar K, Sinnett D, Laprise C, Raby BA, Pastinen T, Naumova AK. Allele-specific chromatin remodeling in the zbp2/gsdmb/ormdl3 locus associated with the risk of asthma and autoimmune disease. *Am J Hum Genet*. 2009; 85:377–393. [PubMed: 19732864]
39. Berlivet S, Moussette S, Ouimet M, Verlaan DJ, Koka V, Al Tuwaijri A, Kwan T, Sinnett D, Pastinen T, Naumova AK. Interaction between genetic and epigenetic variation defines gene expression patterns at the asthma-associated locus 17q12-q21 in lymphoblastoid cell lines. *Hum Genet*. 2012; 131:1161–1171. [PubMed: 22271045]
40. Verlaan DJ, Ge B, Grundberg E, Hoberman R, Lam KC, Koka V, Dias J, Gurd S, Martin NW, Mallmin H, Nilsson O, Harmsen E, Dewar K, Kwan T, Pastinen T. Targeted screening of cis-regulatory variation in human haplotypes. *Genome research*. 2009; 19:118–127. [PubMed: 18971308]
41. Stranger BE, Nica AC, Forrest MS, Dimas A, Bird CP, Beazley C, Ingle CE, Dunning M, Flicek P, Koller D, Montgomery S, Tavaré S, Deloukas P, Dermitzakis ET. Population genomics of human gene expression. *Nature genetics*. 2007; 39:1217–1224. [PubMed: 17873874]
42. Dixon AL, Liang L, Moffatt MF, Chen W, Heath S, Wong KC, Taylor J, Burnett E, Gut I, Farrall M, Lathrop GM, Abecasis GR, Cookson WO. A genome-wide association study of global gene expression. *Nat Genet*. 2007; 39:1202–1207. [PubMed: 17873877]
43. Murphy A, Chu JH, Xu M, Carey VJ, Lazarus R, Liu A, Szeffler SJ, Strunk R, Demuth K, Castro M, Hansel NN, Diette GB, Vonakis BM, Adkinson NF Jr, Klanderman BJ, Senter-Sylvia J, Ziniti J, Lange C, Pastinen T, Raby BA. Mapping of numerous disease-associated expression polymorphisms in primary peripheral blood cd4+ lymphocytes. *Human molecular genetics*. 2010; 19:4745–4757. [PubMed: 20833654]
44. Hao K, Bosse Y, Nickle DC, Pare PD, Postma DS, Laviolette M, Sandford A, Hackett TL, Daley D, Hogg JC, Elliott WM, Couture C, Lamontagne M, Brandsma CA, van den Berge M, Koppelman G, Reicin AS, Nicholson DW, Malkov V, Derry JM, Suver C, Tsou JA, Kulkarni A, Zhang C, Vessey R, Opitck GJ, Curtis SP, Timens W, Sin DD. Lung eqtls to help reveal the molecular underpinnings of asthma. *PLoS Genet*. 2012; 8:e1003029. [PubMed: 23209423]
45. Miller M, Rosenthal P, Beppu A, Mueller JL, Hoffman HM, Tam AB, Doherty TA, McGeough MD, Pena CA, Suzukawa M, Niwa M, Broide DH. Ormdl3 transgenic mice have increased airway remodeling and airway responsiveness characteristic of asthma. *Journal of immunology*. 2014; 192:3475–3487.







**Figure 1. Manhattan plot of meta-analysis results**

Manhattan plot showing the summary meta-analysis results of Hartford-PR, GALA I, GALA II, and HCHS/SOL. The chromosomal position of each SNP is displayed along the X-axis and the negative logarithm of the association  $P$ -value is displayed on the Y-axis. The blue line represents the suggestive significance line ( $P < 1 \times 10^{-5}$ ). The red line represents the genome-wide significance line ( $P < 5 \times 10^{-8}$ ).





**Figure 2. Forest plots of odds ratio and 95% confidence interval for the association with asthma**  
Forest plots for rs907092, the most significant SNP in the meta-analysis.



**Figure 3. Results of the meta-analysis (Hartford-PR, GALA I, GALA II and HCHS/SOL) on the chromosome 17 region**

The relative location of genes and the direction of transcription are shown in the lower portion of the figure, and the chromosomal position is shown on the  $x$  axis. The light blue line shows the recombination rate across the region (right  $y$  axis), and the left  $y$  axis shows the significance of the associations. The purple diamond shows the  $P$ -value for rs907092 that is the most significant SNP in the meta-analysis (Hartford-PR, GALA I, GALA II and HCHS/SOL). The circles show the  $P$ -values for all other SNPs and are color coded according to the level of LD with rs907092 in the 1000 Genome Project Admixed American (AMR) population.

**Table 1**

Summary of characteristics of participants included in the meta-analysis

	<b>Hartford-PR (n=948)</b>	<b>GALA I (n=437)</b>	<b>GALA II (n=1,786)</b>	<b>HCHS/SOL (n=1,866)</b>
Age in years (mean $\pm$ standard deviation)	10.0 $\pm$ 2.7	18.2 $\pm$ 9.5	12.9 $\pm$ 3.4	48.2 $\pm$ 14.0
Male gender (n, %)	495(52.2)	187(42.8)	907(50.8)	784(42.0)
Asthma (n, %)	523(55.2)	251(57.4)	892(49.9)	478(25.6)
Study sites	Hartford (CT) and San Juan (PR)	Puerto Rico and New York (NY)	Chicago (IL), Bronx (NY), Houston (TX), San Francisco (CA) and Puerto Rico	Bronx (NY), Chicago (IL), Miami (FL) and San Diego (CA)
Genotyping platform	Illumina 2.5M	Affymetrix 6.0 GeneChip	Axiom <sup>®</sup> LAT1 array (Affymetrix)	Illumina HCHS/SOL custom 15041502 B3 array

Table 2

Results of the meta-analysis of GWAS of asthma in Puerto Ricans, for SNPs associated with asthma in two previous meta-analysis in Europeans (by the GABRIEL consortium) and ethnically diverse North Americans (by the EVE consortium)

SNP	Chr	Position	Ref	Alt	Gene	In cited paper		Hartford-PR			GALA I			GALA II			HCHS/SOL			Combined	
						OR	P-value	AAF	OR	P-value	AAF	OR	P-value	AAF	OR	P-value	AAF	OR	P-value	OR	P-value
Loci reported by the GABRIEL consortium in Moffatt M.F. et al. <i>N Engl J Med</i> 2010; 363:1211–21																					
rs3771166	2	102986222	G	A	<i>IL18R1</i>	0.85*	0.45	0.90	0.500	0.44	0.75	0.114	0.43	0.85	0.018	0.46	1.14	0.110	0.94	0.111	
rs9273349	6	32625869	A	G	<i>HLA-DQ</i>	1.14	0.61	0.98	0.860	NA	NA	NA	NA	NA	NA	0.62	1.22	0.016	1.13	0.079	
rs1342326	9	6190076	A	C	<i>IL33</i>	1.27	0.21	0.90	0.403	0.26	1.12	0.571	0.22	1.05	0.528	0.20	1.13	0.213	1.05	0.376	
rs744910	15	67446785	G	A	<i>SMAD3</i>	0.89	0.43	0.88	0.175	0.42	1.01	0.981	0.45	0.93	0.269	0.45	1.05	0.572	0.95	0.326	
rs2305480	17	38062196	G	A	<i>GSDMB</i>	0.76	0.31	0.62	2.0×10 <sup>−6</sup>	0.31	0.54	0.003	0.32	0.71	6.7×10 <sup>−6</sup>	0.34	0.86	0.071	0.72	7.4×10 <sup>−12</sup>	
rs3894194	17	38121993	G	A	<i>GSDMA</i>	1.26	0.37	1.23	0.034	0.39	0.92	0.648	0.40	1.23	0.002	0.39	0.94	0.406	1.11	0.023	
rs2284033	22	37534034	G	A	<i>IL2RB</i>	0.92	0.41	0.91	0.272	0.39	1.09	0.638	0.40	1.17	0.024	0.42	1.00	0.997	1.04	0.289	
rs2073643	5	131723288	T	C	<i>SLC22A5</i>	0.89	0.50	0.96	0.668	0.48	1.18	0.356	0.50	0.93	0.257	0.51	1.01	0.856	0.98	0.551	
rs1295686	5	131995843	T	C	<i>IL13</i>	0.85	0.59	1.01	0.943	0.62	0.75	0.136	0.62	0.90	0.159	0.61	1.07	0.398	0.96	0.408	
rs11071559	15	61069988	C	T	<i>RORA</i>	0.88	0.26	1.07	0.525	0.25	1.69	0.012	0.24	0.90	0.193	0.24	0.85	0.087	0.96	0.486	
Replicated genes reported by the EVE consortium in Torgerson D.G. et al. <i>Nat. Genet.</i> 2011; 43(9):887–92																					
rs1102000	1	158932907	T	C	<i>PYHIN1</i>	0.76#	0.08	1.09	0.651	0.06	0.50	0.061	NA	NA	NA	0.07	1.21	0.986	0.93	0.611	
rs10173081	2	102957348	C	T	<i>IL1RL1</i>	0.83	0.19	0.84	0.132	0.16	0.85	0.519	0.26	0.78	0.010	0.18	0.93	0.506	0.84	0.004	
rs1837253	5	110401872	C	T	<i>TSLP</i>	0.84	0.25	0.73	0.004	0.23	0.65	0.041	0.27	0.83	0.014	0.26	0.96	0.701	0.82	2.5×10 <sup>−4</sup>	
rs2381416	9	6193455	A	C	<i>IL33</i>	1.18	0.38	0.98	0.847	0.39	1.00	0.980	0.37	1.09	0.249	0.35	1.13	0.157	1.07	0.155	
rs11078927	17	38064405	C	T	<i>GSDMB</i>	0.79	0.30	0.61	1.4×10 <sup>−6</sup>	0.31	0.55	0.003	0.31	0.71	7.9×10 <sup>−6</sup>	0.34	0.86	0.079	0.72	8.1×10 <sup>−12</sup>	

Ref: reference allele; Alt: alternative allele; AAF: alternative allele frequency; OR: odds ratio.

<sup>\*</sup> This column is childhood onset odds ratio in Moffatt M.F. et al. *N Engl J Med* 2010; 363:1211–21;

<sup>#</sup> This column is overall odds ratio in Torgerson D.G. et al. *Nat. Genet.* 2011; 43(9):887–92.

Human Genome version: hg19