



Published in final edited form as:

Knee Surg Sports Traumatol Arthrosc. 2018 April ; 26(4): 1137–1144. doi:10.1007/s00167-017-4534-6.

Hip External Rotation Strength Predicts Hop Performance after Anterior Cruciate Ligament Reconstruction

Paul W. Kline, PT, DPT^a, Jeremy Burnham, MD^b, Michael Yonz, MD^c, Darren Johnson, MD^c, Mary Lloyd Ireland, MD^c, and Brian Noehren, PT, PhD^d

^aDepartment of Rehabilitation Sciences, University of Kentucky

^bCenter for Sports Medicine, University of Pittsburgh Medical Center

^cDepartment of Orthopedic Surgery and Sports Medicine, University of Kentucky

^dDivision of Physical Therapy, University of Kentucky

Abstract

Purpose—Quadriceps strength and single-leg hop performance are commonly evaluated prior to return to sport after anterior cruciate ligament reconstruction (ACLR). However, few studies have documented potential hip strength deficits after ACLR, or ascertained the relative contribution of quadriceps and hip strength to hop performance.

Methods—Patients cleared for return to sports drills after ACLR were compared to a control group. Participants' peak isometric knee extension, hip abduction, hip extension, and hip external rotation (HER) strength was measured. Participants also performed single-leg hops, timed hops, triple hops, and crossover hops. Between-limb comparisons for the ACLR to control limb and the non-operative limb were made using independent two-sample and paired-sample t-tests. Pearson's correlations and stepwise multiple linear regression were used to determine the relationships and predictive ability of limb strength, graft type, sex, and limb dominance to hop performance.

Results—Sixty-five subjects, 20 ACLR (11F, Age: 22.8(15–45) years, 8.3±2 months post-op, Mass: 70.47±12.95 kg, Height: 1.71±0.08 m, Tegner: 5.5 (3–9)) and 45 controls (22F, Age: 25.8(15–45) years, Mass: 74.0±15.2 kg, Height: 1.74±0.1 m, Tegner: 6 (3–7)) were tested. Knee extension (4.4±1.5 vs 5.4±1.8 N/kg, $p=0.02$), HER (1.4±0.4 vs 1.7±0.5 N/kg, $p=0.04$), single-leg hop (146±37 vs 182±38 % limb length, $p<0.01$), triple hop (417±106 vs 519±102 % limb length, $p<0.01$), timed hop (3.3±2.0 vs 2.3±0.6 s, $p<0.01$), and crossover hop (364±107 vs 446±123 % limb length, $p=0.01$) were significantly impaired in the operative versus control subject limbs. Similar deficits existed between the operative and non-operative limbs. Knee extension and HER strength were significantly correlated to each of the hop tests, but only HER significantly predicted hop performance.

Please address correspondence to: Brian Noehren, PT, PhD, 900 S Limestone, Room 204D, Lexington, KY 40536-0200, Phone: (859) 218-0581, Fax: (859) 323-6003, b.noehren@uky.edu.

CONFLICT OF INTEREST: The authors declare that they have no conflict of interest.

ETHICAL APPROVAL: All procedures performed in studies involving human participants were in accordance with the ethical standards of the institutional and/or national research committee and with the 1964 Helsinki declaration and its later amendments or comparable ethical standards.

INFORMED CONSENT: Informed consent was obtained from all individual participants included in this study.

Conclusions—After ACLR, patients have persistent HER strength, knee extension strength, and hop test deficits in the operative limb compared to the control and non-operative limbs, even after starting sport specific drills. Importantly, HER strength independently predicted hop performance. Based on these findings, to resolve between-limb deficits in strength and hop performance clinicians should include HER strengthening exercises in postoperative rehabilitation.

Keywords

ACL; hip strength; rehabilitation; quadriceps

INTRODUCTION

Between 130,000 and 175,000 anterior cruciate ligament reconstructions (ACLRs) are performed annually in the United States with only 44–55% of all patients returning to competitive sport [1,2,23]. Of those who return to sport (RTS), 11.1–29.5% experience a second ACL injury in the same or contralateral knee [42]. Poor RTS outcomes have generated significant discussion regarding the criteria used to assess athlete readiness for RTS after ACLR [1,22,24].

Of the objective RTS criteria described, quadriceps strength and single leg hop tests are the most commonly assessed [22]. Quadriceps muscle weakness following ACLR has been well documented and is a significant risk factor for re-injury after RTS [9,18,33]. Furthermore, athletes achieving 90% symmetry in quadriceps strength and single leg hop testing compared to the non-injured limb were reinjured less frequently [9]. Despite the recommended utilization of quadriceps strength and hop testing in RTS decision-making, there appears to be a minimal relationship between the two measures [10,37]. A lack of relationship between quadriceps strength and hop testing suggests that the two measures assess different constructs in recovery from ACLR and highlights the need to explore the contribution of weakness in other muscle groups, such as the hip, to hop performance.

While less commonly studied than quadriceps strength, hip muscle weakness has also been described after ACLR [5,14,33]. Interestingly, several prospective studies have linked hip dysfunction to knee biomechanics associated with increased incidence of lower extremity injury [7,12,13,39]. Despite these findings, hip muscle strength is not often used as a RTS criterion or evaluated in terms of ACL re-injury risk. Likewise, the relationships between hip strength and common RTS criteria such as single leg hop testing have yet to be elucidated. Given the contribution of the hip musculature in both propulsion and eccentric control during single leg landings, deficits in hip strength may play a role in impaired hop performance after ACLR [4,13].

The purpose of this study was to compare the isometric hip and quadriceps muscle strength and single leg hop test performance of the ACLR extremity to a control group, as well as to the non-operative extremity. This study also sought to examine the relationship between hip and quadriceps muscle strength with performance on single leg hop tests. It was hypothesized that ACLR patients would have persistent muscle strength and hop test deficits, and that hip muscle strength would be more closely related to hop test performance than quadriceps strength. Additionally, it was hypothesized that hip muscle strength would

better predict hop test performance than quadriceps strength, sex, graft type, and limb dominance.

MATERIALS AND METHODS

ACL Reconstruction Patients and Controls

Patients included in the ACLR group were recruited from eligible patients at the University's outpatient sports medicine clinic. Inclusion criteria included: (1) at least six months status post ACL reconstruction with hamstring or bone-patellar tendon-bone (BPTB) autograft, (2) no injury to the ipsilateral or contralateral limb in the prior 3 months, (3) no previous history of injury or surgery to the contralateral limb that may affect hip or knee function, and (4) cleared for return to sport drills and sport-specific training by their physical therapist and surgeon. Patients were excluded if they had any of the following: (1) a history of other ligamentous injuries to either knee, (2) knee effusion in either knee, (3) positive Lachman's test in either knee, or (4) positive pivot shift in either knee.

Participants included in the control group were recruited from a sample of convenience. Flyers and recruitment emails were distributed amongst university classes and throughout the community. To be included, all participants were in good general health and met the following inclusion criteria: (1) age between 18–45 years, (2) currently free of any trunk, hip, or knee injuries within the last three months, and (3) no previous history of injury or surgery that may affect their trunk, hip, or knee function. At the time of data collection, all participants completed a Tegner activity scale to quantify their current physical activity level and declared their pre-injury limb dominance.

Strength Testing

Participants completed a series of isometric lower extremity strength tests: hip abduction, hip extension, hip external rotation, and knee extension. Tests were performed by two male assessors using a hand-held dynamometer (Lafayette Instruments, Lafayette, IN) secured by a stabilization strap, as previously described and validated [3,41]. One practice and three experimental trials were performed for five seconds, with 15 seconds of rest between contractions. The average of the three experimental trials was used for calculations. To allow for comparison between groups, the experimental trials were normalized to body mass by dividing the strength value by the subject's weight in kilograms. Hip abduction strength was tested with the subject lying in the sidelying position. Hip extension strength was tested with participants in the prone position and the knee flexed to 90°. Hip external rotation strength was tested with participants in the seated position, with the knee and hip flexed to 90°. Knee extension strength was tested with the participant in the sitting position, the examined thigh parallel to the floor, and the leg hanging off the table in a vertical position with the knee flexed to 90°.

Single Leg Hop Testing

Participants in each group performed a series of single leg hop tests as described and validated [17,21,25,35,36]. This battery of hop tests included a single leg hop for distance, a timed 6-meter hop, a triple hop for distance, and a crossover hop for distance. For all of the

tests, the participant was required to start from a resting single leg stance. ACLR participants performed with their non-operative limbs first to prevent inadvertently biasing the performance of the non-operative limb to match the performance of the operative limb. One trial hop followed by two measured hops were performed for each test [35]. The average value of the measured hops was used for later calculations. To allow for comparison between groups, hop distances were normalized to limb length by dividing the distance by limb length as measured from the anterior superior iliac spine to the medial malleolus [25]. The Institutional Review Board of the University of Kentucky approved this study (13-0326-P1H).

Statistical Analysis

Comparisons between the operative and non-operative limb for the ACLR group were made using two-tailed paired samples t-tests to compare knee strength, hip strength, and single leg hop performance. To compare the ACLR operative limb with the control group and for comparisons of sex, limb dominance, and graft type in the ACLR group, independent two-sample t-tests were utilized. As an ordinal variable, the Tegner activity scale was compared using a Mann-Whitney U test. Muscle strength and hop performance variables identified as significantly different between-groups or between-limbs were included in subsequent correlational analyses. Pearson's product moment correlation coefficients were calculated to assess the relationship between limb strength and hop performance. Subsequently, only those relationships which were significantly correlated were entered into a stepwise multiple linear regression along with sex, limb dominance, and graft type to determine the predictability of hip and knee strength measures on single leg hop performance in the ACLR operative limb. PASW Statistics Version 18.0 (SPSS Inc, Chicago, IL, USA) was utilized for all limb comparisons, correlations, and linear regression analyses. Statistical significance was defined as $p < 0.05$. To detect an effect size of 0.5 at $\alpha = .05$ and $\beta = 0.8$, a sample size calculation revealed a need for a minimum of 17 subjects per group.

RESULTS

A total of 65 participants (20 ACLR, 45 controls) completed the study. No significant differences in mean age, height, weight, or Tegner activity level between the ACLR and Control groups were present at the time of testing (Table 1).

There was a significant difference between the ACLR operative limb and control limb in hip external rotation strength, knee extension strength, single-leg hop, timed hop, triple hop, and crossover hop, with the ACLR operative limb significantly weaker and demonstrating poorer performance in each hop test (Table 2). There were no significant differences in hip abduction strength or hip extension strength. Similar results were observed when comparing the ACLR operative to the non-operative limb (Table 3).

Males significantly outperformed females in all four hop tests in the ACLR limb. There were no significant differences when the ACLR group was stratified by injury to the dominant or non-dominant limb. Lastly, subjects with BPTB autograft demonstrated greater hip extension strength, hip external rotation strength, hip abduction strength, and single leg hop

performance compared to subjects with hamstring autograft. For complete results by sex, limb dominance, and graft type, refer to Table 4.

Significant correlations were found between hip external rotation and knee extension strength, and performance on all hop tests (Table 5). Stepwise multiple linear regression models revealed hip external rotation as the sole predictor of hop performance (single leg hop: $b = 0.833$, $p = .000$; triple hop: $b = 2.23$, $p = .000$; timed hop $b = -0.034$, $p = .007$; crossover hop: $b = 2.37$, $p = .000$). The R^2 of the models were 0.56, 0.48, 0.30, and 0.56, respectively.

DISCUSSION

The most important finding of the present study was the presence of isometric hip external rotation weakness which predicted single leg hop performance independent of knee extension strength after ACLR. Despite recommendations that hip strength be assessed after ACLR, little is known about hip muscle weakness following ACL reconstruction [6,11]. The results of the current study partially agree with previous investigations of hip strength impairments after ACLR, which also found no differences in hip abduction strength in a cohort of females 7 months post-ACLR [28]. However, in contrast, the current study found that ACLR subjects had significantly weaker hip external rotation strength than controls while the previous study reported no significant differences [28]. Differences in hip external rotation strength between this and the former study may be partially due to the mixed gender cohort and lower Tegner score at the time of testing in the current cohort compared to the previous study (5.7 vs 6.5). Additionally, one previous study also reported no difference in isometric hip external rotation strength in the ACLR limb compared to a control group, but the participants in this study were greater than 3 years post-ACLR, as compared to 8 months post-ACLR in the current study [5]. As such, a direct comparison between the two studies is difficult, and differences are likely due to the variability in time points used for testing. Nonetheless, these data suggest that hip external rotation strength deficits are present at time of return to sport after ACLR and represent a potential area for additional intervention during postoperative rehabilitation.

The current study's finding of reduced hip external rotation strength in the ACLR group is notable in light of a several prospective studies linking hip muscle function to ACL injury risk. For instance, Paterno et al. identified reduced contralateral hip external rotation torque production as a significant predictor of a second ACL tear [32]. Additionally, Khayambashi et al. demonstrated that hip external rotation strength independently predicted non-contact primary ACL injury in a large prospective study of male and female competitive athletes [15]. The findings of this study extend those of previous studies by providing evidence that hip external rotation strength remains impaired following rehabilitation, possibly contributing to impaired performance and heightened injury risk. These results suggest the need for hip external rotation strengthening exercises during rehabilitation after ACLR. To date, only one study has evaluated the efficacy of an isolated hip strengthening intervention during post-ALCR rehabilitation [8]. This study demonstrated minimal differences in 3-month knee extension range of motion, pain rating, and International Knee Documentation Committee scores compared to the group that did not receive early hip strengthening during

rehabilitation. However, this study did not assess hip strength at any time during the study to determine if baseline impairments in hip strength were present and/or were improved after the intervention [8]. The lack of objective hip strength assessment makes it difficult to assess if any improvements in hip strength were achieved, possibly accounting for lack of significant findings. Based on data presented in the current study, additional investigations into the role of hip external rotation strengthening exercises during recovery from ACLR on subsequent sport performance and injury are needed.

The findings of significantly reduced knee extension strength and single leg hop performance compared to controls are consistent with previous studies of knee extension strength deficits and hop performance at several time points after ACLR [19,30,31,33,34]. Recovery of quadriceps strength has been cited as an important factor in achieving a successful outcome after ACLR [16,34,38]. Assessments of quadriceps strength and hop performance are commonly performed to provide objective criteria for return to sport [24,35]. However, the relationship between quadriceps strength and single leg hop performance is variable, suggesting that other factors contribute to improved hop performance [11]. In the current study, while both knee extension strength and hip external rotation strength were significantly related to hop performance, the relationship with hip external rotation strength was the strongest (Table 5). A recent review of the measurement properties of single leg hop tests noted limited and conflicting evidence regarding the tests' abilities to predict injury [10]. It should be noted, however, that the value of single leg hop assessments in RTS testing may be from these tests serving as valuable benchmarks for recovery from ACLR. Of the single leg hop tests utilized in the current study, the single hop for distance is the most studied and demonstrates good discriminative validity in males after ACLR [10,26]. Additionally, the single leg hop for distance is responsive to improvements in performance after ACLR [10,40]. Thus, improvement in single hop performance during rehabilitation from ACLR may further determine the degree of recovery achieved. Without additional evidence on how hop test performance contributes to future injury risk or readiness for RTS after ACLR, it is difficult to derive absolute meaning from observed asymmetries for an individual athlete. However, the minimal equipment demands associated with hop testing, the discriminative validity of the single hop test after ACLR, responsiveness to rehabilitation, and the inclusion of hop testing for limb symmetry in successful RTS testing batteries suggest there is clinical utility in administering single leg hop testing in patients after ACLR [9,10,18].

The results of this study highlight the need to address hip external rotation strength deficits during rehabilitation after ACLR. It was found that when knee extension strength, hip external rotation strength, sex, graft type, and injury to dominant or non-dominant limb were entered into a regression model, only hip external rotation strength was a significant predictor of hop performance and independently predicted between 30 to 56% of the variance in performance. Hopping for maximum distance demands large amounts of muscle power to propel the body forward and to control the landing. As one of the most powerful muscles in the human body, and a significant contributor to trunk control during dynamic lower extremity tasks [20,29], the gluteus maximus is critical for performance of the hopping tasks. This may explain why patients who have sufficient postoperative quadriceps strength recovery continue to demonstrate asymmetries in hop testing performance, and

points to possible global muscle strength impairment in the ACLR limb. Active hip external rotation, as tested in this study, consists mostly of gluteus maximus recruitment [20,27]. Other muscles commonly associated with hip external rotation, such as the piriformis and short external rotators, have little or no effect on external rotation when the hip is flexed to 90 degrees [27]. Interestingly, despite differences observed in hip external rotation strength, hip extension strength was not significantly different between the ACLR, non-operative limb, or control group. Testing was performed in prone with the knee flexed to 90° and the lumbar spine stabilized with a strap to limit the contribution of the hamstrings and lumbar extensors, respectively. However, contributions from these muscle groups may have masked gluteal weakness during hip extension. Future work should investigate the efficacy of hip strengthening intervention on reducing biomechanical risk factors for second ACL injury, rate of successful return to sport, and the role of improved hip strength on psychological factors related to recovery after ACLR. These studies will further clarify the significance of proximal weakness and identify the most successful means of intervention to improve hip strength in this population.

There are several limitations of the current study. First, the study was cross-sectional and thus potential associations between muscle weakness or hop performance with future injury risk cannot be made. Additionally, hip strength was not assessed pre-operatively or at prior time points post-operatively so baseline differences cannot be accounted for nor can the time of onset of hip external rotation weakness be established. Lastly, although isometric testing is the most easily reproduced clinically, it does not reflect how these muscles perform during dynamic activities like hopping.

The findings of this study showed that isometric hip external rotation weakness is present after ACLR and is predictive of single leg hop performance. Based on these findings, interventions to increase hip external rotation strength should be included as part of rehabilitation after ACLR in order to achieve better and potentially more symmetrical single leg hop performance.

CONCLUSIONS

Patients after ACLR have significant deficits in hip external rotation strength, knee extension strength, and single leg hop performance with hip external rotation strength independently predicting single leg hop performance. Although quadriceps strengthening should continue to be an important component of rehabilitation after ACLR, patients may also benefit from exercises to improve hip external rotation strength to facilitate better dynamic limb function.

Acknowledgments

FUNDING: This study was funded by the National Institute of Arthritis and Musculoskeletal and Skin Diseases of the National Institutes of Health (grant number K23AR062069) and by the National Center for Advancing Translational Sciences of the National Institutes of Health (grant number TL1TR00015).

LIST OF ABBREVIATIONS

ACLR anterior cruciate ligament reconstruction

HER	hip external rotation
RTS	return to sport

References

1. Anderson MJ, Browning WM 3rd, Urband CE, Kluczynski MA, Bisson LJ. A systematic summary of systematic reviews on the topic of the anterior cruciate ligament. *Orthop J Sports Med.* 2016; 4(3) 2325967116634074.
2. Buller LT, Best MJ, Baraga MG, Kaplan LD. Trends in anterior cruciate ligament reconstruction in the united states. *Orthop J Sports Med.* 2015; 3(1) 2325967114563664.
3. Burnham JM, Yonz MC, Robertson KE, McKinley R, Wilson BR, Johnson DL, Ireland ML, Noehren B. Relationship of hip and trunk muscle function with single leg step-down performance: implications for return to play screening and rehabilitation. *Phys Ther Sport.* 2016; 22:66–73. [PubMed: 27592407]
4. Comyns T, Kenny I, Scales G. Effects of a low-load gluteal warm-up on explosive jump performance. *J Hum Kinet.* 2015; 46:177–187. [PubMed: 26240661]
5. Dalton EC, Pfile KR, Weniger GR, Ingersoll CD, Herman D, Hart JM. Neuromuscular changes after aerobic exercise in people with anterior cruciate ligament-reconstructed knees. *J Athl Train.* 2011; 46(5):476–483. [PubMed: 22488134]
6. Di Stasi S, Myer GD, Hewett TE. Neuromuscular training to target deficits associated with second anterior cruciate ligament injury. *J Orthop Sports Phys Ther.* 2013; 43(11):777–792. A771-711. [PubMed: 24175599]
7. Frank B, Bell DR, Norcross MF, Blackburn JT, Goerger BM, Padua DA. Trunk and hip biomechanics influence anterior cruciate loading mechanisms in physically active participants. *Am J Sports Med.* 2013; 41(11):2676–2683. [PubMed: 23884306]
8. Garrison JC, Bothwell J, Cohen K, Conway J. Effects of hip strengthening on early outcomes following anterior cruciate ligament reconstruction. *Int J Sports Phys Ther.* 2014; 9(2):157–167. [PubMed: 24790777]
9. Grindem H, Snyder-Mackler L, Moksnes H, Engebretsen L, Risberg MA. Simple decision rules can reduce reinjury risk by 84% after acl reconstruction: the delaware-oslo acl cohort study. *Br J Sports Med.* 2016; 50(13):804–808. [PubMed: 27162233]
10. Hegedus EJ, McDonough S, Bleakley C, Cook CE, Baxter GD. Clinician-friendly lower extremity physical performance measures in athletes: a systematic review of measurement properties and correlation with injury, part 1. The tests for knee function including the hop tests. *Br J Sports Med.* 2015; 49(10):642–648. [PubMed: 25497489]
11. Hewett TE, Di Stasi SL, Myer GD. Current concepts for injury prevention in athletes after anterior cruciate ligament reconstruction. *Am J Sports Med.* 2013; 41(1):216–224. [PubMed: 23041233]
12. Hollman JH, Ginos BE, Kozuchowski J, Vaughn AS, Drause DA, Youdas JW. Relationships between knee valgus, hip-muscle recruitment during a single-limb step-down. *Journal of Sport Rehabilitation.* 2009; 18(1):104–117. [PubMed: 19321910]
13. Jacobs CA, Uhl TL, Mattacola CG, Shapiro R, Rayens WS. Hip abductor function and lower extremity landing kinematics: sex differences. *J Athl Train.* 2007; 42(1):76–83. [PubMed: 17597947]
14. Karanikas K, Arampatzis A, Bruggemann GP. Motor task and muscle strength followed different adaptation patterns after anterior cruciate ligament reconstruction. *Eur J Phys Rehabil Med.* 2009; 45(1):37–45. [PubMed: 18987563]
15. Khayambashi K, Ghoddosi N, Straub RK, Powers CM. Hip muscle strength predicts noncontact anterior cruciate ligament injury in male and female athletes: a prospective study. *Am J Sports Med.* 2016; 44(2):355–361. [PubMed: 26646514]
16. Kline PW, Johnson DL, Ireland ML, Noehren B. Clinical predictors of knee mechanics at return to sport after acl reconstruction. *Med Sci Sports Exerc.* 2016; 48(5):790–795. [PubMed: 26694845]
17. Kockum B, Heijne AI. Hop performance and leg muscle power in athletes: reliability of a test battery. *Phys Ther Sport.* 2015; 16(3):222–227. [PubMed: 25891995]

18. Kyritsis P, Bahr R, Landreau P, Miladi R, Witvrouw E. Likelihood of acl graft rupture: not meeting six clinical discharge criteria before return to sport is associated with a four times greater risk of rupture. *Br J Sports Med.* 2016; 50(15):946–951. [PubMed: 27215935]
19. Lepley LK. Deficits in quadriceps strength and patient-oriented outcomes at return to activity after acl reconstruction: a review of the current literature. *Sports Health.* 2015; 7(3):231–238. [PubMed: 26131300]
20. Lieberman DE, Raichlen DA, Pontzer H, Bramble DM, Cutright-Smith E. The human gluteus maximus and its role in running. *J Exp Biol.* 2006; 209(Pt 11):2143–2155. [PubMed: 16709916]
21. Logerstedt D, Grindem H, Lynch A, Eitzen I, Engebretsen L, Risberg MA, Axe MJ, Snyder-Mackler L. Single-legged hop tests as predictors of self-reported knee function after anterior cruciate ligament reconstruction: the delaware-oslo acl cohort study. *Am J Sports Med.* 2012; 40(10):2348–2356. [PubMed: 22926749]
22. Makhni EC, Crump EK, Steinhaus ME, Verma NN, Ahmad CS, Cole BJ, Bach BR Jr. Quality and variability of online available physical therapy protocols from academic orthopaedic surgery programs for anterior cruciate ligament reconstruction. *Arthroscopy.* 2016; 32(8):1612–1621. [PubMed: 27032604]
23. Mall NA, Chalmers PN, Moric M, Tanaka MJ, Cole BJ, Bach BR Jr, Paletta GA Jr. Incidence and trends of anterior cruciate ligament reconstruction in the united states. *Am J Sports Med.* 2014; 42(10):2363–2370. [PubMed: 25086064]
24. Mueller LM, Bloomer BA, Durall CJ. Which outcome measures should be utilized to determine readiness to play after acl reconstruction? *J Sport Rehabil.* 2014; 23(2):158–164. [PubMed: 23981536]
25. Munro AG, Herrington LC. Between-session reliability of four hop tests and the agility t-test. *J Strength Cond Res.* 2011; 25(5):1470–1477. [PubMed: 21116200]
26. Myer GD, Schmitt LC, Brent JL, Ford KR, Barber Foss KD, Scherer BJ, Heidt RS Jr, Divine JG, Hewett TE. Utilization of modified nfl combine testing to identify functional deficits in athletes following acl reconstruction. *J Orthop Sports Phys Ther.* 2011; 41(6):377–387. [PubMed: 21289456]
27. Neumann DA. Kinesiology of the hip: a focus on muscular actions. *J Orthop Sports Phys Ther.* 2010; 40(2):82–94. [PubMed: 20118525]
28. Noehren B, Abraham A, Curry M, Johnson D, Ireland ML. Evaluation of proximal joint kinematics and muscle strength following acl reconstruction surgery in female athletes. *J Orthop Res.* 2014; 32(10):1305–1310. [PubMed: 25044305]
29. Norris B, Trudelle-Jackson E. Hip- and thigh-muscle activation during the star excursion balance test. *J Sport Rehabil.* 2011; 20(4):428–441. [PubMed: 22012497]
30. Pairo de Fontenay B, Argaud S, Blache Y, Monteil K. Contralateral limb deficit seven months after acl-reconstruction: an analysis of single-leg hop tests. *Knee.* 2015; 22(4):309–312. [PubMed: 25981951]
31. Palmieri-Smith RM, Lepley LK. Quadriceps strength asymmetry after anterior cruciate ligament reconstruction alters knee joint biomechanics and functional performance at time of return to activity. *Am J Sports Med.* 2015; 43(7):1662–1669. [PubMed: 25883169]
32. Paterno MV, Schmitt LC, Ford KR, Rauh MJ, Myer GD, Huang B, Hewett TE. Biomechanical measures during landing and postural stability predict second anterior cruciate ligament injury after anterior cruciate ligament reconstruction and return to sport. *Am J Sports Med.* 2010; 38(10):1968–1978. [PubMed: 20702858]
33. Petersen W, Taheri P, Forkel P, Zantop T. Return to play following acl reconstruction: a systematic review about strength deficits. *Arch Orthop Trauma Surg.* 2014; 134(10):1417–1428. [PubMed: 25091127]
34. Pietrosimone BG, Lepley AS, Ericksen HM, Gribble PA, Levine J. Quadriceps strength and corticospinal excitability as predictors of disability after anterior cruciate ligament reconstruction. *J Sport Rehabil.* 2013; 22(1):1–6. [PubMed: 22951289]
35. Reid A, Birmingham TB, Stratford PW, Alcock GK, Giffin JR. Hop testing provides a reliable and valid outcome measure during rehabilitation after anterior cruciate ligament reconstruction. *Phys Ther.* 2007; 87(3):337–349. [PubMed: 17311886]

36. Reinke EK, Spindler KP, Lorrington D, Jones MH, Schmitz L, Flanigan DC, An AQ, Quiram AR, Preston E, Martin M, Schroeder B, Parker RD, Kaeding CC, Borzi L, Pedroza A, Huston LJ, Harrell FE Jr, Dunn WR. Hop tests correlate with ikdc and koos at minimum of 2 years after primary acl reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2011; 19(11):1806–1816. [PubMed: 21445595]
37. Ross MD, Irrgang JJ, Denegar CR, McCloy CM, Unangst ET. The relationship between participation restrictions and selected clinical measures following anterior cruciate ligament reconstruction. *Knee Surg Sports Traumatol Arthrosc.* 2002; 10(1):10–19. [PubMed: 11819015]
38. Schmitt LC, Paterno MV, Hewett TE. The impact of quadriceps femoris strength asymmetry on functional performance at return to sport following anterior cruciate ligament reconstruction. *J Orthop Sports Phys Ther.* 2012; 42(9):750–759. [PubMed: 22813542]
39. Stickler L, Finley M, Gulgin H. Relationship between hip and core strength and frontal plane alignment during a single leg squat. *Physical Therapy in Sport.* 2015; 16(1):66–71. [PubMed: 25070759]
40. Svensson M, Sernert N, Ejerhed L, Karlsson J, Kartus JT. A prospective comparison of bone-patellar tendon-bone and hamstring grafts for anterior cruciate ligament reconstruction in female patients. *Knee Surg Sports Traumatol Arthrosc.* 2006; 14(3):278–286. [PubMed: 16292682]
41. Thorborg K, Bandholm T, Holmich P. Hip- and knee-strength assessments using a hand-held dynamometer with external belt-fixation are inter-tester reliable. *Knee Surg Sports Traumatol Arthrosc.* 2013; 21(3):550–555. [PubMed: 22773065]
42. Wiggins AJ, Grandhi RK, Schneider DK, Stanfield D, Webster KE, Myer GD. Risk of secondary injury in younger athletes after anterior cruciate ligament reconstruction: a systematic review and meta-analysis. *Am J Sports Med.* 2016; 44(7):1861–1876. [PubMed: 26772611]

Table 1**Subject Demographics**

	Number	Age (years)	Sex	Weight (kg)	Injured Side	Follow-up (months)	Graft Type	Tegner
ACL	20	22.8 (15–45)	11 female, 9 male	70.5±12.9	8 left, 12 right	8.3 (6–14)	8 Hamstring Autograft, 12 BPTB Autograft	5.5 (3–9)
Controls	45	25.8 (15–45)	22 female, 23 male	74.0±15.2	-	-	-	6 (3–7)
P-value	-	0.097	0.649	0.370	-	-	-	0.761

- Not applicable

ACL: Anterior cruciate ligament reconstruction

BPTB: Bone patellar tendon bone

Data presented as mean ± standard deviation or mean (range).

Tegner presented as median (range).

Table 2
Comparison of Muscle Strength and Hop Test Performance Between ACLR and Control Group Limbs

Test	ACLR Mean	Controls Mean	% Difference	95% CI	P-value
Hip Extension	2.9 ± 1.1	3.1 ± 1.0	-6.7	-0.8, 0.4	n.s.
Hip External Rotation	1.4 ± 0.4	1.7 ± 0.5	-19.4	-0.5, -0.1	0.04 *
Hip Abduction	4.1 ± 0.7	4.3 ± 1.0	-4.8	-0.6, 0.4	n.s.
Knee Extension	4.4 ± 1.5	5.4 ± 1.8	-20.4	-1.9, -0.2	0.02 *
Single Leg Hop	146 ± 37	182 ± 38	-22	-56, -16	<0.01 *
Timed Hop (seconds)	3.3 ± 2.0	2.3 ± 0.6	35.7	0.4, 1.6	<0.01 *
Triple Hop	417 ± 106	519 ± 102	-21.8	-156, -46	<0.01 *
Crossover Hop	364 ± 107	446 ± 123	-20.2	-147, -18	0.01 *

Strength tests reported in N/kg. Hop tests reported as % of limb length, unless otherwise specified.
95% CI: 95% Confidence Interval for mean difference between ACLR and Controls.

* Statistically significant

± Standard deviation

ACLR: Anterior cruciate ligament reconstruction

Note: Statistically significant differences in bold; Hop test results normalized for limb length.

Table 3
Between-Limb Comparison of Muscle Strength and Hop Test Performance in ACLR Group

Test	ACLR Extremity		Non-operative Extremity		% Difference	95% CI	P-value
	Mean	SD	Mean	SD			
Hip Extension	195.4 ± 70.7		193.6 ± 44.6		0.9	-21.7, 25.3	n.s.
Hip External Rotation	101.6 ± 33.7		114.4 ± 36.7		-11.9	-25.6, -0.1	0.05 *
Hip Abduction	287.8 ± 56.5		287.1 ± 59.1		0.2	-22.9, 24.4	n.s.
Knee Extension	308.0 ± 120.1		366.4 ± 119.4		-17.3	-103.2, -13.6	0.01 *
Single Leg Hop	132 ± 37		162 ± 34		-20.4	-37, -22	<0.01 *
Timed Hop (seconds)	3.3 ± 2.0		2.5 ± 0.9		27.6	0.2, 1.5	0.02 *
Triple Hop	379 ± 105		452 ± 108		-17.6	-95, -51	<0.01 *
Crossover Hop	331 ± 107		390 ± 108		-16.4	-79, -38	<0.01 *

Strength tests reported in Newtons; Hop tests reported in cm, unless otherwise specified.

95% CI: 95% Confidence Interval for mean difference between ACLR and Non-operative extremity.

* Statistically significant

± Standard deviation

ACLR: Anterior cruciate ligament reconstruction

Note: Statistically significant differences in bold.

Table 4

Comparison of Muscle Strength and Hop Test Performance of the ACLR extremity in the ACLR Group Stratified by Sex, Limb Dominance, and Graft Type.

Test	Male	Female	P-value	Dominant Limb Injured	Non-dominant Limb Injured	P-value	BPTB Autograft	Hamstring Autograft	P-value
Hip Extension	2.7 ± 1.2	3.0 ± 1.0	n.s.	3.0 ± 1.0	2.7 ± 1.2	n.s.	3.3 ± 1.1	2.3 ± 0.7	0.03 *
Hip External Rotation	1.6 ± 0.5	1.4 ± 0.4	n.s.	1.5 ± 0.4	1.4 ± 0.5	n.s.	1.7 ± 0.4	1.1 ± 0.3	0.01 *
Hip Abduction	4.3 ± 0.8	4.0 ± 0.5	n.s.	4.3 ± 0.7	3.9 ± 0.6	n.s.	4.4 ± 0.7	3.8 ± 0.4	0.04 *
Knee Extension	4.5 ± 1.6	4.3 ± 1.4	n.s.	4.3 ± 1.6	4.4 ± 1.4	n.s.	4.8 ± 1.7	3.8 ± 0.8	n.s.
Single Leg Hop	164 ± 36	130 ± 32	0.04 *	148 ± 41	142 ± 34	n.s.	159 ± 36	126 ± 30	0.04 *
Timed Hop	2.3 ± 0.7	4.2 ± 2.3	0.03 *	3.5 ± 2.3	3.1 ± 1.6	n.s.	2.6 ± 1.2	4.4 ± 2.5	n.s.
Triple Hop	479 ± 95	367 ± 89	0.02 *	427 ± 126	406 ± 81	n.s.	453 ± 109	363 ± 79	n.s.
Crossover Hop	429 ± 91	305 ± 87	0.01 *	361 ± 123	367 ± 89	n.s.	399 ± 103	303 ± 91	n.s.

Strength tests reported in N/kg. Hop tests reported as % of limb length, except the Timed Hop which is reported in seconds.

* Statistically significant

Data presented as mean ± standard deviation

ACLR: Anterior cruciate ligament reconstruction

BPTB: Bone patellar tendon bone

Table 5

Pearson Correlations Between Limb Strength and Hop Performance of the ACLR Limb

	Single Leg Hop	Triple Hop	Timed Hop	Crossover Hop
Hip External Rotation	0.765 (.000) *	0.714 (.000) *	-0.579 (.007) *	0.766 (.000) *
Knee Extension	0.554 (.011) *	0.513 (.021) *	-0.426 (n.s.)	0.461 (.047) *

Data presented as r-value (p-value).

* statistically significant at $p < 0.05$