

Vitamin D Deficiency in Pediatric Critical Care

J. Dayre McNally^{1,2} Karin Amrein³

¹Division of Critical Care, Department of Pediatrics, Faculty of Medicine, University of Ottawa, Children's Hospital of Eastern Ontario, Ottawa, Canada

²Research Institute, Children's Hospital of Eastern Ontario, Ottawa, Canada

³Division of Endocrinology and Metabolism, Department of Internal Medicine, Medical University of Graz, Austria

Address for correspondence Dayre McNally, MSc, PhD, University of Ottawa, Pediatric Intensivist, Division of Critical Care, Children's Hospital of Eastern Ontario, 401 Smyth Road, Ottawa, Ontario, Canada, K1H 8L1 (e-mail: dmcnally@cheo.on.ca).

J Pediatr Intensive Care 2016;5:142–153.

Abstract

Vitamin D deficiency (VDD) is a well-established cause of pediatric bone and muscle disease. In addition, a role has been recognized for vitamin D in the health and stress response of other organs, including the cardiovascular, immune, and respiratory systems. As these organs are central to the development of and recovery from critical illness, VDD has been hypothesized to be a modifiable risk factor for ICU outcome. Over the past 5 years, a growing number of adult and pediatric critical care studies have investigated the prevalence of VDD and its association with illness severity and outcome. The adult studies have recently been synthesized in systematic reviews, with results that convincingly suggest the need for trials to determine whether optimization of vitamin D status improves outcome. In contrast, the pediatric ICU and related literature has not been similarly synthesized. The goal of this review is to describe vitamin D metabolism, known biological mechanisms, potential role in pathophysiology, and summarize the available pediatric intensive care unit (PICU) studies reporting on prevalence of VDD deficiency and its association with outcome. The problems with currently approved supplementation approaches and alternative strategies are discussed, including evidence from available RCTs in adult ICU. Altogether the results suggest that critically ill children are at risk for VDD, and that VDD appears to be associated with a worse clinical course. Clinical trials evaluating novel approaches to testing for and supplementing vitamin D require exploration.

Keywords

- pediatric critical care
- endocrinology
- vitamin D

Introduction

Severe vitamin D deficiency (VDD) is a well-established cause of pediatric disease, including hypocalcemia, skeletal abnormalities, and stunted growth.^{1–3} Although severe deficiency is now rare, many adults and children have subclinical VDD that does not manifest classic symptoms. Concern has been expressed about subclinical VDD as research suggests that it could predispose patients to or impair recovery from a variety of neurologic, cardiovascular, respiratory, and immune disorders.^{4–6} As these organs are central to the development and

recovery from critical illness, VDD has been hypothesized to be a modifiable risk factor for worse outcomes in intensive care unit (ICU).⁷

Over the past 5 years there has been a rapid growth of ICU literature on vitamin D. Dozens of adult studies have convincingly demonstrated associations between VDD and illness severity, patient morbidity, and mortality. This work has culminated in book chapters,⁸ narrative reviews,^{9–13} systematic reviews,^{14–16} pilot clinical trials,^{17–22} and a phase III randomized clinical trials (RCT) suggesting safety and clinical efficacy of high-dose supplementation.²³ In contrast, there

received
September 30, 2015
accepted after revision
October 22, 2015
published online
May 4, 2016

Issue Theme Endocrinology in Pediatric Critical Care; Guest Editors: Kusum Menon, MSc, MD, and Dayre McNally, MSc, PhD

Copyright © 2016 by Georg Thieme Verlag KG, Stuttgart · New York

DOI <http://dx.doi.org/10.1055/s-0036-1583285>.
ISSN 2146-4618.

have been relatively few attempts to synthesize the pediatric literature.²⁴ This review summarizes the pediatric intensive care unit (PICU) literature on VDD, discusses problems with approved supplementation regimens, and presents alternative strategies for evaluation as part of clinical trials.

Vitamin D Deficiency

What Is Vitamin D?

The term *vitamin D* refers to two molecules: ergocalciferol (D2) and cholecalciferol (D3). Both forms can be attained through naturally occurring dietary sources, fortified foods, and pharmaceutical supplements. Of the two, only cholecalciferol can be synthesized in the skin from 7-dehydrocholesterol following ultraviolet (UV) irradiation. Both D2 and D3 are biologically inert, requiring two separate hydroxylations to become active. The initial hydroxylation is performed by liver cytochrome P450 enzymes (25-hydroxylases), producing the prehormone calcidiol or 25-hydroxyvitamin D (25OHD). Calcidiol is the major vitamin D metabolite in the body (nanomolar blood concentrations) and circulates 99% bound to vitamin D-binding protein (VDBP) and albumin. A subsequent nonliver hydroxylation generates the active hormone called *calcitriol* (1,25 dihydroxyvitamin D or 1,25 (OH)₂D) (►Fig. 1).

Vitamin D Parathyroid Renal Axis

The best understood function of vitamin D is the maintenance of blood and body calcium levels²⁵ (see ►Fig. 1). When blood calcium falls, the parathyroid gland is stimulated and increases parathyroid hormone (PTH) secretion. Higher PTH levels prompt the kidneys to increase hydroxylase activity converting more available calcidiol to active hormone. The active hormone is circulated in the blood to the bone, gut, and kidneys where it binds to vitamin D receptor (VDR) and translocates to the nucleus. The calcitriol:VDR complex interacts with other proteins triggering changes in transcription of hundreds of genes. These changes work to restore calcium homeostasis through increased enteral absorption, decreased renal secretion, and modulation of osteoclast and osteoblast activity.

Nonclassic Functions of Vitamin D

It is now well recognized that dozens of cells outside of the bone, intestine, and kidneys have active VDRs, including myocytes, endothelial cells, white blood cells, pancreatic islets, and skeletal muscle.^{26,27} Additional studies have also shown that many cells remote from the kidneys produce α hydroxylases and generate calcitriol from calcidiol for both autocrine and paracrine purposes. Laboratory studies involving both cell lines and animal models have convincingly demonstrated that depletion of vitamin D negatively impacts normal development and stress response. Finally, in addition to altering transcription, it is now evident that vitamin D has nongenomic mechanisms where it can rapidly (within minutes) alter cell functioning through signaling pathways at the plasma membrane (caveolae) or t-tubules.^{27–30}

Definition of Vitamin D Status

Calcidiol (25OHD) levels are widely regarded as the best indicator of vitamin D status. Although variability exists, 50 nmol/L is widely used as the threshold to define deficiency, with 25 or 30 nmol/L representing severe deficiency (►Fig. 2). Some agencies also recognize an additional state of “insufficiency” with adequate or normal status achieved when levels exceed 75 or 80 nmol/L.^{31,32} These thresholds are based on biochemical indicators of axis stress and values below which disease predisposition rises. In brief, when 25OHD falls to 50 nmol/L, maintenance of circulating calcitriol levels requires elevation of serum PTH and increased renal hydroxylase activity.^{33,34} As 25OHD falls to 30 nmol/L renal production of calcitriol declines and healthy individuals can develop electrolyte disturbances and clinically evident musculoskeletal pathology.^{34–36} These thresholds apply specifically to bone health; the 25OHD concentrations required for optimal non-renal hydroxylase activity have not been defined.

Explanation of the Potential Role for Vitamin D in ICU Pathophysiology

The biological mechanisms through which VDD may contribute to primary or secondary ICU pathophysiology have been presented in full elsewhere^{11,37}; however, a few examples are discussed here.

Hypocalcemia

Hypocalcemia occurs frequently during pediatric critical illness, and both its occurrence and administration of parenteral calcium have been associated with increased morbidity and mortality.^{38–41} Normal intra- and extracellular calcium homeostasis is important as it initiates and propagates nerve conduction, muscle contraction, and contributes to signal transduction. In addition to adult critical care studies, pediatric work has shown that patients with low calcium are more likely to have abnormalities of the vitamin D axis, including low 25OHD, hypoparathyroidism, and/or renal dysfunction.^{38,39,42}

Immune Dysfunction

Pediatric critical illness is frequently accompanied by an intense inflammatory response.^{43,44} VDRs have been identified on all major immune cell types, and current evidence suggests that proper signaling through these receptors is essential for modulating the immune response. For example, vitamin D has been demonstrated to inhibit antigen-induced T-cell proliferation, antagonize the proinflammatory Th1 response, alter gene expression of adhesion factors, and decrease neutrophil adherence and chemotaxis.^{45–47} Important functions in innate immunity have also been identified. For example, activation of the VDR is important for transcription and translation of antimicrobial peptides (e.g., cathelicidin).^{48–50} Cathelicidin levels have been correlated with both infectious disease outcomes and vitamin D concentrations.^{50–52} Pediatric observational studies have linked 25OHD levels and polymorphisms of both VDR and VDBP to various autoimmune and infectious conditions, including RSV, diabetes, and asthma exacerbations.^{53–55}




Fig. 1 Metabolism of vitamin D. Vitamin D₃ can be synthesized in human skin following exposure to ultraviolet light. Both vitamins D₃ and D₂ can also be acquired from consumption of specific foods, fortified foods, and pharmaceutical supplements. Both forms are transported to the liver where CYP450 enzymes generate the 25 hydroxyvitamin D metabolite and release it into the circulation. A second hydroxylation is required to produce the active hormone calcitriol. Circulating calcitriol is primarily produced in the kidney by an α hydroxylase that is regulated by parathyroid hormone (positively) and fibroblast growth factor (negatively). Multiple cells also synthesize their own hydroxylase and can locally convert 25OHD for autocrine and paracrine use. Once activated, there are multiple mechanisms through which vitamin D can alter cell function. It is well known that calcitriol binding leads to nuclear translocation of the VDR where it works with transcription factors to upregulate and downregulate transcription of genes. In addition, it has also been established that calcitriol can elicit several rapid nongenomic responses from the plasma membrane (or t-tubules) through binding to the VDR or alternative calcitriol receptors (e.g., MARRS). Abbreviations: Ca²⁺, ionized calcium; GPCR, G protein coupled receptor; MAPK, MAP kinase; MARRS, membrane-associated rapid-response steroid binding; PDIA3, protein disulfide isomerase family A, member 3; PKC, protein kinase C; PLA, phospholipase A; PLC, phospholipase C; PTH, parathyroid hormone; RXR, retinoid-X receptor; UV, ultraviolet; VDR, vitamin D receptor.

Cardiovascular Dysfunction

Cardiovascular dysfunction is common in PICU, with patients frequently receiving fluid and inotropes to support cardiac output.⁵⁶ In addition to indirect effects through calcium,

vitamin D influences myocyte structure and function via gene and protein expression.²⁶ Additional research has shown that myocyte contractility can be favorably altered within minutes by 1,25(OH)₂D supplementation, mediated

through signal transduction pathways, enzymatic reactions, and ion channels.^{28–30} Pediatric clinical studies support these laboratory observations, demonstrating that children with rickets frequently have subclinical cardiac dysfunction along with case reports/series suggesting cardiomyopathy secondary to severe VDD.^{57–61} A recent pilot RCT evaluating stable children with heart failure demonstrated improved heart function scores and echocardiographic findings with high-dose vitamin D (2,000 IU/d) when compared with placebo.⁶²

Muscle Weakness and Rehabilitation

ICU-acquired weakness is a well-recognized consequence of critical illness, and contributes to mortality, morbidity, worse functional outcomes, and quality of life.^{63–66} A significant body of observational research on children and adults has clearly demonstrated that severe VDD can cause muscle pathology and clinically relevant weakness.^{67–71} Recently, studies have demonstrated potentially long-lasting effects of high-dose vitamin D on body lean muscle mass in infants and young children.^{72,73} Further, a recent RCT evaluating administration of 540,000 IU to critically ill adults demonstrated that patients with 25OHD levels between 30 and 50 nmol/L who received study drug had improved grip strength and physical components of the SF-12.²³

Prevalence of Vitamin D Deficiency in PICU

Vitamin D Deficiency in PICU

Prior to 2012, little attention was paid to vitamin D status during pediatric critical illness, with only three small studies reporting 25OHD levels on a total of 88 children.^{39,74,75} Since 2012 there have been 10 observational studies evaluating vitamin D status in general PICU populations,^{76–85} two focused on postoperative congenital heart disease (CHD),^{86,87} and a pilot study of high-dose vitamin D in children with burns.⁸⁸ The 10 observational PICU studies originate from six countries and evaluate 2,000 children. Almost all studies were single center, collected only one blood sample (at or near admission), and evaluated vitamin D status using total 25OHD. Further, with the exception of Rey and colleagues,⁷⁹ all studies defined VDD using the 50 nmol/L threshold. As shown in ►Fig. 3, prevalence of VDD ranged from 28 to 84%, suggesting significant risk for VDD in the PICU. Three of the studies also sought to determine whether 25OHD levels were lower in critically ill children compared with other children. Hebbar and colleagues showed statistically lower 25OHD levels compared with a group of unwell children receiving care at the same hospital (99 vs. 46 nmol/L).⁸¹ Similarly, in their investigation of PICU patients with sepsis, Ponnarmeni et al demonstrated significantly lower 25OHD levels compared with healthy controls (49.25 vs. 68.7 nmol/L).⁸⁵

In addition to studies on the general PICU populations, there have been two recent observational studies focused on children with CHD.^{86,87} In a secondary analysis, Graham and colleagues reported that 84% of neonates undergoing cardiac surgery had VDD postoperatively. In a prospective Canadian study, mean immediate postoperative vitamin D status was 35 nmol/L (85% VDD). In the pilot RCT of high-dose vitamin D

(100 IU/kg) in severe pediatric burn injury, the mean 25OHD concentration was 50 nmol/L.⁸⁸

Change in 25OHD Levels during Hospital Admission

Available PICU literature evaluates VDD at admission but does not address how vitamin D status changes throughout the hospital stay. Only two of the pediatric studies evaluated 25OHD levels at more than one time in the PICU. The Canada multicenter study did not observe differences in group 25OHD levels between days 1 and 2,⁷⁶ and the prospective longitudinal study of children undergoing CHD surgery demonstrated that 25OHD levels remained stable over a 48-hour period following separation from cardiopulmonary bypass.⁸⁶ Studies from related populations evaluating 25OHD over a longer period provide more useful and concerning findings. In a mixed population of hospitalized children (ward/PICU) Dayal and colleagues observed a significant decline in 25OHD from admission (72 nmol/L) to discharge (49 nmol/L).⁸⁹ Similar adult studies by Higgins et al⁹⁰ and Yi et al⁹¹ reported a significant decline in 25OHD levels over 10- and 35-day study periods, respectively. The placebo group in a phase III RCT of high-dose vitamin D in critically ill adults found no significant change in 25OHD (mean, 35 nmol/L) over the first 30 days.²³

Vitamin D Deficiency and Critical Illness

Similar to the general population, reduced consumption of specific foods and inadequate UV exposure (sun avoidance, sunscreen, skin melanin content, high latitude) predisposes many children to preillness VDD. It is also increasingly evident that critical illness and related interventions (surgery, fluids, extracorporeal membrane oxygenation [ECMO], cardiopulmonary bypass, plasma exchange) may significantly reduce vitamin D levels.⁸⁶ Disruption of the hepatic, parathyroid and renal organ function (reducing hydroxylation) and greater tissue requirements during catabolism may also reduce 25OHD. Acute changes in 25OHD may be more harmful than chronically low levels as the endocrine axis and individual cells require time to adjust to new steady-state levels.

Association of Vitamin D Status with Illness Severity and Clinical Outcome

Adult ICU Observational Studies

Since 2009, dozens of observational studies have investigated the role of vitamin D in more than 10,000 critically ill adults. Many of these studies estimate concerning VDD rates and report associations with illness severity, organ dysfunction, complications, clinical course, and/or mortality.^{7,12–15,19,29–31} In 2014, two systematic reviews synthesized the available literature and calculating an almost twofold greater risk of in-hospital death in VDD.^{15,16} Further meta-analysis demonstrated statistically significant differences in sepsis,¹⁵ culture positive infection,¹⁵ and a descriptive analysis suggested an association with increased length of stay.¹⁶ Three large studies, linking hospital laboratory and




Fig. 2 Example thresholds for vitamin D status. Vitamin D thresholds suggested by various agencies and societies to define vitamin D deficiency (dark bar), vitamin D insufficiency (gray bar), and sufficiency (white bar). The 25 hydroxyvitamin D levels suggested to be associated with adverse events (AE) or toxicity are shown to the right of the bar. Abbreviations: AAP, American Academy of Pediatrics; CPS, Canadian Pediatric Society; ES, Endocrine Society; IOM, Institute of Medicine; NR, not reported.

admission data, determined preillness vitamin D levels to be associated with in-hospital mortality.^{92–94} Given the convincing quantity and quality of the observational evidence, multiple adult research groups have proceeded to clinical trials.

PICU Observational Studies

In addition to describing vitamin D status, all the general PICU studies assessed associations between 25OHD and at least one marker of illness severity or clinical outcome. As shown in **Table 1**, there is considerable variability in the number and type of measures reported. This lack of consistency, combined with the application of different measurement tools and considerable differences in sample size, makes it challenging to interpret and synthesize the findings. Mortality, the clinical outcome arguably of greatest importance and least subjectivity was reported in six publications. Unfortunately, no individual PICU study was sufficiently powered to investigate an association with mortality. However, it is worth noting that with the exception of the study by Rippel and

colleagues,⁷⁷ the five remaining studies all reported higher mortality rates in the VDD groups. With the cumulative number of children approaching 2,000, it would be worthwhile combining individual study findings as part of a systematic review and meta-analysis. Of the remaining associations investigated, the two most common included illness severity ($n = 10$) and cardiovascular support ($n = 8$). Overall, an association between illness severity and VDD is strongly suggested as four studies reported statistically significant associations and three studies calculated trends toward statistical significance. Similarly, among the eight studies evaluating cardiovascular outcomes, four reported significantly more cardiovascular support in at least one analysis and two others reported clinically meaningful differences that did not achieve statistical significance. A significant association between 25OHD levels and cardiovascular support was also reported in both studies focused on postoperative CHD patients.^{86,87} It is worth commenting on the findings from a few specific studies. First, the lone multicenter study




Fig. 3 Prevalence of vitamin D receptor (VDD) in PICU observational studies. This figure shows the prevalence of VDD in the 10 general PICU observational studies. With the exception of Rey and colleagues (37.5 nmol/L), VDD was defined using the 50 nmol/L threshold.

Table 1 Published PICU observational studies reporting on vitamin D status and outcomes

	McNally 2012	Madden 2012	Rippel 2012	Rey 2014	Ayulo 2014	Hebbar 2014	Prasad 2015	Onwuneme 2015	Ebenezer 2015	Ponnarneni 2015
Sample size	326	511	316	156	216	61	80	120	54	120
Location	Canada	USA	Australia	Spain	USA	USA	India	Ireland	India	India
Age (eligibility)	0–17 y	< 21 y	NR	< 16 y	1–21 y	0–18 y	2 m–12 y	< 12 y	All	1–12 y
Illness severity and outcome measures										
Inotropes/catecholamines	+	+	+	+			+	+	+	+
Calcium/phosphate	+	+	+		+		+		+	+
Fluid requirements	+							+		
Ventilation	+	+	+	+			+	+	+	+
Admission severity score	+	+	+	+	+	+	+	+	+	+
Organ dysfunction						+		+		+
LOS—PICU	+		+	+			+	+	+	+
LOS—hospital			+				+			
Mortality	+		+		+		+		+	+
Coagulopathy/platelets				+			+	+		
Renal failure/dialysis								+		
Hypoglycemia/steroid							+			
Infectious			+			+	+	+	+	+
Inflammatory marker				+				+	+	

Abbreviations: LOS, length of stay; PICU, pediatric intensive care unit.

demonstrated that VDD was associated with a significantly increased length of PICU stay (2 days) after adjusting for important covariates including PRISM (Pediatric Risk of Mortality Score).⁷⁶ Second, an Irish study focused on PICU patients with sepsis and the authors reported significantly higher rates of culture proven sepsis in the VDD group.⁸³ Finally, although the cumulative evidence supports a role for VDD in PICU, it cannot be overlooked that two moderately sized studies failed to find associations with most clinical outcome measures.^{77,85} Based on existing work, it would appear that a large multicenter study enrolling a thousand patients, and correcting for important confounders, would be required to further solidify the relationship between 25OHD levels and clinical course. As this observational study would still leave the more important question unanswered, it might be more appropriate to use the resources for a large multicenter interventional.

Approved Vitamin D Supplementation Regimens

Usual Care

Because of the known negative health consequences of VDD, it is recommended that children intake a minimum quantity of vitamin D. The recommended daily allowance (RDA) or adequate intake (AI) suggested by the Institute of Medicine (IOM) and supported by Health Canada are 400 IU for infants and 600 IU for older children.⁹⁵ Slightly higher doses have been suggested by the Canadian Pediatric Society (CPS) for infants living in Northern Canada (800 IU).⁹⁶ Presently, no standard of care for vitamin D supplementation has been established during or following pediatric critical illness. If ordered at all, a daily dose of vitamin D between 400 and 800 IU is provided enterally or with total parenteral nutrition, but it is well recognized that these doses can take 2 or more months to restore vitamin D status in stable, otherwise healthy children.^{73,97} As previously discussed, available evidence on hospitalized and critically ill patients suggests that vitamin D levels generally remain constant or fall over time with usual care.^{89,90} As this approach leaves critically ill patients in a VDD state over an extended period, alternative dosing regimens need consideration.

Approved High-Dose Regimens

In addition to RDA, the IOM provides a higher age-specific dose called the tolerable upper intake level (UL), which ranges from 1,000 to 4,000 IU.⁹⁵ When given daily, this dose is intended to gradually elevate vitamin D into the high normal range, while safely avoiding toxicity. The UL provided by the IOM was based on evidence from healthy children, and serves as a starting point for research on high-risk populations. A recent systematic review of pediatric clinical trials evaluated dosing in the 1,000 to 4,000 IU range, demonstrating that most patients would not achieve normal levels until after a month of treatment.⁹⁸ Further concerning, this systematic review also determined that children who are unwell may have a blunted 25OHD response suggesting that normalization of vitamin D status in deficient critically ill patients using

a daily dosing approach may be even slower. A recent pilot RCT evaluating a regimen of 800 IU + 100 IU/kg in pediatric burn patients confirmed these concerns showing that half of the patients remained VDD at the projected midpoint of their hospital admission.⁸⁸ Altogether, it appears that approved high-dose daily regimens will not restore vitamin D status in a time frame beneficial for the critically ill child.

Alternative Vitamin D Supplementation Strategies

Loading Dose Vitamin D (Stoss Therapy)

The supplementation strategy most appropriate for the ICU setting is loading dose (or Stoss therapy) where 40,000 to 600,000 IU are provided as single or divided doses. A recent systematic review sought to understand 25OHD response to loading dose therapy in children.⁹⁸ Although dozens of studies administering loading dose therapy were identified, none included critically ill children. Evaluation of the change in levels of 25OHD demonstrated that vitamin D levels can be normalized within 48 hours, but that weight-based dosing was necessary to avoid both under- and overdosing. Ultimately, 10,000 IU/kg was proposed as the dose for consideration in future trials.

Trials of High-Dose Vitamin D in Related Populations

There is some clinical trial evidence to suggest that loading therapy may improve outcomes in related non-PICU populations. Primary among the evidence is VITdAL-ICU trial, where 540,000 IU (enteral) and placebo were compared in critically ill adults. Although there was no difference in length of stay (primary objective), there was a clinically meaningful difference in mortality (43 vs. 35%, $p = 0.09$).²³ Further subgroup analysis suggested an interaction between outcome and baseline 25(OH)D level, as (1) participants with baseline 25(OH)D under 30 nmol/L had a statistically significant mortality reduction (50 vs. 35%, $p = 0.02$), and (2) participants with baseline 25(OH)D between 30 and 50 nmol/L showing improved grip strength and physical components of the SF-12 questionnaire. Two pediatric studies evaluating loading dose therapy in ill populations have also suggested benefit. First, an RCT of 450 young children presenting with acute lower respiratory tract infection demonstrated that a 100,000-IU enteral dose reduced repeat episodes.⁹⁹ Second, a recent systematic review of pediatric clinical trials of vitamin D in asthma demonstrated that high-dose vitamin D supplementation, including a 60,000-IU monthly loading regimen, reduced asthma exacerbations by 50%.^{100,101}

Safety of High-Dose Vitamin D

As an inexpensive and simple intervention, it would be understandable for PICUs to consider overlooking the lack of high-quality clinical trial evidence and begin implementation of high-dose therapy. Preventing this are the concerns spawned from case reports, case series, and a small number of clinical trials establishing that inappropriate vitamin D intake, resulting in supraphysiological 25OHD

levels, can cause toxicity.^{102,103} Vitamin D toxicity is characterized by hypercalcemia or hypercalciuria, with the classic symptoms (lethargy, abdominal pain, anorexia, constipation, polyuria, and nocturia) directly attributable to these abnormalities. With prolonged states of hypercalcemia and hypercalciuria, children are at risk for developing nephrocalcinosis and other renal pathology. However, there is considerable evidence to suggest that loading dose therapy, designed with the goal of achieving normal vitamin D levels, will be safe. First, studies in healthy children receiving regimens approximating the UL did not demonstrate toxicity with cumulative dosing approximating 10,000 IU/kg.^{73,97,104} Second, a review of pediatric nephrocalcinosis studies identified that cases attributed to vitamin D occur in the context of one or more doses of greater than 600,000 IU administered to healthy children or those with genetic abnormalities of VDR.^{105–109} Third, an adverse event analysis performed as part of a systematic review of high-dose pediatric trials did not find evidence of toxicity until dosing exceeded 400,000 IU. Fourth, the VITdAL-ICU study did not demonstrate increased hypercalcemia, hypercalciuria, or nephrocalcinosis in the group of vitamin D–deficient critically ill adults that received 540,000 IU.²³ Fifth, pediatric case reports and case series of clinical and subclinical cardiac dysfunction secondary to VDD describe improvements in patient status with both gradual and rapid restoration of vitamin D levels.^{57–61,110} Finally, pediatric RCTs evaluating loading dose therapy in healthy and stable unwell pediatric populations (e.g., pneumonia, asthma) have not suggested safety con-

cerns.^{99–101,111} Altogether, these findings strongly suggest that rapid normalization of vitamin D status in critically ill children could have more benefit than harm, but further study is necessary to confirm this.

Controversies and Knowledge Gaps

Are There Better Markers for Vitamin D Status for ICU?

Given the high incidence of hepatic, renal, and parathyroid organ dysfunction, it has been suggested that calcitriol might represent a superior marker of vitamin D status in critically ill patients. Two adult studies have been published to date evaluating this question, with neither showing calcitriol to be superior or additive to calcidiol in the relationship with illness severity or clinical outcome.^{112,113} A large pediatric study published in 2015 further demonstrated that although associations with interventions and clinical course were evident, they disappeared once 25OHD was considered.¹¹⁴ More recently, groups have suggested that bioavailable 25OHD might be a better measure, given alterations in VDBP and albumin levels during critical illness. The limited adult and pediatric work in this area have not demonstrated total 25OHD to be inferior,^{112,115} but further study is warranted.

Uncertainty Surrounding the Best Dosing Regimen

To date, seven pilot and one phase III trials have been published in various adult ICU settings (►Table 2). With the exception of the pilot performed prior to the phase III VITdAL-ICU, none of studies used the same combination of

Table 2 Vitamin D dosing regimens considered to date as part of clinical trials in ICU

	Population	n	Metabolite	Dose	Frequency	Route	Usual care arm	Comment
Pediatric—pilot								
Gottschlich, 2015	Burns	18	Ergocalciferol	100 IU/kg	Daily	PO	Yes	
Gottschlich, 2015	Burns	15	Cholecalciferol	100 IU/kg	Daily	PO	Yes	
Adult—pilot								
Ingels, 2010	General	11	Calcidiol	8,000 IU load, 600 IU/d	10 d	IV	Yes	Abstract only
Mata-Granados, 2010	Sepsis	11	Calcidiol	60,000 IU	Days: 0, 4	PO	Yes	Non-RCT
Mata-Granados, 2010	Sepsis	10	Calcitriol	2.0 mcg	Days: 0, 2, 4, 6	IV	Yes	
Amrein, 2011	General, VDD	10	Cholecalciferol	540,000 IU	Once	PO	Yes	
Leaf, 2014	Severe sepsis	36	Calcitriol	2.0 µg	Once	IV	Yes	
Nair, 2015	Sepsis/SIRS	25	Cholecalciferol	150,000 IU	Once	IM	No	
Nair, 2015	Sepsis/SIRS	25	Cholecalciferol	300,000 IU	Once	IM	No	
Dickerson, 2015	Trauma	16	Ergocalciferol	50,000 IU	Once weekly	PO	No	
Dickerson, 2015	Trauma	18	Ergocalciferol	50,000 IU	Twice weekly	PO	No	
Dickerson, 2015	Trauma	31	Ergocalciferol	50,000 IU	Thrice weekly	PO	No	
Rousseau, 2015	Burn	13	Cholecalciferol	200,000 IU	Q3m	IM	Yes	
Amrein, 2014	General	249	Cholecalciferol	540,000 IU then 90,000 IU	Once	Once + per month	Yes	Phase III RCT

Abbreviations: ICU, intensive care unit; IM, intramuscular; IV, intravenous; PO, oral; RCT, randomized clinical trial; SIRS, systemic inflammatory response syndrome; VDD, vitamin D deficiency.

metabolite, dose, frequency, and route. Demonstrating the variability, the metabolites selected in nonplacebo arms included cholecalciferol ($n = 5$), ergocalciferol ($n = 3$), calcidiol ($n = 2$), and calcitriol ($n = 2$). Within the D3 and D2 arms, there were five doses (50,000–540,000 IU) and four frequency options. Importantly, roughly half of the arms utilized intramuscular administration, which has the disadvantage of a longer time to reach peak effect and lower final 25OHD concentration. Although calcidiol is a reasonable option for supplementation, there are valid arguments against the use of calcitriol, the active hormone: (1) increased expense relative to cholecalciferol, (2) knowledge that calcitriol levels do not correlate better with clinical outcome, and (3) observations that loading therapy with D3 or 25OHD results in a rapid rise in calcitriol (two to three fold). Finally, the suggestive results of the VITdAL-ICU trial support further evaluation of enteral loading dose cholecalciferol, but given that many ICU patients cannot receive enteral medications for prolonged periods, it would be reasonable to also explore an IV preparation of cholecalciferol or 25OHD.

Potential Impact and Next Steps

Critical illness occurs in millions of children worldwide every year. In addition to death, these children are at risk for significant morbidity due to organ dysfunction, prolonged periods of rehabilitation, and chronic disease. Given the potential importance of vitamin D to the health and stress response of multiple organ systems, the high VDD rate reported in many ICUs should be of concern. Cumulative observational and clinical trial data suggest that rapid normalization of vitamin D status could represent a simple, inexpensive, and safe means of improving outcomes and reducing health care spending. Unfortunately many of the approved daily dosing regimens for vitamin D can require months to restore levels. Loading therapy represents a more appropriate approach for restoring vitamin D status in critically ill children. Unfortunately, there have been no studies of loading therapy in the PICU setting. Consequently, despite significant literature suggesting VDD to be a modifiable risk factor in critical illness, there is no robust evidence to inform us on how to rapidly normalize levels or the true benefits or risks. RCTs including dose-evaluation pilots and phase III RCTs will be required.

Funding

None.

Conflict of interest

None.

References

- 1 Ward LM, Gaboury I, Ladhani M, Zlotkin S. Vitamin D-deficiency rickets among children in Canada. *CMAJ* 2007;177(2):161–166
- 2 Mehrotra P, Marwaha RK, Aneja S, et al. Hypovitaminosis d and hypocalcemic seizures in infancy. *Indian Pediatr* 2010;47(7):581–586
- 3 Basatemur E, Sutcliffe A. Incidence of hypocalcemic seizures due to vitamin D deficiency in children in the United Kingdom and Ireland. *J Clin Endocrinol Metab* 2015;100(1):E91–E95
- 4 Ginde AA, Mansbach JM, Camargo CA Jr. Vitamin D, respiratory infections, and asthma. [Review] [53 refs] *Curr Allergy Asthma Rep* 2009;9(1):81–87
- 5 Kim SY. The pleiomorphic actions of vitamin D and its importance for children. *Ann Pediatr Endocrinol Metab* 2013;18(2):45–54
- 6 Zittermann A, Gummert JF. Nonclassical vitamin D action. *Nutrients* 2010;2(4):408–425
- 7 Lee P, Nair P, Eisman JA, Center JR. Vitamin D deficiency in the intensive care unit: an invisible accomplice to morbidity and mortality? *Intensive Care Med* 2009;35(12):2028–2032
- 8 Youssef D, Amrein K, Schnedl C, Dobnig H, Peiris AN. Intensive Care and Vitamin D status. *Diet and Nutrition in Critical Care*. Vol 1. New York, NY: Springer; 2014:16
- 9 Amrein K, Venkatesh B. Vitamin D and the critically ill patient. *Curr Opin Clin Nutr Metab Care* 2012;15(2):188–193
- 10 Christopher KB. Vitamin D supplementation in the ICU patient. *Curr Opin Clin Nutr Metab Care* 2015;18(2):187–192
- 11 Lee P. Vitamin D metabolism and deficiency in critical illness. *Best Pract Res Clin Endocrinol Metab* 2011;25(5):769–781
- 12 Amrein K, Christopher KB, McNally JD. Understanding vitamin D deficiency in intensive care patients. *Intensive Care Med* 2015;41(11):1961–1964
- 13 Williams S, Heuberger R. Outcomes of vitamin D supplementation in adults who are deficient and critically ill: a review of the literature. *Am J Ther* 2015
- 14 Zajic P, Amrein K. Vitamin D deficiency in the ICU: a systematic review. *Minerva Endocrinol* 2014;39(4):275–287
- 15 de Haan K, Groeneveld AB, de Geus HR, Egal M, Struijs A. Vitamin D deficiency as a risk factor for infection, sepsis and mortality in the critically ill: systematic review and meta-analysis. *Crit Care* 2014;18(6):660
- 16 Zhang YP, Wan YD, Sun TW, Kan QC, Wang LX. Association between vitamin D deficiency and mortality in critically ill adult patients: a meta-analysis of cohort studies. *Crit Care* 2014;18(6):684
- 17 Amrein K, Sourij H, Wagner G, et al. Short-term effects of high-dose oral vitamin D3 in critically ill vitamin D deficient patients: a randomized, double-blind, placebo-controlled pilot study. *Crit Care* 2011;15(2):R104–R104
- 18 Mata-Granados JM, Vargas-Vassero J, Ferreira-Vera C, Luque de Castro MD, Pavón RG, Quesada Gómez JM. Evaluation of vitamin D endocrine system (VDES) status and response to treatment of patients in intensive care units (ICUs) using an on-line SPE-LC-MS/MS method. *J Steroid Biochem Mol Biol* 2010;121(1-2):452–455
- 19 Nair P, Venkatesh B, Lee P, et al. A randomized study of a single dose of intramuscular cholecalciferol in critically ill adults. *Crit Care Med* 2015;43(11):2313–2320
- 20 Dickerson RN, Berry SC, Ziebarth JD, et al. Dose-response effect of ergocalciferol therapy on serum 25-hydroxyvitamin D concentration during critical illness. *Nutrition* 2015;31(10):1219–1223
- 21 Leaf DE, Raed A, Donnino MW, Ginde AA, Waikar SS. Randomized controlled trial of calcitriol in severe sepsis. *Am J Respir Crit Care Med* 2014;190(5):533–541
- 22 Aminmansour B, Nikbakht H, Ghorbani A, et al. Comparison of the administration of progesterone versus progesterone and vitamin D in improvement of outcomes in patients with traumatic brain injury: a randomized clinical trial with placebo group. *Adv Biomed Res* 2012;1:58
- 23 Amrein K, Schnedl C, Holl A, et al. Effect of high-dose vitamin D3 on hospital length of stay in critically ill patients with vitamin D deficiency: the VITdAL-ICU randomized clinical trial. *JAMA* 2014;312(15):1520–1530

- 24 Abou-Zahr R, Kandil SB. A pediatric critical care perspective on vitamin D. *Pediatr Res* 2015;77(1-2):164–167
- 25 Shoback D. Clinical practice. Hypoparathyroidism. *N Engl J Med* 2008;359(4):391–403
- 26 Nibbelink KA, Tishkoff DX, Hershey SD, Rahman A, Simpson RU. 1,25(OH)₂-vitamin D₃ actions on cell proliferation, size, gene expression, and receptor localization, in the HL-1 cardiac myocyte. *J Steroid Biochem Mol Biol* 2007;103(3-5):533–537
- 27 Haussler MR, Jurutka PW, Mizwicki M, Norman AW. Vitamin D receptor (VDR)-mediated actions of 1 α ,25(OH)₂ vitamin D₃: genomic and non-genomic mechanisms. *Best Pract Res Clin Endocrinol Metab* 2011;25(4):543–559
- 28 Green JJ, Robinson DA, Wilson GE, Simpson RU, Westfall MV. Calcitriol modulation of cardiac contractile performance via protein kinase C. *J Mol Cell Cardiol* 2006;41(2):350–359
- 29 Santillán GE, Vazquez G, Boland RL. Activation of a beta-adrenergic-sensitive signal transduction pathway by the secosteroid hormone 1,25-(OH)₂-vitamin D₃ in chick heart. *J Mol Cell Cardiol* 1999;31(5):1095–1104
- 30 Tishkoff DX, Nibbelink KA, Holmberg KH, Dandu L, Simpson RU. Functional vitamin D receptor (VDR) in the t-tubules of cardiac myocytes: VDR knockout cardiomyocyte contractility. *Endocrinology* 2008;149(2):558–564
- 31 Holick MF, Binkley NC, Bischoff-Ferrari HA, et al; Endocrine Society. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab* 2011;96(7):1911–1930
- 32 Wagner CL, Greer FR; American Academy of Pediatrics Section on Breastfeeding; American Academy of Pediatrics Committee on Nutrition. Prevention of rickets and vitamin D deficiency in infants, children, and adolescents. *Pediatrics* 2008;122(5):1142–1152
- 33 Willett AM. Vitamin D status and its relationship with parathyroid hormone and bone mineral status in older adolescents. *Proc Nutr Soc* 2005;64(2):193–203
- 34 Christensen MH, Lien EA, Hustad S, Almås B. Seasonal and age-related differences in serum 25-hydroxyvitamin D, 1,25-dihydroxyvitamin D and parathyroid hormone in patients from Western Norway. *Scand J Clin Lab Invest* 2010;70(4):281–286
- 35 Docio S, Riancho JA, Pérez A, Olmos JM, Amado JA, González-Macías J. Seasonal deficiency of vitamin D in children: a potential target for osteoporosis-preventing strategies? *J Bone Miner Res* 1998;13(4):544–548
- 36 Thacher TD, Fischer PR, Isichei CO, Pettifor JM. Early response to vitamin D₂ in children with calcium deficiency rickets. *J Pediatr* 2006;149(6):840–844
- 37 Lee P, Eisman JA, Center JR. Vitamin D deficiency in critically ill patients. *N Engl J Med* 2009;360(18):1912–1914
- 38 Cardenas-Rivero N, Chernow B, Stoiko MA, Nussbaum SR, Todres ID. Hypocalcemia in critically ill children. *J Pediatr* 1989;114(6):946–951
- 39 Gauthier B, Trachtman H, Di Carmine F, et al. Hypocalcemia and hypercalcitoninemia in critically ill children. *Crit Care Med* 1990;18(11):1215–1219
- 40 Iqbal M, Rehmani R, Hijazi M, Abdulaziz A, Kashif S. Hypocalcemia in a Saudi intensive care unit. *Ann Thorac Med* 2008;3(2):57–59
- 41 Dyke PC II, Yates AR, Cua CL, et al. Increased calcium supplementation is associated with morbidity and mortality in the infant postoperative cardiac patient. *Pediatr Crit Care Med* 2007;8(3):254–257
- 42 Broner CW, Stidham GL, Westenkirchner DF, Tolley EA. Hypomagnesemia and hypocalcemia as predictors of high mortality in critically ill pediatric patients. *Crit Care Med* 1990;18(9):921–928
- 43 Kozik DJ, Tweddell JS. Characterizing the inflammatory response to cardiopulmonary bypass in children. *Ann Thorac Surg* 2006;81(6):S2347–S2354
- 44 Jagers J, Lawson JH. Coagulopathy and inflammation in neonatal heart surgery: mechanisms and strategies. *Ann Thorac Surg* 2006;81(6):S2360–S2366
- 45 Baekke F, Gysemans C, Korf H, Mathieu C. Vitamin D insufficiency: implications for the immune system. *Pediatr Nephrol* 2010;25(9):1597–1606
- 46 Rigby WF, Denome S, Fanger MW. Regulation of lymphokine production and human T lymphocyte activation by 1,25-dihydroxyvitamin D₃. Specific inhibition at the level of messenger RNA. *J Clin Invest* 1987;79(6):1659–1664
- 47 Bhalla AK, Amento EP, Serog B, Glimcher LH. 1,25-Dihydroxyvitamin D₃ inhibits antigen-induced T cell activation. *J Immunol* 1984;133(4):1748–1754
- 48 Hata TR, Kotol P, Jackson M, et al. Administration of oral vitamin D induces cathelicidin production in atopic individuals. *J Allergy Clin Immunol* 2008;122(4):829–831
- 49 Gombart AF, Borregaard N, Koeffler HP. Human cathelicidin antimicrobial peptide (CAMP) gene is a direct target of the vitamin D receptor and is strongly up-regulated in myeloid cells by 1,25-dihydroxyvitamin D₃. *FASEB J* 2005;19(9):1067–1077
- 50 Jeng L, Yamshchikov AV, Judd SE, et al. Alterations in vitamin D status and anti-microbial peptide levels in patients in the intensive care unit with sepsis. *J Transl Med* 2009;7:28–28
- 51 Gombart AF, Bhan I, Borregaard N, et al. Low plasma level of cathelicidin antimicrobial peptide (hCAP18) predicts increased infectious disease mortality in patients undergoing hemodialysis. *Clin Infect Dis* 2009;48(4):418–424
- 52 Leaf DE, Croy HE, Abrahams SJ, Raed A, Waikar SS. Cathelicidin antimicrobial protein, vitamin D, and risk of death in critically ill patients. *Crit Care* 2015;19:80
- 53 McNally JD, Sampson M, Matheson LA, Hutton B, Little J. Vitamin D receptor (VDR) polymorphisms and severe RSV bronchiolitis: a systematic review and meta-analysis. *Pediatr Pulmonol* 2014;49(8):790–799
- 54 Brehm JM, Schuermann B, Fuhlbrigge AL, et al; Childhood Asthma Management Program Research Group. Serum vitamin D levels and severe asthma exacerbations in the Childhood Asthma Management Program study. *J Allergy Clin Immunol* 2010;126(1):52–8.e5
- 55 Maxwell CS, Carbone ET, Wood RJ. Better newborn vitamin D status lowers RSV-associated bronchiolitis in infants. *Nutr Rev* 2012;70(9):548–552
- 56 Menon K, Ward RE, Lawson ML, Gaboury I, Hutchison JS, Hébert PC; Canadian Critical Care Trials Group. A prospective multicenter study of adrenal function in critically ill children. *Am J Respir Crit Care Med* 2010;182(2):246–251
- 57 Olgun H, Ceviz N, Ozkan B. A case of dilated cardiomyopathy due to nutritional vitamin D deficiency rickets. *Turk J Pediatr* 2003;45(2):152–154
- 58 Price DI, Stanford LC Jr, Braden DS, Ebeid MR, Smith JC. Hypocalcemic rickets: an unusual cause of dilated cardiomyopathy. *Pediatr Cardiol* 2003;24(5):510–512
- 59 Verma S, Khadwal A, Chopra K, Rohit M, Singhi S. Hypocalcemia nutritional rickets: a curable cause of dilated cardiomyopathy. *J Trop Pediatr* 2011;57(2):126–128
- 60 Maiya S, Sullivan I, Allgrove J, et al. Hypocalcaemia and vitamin D deficiency: an important, but preventable, cause of life-threatening infant heart failure. *Heart* 2008;94(5):581–584
- 61 Uysal S, Kalayci AG, Baysal K. Cardiac functions in children with vitamin D deficiency rickets. *Pediatr Cardiol* 1999;20(4):283–286
- 62 Shedeed SA. Vitamin D supplementation in infants with chronic congestive heart failure. *Pediatr Cardiol* 2012;33(5):713–719
- 63 Banwell BL, Mildner RJ, Hassall AC, Becker LE, Vajsaar J, Shemie SD. Muscle weakness in critically ill children. *Neurology* 2003;61(12):1779–1782
- 64 Williams S, Horrocks IA, Ouvrier RA, Gillis J, Ryan MM. Critical illness polyneuropathy and myopathy in pediatric intensive care: a review. *Pediatr Crit Care Med* 2007;8(1):18–22

- 65 Kress JP, Hall JB. ICU-acquired weakness and recovery from critical illness. *N Engl J Med* 2014;370(17):1626–1635
- 66 Batt J, dos Santos CC, Cameron JL, Herridge MS. Intensive care unit-acquired weakness: clinical phenotypes and molecular mechanisms. *Am J Respir Crit Care Med* 2013;187(3):238–246
- 67 Al-Said YA, Al-Rached HS, Al-Qahtani HA, Jan MMS. Severe proximal myopathy with remarkable recovery after vitamin D treatment. *Can J Neurol Sci* 2009;36(3):336–339
- 68 Al-Said YA, Al-Rached HS, Al-Qahtani HA, Jan MM. Severe proximal myopathy with remarkable recovery after vitamin D treatment. *Can J Neurol Sci* 2009;36(3):336–339
- 69 Ward KA, Das G, Berry JL, et al. Vitamin D status and muscle function in post-menarchal adolescent girls. *J Clin Endocrinol Metab* 2009;94(2):559–563
- 70 van der Heyden JJ, Verrips A, ter Laak HJ, Otten B, Fiselier T. Hypovitaminosis D-related myopathy in immigrant teenagers. *Neuropediatrics* 2004;35(5):290–292
- 71 Torun E, Genç H, Gönüllü E, Akovali B, Ozgen IT. The clinical and biochemical presentation of vitamin D deficiency and insufficiency in children and adolescents. *J Pediatr Endocrinol Metab* 2013;26(5-6):469–475
- 72 Hazell TJ, Gallo S, Berzina I, Vanstone CA, Rodd C, Weiler HA. Plasma 25-hydroxyvitamin D, more so than its epimer, has a linear relationship to leaner body composition across infancy in healthy term infants. *Appl Physiol Nutr Metab* 2014;39(10):1137–1143
- 73 Gallo S, Comeau K, Vanstone C, et al. Effect of different dosages of oral vitamin D supplementation on vitamin D status in healthy, breastfed infants: a randomized trial. *JAMA* 2013;309(17):1785–1792
- 74 McNally JD, Leis K, Matheson LA, Karuananyake C, Sankaran K, Rosenberg AM. Vitamin D deficiency in young children with severe acute lower respiratory infection. *Pediatr Pulmonol* 2009;44(10):981–988
- 75 Gottschlich MM, Mayes T, Khoury J, Warden GD. Hypovitaminosis D in acutely injured pediatric burn patients. *J Am Diet Assoc* 2004;104(6):931–941, quiz 1031
- 76 McNally JD, Menon K, Chakraborty P, et al; Canadian Critical Care Trials Group. The association of vitamin D status with pediatric critical illness. *Pediatrics* 2012;130(3):429–436
- 77 Rippel C, South M, Butt WW, Shekerdemian LS. Vitamin D status in critically ill children. *Intensive Care Med* 2012;38(12):2055–2062
- 78 Madden K, Feldman HA, Smith EM, et al. Vitamin D deficiency in critically ill children. *Pediatrics* 2012;130(3):421–428
- 79 Rey C, Sánchez-Arango D, López-Herce J, et al. Vitamin D deficiency at pediatric intensive care admission. *J Pediatr (Rio J)* 2014;90(2):135–142 (Versão em Português)
- 80 Ayulo MJr, Katyal AG, Agarwal Ch, et al. The prevalence of vitamin D deficiency and its relationship with disease severity in an urban pediatric critical care unit. *Endocr Regul* 2014;48(2):69–76
- 81 Hebbar KB, Wittkamp M, Alvarez JA, McCracken CE, Tangpricha V. Vitamin D. Vitamin D deficiency in pediatric critical illness. *J Clin Transl Endocrinol* 2014;1(4):170–175
- 82 Prasad S, Raj D, Warsi S, Chowdhary S. Vitamin D. Vitamin D deficiency and critical illness. *Indian J Pediatr* 2015;82(11):991–995
- 83 Onwuneme C, Carroll A, Doherty D, et al. Inadequate vitamin D levels are associated with culture positive sepsis and poor outcomes in paediatric intensive care. *Acta Paediatr* 2015;104(10):e433–e438
- 84 Ebenezer K, Job V, Antonisamy B, Dawodu A, Manivachagan MN, Steinhoff M. Serum vitamin D status and outcome among critically ill children admitted to the pediatric intensive care unit in South India. *Indian J Pediatr* 2015
- 85 Ponnarmeni S, Kumar Angurana S, Singhi S, et al. Vitamin D deficiency in critically ill children with sepsis. *Paediatr Int Child Health* 2016;36(1):15–21
- 86 McNally JD, Menon K, Chakraborty P, et al. Impact of anesthesia and surgery for congenital heart disease on the vitamin D status of infants and children: a prospective longitudinal study. *Anesthesiology* 2013;119(1):71–80
- 87 Graham EM, Taylor SN, Zyblewski SC, et al. Vitamin D status in neonates undergoing cardiac operations: relationship to cardiopulmonary bypass and association with outcomes. *J Pediatr* 2013;162(4):823–826
- 88 Gottschlich MM, Mayes T, Khoury J, Kagan RJ. Clinical Trial of Vitamin D2 vs D3 Supplementation in Critically Ill Pediatric Burn Patients. *JPEN J Parenter Enteral Nutr* 2015
- 89 Dayal D, Kumar S, Sachdeva N, Kumar R, Singh M, Singhi S. Fall in vitamin D levels during hospitalization in children. *Int J Pediatr* 2014;2014:291856
- 90 Higgins DM, Wischmeyer PE, Queensland KM, Sillau SH, Sufit AJ, Heyland DK. Relationship of vitamin D deficiency to clinical outcomes in critically ill patients. *JPEN J Parenter Enteral Nutr* 2012;36(6):713–720
- 91 Yi HJ, Jeong JH, Jin ES, Shin IY, Hwang HS, Moon SM. Evaluation of vitamin D level in patients from neurosurgical intensive care unit. *Neural Regen Res* 2013;8(16):1528–1534
- 92 Lange N, Litonjua AA, Gibbons FK, Giovannucci E, Christopher KB. Pre-hospital vitamin D concentration, mortality, and bloodstream infection in a hospitalized patient population. *Am J Med* 2013;126(7):640.e19–640.e27
- 93 Amrein K, Litonjua AA, Moromizato T, et al. Increases in pre-hospitalization serum 25(OH)D concentrations are associated with improved 30-day mortality after hospital admission: A cohort study. *Clin Nutr* 2015
- 94 Braun A, Chang D, Mahadevappa K, et al. Association of low serum 25-hydroxyvitamin D levels and mortality in the critically ill. *Crit Care Med* 2011;39(4):671–677
- 95 Ross AC, Manson JE, Abrams SA, et al. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab* 2011;96(1):53–58
- 96 Godel JC. Vitamin D supplementation: recommendations for Canadian mothers and infants. *Paediatr Child Health (Oxford)* 2007;12(7):583–598
- 97 Holmlund-Suila E, Viljakainen H, Hytinen T, Lamberg-Allardt C, Andersson S, Mäkitie O. High-dose vitamin D intervention in infants—effects on vitamin D status, calcium homeostasis, and bone strength. *J Clin Endocrinol Metab* 2012;97(11):4139–4147
- 98 McNally JD, Iliriani K, Pojsupap S, et al. Rapid normalization of vitamin D levels: a meta-analysis. *Pediatrics* 2015;135(1):e152–e166
- 99 Manaseki-Holland S, Qader G, Isaq Masher M, et al. Effects of vitamin D supplementation to children diagnosed with pneumonia in Kabul: a randomised controlled trial. *Trop Med Int Health* 2010;15(10):1148–1155
- 100 Pojsupap S, Iliriani K, Sampaio TZ, et al. Efficacy of high-dose vitamin D in pediatric asthma: a systematic review and meta-analysis. *J Asthma* 2014:1–9
- 101 Yadav M, Mittal K. Effect of vitamin D supplementation on moderate to severe bronchial asthma. *Indian J Pediatr* 2014;81(7):650–654
- 102 Joshi R. Hypercalcemia due to hypervitaminosis D: report of seven patients. *J Trop Pediatr* 2009;55(6):396–398
- 103 Markestad T, Hesse V, Siebenhuner M, et al. Intermittent high-dose vitamin D prophylaxis during infancy: effect on vitamin D metabolites, calcium, and phosphorus. *Am J Clin Nutr* 1987;46(4):652–658
- 104 Lewis RD, Laing EM, Hill Gallant KM, et al. A randomized trial of vitamin D₃ supplementation in children: dose-response effects on vitamin D metabolites and calcium absorption. *J Clin Endocrinol Metab* 2013;98(12):4816–4825

- 105 Ammenti A, Pelizzoni A, Cecconi M, Molinari PP, Montini G. Nephrocalcinosis in children: a retrospective multi-centre study. *Acta Paediatr* 2009;98(10):1628–1631
- 106 Rönnefarth G, Misselwitz J. Nephrocalcinosis in children: a retrospective survey. Members of the Arbeitsgemeinschaft für pädiatrische Nephrologie. *Pediatr Nephrol* 2000;14(10–11):1016–1021
- 107 Moncrieff MW, Chance GW. Nephrotoxic effect of vitamin D therapy in vitamin D refractory rickets. *Arch Dis Child* 1969;44(237):571–579
- 108 Mantan M, Bagga A, Viridi VS, Menon S, Hari P. Etiology of nephrocalcinosis in northern Indian children. *Pediatr Nephrol* 2007;22(6):829–833
- 109 Alon US. Nephrocalcinosis. *Curr Opin Pediatr* 1997;9(2):160–165
- 110 Kosecik M, Ertas T. Dilated cardiomyopathy due to nutritional vitamin D deficiency rickets. *Pediatr Int* 2007;49(3):397–399
- 111 Manaseki-Holland S, Maroof Z, Bruce J, et al. Effect on the incidence of pneumonia of vitamin D supplementation by quarterly bolus dose to infants in Kabul: a randomised controlled superiority trial. *Lancet* 2012;379(9824):1419–1427
- 112 Quraishi SA, Bittner EA, Blum L, McCarthy CM, Bhan I, Camargo CA Jr. Prospective study of vitamin D status at initiation of care in critically ill surgical patients and risk of 90-day mortality. *Crit Care Med* 2014;42(6):1365–1371
- 113 Nair P, Lee P, Reynolds C, et al. Significant perturbation of vitamin D-parathyroid-calcium axis and adverse clinical outcomes in critically ill patients. *Intensive Care Med* 2013;39(2):267–274
- 114 Dayre McNally J, Menon K, Lawson ML, Williams K, Doherty DR. 1,25 dihydroxyvitamin D levels in pediatric intensive care units: risk factors and association with clinical course. *J Clin Endocrinol Metab* 2015;100(8):2942–5
- 115 Madden K, Feldman HA, Chun RF, et al. Critically ill children have low vitamin d-binding protein, influencing bioavailability of vitamin D. *Ann Am Thorac Soc* 2015;12(11):1654–1661