



Published in final edited form as:

Surg Endosc. 2020 July ; 34(7): 3011–3019. doi:10.1007/s00464-019-07097-y.

Surgeon Delivered Laparoscopic Transversus Abdominis Plane Blocks are Non-Inferior to Anesthesia Delivered Ultrasound Guided Transversus Abdominis Plane Blocks:

A Blinded, Randomized Non-Inferiority Trial

Daniel J Wong, MD¹, Thomas Curran, MD MPH¹, Vitaliy Y Poylin, MD¹, Thomas E Cataldo, MD¹

¹Division of Colon & Rectum Surgery, Beth Israel Deaconess Medical Center, Harvard Medical School, Boston, MA 02215

Abstract

Background—The transversus abdominis plane (TAP) block is an important non-narcotic adjunct for post-operative pain control in abdominal surgery. Surgeons can use laparoscopic guidance for TAP block placement (LTAP), however, direct comparisons to conventional ultrasound guided TAP (UTAPs) have been lacking. The aim of this study is to determine if surgeon placed LTAPs were non-inferior to anesthesia placed UTAPs for post-operative pain control in laparoscopic colorectal surgery.

Methods—This was a prospective, randomized, patient and observer blinded parallel-arm non-inferiority trial conducted at a single tertiary academic center between 2016–2018 on adult patients undergoing laparoscopic colorectal surgery. Narcotic consumption and pain scores were compared for LTAP vs. UTAP for 48 hours post-operatively. The trial was registered at clinicaltrials.gov Identifier [NCT03577912](https://clinicaltrials.gov/ct2/show/study?term=NCT03577912).

Results—60 patients completed the trial (31 UTAP, 29 LTAP) of which 25 patients were female (15 UTAP, 10 LTAP) and the mean ages (SD) were 60.0 (13.6) and 61.5 (14.3) in the UTAP and LTAP groups, respectively. There was no significant difference in post-operative narcotic consumption between UTAP and LTAP at the time of PACU discharge (median [IQR] milligrams of morphine, 1.8 [0–4.5] UTAP vs 0 [0–8.7] LTAP $P=.32$), 6 hours post-operatively (5.4 [1.8–17.1] UTAP vs 3.6 [0–12.6] LTAP $P=.28$) at 12 hours post-operatively (9.0 [3.6–29.4] UTAP vs 7.2 [0.9–22.5] LTAP $P=.51$) at 24 hours post-operatively (9.0 [3.6–29.4] UTAP vs 7.2 [0.9–22.5] LTAP $P=.63$) and 48 hours post-operatively 39.9 [7.5–70.2] UTAP vs 22.2 [7.5–63.8] LTAP $P=.41$). Patient reported pain scores as well as pre-, intra-, and post-operative course were similar between groups. Non-inferiority criteria were met at all post-op time points up to and including 24 hours but not at 48 hours.

Corresponding Author/Reprints: Thomas Cataldo MD, FACS, FASCRS, Beth Israel Deaconess Medical Center, 330 Brookline Avenue, Gryzmish Building 6th floor, Boston, MA 02215, Phone: (617) 667-4159 Fax: (617) 667-2978, Tcatald1@bidmc.harvard.edu

Disclosures

Dr. Wong reports no conflict of interest.

Dr. Curran reports personal fees from KCI Inc, outside the scope of the submitted work.

Dr. Poylin reports no conflict of interest.

Dr. Cataldo reports no conflict of interest.

Conclusions—Surgeon delivered LTAPs are safe, effective and non-inferior to anesthesia administered UTAPs in the immediate post-operative period.

Keywords

TAP block; Laparoscopy; Surgeon; non-inferior

Introduction

Enhanced recovery after surgery (ERAS) programs improve post-operative outcomes and reduces cost in colorectal surgery.[1–3] A multimodal, opiate sparing pain regimen is a central component of ERAS protocols.[4] Benefits of reducing opiates include decreased post-operative nausea, vomiting, and sedation as well as reducing time to return of bowel function and expediting patient recovery.[5] A reduction in opiate consumption postoperatively is particularly relevant in light of the ongoing opiate epidemic.

The transversus abdominis plane (TAP) block, whereby local anesthetic is injected between the internal oblique and transversus abdominis fascia to facilitate blockade of somatic afferents nerves, is an effective, non-narcotic adjunct incorporated into many ERAS protocols. Numerous studies have shown that the TAP block is effective in reducing post-operative pain, opiate requirement, time to return of bowel function and length of stay (LOS).[6–9] The efficacy of the TAP block has been established across a number of other surgical disciplines as well including general, gynecologic, and urologic surgery.[6, 10–12] Correct positioning of the needle in the TAP plane is essential to the safety and efficacy of the procedure. The high rate of peritoneal needle placement during TAP blocks when solely anatomic landmarks and tactile sensation were employed for delivery led to ultrasound guidance becoming standard.[13] Currently, in many institutions anesthesiologists administer the TAP block under ultrasound guidance (UTAP) immediately prior to or following the operation. This requires an additional provider, time, and potentially an avoidable professional fee.

Recently, there have been several reports of localizing the TAP injection using laparoscopic guidance.[14–17] Laparoscopically guided TAP blocks (LTAPs) have the potential to decrease resource utilization including time, provider, and cost. The comparative efficacy of surgeon delivered LTAPs versus anesthesia delivered UTAPs has only twice been studied in colorectal surgery; employing different delivery techniques and in populations with unclear generalizability to the average colorectal surgery practice.[15, 18]

Therefore we aimed to address this gap in the current understanding of the relative efficacy of LTAP and UTAP through a prospective trial. We hypothesized that surgeon delivered LTAPs would be non-inferior to anesthesia delivered UTAPs in terms of post-operative pain control. To test this, we conducted a prospective, randomized, patient and observer blinded non-inferiority trial using post-operative opiate consumption as our primary outcome.

Materials and Methods

The study design was a prospective, randomized, observer and patient-blinded non-inferiority trial with 2 parallel arms. Approval was obtained from the institutional review board of the Beth Israel Deaconess Medical Center before patient enrollment (2014P-000347). This trial was registered at [clinicaltrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT03577912) NCT03577912.

Participants

Adult (>18 years old) patients undergoing colorectal resection in the Division of Colorectal Surgery from March 2016 and April 2018 were eligible. Initially open and laparoscopic patients were eligible for enrollment, however, the rate of intended open cases was too low, thus only laparoscopic segmental colectomies without preoperative intended stoma were recruited. Those patients with allergy to bupivacaine, preoperative chronic narcotic usage or chronic pain syndrome were excluded from enrollment. Patients whose anatomy was thought to preclude effective placement of the TAP block, such as those with prior complex abdominal wall reconstruction were excluded. Prior abdominal surgery in itself was not an exclusion criteria. Post-enrollment, those patients who required an unplanned return to the operating room thus incurring additional incisional pain were also excluded.

Randomization and Treatment Allocation

Patients were randomized on the day of surgery. The randomization was performed at the beginning of the study by computer algorithm from a non-affiliated statistician with a goal 1:1 patient allocation. Pre-sealed, opaque envelopes were held by independent research staff and delivered to the OR on the day of surgery and opened after induction of anesthesia.

Blinding

The surgeons, regional block team, and operating room staff were not blinded to the treatment. Patients as well as independent research staff who collected pain scores and post-operative nursing teams were all blinded to the treatment received. There were no differences in patient bandages.

Interventions

Two staff surgeons (T.C. & V.P.) performed laparoscopic colorectal surgery using standard port placement with midline specimen extraction. TAP blocks were administered either by the surgeon with laparoscopic guidance or by the anesthesia regional block team under ultrasound guidance. In both groups, the TAP block consisted of 0.25% bupivacaine with 1:100,000 epinephrine with a volume of 1cc/kg up to 60cc which was divided equally and injected bilaterally.

Surgeon LTAPs were performed under direct visualization with the laparoscope just prior to closure of the fascia. With the abdomen insufflated, the operating surgeon palpated the lateral border of the rectus to ensure adequate lateral placement and then a 19 gauge needle was inserted percutaneously just inferior to the costal margin and as lateral as possible within the draped surgical field. Using the laparoscope to ensure that the peritoneum is not penetrated, the needle is advanced through the internal and external obliques (Fig 2a) and a

small amount of anesthetic is injected into the TAP plane. The spreading of the bulge along the TAP plane is confirmed visually prior to the injection of the whole amount of anesthetic (Fig. 2b). The same process was repeated on the contralateral side. After bilateral injection, the operation was completed in standard form including maturation of an ostomy, if necessary.

Anesthesia UTAPs were administered by the regional block team under ultrasound guidance (body habitus dictated linear high frequency 12 MHz or curvilinear abdominal 6Mhz probe use) using a 22g needle and standard lateral approach immediately following removal of the surgical drapes but prior to patient emergence from general anesthesia.

Post-operative care

All patients received non-narcotic analgesics including acetaminophen and ketorolac as well as anti-emetic medication as part of a standardized ERAS protocol. Patients received an opiate patient-controlled analgesia (PCA) pump in the post-anesthesia care unit (PACU). Rescue boluses of narcotic were administered at discretion of PACU staff. Once tolerating a regular diet, patients were transitioned to an oral narcotic with ongoing utilization of non-narcotic analgesics.

Data Collection

Patient, operative, and post-operative characteristics were collected from the patients' medical records by independent nursing staff and recorded using a standardized data collection tool. A blinded member of the research staff assessed subjective pain scores on a scale from 0–10 in the PACU, 6, 12, 24, and 48 hours post-operatively. Post-operative nausea or vomiting (PONV) was also recorded.

Outcomes measured

The primary outcome was cumulative post-operative opiate requirement. Opiate requirement was tracked from admission to PACU, at 6, 12, 24, and 48 hours post-operatively and standardized across pain regimens using milligrams of oral morphine equivalents. Secondary outcomes included subjective pain scores both at rest and with activity as assessed by blinded clinical research staff. Post-operative complications were graded using the modified Clavien-Dindo classification for surgical complications.[19]

Sample Size Calculation and Statistical Analysis

This study was designed as a non-inferiority trial.[20] Historical data on the opiate sparing effect of a TAP block demonstrated a mean 58mg decrease in oral morphine usage over 24hrs in the TAP block group.[21] A non-inferiority margin of 15mg of morphine was selected. Using 80% power and a two-sided significance level of 0.05, power calculations necessitated 27 patients per arm of the trial. As the primary outcome is achieved within 48 hours of surgery with a very low mortality rate, it was not expected that any patients would be lost to follow up. However, given that data collection may be imperfect, we increased the sample size by approximately 10% to account for this with a goal of 30 patients per arm.

Categorical data were analyzed with Fisher's exact or chi-square tests. Continuous variables were tested with unpaired two-tailed t-tests if normally distributed or using the Mann-Whitney U test if not normally distributed. Significance was considered as $p < 0.05$ unless otherwise stated. Non-inferiority was established when the 95% confidence interval of the differences of means fell within the pre-specified non-inferior margin. All analyses were performed using GraphPad Prism (version 7.03, La Jolla, CA).

Results

Study Population

80 patients were consented during the enrollment period between April 2016 and April 2018 and 64 underwent randomization (Fig. 1). The primary reason for lack of randomization among those consented ($N = 16$) was the inability to schedule the research personnel to perform independent blinded assessments. Of the 64 patients randomized on the day of surgery, 4 patients were excluded from the final analysis. One patient did not have the anticipated procedure and did not require a midline abdominal incision. One patient was re-intubated and unavailable for pain assessment. One patient could not have their blinded assessment completed due to a scheduling conflict, and one patient could not receive standardized TAP block due to drug shortage.

Patient Characteristics

There were no significant differences between groups in age, sex, body mass index (BMI), rate of prior abdominal surgery, diabetes, or pre-operative narcotic usage (Table 1). Nor were there any significant differences in indication for colectomy.

Operative and Post-operative Course

No significant differences in operative details were observed between groups (Table 2). Case time, as defined by "room in" to "room out" times, was similar as was estimated blood loss (EBL). There were no conversions to open surgery. There were no significant differences in rate of stoma creation or anatomic location of resection. Similarly there were no significant differences in post-operative course including frequency of PONV, time to oral narcotic consumption (indicating tolerance of a regular diet), or time to discharge (Table 2). No adverse events such as hematoma or inadvertent bowel or solid organ puncture related to the TAP block occurred. There were 3 and 5 Clavien-Dindo grade I-II complications and 2 and 1 Grade III-V complication in the UTAP and LTAP groups respectively which were not significantly different.

Post-operative narcotic consumption

Post-operative narcotic requirement administration was recorded from admission to the PACU until 48 hours following the operation. Intravenous and oral doses were converted to oral morphine equivalents. Cumulative opiate consumption at the time of PACU discharge (median [IQR] milligrams of morphine, 1.8 [0–4.5] UTAP vs 0 [0–8.7] LTAP $P = .32$), 6 hours post-operatively (5.4 [1.8–17.1] UTAP vs 3.6 [0–12.6] LTAP $P = .28$) at 12 hours post-operatively (9.0 [3.6–29.4] UTAP vs 7.2 [0.9–22.5] LTAP $P = .51$) at 24 hours post-operatively (9.0 [3.6–29.4] UTAP vs 7.2 [0.9–22.5] LTAP $P = .63$) and 48 hours post-

operatively 39.9 [7.5–70.2] UTAP vs 22.2 [7.5–63.8] LTAP $P=.41$) (Fig. 3a). There were no significant differences at any time point.

The upper boundary (favoring anesthesia delivered UTAP) of the 95% confidence interval of the difference of mean opiate consumption between groups was 3.3 in the PACU, 2.6 at 6 hours post-op, 7.1 at 12 hours post-op, 12.6 at 24 hours, 18.1 at 48 hours. Since the pre-determined non-inferiority margin was 15mg of morphine, surgeon delivered LTAP met the non-inferiority criteria at all points up to and including 24 hours but not at 48 hours (Fig. 3b).

Pain scores

Patient reported pain scores at rest (RPS) and with motion (MPS) were recorded on a scale from 0–10 in the PACU and at 6, 12, 24, and 48hrs post-operatively (Fig. 4a). Mean resting pain score \pm standard deviation (SD) at 48 hours was 2.8 ± 2.7 and 1.4 ± 1.6 ($P=0.048$) for anesthesia delivered UTAP and surgeon delivered LTAP, respectively (Fig. 4b). There were no other significant differences at any other time point.

Discussion

In this prospective, randomized, non-inferiority trial comparing LTAP vs. UTAP, we found that surgeon delivered LTAPs were non-inferior to anesthesia delivered UTAPs in the first 24 hours post-operatively with regard to cumulative narcotic requirement in these similar groups. In the 48 hours following surgery, there were no significant differences in narcotic requirement or patient reported pain scores (except at the 48hr resting pain score which favored surgeon delivered LTAPs). Therefore, we conclude that surgeon delivered LTAPs are non-inferior to anesthesia delivered UTAPs in patients undergoing laparoscopic colorectal resections.

Over the past two decades, TAP blocks have been shown to reduce post-operative pain and narcotic requirement in patients undergoing a wide range of surgery requiring abdominal incision.[6, 7, 22, 23] A number of studies have also demonstrated that TAP blocks reduce time to discharge prompting the inclusion of the TAP block into many ERAS protocols.[8, 24] This effect persists in newer studies using liposomal formulations of the local anesthetic.[25, 26] Moreover, the growing body of evidence surrounding TAP block and its inclusion in individual society ERAS guidelines demonstrate that TAP blocks are a reliable and broadly applicable non-narcotic adjunct to post-operative pain management.[4, 7]

Laparoscopically guided TAP blocks have been gaining popularity in recent years and several studies have demonstrated their efficacy, however, direct comparisons between LTAPs and UTAPs are lacking. [14–18] One previous study examined LTAP vs. UTAP in Korean patients undergoing colorectal resections demonstrated that LTAP was non-inferior although the primary outcome was subjective pain scores. Our findings build upon those of Park and colleagues in several important ways.[15] First, these authors employ an intra-abdominal approach to LTAP using a laparoscopic needle rather than a percutaneous technique, which requires specialized equipment for administration. Second, our patients

more accurately reflect the United States population with a mean BMI of 29.3 as compared to their cohort with a mean BMI of 24.

A recent study by Zaghiyan & colleagues examined a protocol similar to our own for LTAPs in comparison to anesthesia delivered UTAPs and patients receiving no TAP.[18] This study found that LTAPs were superior to UTAPs, however, in this study UTAPs had no benefit over the no TAP group in the primary endpoint of 24 hour morphine consumption calling into question the efficacy of the UTAPs delivered in this study. Similar to the study by Park and colleagues, average BMI in the Zaghiyan trial was between 23.6–25.6. Nevertheless the superiority of LTAP over UTAP in their study reinforces the results of this study that LTAP is, at the least, non-inferior to UTAP.

There was no difference in length of stay, operating room time, or other operative or post-operative parameters between UTAP and LTAP. Anecdotally we feel that surgeon administered LTAPs decrease operating room time as compared to a UTAP when performed before the patient leaves the operating room. However, our study did not seek to compare operative time and the large standard deviation in case time precluded identification of differences in this context. Additionally, while billing practices and compensation rates vary, we suggest that performing surgeon delivered TAP blocks as part of the surgery avoids potential fees involved with a separate operator and additional required equipment.

A strength of this study was its performance in the context of our ERAS protocol with both narcotic and non-narcotic pain medication used at the patient's discretion. The primary outcome in this study was post-operative opioid requirement as compared to other studies that have focused primarily on patient reported pain scores. [10, 15] We felt that an emphasis on the objective measure of post-operative opiate requirement was more generalizable, and thus we chose that value as our primary outcome. Opiate requirement differs among patients both in effective dose and agent and this variation is reflected in the wide range of dosage used by patients in this study. While lacking some of the uniformity of other studies, we believe that this “real-world” method further emphasizes the generalizability of these findings in modern colorectal surgery.

This study has several limitations. There was no placebo arm in this trial, which would have strengthened conclusions about the efficacy of LTAP and UTAP, however, given the widespread adoption of TAP block in abdominal surgery our institutional review board felt that excluding patients from a proven non-narcotic adjunct was not ethical. Another limitation is the employment of opiate patient controlled analgesia (PCA) in the initial post-operative period instead of an immediate transition to oral pain medication which is the case in many ERAS protocols. There are a range of doses, routes of administration (single vs. multiple injections and continuous infusion), and timing used for TAP blocks. This study specifically compares bupivacaine with epinephrine in one subcostal injection site bilaterally at the end of the operation in an academic teaching hospital. Therefore our findings may not be generalizable to other techniques, formulations of local anesthetic, or healthcare settings. [23, 28] All patients underwent peri-umbilical midline extraction of the specimen in this study and thus generalizability to other extraction sites is limited. Cadaver studies with single subcostal TAP injections have demonstrated broad spread along the abdominal wall

encompassing multiple thoracic segmental nerves, however, this technique may not be adequate for Pfannenstiel incisions.[28] Finally the proportion of patients who had undergone previous abdominal surgery in the LTAP group was nearly twice that of the UTAP group, which could have resulted in potential confounding with patient opiate consumption and pain scores.

Conclusion

Multi-modal adjuncts to post-operative pain management have become the standard of care in colorectal surgery. While there remains debate about which method or medication is best for patients, our results demonstrate that surgeon delivered LTAP is safe, simple, effective and non-inferior to anesthesia delivered UTAP. This method should be considered in all patients undergoing laparoscopic colorectal surgery where an ultrasound guided TAP block is planned.

Acknowledgments

The authors thank Shaelah Huntington and Dr. Marc Shnider for their contributions. This work in part was supported by NIH 5T32HL007734 fellowship funding to Dr. Wong. This work was conducted with support from Harvard Catalyst | The Harvard Clinical and Translational Science Center (National Center for Advancing Translational Sciences, National Institutes of Health Award UL1 TR001102) and financial contributions from Harvard University and its affiliated academic healthcare centers. The content is solely the responsibility of the authors and does not necessarily represent the official views of Harvard Catalyst, Harvard University and its affiliated academic healthcare centers, or the National Institutes of Health.

This work in part was supported by NIH 5T32HL007734 fellowship funding to Dr. Wong and by The Harvard Clinical and Translational Science Center (National Center for Advancing Translational Sciences, National Institutes of Health Award UL1 TR001102) and financial contributions from Harvard University and its affiliated academic healthcare centers.

References

- Spanjerberg W, Reurings J, Keus F, Cjhm VL (2011) Fast track surgery versus conventional recovery strategies for colorectal surgery. *Cochrane Database Syst Rev* 10.1002/14651858.CD007635.pub2. www.cochranelibrary.com
- Zhuang C-L, Ye X-Z, Zhang X-D, et al. (2013) Enhanced Recovery After Surgery Programs Versus Traditional Care for Colorectal Surgery. *Dis Colon Rectum* 56:667–678. 10.1097/DCR.0b013e3182812842 [PubMed: 23575408]
- Lee L, Mata J, Ghitulescu GA, et al. (2015) Cost-effectiveness of enhanced recovery versus conventional perioperative management for colorectal surgery. *Ann Surg* 262:1026–1033. 10.1097/SLA.0000000000001019 [PubMed: 25371130]
- Carmichael JC, Keller DS, Baldini G, et al. (2017) Clinical Practice Guidelines for Enhanced Recovery After Colon and Rectal Surgery From the American Society of Colon and Rectal Surgeons and Society of American Gastrointestinal and Endoscopic Surgeons. *Dis Colon Rectum* 60:761–784. 10.1097/DCR.0000000000000883 [PubMed: 28682962]
- Wick EC, Grant MC, Wu CL (2017) Postoperative multimodal analgesia pain management with nonopioid analgesics and techniques a review. *JAMA Surg* 152:691–697. 10.1001/jamasurg.2017.0898 [PubMed: 28564673]
- Hain E, Maggiori L, Prost à la Denise J, Panis Y (2018) Transversus abdominis plane (TAP) block in laparoscopic colorectal surgery improves postoperative pain management: a meta-analysis. *Color Dis* 20:279–287. 10.1111/codi.14037
- Johns N, O'Neill S, Ventham NT, et al. (2012) Clinical effectiveness of transversus abdominis plane (TAP) block in abdominal surgery: A systematic review and meta-analysis. *Color Dis* 14:635–642. 10.1111/j.1463-1318.2012.03104.x

8. Favuzza J, Brady K, Delaney CP (2013) Transversus abdominis plane blocks and enhanced recovery pathways: Making the 23-h hospital stay a realistic goal after laparoscopic colorectal surgery. *Surg Endosc Other Interv Tech* 27:2481–2486. 10.1007/s00464-012-2761-y
9. Pirrera B, Alagna V, Lucchi A, et al. (2017) Transversus abdominis plane (TAP) block versus thoracic epidural analgesia (TEA) in laparoscopic colon surgery in the ERAS program. *Surg Endosc Other Interv Tech* 1–7. 10.1007/s00464-017-5686-7
10. Elkassabany N, Ahmed M, Malkowicz SB, et al. (2013) Comparison between the analgesic efficacy of transversus abdominis plane (TAP) block and placebo in open retropubic radical prostatectomy: A prospective, randomized, double-blinded study. *J Clin Anesth* 25:459–465. 10.1016/j.jclinane.2013.04.009 [PubMed: 23965191]
11. Jadon A, Jain P, Chakraborty S, et al. (2018) Role of ultrasound guided transversus abdominis plane block as a component of multimodal analgesic regimen for lower segment caesarean section: A randomized double blind clinical study. *BMC Anesthesiol* 18:1–7. 10.1186/s12871-018-0512-x [PubMed: 29298664]
12. Gao T, Zhang JJ, Xi FC, et al. (2017) Evaluation of Transversus Abdominis Plane (TAP) block in hernia surgery. *Clin J Pain* 33:369–375. 10.1097/AJP.0000000000000412 [PubMed: 27518492]
13. McDermott G, Korba E, Mata U, et al. (2012) Should we stop doing blind transversus abdominis plane blocks? *Br J Anaesth* 108:499–502. 10.1093/bja/aer422 [PubMed: 22236911]
14. Favuzza J, Delaney CP (2013) Laparoscopic-guided transversus abdominis plane block for colorectal surgery. *Dis Colon Rectum* 56:389–91. 10.1097/DCR.0b013e318280549b [PubMed: 23392158]
15. Park SY, Park JS, Choi GS, et al. (2017) Comparison of Analgesic Efficacy of Laparoscope-Assisted and Ultrasound-Guided Transversus Abdominis Plane Block after Laparoscopic Colorectal Operation: A Randomized, Single-Blind, Non-Inferiority Trial. *J Am Coll Surg* 225:403–410. 10.1016/j.jamcollsurg.2017.05.017 [PubMed: 28610880]
16. Fields AC, Gonzalez DO, Chin EH, et al. (2015) Laparoscopic-Assisted Transversus Abdominis Plane Block for Postoperative Pain Control in Laparoscopic Ventral Hernia Repair: A Randomized Controlled Trial. *J Am Coll Surg* 221:462–469. 10.1016/j.jamcollsurg.2015.04.007 [PubMed: 26206644]
17. Elamin G, Waters PS, Hamid H, et al. (2015) Efficacy of a Laparoscopically Delivered Transversus Abdominis Plane Block Technique during Elective Laparoscopic Cholecystectomy: A Prospective, Double-Blind Randomized Trial. *J Am Coll Surg* 221:335–344. 10.1016/j.jamcollsurg.2015.03.030 [PubMed: 25899736]
18. Zaghiyan KN, Mendelson BJ, Eng MR, et al. (2019) Randomized Clinical Trial Comparing Laparoscopic Versus Ultrasound-Guided Transversus Abdominis Plane Block in Minimally Invasive Colorectal Surgery. *Dis Colon Rectum* 62:203–210. 10.1097/DCR.0000000000001292 [PubMed: 30540660]
19. Dindo D, Demartines N, Clavien PA (2004) Classification of surgical complications: A new proposal with evaluation in a cohort of 6336 patients and results of a survey. *Ann Surg* 240:205–213. 10.1097/01.sla.0000133083.54934.ae [PubMed: 15273542]
20. Schumi J, Wittes JT (2011) Through the looking glass: understanding non-inferiority. *Trials* 12:106. 10.1186/1745-6215-12-106 [PubMed: 21539749]
21. McDonnell JG, O'Donnell B, Curley G, et al. (2007) The analgesic efficacy of transversus abdominis plane block after abdominal surgery: A prospective randomized controlled trial. *Anesth Analg* 104:193–197. 10.1213/01.ane.0000250223.49963.0f [PubMed: 17179269]
22. Siddiqui MRS, Sajid MS, Uncles DR, et al. (2011) A meta-analysis on the clinical effectiveness of transversus abdominis plane block. *J Clin Anesth* 23:7–14. 10.1016/j.jclinane.2010.05.008 [PubMed: 21296242]
23. Ma N, Duncan JK, Scarfe AJ, et al. (2017) Clinical safety and effectiveness of transversus abdominis plane (TAP) block in post-operative analgesia: a systematic review and meta-analysis. *J Anesth* 31:432–452. 10.1007/s00540-017-2323-5 [PubMed: 28271227]
24. Keller DS, Ermlich BO, Delaney CP (2014) Demonstrating the benefits of transversus abdominis plane blocks on patient outcomes in laparoscopic colorectal surgery: Review of 200 consecutive cases. *J Am Coll Surg* 219:1143–1148. 10.1016/j.jamcollsurg.2014.08.011 [PubMed: 25442068]

25. Stokes AL, Adhikary SD, Quintili A, et al. (2017) Liposomal Bupivacaine Use in Transversus Abdominis Plane Blocks Reduces Pain and Postoperative Intravenous Opioid Requirement after Colorectal Surgery. *Dis Colon Rectum* 60:170–177. 10.1097/DCR.0000000000000747 [PubMed: 28059913]
26. Torgeson M, Kileny J, Pfeifer C, et al. (2018) Conventional Epidural vs Transversus Abdominis Plane Block with Liposomal Bupivacaine: A Randomized Trial in Colorectal Surgery. *J Am Coll Surg* 10.1016/j.jamcollsurg.2018.04.021
27. Bliss LA, Maguire LH, Chau Z, et al. (2015) Readmission After Resections of the Colon and Rectum. *Dis Colon Rectum* 58:1164–1173. 10.1097/DCR.0000000000000433 [PubMed: 26544814]
28. Barrington MJ, Ivanusic JJ, Rozen WM, Hebbard P (2009) Spread of injectate after ultrasound-guided subcostal transversus abdominis plane block: A cadaveric study. *Anaesthesia* 64:745–750. 10.1111/j.1365-2044.2009.05933.x [PubMed: 19624629]




FIGURE 1. CONSORT flow diagram of patients in trial.

Abbreviations: UTAP (Ultrasound guided transversus abdominis block) LTAP (laparoscopic guided transversus abdominis block)




FIGURE 2. Surgeon delivered laparoscopically guided TAP block.

(A) Laparoscopic view of abdominal wall with circle highlighting initial bulge of injection within the TAP plane. Laparoscope is used to ensure peritoneum is not entered with needle.
(B) Laparoscopic view of anesthetic spreading along TAP plane following injection.




FIGURE 3. Post-operative narcotic consumption.

(A) No significant differences in opiate requirement between anesthesia administered ultrasound guided TAP block (UTAP) and surgeon administered laparoscopic TAP block (LTAP) Bars represent mean opiate consumption \pm SD. (B) Graphic representation of 95% CIs of difference of means of opiate requirements in relation to 15mg non-inferiority margin. Numbers indicate bounds of 95% CI. * signifies met non-inferiority criteria at indicated time point. Abbreviations: PACU (Post-anesthesia care unit) mg (milligrams)




FIGURE 4. Post-operative pain scores.

No differences in patient reported pain scores at rest (a) or in motion (b) between UTAP and LTAP groups. Bars represent mean scores \pm SD. * = $p < 0.05$. Abbreviations: UTAP (Ultrasound guided transversus abdominis block) LTAP (laparoscopic guided transversus abdominis block) PACU (Post-anesthesia care unit)

Table 1.

Demographic and Pre-operative Clinical Characteristics.

Characteristic	Ultrasound TAP (N=31)	Laparoscopic TAP (N=29)	P-value
Age, years; mean \pm SD	60.0 \pm 13.6	61.5 \pm 14.3	.68 ^a
Female sex; N (%)	15 (48.4)	10 (34.5)	.31 ^b
BMI; mean \pm SD	28.1 \pm 5.9	30.6 \pm 5.8	.11 ^a
Pre-op narcotic usage; N (%)	0 (0)	2 (6.8)	.23 ^b
Baseline Pain; median (IQR)	0 (0–0)	0 (0–1.5)	.70 ^c
Diabetes; N (%)	3 (9.7)	5 (17.2)	.47 ^b
Prior Abdominal Surgery; N (%)	11 (35.5)	18 (62.1)	.07 ^b
Surgery Indication; N (%)			.39 ^d
Cancer	12 (38.7)	16 (55.1)	
Benign	16 (51.6)	10 (34.4)	
IBD	3 (9.6)	3 (10.3)	

Abbreviations: BMI (Body mass index); Pre-op (pre-operative); IBD (inflammatory bowel disease)

^{a)} Statistical analysis performed using unpaired t-test^{b)} statistical analysis performed with Fisher's exact test^{c)} statistical analysis performed using Mann-Whitney U test^{d)} statistical analysis performed with Chi-square all p-values reported as 2-sided

Table 2.

Operative details and post-operative hospital course of patients who completed study.

Characteristic	Ultrasound TAP (n=31)	Laparoscopic TAP (n=29)	P-value
Estimated blood Loss (ml) median (IQR)	50 (25–100)	30 (20–62.5)	.39 ^a
Case time, min.; median (IQR)	157 (120–199)	153 (124.5–198)	.72 ^a
Conversion to open No. (%)	0 (0)	0 (0)	>.99
Stoma creation No. (%)	0 (0)	2 (6.9)	.23
Colectomy type No. (%)			.64 ^b
Right	16 (51.6)	14 (48.3)	
Left	4 (12.9)	2 (6.9)	
Sigmoid	11 (35.5)	13 (44.8)	
PONV No. (%)			
PACU	8 (25.8)	3 (10.3)	.18 ^c
12 hrs post-op	5 (16.1)	5 (17.2)	>.99 ^c
24 hrs post-op	4 (12.9)	3 (10.3)	>.99 ^c
48 hrs post-op	3 (12.5)	1 (4.4)	.61 ^c
Time (hrs) to oral pain medication median (IQR)	25.9 (22.9–33.1)	27.5 (24.3–42.5)	.23 ^a
Time (hrs) to discharge median (IQR)	50.8 (46.4–73.0)	50.4 (46.1–70.0)	.74 ^a

Abbreviations: min (minutes); PONV (Post-operative nausea or vomiting); PACU (Post-anesthesia care unit); post-op (Post-operative); hrs (hours).

^{a)} statistical analysis performed using Mann-Whitney U test^{b)} statistical analysis performed with Chi-square^{c)} Fisher's exact test. All p-values reported as 2 sided.