



Since January 2020 Elsevier has created a COVID-19 resource centre with free information in English and Mandarin on the novel coronavirus COVID-19. The COVID-19 resource centre is hosted on Elsevier Connect, the company's public news and information website.

Elsevier hereby grants permission to make all its COVID-19-related research that is available on the COVID-19 resource centre - including this research content - immediately available in PubMed Central and other publicly funded repositories, such as the WHO COVID database with rights for unrestricted research re-use and analyses in any form or by any means with acknowledgement of the original source. These permissions are granted for free by Elsevier for as long as the COVID-19 resource centre remains active.

Letter to the Editor

Controversy regarding ACE inhibitors / ARBs in COVID-19**Controversias sobre el tratamiento con IECA/ARA-II y COVID-19****To the Editor,**

In the context of the severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) pandemic, the letter published by Gerard J O'Mara in the *British Medical Journal* suggesting that treatment with angiotensin-converting enzyme (ACE) inhibitors and angiotensin II receptor blockers (ARBs) could facilitate coronavirus infection by increasing circulating ACE2 levels, leading to increased viral load and more severe lung injury, has created considerable unease.¹

There is, however, evidence that the interaction between previous versions of coronavirus (SARS-CoV, which, to avoid confusion, we will hereafter refer to as SARS-CoV-1) and the renin-angiotensin-aldosterone system and ACE2 is much more complex. Experimental data from mouse models indicate that losartan might actually protect against coronavirus-induced lung injury.^{2,3}

The SARS-CoV-1 spike protein interacts with ACE2, using it for entry into the host cell, and deregulates this lung protective pathway by reducing ACE2 expression, resulting in unopposed angiotensin II accumulation and tissue injury. This process would explain the proposed protective effect of ARBs against SARS-CoV-1-induced lung injury in mice.²

Additionally, although absence of ACE2 (in knockout mice) protects against infection, it is not clear whether partial reduction of ACE2 levels, achieved through withdrawal of ACE inhibitors or ARBs, would have clinically relevant benefits.

As there are quite a few similarities between SARS-CoV-1 and SARS-CoV-2, particularly in terms of protein binding to ACE2, it is likely that the above findings will also be applicable to the current situation with coronavirus disease 2019 (COVID-19).^{4,5}

We therefore believe that the views expressed by O'Mara¹ should be treated as simple reflections and that caution should be exercised when considering their clinical applicability. The

relationship between the renin-angiotensin-aldosterone system and SARS-CoV-1 pathogenicity, and, by analogy, SARS-CoV-2 pathogenicity in the context of COVID-19, is highly complex. In fact, basic research suggests that ACE inhibitors, and ARBs in particular, have a protective role.

In this context, thus, there is no basis for discontinuing existing treatments with ACE inhibitors and ARBs. Moreover, complications due to the indiscriminate discontinuation of these drugs could have far more serious consequences than many of the surmised adverse effects.

Federico Soria Arcos,^a Antonio Romero Puche,^{b,*} and Tomás Vicente Vera^b

^aServicio de Cardiología, Hospital Santa Lucía, Cartagena, Murcia, Spain

^bServicio de Cardiología, Hospital Reina Sofía, Murcia, Spain

*Corresponding author:

E-mail address: antoniojoserp@hotmail.com (A. Romero Puche).

Available online 15 April 2020

REFERENCES

1. O'Mara GJ. Could ACE inhibitors and particularly ARBs increase susceptibility to COVID-19 infection. *Marzo 2020*. Disponible en: <https://www.bmj.com/content/368/bmj.m406/r-13>. Consultado 15 Mar 2020
2. Imai Y, Kuba K, Rao S, et al. Angiotensin-converting enzyme 2 protects from severe acute lung failure. *Nature*. 2005;436:112–116.
3. Kuba K, Imai Y, Rao S, et al. A crucial role of angiotensin converting enzyme 2 (ACE2) in SARS coronavirus-induced lung injury. *Nat Med*. 2005;11:875–879.
4. Xu X, Chen P, Wang J, et al. Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. *Sci China Life Sci*. 2020;63:457–460.
5. Zhang H, Penninger JM, Li Y, Zhong N, Slutsky AS. Angiotensin-converting enzyme 2 (ACE2) as a SARS-CoV-2 receptor: molecular mechanisms and potential therapeutic target. *Intensive Care Med*. 2020. <http://dx.doi.org/10.1007/s00134-020-05985-9>.