



Published in final edited form as:

Annu Rev Microbiol. 2011 ; 65: 189–213. doi:10.1146/annurev-micro-090110-102946.

The RpoS-Mediated General Stress Response in *Escherichia coli**

Aurelia Battesti, Nadim Majdalani, Susan Gottesman

Laboratory of Molecular Biology, National Cancer Institute, Bethesda, Maryland 20892;

Abstract

Under conditions of nutrient deprivation or stress or as cells enter stationary phase, *Escherichia coli* and related bacteria increase the accumulation of RpoS, a specialized sigma factor. RpoS-dependent gene expression leads to general stress resistance of cells. During rapid growth RpoS translation is inhibited and any RpoS protein that is synthesized is rapidly degraded. The complex transition from exponential growth to stationary phase has been partially dissected by analyzing the induction of RpoS after specific stress treatments. Different stress conditions lead to induction of specific sRNAs that stimulate RpoS translation or to induction progress has led to a better, but still far from complete, understanding of how stresses lead to RpoS induction and what RpoS-dependent genes help the cell deal with the stress.

Keywords

Hfq; (p)ppGpp; RssB; sRNA; ClpXP

INTRODUCTION

Bacteria constantly face changes in their environment, from nutrient starvation to variations in temperature, osmolarity, or pH. To adapt to or resist these changing conditions, they have developed various responses. Many of these adaptation response pathways are quite specific, involving a given regulator and a set of related genes, termed a regulon, that help the cell survive the original stress. The genes induced as a response to the stress include those necessary for repairing the damage or returning the cell to homeostasis. The ability of the bacteria to adapt to stress provides evidence of the inducible nature of the response—a pretreatment of cells with a nonlethal stress can improve resistance to a later, otherwise lethal, stress, and this adaptation requires new protein synthesis.

The alternative to these very targeted responses to specific stresses are more global changes in metabolism and gene expression that provide protection from many types of stress. Generally, these global responses have been identified under conditions of nutrient

*This is a work of the U.S. Government and is not subject to copyright protection in the United States.

battestia@mail.nih.gov.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

depletion, for instance, as cells enter stationary phase. In extreme cases, some bacteria, including *Bacillus subtilis* and *Myxococcus*, can differentiate into a stress-resistant spore in response to nutrient depletion.

For *Escherichia coli* and its relatives, a highly resistant state is achieved by triggering a global stress response dependent on the alternative sigma factor RpoS. This response allows cells to become more resistant not only to the stress that they first encounter but also to other stressful treatments. For instance, cells starved for carbon become resistant to hydrogen peroxide, high temperature, and low pH. This cross-protection phenomenon is typical of general stress responses and contrasts with specific stress responses that deal only with consequences of the inducing stress.

As with other sigma factors, RpoS interacts with the core RNA polymerase (RNAP) and controls the expression of a specific but large set of genes. Directly or indirectly, RpoS regulates 10% of the *E. coli* genome (approximately 500 genes) (157). The RpoS response has a number of key characteristics:

- Induction of the RpoS response is reflected in the rapid increase in levels of RpoS, which are very low in exponentially growing cells.
- Regulation of RpoS is at multiple levels [transcription, translation, degradation, and regulation of activity (see below)].
- A vast number of stresses affect RpoS accumulation, by feeding into one or more of these levels of regulation.
- Some of the genes induced as part of the RpoS response are truly RpoS specific; others are also expressed by the vegetative sigma factor, RpoD (σ^{70}), under some specific growth conditions.

In this review, we examine the complex regulatory circuits that center around the regulon controlled by RpoS, the stationary phase/stress sigma factor. The literature on this subject is vast and complex. We focus on work in *E. coli* and, to some extent, in *Salmonella*; we are necessarily selective even for these organisms. Among the questions we attempt to answer are the following: What are the signaling cascades leading to RpoS induction? Can the stresses that lead to induction be matched to genes in the RpoS-dependent response that help to deal with the stress? How does the RpoS-dependent response to a stress compare with the RpoS-independent, specific responses known for the same stress? Other reviews that examine the RpoS response and its regulation include References 52 and 53.

Discovering Stress Regulons and the RpoS-Dependent General Stress Response

Two general approaches were important to initially define regulons, including the RpoS regulon. Two-dimensional gels, first used for this purpose by Neidhardt and coworkers (79), made it possible to begin to define the proteins induced under a given stress treatment or affected by a given regulatory mutation. Such studies in the late 1980s and early 1990s allowed Martin and coworkers (49) to define a set of protein spots on two-dimensional gels that were common to starvation for carbon (C), phosphate (P), and nitrogen (N).

The second approach made use of transposable reporter systems—generally the *lacZ* gene, without a promoter, within an engineered transposable element such as the Mu phage. Transposition of the element containing *lacZ* led to isolation of a set of *lac* fusions to the promoters of random genes, which could then be queried on indicator plates for responses to a given treatment. These systems, developed by Casadaban & Cohen (25), were first used to define the SOS regulon (67) and then for many other stresses (reviewed in Reference 128). Studies of this sort identified genes induced by C starvation, for instance, that turned out to be induced by other stresses as well, helping to define the RpoS regulon (75). Mutations characterized by their effects on a particular phenotype or fusion were shown to be in a common gene, now called *rpoS*, encoding an alternative sigma factor referred to as RpoS (to be used here) or σ^{38} [see multiple alternative names listed in EcoCyc: *katF*, *appR*, *otsX*, *csi2*, *nur* (68)].

Untangling the Regulatory Cascades for the RpoS Regulon

Understanding any regulatory cascade requires defining the signals and mechanisms that lead to induction of the response, as well as an understanding of the outcome of induction (i.e., Which genes/functions are expressed and what do they do?) and an understanding of how the cell recovers from induction.

Induction: increasing RpoS levels and activity.—Under optimal laboratory growth conditions, RpoS levels are very low and increase as cells enter stationary phase. This is achieved by a combination of regulatory mechanisms. The default situation for RpoS, during exponential growth, is for synthesis to be low, because, even if transcription occurs, translation is shut off; in addition, any RpoS that is made is rapidly degraded. Thus, the inducing signals that increase RpoS in stationary phase or in response to a variety of stresses are designed to overcome these negative locks on RpoS accumulation.

The induction of RpoS synthesis and downregulation of degradation in response to starvation (for C, P, or Mg) or unfavorable conditions (e.g., high or low pH, high or low temperature, high or low osmolarity, DNA damage) require that the cell has a way of sensing and transducing the starvation or stress. It is clear that induction of the RpoS system takes advantage of many of the regulatory cascades that have been studied in other contexts. Among the most prominent of these collaborating systems is the stringent response that uses (p)ppGpp as a molecule for signaling a wide variety of starvation conditions. As (p)ppGpp increases, it promotes increased *rpoS* transcription and translation, inhibits RpoS degradation, and also improves RpoS activity. The Rcs phosphorelay and the PhoPQ two-component system also contribute to RpoS induction.

Some regulatory cascades, important for specific stress responses, seem to play more antagonistic roles for the RpoS regulon. Cyclic AMP (cAMP) and catabolite response protein (CRP), for instance, primarily have negative effects on RpoS. The ArcAB phosphorelay negatively regulates RpoS at multiple levels.

Outcome: expression of RpoS-dependent genes.—When the RpoS system is induced, the cell is resistant to a wide range of stress and starvation treatments. Cross-resistance suggests that all (or many) RpoS-dependent genes are induced in response to each

of these stresses. C starvation, for instance, leads to protection from low pH, oxidative stress, high temperature, and osmotic shock. If resistance to each of these depends on different output genes, we would argue that core resistance genes must all be induced without further specific input. In fact, microarrays carried out under a variety of conditions do support the idea that there is a common core of genes for which induction of RpoS is sufficient to increase transcription (157). Presumably, the RpoS-dependent genes necessary for cross-resistance all reside in this common core of genes. Which gene(s) provide resistance to which stress are known in only a subset of cases, however.

Many other genes are induced in an RpoS-dependent manner, but only after a specific stress (157). Why are they not induced when RpoS is present? Two general mechanisms can be imagined. The simplest is that transcription of these genes requires a combination of inputs—RpoS plus something else (a small molecule, an additional transcriptional activator, or inactivation of a repressor in response to the specific inducing signal). *csiD*, for instance, requires cAMP and CRP, in addition to RpoS; Fis is necessary for the RpoS-dependent transcription of *proP* (87, 159). In both cases, binding sites for the auxiliary regulators have been defined in the promoters of the RpoS-dependent genes. The second possibility is that some genes require higher RpoS levels than others do to be transcribed and that the levels of RpoS or the ability to compete with other sigma factors varies after different inducing treatments. This would imply a nested set of gene induction, with the strongest inducing treatment leading to induction of everything and weaker ones leading to a subset of these genes. Such is not what was seen in arrays (157), but certainly some sort of hierarchy is likely to exist.

Among the genes that are induced in response to RpoS, many but not all are also transcribed by RpoD in vitro, or in vivo under certain conditions (see, for instance, References 31 and 138). During exponential growth, when RpoS is not induced, the cell responds to specific stresses with induction of genes tailored to recovery from that stress. Some genes necessary for these specific stress responses are also induced in stationary phase by RpoS; one example is *dps* (3, 48, 88). How then is specificity achieved? Although the answers vary for different genes, global regulators such as Lrp and H-NS seem to play particularly important roles in enforcing RpoS-dependent regulation by repressing RpoD-dependent transcription (10, 31).

Recovery: returning to equilibrium.—For most stress responses, the process of reversing induction and returning cells to growth is as important as the induction process, but this process has been less studied. A variety of experiments as well as analysis of *E. coli* isolated from the environment suggest that there are many conditions under which the cell accumulates mutations that inactivate or downregulate *rpoS* (70, 137, 163). The isolation of these mutants certainly suggests that RpoS is not always good for the cell and supports the importance of recovery mechanisms.

REGULATING RpoS: MAJOR MECHANISMS OF REGULATION

Alternative sigma factors such as RpoS are generally under tight negative regulation, so that they do not compete for core RNAP under conditions when they are not needed. This is

certainly true for RpoS; it is barely detectable under rapid growth conditions and accumulates to high levels in response to a variety of stress treatments. We describe the general mechanisms for regulating RpoS here; some mechanisms are developed in more detail below in the context of the specific regulators and the signaling cascades in which they participate. Overexpression of RpoS in exponential phase cells is not sufficient for full induction of many RpoS-dependent genes (72), suggesting additional controls on how RpoS is able to compete effectively with RpoD.

Transcription of *rpoS*

In *E. coli* and many other species, the gene encoding RpoS is just downstream of *nlpD*, encoding an outer membrane lipoprotein of unknown function. Although a promoter upstream of *nlpD* may provide some transcription into *rpoS*, the major *rpoS* transcript initiates at a promoter internal to *nlpD*, starting 567 nucleotides (nt) upstream of the AUG for *rpoS* (Figure 1). This long 5' untranslated region (UTR) is critical for translational regulation. Regulation at the level of transcription is not dramatic compared with effects on translation and protein turnover (76). In addition, interpretation of some results based on reporter fusions or mRNA levels may be complicated by effects of translational regulation on mRNA stability and should be revisited (89).

In two cases, the protein regulators bind directly to the *rpoS* promoter. In *Salmonella enterica*, Fis, abundant during exponential growth and low during stationary phase, negatively regulates *rpoS* transcription ninefold during exponential growth, dependent on an upstream binding site (57). Another negative regulator of *rpoS*, ArcA-P, the response regulator of the two-component ArcB/ArcA system, had a three- to fourfold effect on *rpoS* transcription in exponential phase (92; also see Reference 69 for a review). For other regulators, direct binding remains to be shown. cAMP and CRP negatively regulate an *rpoS-lacZ* transcriptional fusion, and higher levels of RpoS are found in exponentially growing cells in rich medium in the absence of cAMP (76). The two-component BarA-UvrY system has been reported to positively regulate *rpoS* transcription, although the binding site has not been determined (100).

Levels of *rpoS* mRNA rise after (p)ppGpp induction, suggesting a stimulatory effect of (p)ppGpp on *rpoS* mRNA transcription or stability (36, 144, 145). However, studies of transcriptional fusions suggest that these effects may be at the level of mRNA stability or elongation, rather than initiation (56, 74). Some of the reported effects of (p)ppGpp on mRNA levels could be indirect, for instance, via the (p)ppGpp-dependent accumulation of polyphosphate, which is also associated with induction of *rpoS* (127).

Translational Regulation: Critical Roles for sRNAs

The long 5' UTR of the *rpoS* transcript folds into a stem-loop that occludes the ribosome-binding site and minimizes translation of *rpoS* (21, 34, 97) (Figure 1). This inhibitory structure is overcome by *trans*-encoded small RNAs (sRNAs) that stimulate *rpoS* translation. The sRNAs require the RNA chaperone protein Hfq for their action (reviewed in Reference 19). Hfq both stabilizes the sRNAs in vivo and promotes pairing with mRNA targets in vitro.

Although Hfq is used by many sRNAs in *E. coli*, the first observed phenotypes of *hfq* mutants in *E. coli* and *Salmonella* are attributable to very low levels of RpoS (20, 98).

Three Hfq-dependent sRNAs (DsrA, RprA, and ArcZ) stimulate *rpoS* translation. All operate by a similar mechanism, pairing between complementary nucleotides in the sRNA and the 5' UTR of the mRNA to open the hairpin, thus freeing the ribosome-binding site (82, 84, 86). The major effect of pairing is to allow translation; an associated, and possibly indirect, effect is to stabilize the *rpoS* mRNA (89). Hfq requires an (AAN)₄ repeat far upstream within the *rpoS* 5' UTR for sRNAs to activate translation effectively (131, 132). A fourth sRNA, OxyS, negatively regulates *rpoS* translation, likely by titrating Hfq, as no pairing could be detected (59, 165; K. Moon & S. Gottesman, manuscript in preparation). These sRNAs are synthesized in response to different stresses under the control of different regulators, allowing the bacterium to integrate and respond to numerous stress signals for control of *rpoS* translation.

The essential single-stranded endonuclease RNase E appears to be a major player in degrading *rpoS* mRNA. However, inactivating RNase E is not sufficient for full translation, consistent with a critical role of sRNAs in opening up the inhibitory hairpin to allow translation (89).

Interestingly, a strain deleted for all three positively regulating sRNAs has higher expression of a translational fusion to *rpoS* than a strain deleted for *hfq*, suggesting that either Hfq has some direct regulatory activity on the *rpoS* mRNA or that there is at least one as yet undetected sRNA that positively regulates *rpoS* translation (86). A number of reports hint at other Hfq-dependent activators. For instance, deletion of the gene for the sRNA GcvB leads to lower levels of RpoS (61), although overexpression of GcvB did not increase expression of an *rpoS-lacZ* translational fusion (86), suggesting either indirect activation or regulation of RpoS by a mechanism not detected by the reporter fusion (for instance, dependency on the 3' end of the *rpoS* gene). The transcriptional regulator LrhA represses *rpoS* translation in an Hfq-dependent manner (108) by an as yet unknown mechanism. Constitutive expression of the *E. coli* Pho regulon owing to deletion of the *pst* genes also upregulates RpoS, dependent on both Hfq and the PhoB/R two-component system, suggesting yet another sRNA (121) possibly encoded in an intergenic region of the *pst* operon (124).

A number of other proteins have been implicated in modulating *rpoS* translational regulation or in regulating the stability of the *rpoS* message. How directly these act remains to be demonstrated. RNase III, a double-stranded endonuclease encoded by the *rnc* gene, cleaves the *rpoS* mRNA within the inhibitory hairpin. *rnc* mutants have higher levels of *rpoS* mRNA and RpoS protein, suggesting that RNase III is important to maintain low RpoS levels (118) (Figure 1). Binding of DsrA or RprA stimulates *rpoS* translation in a manner that appears to be additive to and independent of the effect of RNase III (89), although DsrA binding does change the RNase III cleavage site (89, 118). It is not yet clear whether specific signals modulate RNase III cleavage of the *rpoS* mRNA. One possible signal is temperature; at high temperature (43.5°C), *rnc* mutants no longer affect *rpoS* mRNA stability (89). The secondary structure elements necessary for this double-stranded endonuclease to cut may be disrupted

at high temperature, preventing cleavage. If so, melting of the hairpin may contribute to the known induction of RpoS at high temperature by overcoming RNase III action (63).

The cold-shock DEAD-box protein CsdA is an RNA helicase that binds to the *rpoS* message. Deletion of *csdA* leads to a decrease in RpoS amounts at 24°C compared with those found at 37°C (119). Because *rpoS* translation at low temperature is primarily dependent on DsrA (130), CsdA may aid in unwinding the *rpoS* secondary structure to facilitate DsrA annealing (119).

Overexpression of CspC or CspE, constitutively produced members of the CspA family of RNA-binding proteins, increases RpoS levels, apparently by stabilizing *rpoS* mRNA. Deletion of *cspC* lowers expression of an *rpoS-lacZ* translational fusion modestly (30), and deletion of both *cspC* and *cspE* impairs osmotic induction of RpoS-dependent genes (109). It is not clear whether these proteins function only under conditions of high osmolarity. Similar to the models for CsdA function, one may imagine that these proteins help to open up the inhibitory hairpin at high osmolarity, contributing to the increased ability of DsrA and other sRNAs to activate translation under these conditions (81). Overall, the results suggest a possible role for CspC and CspE as alternative RNA chaperones affecting *rpoS* mRNA structure and/or stability.

Deletion of *hupA* and *hupB*, encoding the subunits of HU, significantly decreases *rpoS* translation. Because HU can bind tightly to the *rpoS* 5' UTR in vitro, this may reflect a direct effect on *rpoS* mRNA structure (7).

Most studies of translation of *rpoS* have focused on the role of the 5' UTR and how its inhibitory structure is overcome. The translational fusions used to study *rpoS* have also been constructed to focus on the same regions. The translation of the *rpoS* ORF, downstream of the region studied in these fusions, may be subject to additional controls that are not yet understood (56, 115, 156).

Regulated Degradation of RpoS: Critical Roles for RssB and Antiadaptors

Targeted degradation of RpoS is one of the major examples of regulated proteolysis in *E. coli*. RpoS is rapidly degraded during exponential growth under optimal conditions; degradation stops or slows and RpoS accumulates rapidly after stress treatments (starvation for C, Mg, and P) and in stationary phase (76, 85). When cells return to rapid growth, degradation helps reduce the level of RpoS.

RpoS is degraded by the ClpXP ATP-dependent protease (125). However, RpoS is not recognized directly by ClpX but needs to be delivered to the protease by an adaptor protein called RssB or SprE in *E. coli*, MviA in *S. typhimurium*, or ExpM in *Erwinia carotovora* (4, 11, 71, 96, 112, 168). RssB is not degraded, but it is recycled, promoting degradation of multiple molecules of RpoS (168).

The regulation of RpoS proteolysis during growth phase depends on the stoichiometry and activity of a number of different proteins. ClpXP expression does not change significantly with growth phase, although under C starvation, the availability of ClpXP may be limited if

it is actively degrading mistranslated and misfolded proteins (44, 125) or its activity may be lower if ATP levels drop (51).

The limiting component for degradation of RpoS is RssB. The level of RssB is very low (13). If overproduced, it can lead to RpoS degradation even in stationary phase, and it can act as an antisigma factor for RpoS in cells in which RpoS degradation is artificially blocked by mutations in *clpX* or *clpP* (13, 112, 169). Whether this antisigma activity would ever be significant in wild-type cells is not clear. *rssB* is the second gene in an operon with another gene, *rssA*. Although RssA has phospholipase activity, no role for it in modulating RssB activity or synthesis has been reported (51, 114, 120).

RssB expression levels increase modestly during entry into stationary phase in an RpoS-dependent fashion (47, 114, 120). The modest induction of RssB in stationary phase has been suggested to be important for the rapid destruction of RpoS when cells exit from stationary phase, although this remains to be demonstrated. Another possibility is that RssB has another function during stationary phase as suggested by the recent demonstration that RssB promotes the association of polyA polymerase with the RNA degradosome during stationary phase, independent of RpoS (23, 24).

RssB is a response regulator that can be phosphorylated, but the role of RssB phosphorylation in RpoS degradation is not fully understood. In vitro, phosphorylation improves the interaction between RssB and RpoS and stimulates RpoS degradation (12, 14, 71, 94, 168). This suggests that phosphorylated RssB is the active form that degrades RpoS and dephosphorylation of RssB under some stress conditions might be the signal for stabilization of RpoS. However, cells carrying mutations in the site of phosphorylation on RssB still degrade RpoS and respond to starvation or stationary-phase growth signals to stabilize RpoS, suggesting that RssB phosphorylation is not essential for regulated degradation (108). Nonetheless, it remains possible that phosphorylation helps to modulate RssB activity or is important for alternative RssB activities. Although no dedicated phosphatase or histidine kinase has been found, the two-component system ArcA/B and small-molecule P donor acetyl phosphate have been implicated in RssB phosphorylation (14, 92).

Recently, three new proteins able to modulate RssB activity have been identified and termed Ira for inhibitor of RssB activity (15, 17). Each of these three proteins, IraP, IraM, and IraD, interacts with RssB and prevents RpoS degradation in vivo and in vitro; we refer to this class of proteins as antiadaptors. Although they share a similar function, they do not share any sequence similarity. The expression of these proteins under specific stress conditions provides an explanation for how RpoS degradation is regulated in response to stress (Figure 1).

Regulation of RpoS activity

The RpoS regulon is large and complex, including genes that are read only by RpoS, genes transcribed by both RpoS and RpoD, and genes in which auxiliary factors activate RpoS-dependent transcription. Proper expression of RpoS-dependent genes has been studied by

examining the rules for promoter recognition by RpoS as well as factors that contribute to the ability of RpoS to compete successfully for core RNAP.

Promoter recognition by RpoS.—Sigma factors determine promoter recognition, and thus the sequence of the promoter will act to exclude some sigma factors and favor others. However, RpoS is a member of the same family of sigma factors as the vegetative sigma factor RpoD and shows preference for the same consensus promoter elements in vitro as RpoD (46). RpoS is more sensitive to sequences around the -10 promoter region and shows less dependency on -35 regions than does RpoD (reviewed in Reference 151). Many promoters that appear fully RpoS dependent in vivo can be read by RpoD in vitro and vice versa (139). It is difficult to make broad conclusions about RpoS specificity given that each case is different. A combination of favored promoter elements combined with differential sensitivity to repressors and activators poises promoters in vivo so that only when RpoS-containing holoenzyme levels reach a critical level will they fire.

In many genes, the same start point is used by RpoD or RpoS, under different conditions. The promoter for the DNA-binding protein *dps* is repressed by Fis and H-NS, limiting RpoD-holoenzyme access unless the activator OxyR is present. However, in stationary phase, when RpoS levels rise, *dps* is robustly expressed, dependent on RpoS. This is due both to the ability of RpoS-holoenzyme to overcome or compete successfully with H-NS for binding to the promoter and to the decrease in Fis levels in stationary phase (48). The *aidB* gene is expressed in an RpoD-dependent fashion only when the Ada activator is present; RpoS-dependent expression occurs without Ada (73, 153). The promoter for the osmotically inducible *osmY* gene can be read by both RpoD and RpoS holoenzymes but is repressed by IHF, CRP, and Lrp, all blocking access of RpoD but not RpoS (31).

For other genes, alternative promoters allow expression of the same gene dependent on either RpoS or RpoD (8, 18, 54). For instance, *proP*, encoding a transporter for the osmoprotectants proline and glycine betaine, has an osmotically induced RpoD-dependent promoter and a second RpoS-dependent promoter; recognition of the RpoS promoter is significantly helped by binding of Fis nearby (151, 159).

Competition for RNA polymerase core.—All sigma factors compete for the same pool of core RNAP. Thus, anything that affects the binding of RpoD to core polymerase, for instance, necessarily also affects the ability of RpoS to bind core and recognize its promoters. The reverse is true as well; increased RpoS downregulates at least some RpoD promoters, presumably by competing for core RNAP (38).

In in vitro experiments, the affinity of RpoS for core is somewhat less than that of RpoD, the vegetative sigma factor (62, 64). In addition, levels of RpoS are lower than those for RpoD, even in stationary-phase cells. In one study, 7,000 molecules of RpoD were found per cell, significantly in excess of the 2,500 RNAP core complexes, in both exponential and stationary phase. RpoS, undetectable in exponential phase, rose to 1,600 molecules per cell in stationary phase (110). However, under conditions of RpoS accumulation, RpoS is clearly able to acquire core polymerase and read relevant promoters. Ferenci and coworkers (39, 70) have suggested that a balance exists between the advantages of cross-resistance (when RpoS

is active) and the loss of nutritional flexibility associated with the shutdown of many RpoD-dependent pathways (SPANC, or self-preservation and nutritional competence). Not surprisingly, the cell has developed ways to regulate this balance. Crl, Rsd, and 6S RNA have been implicated in helping RpoS effectively compete for core. The pattern of expression of these factors suggests that they reinforce RpoS activity and do not themselves act as part of the signaling cascade for RpoS.

Crl protein is necessary in vivo for maximal expression of RpoS-dependent promoters but does not increase RpoS levels (113). In vitro, Crl helps to stimulate RpoS holoenzyme formation, particularly under limiting levels of sigma. This is presumably particularly important under transition conditions when RpoS levels are first increasing or during recovery from stationary phase (45, 150). Crl is expressed constitutively, possibly so that it is available when RpoS increases. However, Crl was also identified as a substrate for the ClpXP protease (42). Whether degradation of Crl significantly affects its abundance has not been examined; it may be brought to the protease by binding to RpoS. If so, degradation of Crl could be part of the recovery mechanism when rapid growth resumes and RpoS is no longer useful. *crl* mutations also slow RpoS degradation (149); the basis for this is not known and may reflect a lower level of the RpoS-dependent expression of *rssB*. Because RNAP core protects RpoS from degradation in vitro (168), one could imagine a role for Crl as an exchange factor, promoting the ability of RpoS to both bind to and be released from core polymerase.

Unlike Crl, which acts on RpoS holoenzyme, 6S regulatory RNA negatively affects RpoD holoenzyme. 6S regulatory RNA binds to the RpoD holoenzyme and not to RpoS holoenzyme and freezes the holoenzyme in an inactive form (reviewed in Reference 154). 6S RNA is released from RNAP as growth resumes. Because 6S interacts with the holoenzyme, interaction with 6S would not increase the availability of RNAP core for RpoS, but it can lower the relative levels of free RpoD compared with cells deleted for 6S by preventing RpoD recycling.

The in vivo effect of 6S is to increase RpoS-dependent transcription at some promoters and decrease expression of a subset of RpoD-dependent promoters, characterized by extended -10 sequences and poor -35 sequences, possibly because 6S binds to the region of sigma necessary for -35 recognition (26). In competition experiments, 6S provides an advantage in late stationary and under stress conditions such as high pH (147, 148).

Rsd is a protein identified biochemically as a factor that bound RpoD in stationary-phase cells. Rsd interacts with RpoD and interferes with its ability to interact with core polymerase, thus possibly helping RpoS compete for core (162). No phenotype for an *rsd* deletion mutant has been detected in microarrays (93); only when Rsd was overproduced were some RpoS-dependent genes upregulated, consistent with competition. Thus, the physiological role of Rsd is currently unclear.

REGULATORY CASCADES: SIGNALS AND RESPONSES

As described above, many different mechanisms regulate RpoS accumulation and activity, and for each of these mechanisms, there are multiple effectors, each made under particular conditions. In addition, there are many RpoS-dependent genes, some dependent only on whether RpoS is available and some requiring additional inputs. The function and importance of only a handful of these downstream genes is known. In this section, we review a selected group of the known cascades of signaling and the downstream genes implicated in particular resistance pathways. This is not meant to be all-inclusive. In Table 1, a broader list of stresses and the known downstream genes implicated in the RpoS response are listed.

Starvation Responses: (p)ppGpp Unites Starvation Signaling

The small-molecule alarmone (p)ppGpp is the main effector of the stress response that takes place in *E. coli* during starvation. Its accumulation in the cell triggers the inhibition of stable RNA biosynthesis (i.e., rRNA operons and tRNA genes), growth arrest, and the activation of genes needed to resist stress conditions. This response, initially observed during amino acid starvation, was called the stringent response. Today, the term stringent response is commonly used to describe a broad set of stress responses driven by the accumulation of this nucleotide (reviewed in Reference 111).

(p)ppGpp is a unique small-molecule global transcriptional regulator that interacts directly with RNAP (6, 28, 141). In *E. coli*, synthesis of (p)ppGpp is carried out by RelA, during amino acid starvation, and the bifunctional SpoT protein during other stress treatments. RelA, the enzyme mainly responsive for (p)ppGpp synthesis, is associated with the ribosome and is activated upon amino acid starvation (111). SpoT is also responsible for degradation of (p)ppGpp. The effect of (p)ppGpp on RNAP and on gene transcription can be enhanced by DksA, a protein also able to interact with RNAP (104). Accumulation of this nucleotide regulates many genes directly and also affects the expression of other global regulators in the cell such as RpoS. Thus, it is often difficult to discriminate direct from indirect effects of (p)ppGpp.

(p)ppGpp changes promoter activity, inhibiting some classes of promoters and activating others. During rapid growth, a major part of the available RNAP is dedicated to the transcription of stable RNA (reviewed in Reference 105). However, the presence of (p)ppGpp and DksA destabilizes the open complex formed between stable RNA promoters and RNAP holoenzyme (9). As a consequence of inhibiting rRNA transcription, RNAP becomes available for the transcription of lower affinity promoters (those that need a higher amount of available RNAP to be transcribed) (9).

In parallel, (p)ppGpp induces the expression of alternative sigma factors and of the antisigma factor Rsd (33, 36). The increased availability of RNAP core and of the alternative sigma factors, simultaneously with the titration of RpoD by Rsd, favors the transcription of genes by alternative sigma factors, including RpoS (36, 145).

In vitro studies have defined the characteristics of promoters negatively regulated by (p)ppGpp (105). This regulation depends in part on the discriminator sequence of the

promoter, located between the –10 box and the transcriptional start (143). Positively regulated promoters are less well understood: In some cases, data suggest direct effects on RNAP at these promoters (see *iraP* below); in other cases, the inhibition of RNAP from transcribing the extremely active rRNA operons and other stable RNA genes may increase availability for other promoters.

Many of the experiments that tie (p)ppGpp to RpoS were done by studying the properties of strains in which (p)ppGpp cannot be synthesized, as a result of deletions of both *relA* and *spoT*. Under these conditions, very little RpoS is made, and studies have suggested that, in *E. coli*, (p)ppGpp affects RpoS at every level: transcription, translation, proteolysis, and activity. In all cases, there is a positive correlation between (p)ppGpp and RpoS levels and activity, so that these two responses act as two aspects of a more general stress response. Effects on transcription of *rpoS* have been less fully studied and are not further discussed here.

Effects of (p)ppGpp on rpoS translation.—Several studies have shown a regulatory effect of (p)ppGpp on *rpoS* translation (22, 126). In fact, DksA was also implicated in *rpoS* regulation (126, 156) even before the functional link between (p)ppGpp and DksA had been established. The most compelling evidence of a specific mechanism for how (p)ppGpp may improve *rpoS* translation comes from work in *Shigella flexneri*. In *S. flexneri*, DksA has a direct positive effect on *hfq* transcription, which is highly enhanced in the presence of (p)ppGpp. The increased *hfq* transcription, in turn, activates *rpoS* translation, presumably because Hfq is in fact limiting for *rpoS* translation (59, 126; K. Moon & S. Gottesman, manuscript in preparation). It seems likely that this mechanism is conserved in *E. coli*; we have observed twofold-lower levels of Hfq protein in *dksA* mutants (K. Moon & S. Gottesman, manuscript in preparation). Consistent with an effect on Hfq-stimulated translation, the (p)ppGpp effect on *rpoS* translation is abolished in the absence of the inhibitory antisense structure (56).

Effects of (p)ppGpp in regulating RpoS stability: controlling antiadaptor synthesis.—As described above, RpoS is very rapidly degraded under rapid growth conditions. Therefore, even if synthesis is high, the accumulation of RpoS will be limited by degradation. Starvation for C and P, which leads to increased (p)ppGpp, also stabilizes RpoS (76, 108, 133, 158, 164). The recent demonstration that expression of two of the three known antiadaptor proteins, inhibitors of RssB IraP and IraD, is positively regulated by (p)ppGpp provides a clear mechanism explaining at least some of the (p)ppGpp requirement for RpoS accumulation (16, 91).

IraP was named for its role in stabilizing RpoS after P starvation (17); *iraP* is transcriptionally induced after P starvation, and this induction depends on SpoT activity and (p)ppGpp. Because the discriminator region of the *iraP* promoter is important for (p)ppGpp-dependent regulation, it seems likely expression is under the direct control of (p)ppGpp (16). Interestingly, under conditions other than P starvation [amino acid starvation or artificial overproduction of (p)ppGpp], the level of *iraP* mRNA increases but not the level of the protein, suggesting high levels of (p)ppGpp may inhibit IraP translation (16). This further suggests that *iraP* induction would be triggered by a moderate accumulation of (p)ppGpp,

which is consistent with SpoT-dependent *iraP* expression, and not under conditions that induce higher levels of (p)ppGpp.

IraD, identified in a multicopy screen for proteins that stabilize RpoS, stabilizes RpoS after DNA damage; deletions of *iraD* are more sensitive to some types of DNA damage, in an RpoS-dependent fashion (15, 90). IraD also plays a role in stabilizing RpoS as cells enter stationary phase (91). The major *iraD* promoter, P1, is induced during stationary phase, dependent on (p)ppGpp (91). The induction of *iraD* by DNA damage is also dependent on P1; the mechanism of induction is undefined but is not dependent on the SOS regulator LexA (91).

The role of (p)ppGpp in the synthesis of these two antiadaptors provides an explanation for some of the effects of (p)ppGpp on RpoS accumulation. P starvation leads to induction of *iraP*. The end of exponential phase may be the inducing signal for *iraD*; as (p)ppGpp accumulates, *iraP* may also be induced. This leaves a number of aspects of this system unexplained. If the same signal induces both *iraP* and *iraD*, are they really expressed at the same time? For instance, is *iraD* induced under P starvation? If so, why does it not contribute more to stabilization of RpoS (because a deletion of *iraP* eliminates most of the stabilization after P starvation)? If it is not induced, why not? Does it require higher or more sustained levels of (p)ppGpp, or does some other regulator contribute to induction for one or the other of these proteins? A requirement for relatively high levels of (p)ppGpp for RpoS induction was recently shown, although the contributions of IraP and IraD were not specifically tested (146).

Specific Stresses and the Role of sRNAs

In addition to the global role of (p)ppGpp in sensing and transducing starvation signals, RpoS is induced under a large variety of stress treatments that appear to be independent of (p)ppGpp and, in many cases, dependent on expression of sRNAs. In this section, for a selected subset of stress treatments, we try to link what is known about the inducing pathways to the downstream genes that help to protect from the stress. Other stresses known to induce RpoS are listed in Table 1.

DsrA, the RpoS low-temperature response, H-NS, and osmoprotection.—One definition of a stress, as seen by the cell, may come from identifying the downstream genes that provide protection from the stress. For instance, stationary-phase cells may perceive the stress of low temperature similarly to the stress of high osmolarity, as evidenced by the observation that increasing trehalose synthesis provides protection for both.

RpoS accumulates and is active at low temperature, even in exponential phase, and this accumulation reflects increased translation, dependent on the sRNA DsrA (130). No specific low-temperature activator of DsrA is known, and thermocontrol seems to be a combined function of the suboptimal spacer and –10 sequences of the *dsrA* promoter as well as differential stability of the sRNA at different temperatures (116, 117). What is the physiological role of RpoS at low temperature? At least one set of RpoS-dependent genes, *otsAB*, is important at low temperature. The products of *otsA* and *otsB* are necessary for

synthesis of internal trehalose, an osmoprotectant. *E. coli* cells subjected to cold shock accumulate high levels of trehalose but die quickly in the absence of *otsA* (66).

In other experiments, the expression of the *otsAB* operon has been shown to be necessary and sufficient for resistance to high osmolarity in stationary-phase cells. Mutants in *otsAB* are hypersensitive to high osmolarity. Cells deleted for *rpoS* and evolved to grow at high osmolarity in chemostats accumulate insertions that activate the *ots* operon (134). Combined with the observations above, this suggests that adaptation to cold shock is akin to adaptation to high osmolarity and provides a rationale for induction of RpoS at low temperatures as well as at high osmolarity. When cells are exposed to high osmolarity, rapid translational induction of RpoS is primarily dependent on DsrA (81). *otsA* mutants also partially impair the thermotolerance of stationary-phase cells (55). Thus, high levels of internal trehalose may serve to help protect proteins from denaturation at high temperature, low temperature, and high osmolarity.

DsrA also directly negatively regulates the translation of the histone-like protein H-NS, a global regulator of gene expression (77). H-NS has broad effects on transcription. H-NS directly represses RpoD-dependent transcription of many genes with both RpoD- and RpoS-dependent promoters, and RpoS is less sensitive to H-NS repression than RpoD, at least in one tested case (5, 10, 48). The net effect is that H-NS helps make expression of these genes fully RpoS dependent.

In parallel to increasing the RpoS dependency of many promoters, H-NS and related proteins negatively regulate *rpoS* translation by an unknown mechanism (10, 160) and promote its degradation (Figure 1). RpoS is stable in *hns* mutants, suggesting that H-NS may negatively regulate one or more antiadaptors, although which antiadaptors is not yet known (167). In *Salmonella*, an H-NS homolog, StpA, acts to repress the RpoS regulon in exponential growth by repressing expression of the *rssC* antiadaptor. RssC is most similar to the *E. coli* IraM antiadaptor (80). Whether RpoS-inducing signals other than DsrA also negatively regulate H-NS synthesis or activity is not yet known but would not be surprising.

hns mutants are impaired for growth under many conditions; this growth defect is overcome by a second mutation in *rpoS* (10). Because RpoS levels are high in *hns* mutants, this may reflect poor growth due to RNAP core competition, coupled with the higher requirement for RpoD owing to high transcription of genes usually silenced by H-NS.

Acid stress, H-NS, and PhoPQ.—One of the most striking phenotypes of stationary-phase cells is their resistance to low pH. In particular, *E. coli* and *Shigella* become resistant to pH 2.5 when they enter stationary phase or when they are preshocked with an intermediate pH (reviewed in Reference 166). While the acidresistance response is complex and involves many regulators, there is a clear requirement for RpoS for the acid resistance of stationary-phase cells (11, 78). Required for this acid resistance are RpoS-dependent genes (155), including *gadC*, a gene of the glutamate-dependent low-pH-resistance network of genes, *hdeAB*, encoding pH-regulated periplasmic chaperones, and *cfa*, encoding a gene for cyclopropane fatty acid synthesis. *hdeAB* was termed an H-NS-dependent operon; it is strongly repressed by H-NS (161). HdeA binds proteins at low pH, helping to protect

misfolded or damaged proteins from aggregating and then releasing them to refold when the pH returns to neutral (140). The product of the *cfa* gene leads to modification of phospholipids that render the cell able to tolerate low pH (27). In addition to these genes, *gadY*, encoding an sRNA that positively regulates the synthesis of two other regulators of the *gad* genes, GadX and GadW, is also RpoS dependent (103, 142).

The pathway of induction of RpoS by low pH is not fully understood. One component of induction is RpoS stabilization (11), now known to be mediated by the two-component PhoPQ system, the activator of the antiadaptor IraM (15). The importance of coupling PhoPQ to RpoS is demonstrated by the observation that in *Salmonella* another antiadaptor, IraP, is expressed not only from the (p)ppGpp-dependent promoter found in *E. coli* but also from a second PhoPQ-dependent promoter (149). PhoPQ is activated at low Mg and in response to antimicrobial peptides, both encountered in eukaryotic hosts. Another two-component system, EvgA and EvgS, participates in activation of the low-pH response, at least in part by activating PhoPQ and thus increasing synthesis of the IraM antiadaptor (37). PhoP directly and indirectly regulates some of the pH-resistance genes and *phoP* mutants are sensitive to acid stress (170). Some of these effects seem likely to involve the PhoPQ-dependent stabilization of RpoS.

RprA, the Rcs phosphorelay, and biofilm maturation.—An important developmental pathway for *E. coli* and most other bacteria is the development of biofilms. Cells in biofilms are relatively quiescent, and *rpoS* mutants are defective in formation of mature biofilms. Some but not all genes expressed in biofilms are RpoS dependent (1, 32, 60). Thus, it is not surprising that some of the signals leading to RpoS accumulation are also tied to biofilm development.

One regulator of RpoS that may be important in biofilms is the sRNA RprA. RprA is activated by the Rcs phosphorelay (84). The Rcs phosphorelay, important for biofilm formation, is necessary for expression of the genes needed for colanic capsule synthesis in *E. coli* K12, activates a large number of genes, and directly represses others, including genes for flagellar synthesis (reviewed in References 29 and 83). It is not clear which ligand or physiological condition(s) activates the Rcs phosphorelay. However, the genes in the Rcs regulon and the observation that the system is induced by solid surfaces suggest that the Rcs phosphorelay may be important during the development of biofilms, when motility needs to be shut down and capsule accumulates. RpoS is needed for biofilm maturation, and too much or too little RpoS interferes with biofilm development (40). It is attractive to suggest that RprA activation of *rpoS* translation helps to ensure properly timed expression of RpoS during biofilm maturation. Increased osmolarity also induces the Rcs system (43, 129), and in the absence of DsrA, RprA activates *rpoS* translation under osmotic-shock conditions (81).

The interaction of the Rcs phosphorelay and RpoS extends well beyond RprA. There are in fact a number of other genes for which the Rcs phosphorelay may provide regulation, both directly, via RcsB, and indirectly, via stimulation of RpoS. *ftsZ*, encoding the primary septation protein, is synthesized from multiple promoters, among them the RcsB-dependent pZ1 promoter and the RpoS-dependent pQ1 promoter (8, 41). *osmB*, an osmotically

inducible gene encoding a periplasmic protein of unknown function, is transcribed from two promoters, one activated by RcsB and one dependent on RpoS (18). As mentioned below, LrhA represses RpoS, probably via an unidentified sRNA. RcsB also represses *LrhA* expression; thus, in this case, RcsB acts via the LrhA-dependent pathway to stimulate RpoS expression (106).

Another regulatory protein, CsgD, reinforces the connection between RpoS and biofilm development. CsgD, a transcriptional factor important for biofilm development, promotes stabilization of RpoS by increasing synthesis of the IraP antiadaptor (50) (Figure 1). Thus, at the level of both RpoS translation and stabilization, induction of biofilm formation leads to increased RpoS accumulation. CsgD is under complex transcriptional and post-transcriptional control, including a promoter stimulated by RpoS, providing a positive-feedback loop for promoting RpoS accumulation under specific conditions (58, 102).

Too much and too little oxygen: OxyR and ArcB/ArcA negatively regulate RpoS.

—Two sRNAs, OxyS and ArcZ, play opposite roles in regulating RpoS levels as a function of the oxygen/energy status of the cell. Stationary-phase cells are resistant to oxidative damage such as hydrogen peroxide, and one of the first RpoS-dependent genes found was *katE*, encoding a catalase (123). A parallel pathway for resistance to hydrogen peroxide is regulated by OxyR, which in turn activates another catalase gene, *katG* (reviewed in Reference 136). OxyS, an sRNA transcribed in response to OxyR activation, negatively regulates *rpoS* translation (165). Thus, expression of the OxyR response to oxidative damage or the RpoS response may be mutually exclusive, although they share some effector genes, including *dps*. Possibly, the specific OxyR response is preferable when the cell encounters oxidative stress during exponential growth, and the RpoS response is used only when RpoS is induced in response to stationary-phase transition and/or another stress.

ArcZ, the third sRNA activator of RpoS, is negatively regulated by the two-component system ArcB/ArcA (86). ArcZ is processed to a 56-nt-long RNA from the 3' end of the initial 120-nt transcript; the 56-nt form of ArcZ can pair with the same region of the *rpoS* hairpin as DsrA and RprA and activate translation similarly (86). Under aerobic conditions, ArcZ is well expressed, but under anaerobic conditions, ArcA represses ArcZ and thus, indirectly, downregulates *rpoS* translation (86). As with RprA and DsrA, ArcZ sRNA serves as a connector between the general stress response and a more specific response mediated by the ArcB/ArcA two-component system. ArcA and ArcB have previously been implicated in negative regulation of RpoS at the level of transcription (ArcA represses the *rpoS* promoter) and degradation (ArcB-P stimulates RpoS degradation, by increased phosphorylation of RssB) (92). Although the relative contributions of each of these pathways is not yet clear, this concerted downregulation of RpoS by Arc suggests that the cell does not perceive anaerobiosis as a stress or that the specific Arc response is preferable to the RpoS-dependent response.

CONCLUSIONS

The discussion above touches on some of what we now know about the RpoS system, how it is induced, and what genes regulated by RpoS may do for the cell. The combination of years of work on *E. coli* as an experimental system and the advent of genome-wide approaches to monitoring gene expression have left us with a long list of stresses that induce the RpoS system and genes that get induced. However, we are still far from having a complete picture of the network for this response. We study RpoS in the laboratory, under the conditions easiest for us to manipulate. We hope that a full understanding of the functions of downstream genes will allow a better understanding of the stresses that the bacteria has evolved to respond to outside the laboratory.

In the paper that first named RpoS, Lange & Hengge-Aronis (75) noted that many of the resistance phenotypes of the RpoS system were also dealt with by very specific stress systems independent of RpoS. In fact, we now see that, whereas some specific stress systems collaborate and overlap with RpoS (for instance, the Rcs phosphorelay, the PhoPQ two component system, and the (p)ppGpp-dependent stringent response), others act as negative regulators of RpoS or RpoS-dependent genes. Included in the latter are the OxyR-dependent oxidative stress response and the ArcA/ArcB system. Others are likely to be found. For instance, stationary-phase cells show resistance to DNA damage, yet this response is induced independent of the classic LexA-dependent SOS response (90). PhoBR, the two-component system for P starvation, does not contribute significantly to the induction of RpoS after P starvation via IraP (17). What the advantage or disadvantage of keeping these systems distinct (for instance, not simultaneously inducing the specific and the general response for oxidative stress resistance) is not yet clear, but it may reflect only the need to minimize sigma factor competition, allowing either the RpoD-dependent systems or the RpoS-dependent systems to induce fully.

Why does acid stress or P starvation lead to a general stress response, giving overall resistance to many other stresses? Are these conditions usually encountered by the cell either during an infection and development of a biofilm or when cells reach stationary phase? The general stress response may in fact have evolved as a response to stationary phase in which multiple nutrients are limiting and toxic compounds are accumulating. If these stresses are always associated as the end of growth nears, it would make sense for signals of one sort to be interpreted by the cell as evidence for the coming disaster. Inducing a specific stress response may not be feasible when a cell is truly starving. One possible solution for invoking a general rather than a specific response would be if induction of the general stress response is organized such that multiple stresses must coincide for a robust response.

Recent progress has uncovered many of the mechanisms for regulating RpoS rapidly in response to changes in growth conditions. However, new inducing signals continue to be found, and many are not fully understood (Table 1). Unexplored are the physiological roles of most of the RpoS-dependent genes; the advent of genome-wide studies of phenotypes may begin to uncover their functions.

ACKNOWLEDGMENTS

We thank members of our laboratory as well as A. Bougdour, G. Storz, M. Cashel, R. Weisberg, K. Ramamurthi, T. Silhavy, and T. Elliott for their advice, comments on the manuscript, and answers to our questions. We apologize for all the excellent work on RpoS that we were unable to cover in this review. Preparation of this review and research relevant to it in this laboratory were supported by the Intramural Research Program of the National Institutes of Health, National Cancer Institute, Center for Cancer Research.

LITERATURE CITED

- Adams JL, McLean RJ. 1999 Impact of rpoS deletion on *Escherichia coli* biofilms. *Appl. Environ. Microbiol* 65:4285–87 [PubMed: 10473455]
- Almiron M, Link AJ, Furlong D, Kolter R. 1992 A novel DNA-binding protein with regulatory and protective roles in starved *Escherichia coli*. *Genes Dev* 6:2646–54 [PubMed: 1340475]
- Altuvia S, Almiron M, Huisman G, Kolter R, Storz G. 1994 The *dps* promoter is activated by OxyR during growth and by IHF and σ^S in stationary phase. *Mol. Microbiol* 13:265–72 [PubMed: 7984106]
- Andersson RA, Palva ET, Pirhonen M. 1999 The response regulator *expM* is essential for the virulence of *Erwinia carotovora* subsp. *carotovora* and acts negatively on the sigma factor RpoS (σ^S). *Mol. Plant Microbe Interact* 12:575–84 [PubMed: 10478478]
- Arnqvist A, Olsen A, Normark S. 1994 σ^S -dependent growth-phase induction of the *csgBA* promoter in *Escherichia coli* can be achieved in vivo by σ^{70} in the absence of the nucleoid-associated protein H-NS. *Mol. Microbiol* 13:1021–32 [PubMed: 7854117]
- Artsimovitch I, Patlan V, Sekine S, Vassilyeva MN, Hosaka T, et al. 2004 Structural basis for transcription regulation by alarmone ppGpp. *Cell* 117:299 [PubMed: 15109491]
- Balandina A, Claret L, Hengge-Aronis R, Rouviere-Yaniv J. 2001 The *Escherichia coli* histone-like protein HU regulates *rpoS* translation. *Mol. Microbiol* 39:1069–79 [PubMed: 11251825]
- Ballesteros M, Kusano S, Ishihama A, Vicente M, 1998 The *ftsQlp* gearbox promoter of *Escherichia coli* is a major sigma S-dependent promoter in the *ddlB-ftsA* region. *Mol. Microbiol* 30:419–30 [PubMed: 9791185]
- Barker MM, Gaal T, Josaitis CA, Gourse RL. 2001 Mechanism of regulation of transcription initiation by ppGpp. I. Effects of ppGpp on transcription initiation in vivo and in vitro. *J. Mol. Biol* 305:673–88 [PubMed: 11162084]
- Barth M, Marschall C, Muffler A, Fischer D, Hengge-Aronis R. 1995 Role for the histone-like protein H-NS in growth phase-dependent and osmotic regulation of σ^S and many σ^S -dependent genes in *Escherichia coli*. *J. Bacteriol* 177:3455–64 [PubMed: 7768855]
- Bearson SMD, Benjamin WH Jr, Swords WE, Foster JW. 1996 Acid shock induction of RpoS is mediated by the mouse virulence gene *mviA* of *Salmonella typhimurium*. *J. Bacteriol* 178:2572–79 [PubMed: 8626324]
- Becker G, Klauck E, Hengge-Aronis R. 1999 Regulation of RpoS proteolysis in *Escherichia coli*: the response regulator RssB is a recognition factor that interacts with the turnover element in RpoS. *Proc. Natl. Acad. Sci. USA* 96:6439–44 [PubMed: 10339606]
- Becker G, Klauck E, Hengge-Aronis R. 2000 The response regulator RssB, a recognition factor for σ^S proteolysis in *Escherichia coli*, can act like an anti- σ^S factor. *Mol. Microbiol* 35:657–66 [PubMed: 10672187]
- Bouche S, Klauck E, Fischer D, Lucassen M, Jung K, Hengge-Aronis R. 1998 Regulation of RssB-dependent proteolysis in *Escherichia coli*: a role for acetyl phosphate in a response regulator-controlled process. *Mol. Microbiol* 27:787–95 [PubMed: 9515704]
- Bougdour A, Cunning C, Baptiste PJ, Elliott T, Gottesman S. 2008 Multiple pathways for regulation of σ^S (RpoS) stability in *Escherichia coli* via the action of multiple anti-adaptors. *Mol. Microbiol* 68:298–313 [PubMed: 18383615]
- Bougdour A, Gottesman S. 2007 ppGpp regulation of RpoS degradation via an anti-adaptor protein IraP. *Proc. Natl. Acad. Sci. USA* 104:12896–901 [PubMed: 17640895]

17. Bougdour A, Wickner S, Gottesman S. 2006 Modulating RssB activity: IraP, a novel regulator of σ^S stability in *Escherichia coli*. *Genes Dev* 20:884–97 [PubMed: 16600914]
18. Boulanger A, Francez-Charlot A, Conter A, Castanie-Cornet M-P, Cam K, Gutierrez C. 2005 Multi-stress regulation in *Escherichia coli*: expression of *osmB* involves two independent promoters responding either to σ^S or to the RcsCDB His-Asp phosphorelay. *J. Bacteriol* 187:3282–86 [PubMed: 15838058]
19. Brennan RG, Link TM. 2007 Hfq structure, function and ligand binding. *Curr. Opin. Microbiol* 10:125–33 [PubMed: 17395525]
20. Brown L, Elliott T. 1996 Efficient translation of the RpoS sigma factor in *Salmonella typhimurium* requires Host Factor I, an RNA-binding protein encoded by the *hfq* gene. *J. Bacteriol* 178:3763–70 [PubMed: 8682778]
21. Brown L, Elliott T. 1997 Mutations that increase expression of the *rpoS* gene and decrease its dependency on *hfq* function in *Salmonella typhimurium*. *J. Bacteriol* 179:656–62 [PubMed: 9006017]
22. Brown L, Gentry D, Elliott T, Cashel M. 2002 DksA affects ppGpp induction of RpoS at a translational level. *J. Bacteriol* 184:4455–65 [PubMed: 12142416]
23. Carabetta VJ, Mohanty BK, Kushner SR, Silhavy TJ. 2009 The response regulator SprE (RssB) modulates polyadenylation of mRNA stability in *Escherichia coli*. *J. Bacteriol* 191:6812–21 [PubMed: 19767441]
24. Carabetta VJ, Silhavy TJ, Cristea IM. 2010 The response regulator sprE (RssB) is required for maintaining poly(A) polymerase I-degradosome association during stationary phase. *J. Bacteriol* 192:3713–21 [PubMed: 20472786]
25. Casadaban MJ, Cohen SN. 1979 Lactose genes fused to exogenous promoters in one step using a Mu-lac bacteriophage: in vivo probe for transcriptional control sequences. *Proc. Natl. Acad. Sci. USA* 76:4530–33 [PubMed: 159458]
26. Cavanagh AT, Klocko AD, Liu X, Wassarman KM. 2008 Promoter specificity for 6S RNA regulation of transcription is determined by core promoter sequences and competition for region 4.2 of σ^{70} . *Mol. Microbiol* 67:1242–56 [PubMed: 18208528]
27. Chang YY, Cronan JE Jr. 1999 Membrane cyclopropane fatty acid content is a major factor in acid resistance of *Escherichia coli*. *Mol. Microbiol* 33:249–59 [PubMed: 10411742]
28. Chatterji D, Fujita N, Ishihama A. 1998 The mediator for stringent control, ppGpp, binds to the beta-subunit of *Escherichia coli* RNA polymerase. *Genes Cells* 3:279 [PubMed: 9685179]
29. Clarke DJ. 2010 The Rcs phosphorelay: more than just a two-component pathway. *Future Microbiol* 5:1173–84 [PubMed: 20722597]
30. Cohen-Or I, Shenhar Y, Biran D, Ron EZ. 2010 CspC regulates *rpoS* transcript levels and complements *hfq* deletions. *Res. Microbiol* 161:694–700 [PubMed: 20633642]
31. Colland F, Barth M, Hengge-Aronis R, Kolb A. 2000 Sigma factor selectivity of *Escherichia coli* RNA polymerase: role for CRP, IHF and Lrp transcription factors. *EMBO J* 19:3028–37 [PubMed: 10856246]
32. Collet A, Cosette P, Beloin C, Ghigo J-M, Rihouey C, et al. 2008 Impact of *rpoS* deletion on the proteome of *Escherichia coli* grown planktonically and as biofilm. *J. Proteome Res* 7:4659–69 [PubMed: 18826300]
33. Costanzo A, Ades SE. 2006 Growth phase-dependent regulation of the extracytoplasmic stress factor, sigmaE, by guanosine 3',5'-bispyrophosphate (ppGpp). *J. Bacteriol* 188:4627–34 [PubMed: 16788171]
34. Cunning C, Brown L, Elliott T. 1998 Promoter substitution and deletion analysis of upstream region required for *rpoS* translational regulation. *J. Bacteriol* 180:4564–70 [PubMed: 9721296]
35. Demple B, Halbrook J, Linn S. 1983 *Escherichia coli xth* mutants are hypersensitive to hydrogen peroxide. *J. Bacteriol* 153:1079–82 [PubMed: 6337115]
36. Durfee T, Hansen A-M, Zhi H, Blattner FR, Jin DJ. 2008 Transcription profiling of the stringent response in *Escherichia coli*. *J. Bacteriol* 190:1084–96 [PubMed: 18039766]
37. Eguchi Y, Ishii E, Hata K, Utsumi R. 2011 Regulation of acid resistance by connectors of two-component signal transduction systems in *Escherichia coli*. *J. Bacteriol* 193:1222–28 [PubMed: 21193607]

38. Farewell A, Kvint K, Nystrom T. 1998 Negative regulation by RpoS: a case of sigma factor competition. *Mol. Microbiol* 29:1039–51 [PubMed: 9767572]
39. Ferenci T 2005 Maintaining a healthy SPANC balance through regulatory and mutational adaptation. *Mol. Microbiol* 57:1–8 [PubMed: 15948944]
40. Ferrieres L, Thompson A, Clarke DJ. 2009 Elevated levels of σ^S inhibit biofilm formation in *Escherichia coli*: a role for the Rcs phosphorelay. *Microbiology* 155:3544–53 [PubMed: 19696107]
41. Flardh K, Garrido T, Vicente M. 1997 Contribution of individual promoters in the *ddlB-ftsZ* region to the transcription of the essential cell-division gene *ftsZ* in *Escherichia coli*. *Mol. Microbiol* 24:927–36 [PubMed: 9220001]
42. Flynn JM, Neher SB, Kim Y-I, Sauer RT, Baker TA. 2003 Proteomic discovery of cellular substrates of the ClpXP protease reveals five classes of ClpX-recognition signals. *Mol. Cell* 11:671–83 [PubMed: 12667450]
43. Francez-Charlot A, Castanie-Cornet MP, Gutierrez C, Cam K. 2005 Osmotic regulation of the *Escherichia coli* *bdm* (biofilm-dependent modulation) gene by the RcsCDB His-Asp phosphorelay. *J. Bacteriol* 187:3873–77 [PubMed: 15901715]
44. Fredriksson A, Ballesteros M, Peterson CN, Persson O, Silhavy TJ, Nystrom T. 2007 Decline in ribosomal fidelity contributes to the accumulation and stabilization of the master stress response regulator σ^S upon carbon starvation. *Genes Dev* 21:862–74 [PubMed: 17403784]
45. Gaal T, Mandel MJ, Silhavy TJ, Gourse RL. 2006 Crl facilitates RNA polymerase holoenzyme formation. *J. Bacteriol* 188:7966–70 [PubMed: 16980472]
46. Gaal T, Ross W, Estrem ST, Nguyen LH, Burgess RR, Gourse RL. 2001 Promoter recognition and discrimination by σ^S RNA polymerase. *Mol. Microbiol* 42:939–54 [PubMed: 11737638]
47. Gibson KE, Silhavy TJ. 2000 SprE levels are growth phase regulated in a σ^S -dependent manner at the level of translation. *J. Bacteriol* 182:4117–20 [PubMed: 10869095]
48. Grainger DC, Goldberg MD, Lee DJ, Busby SJW. 2008 Selective repression by Fis and H-NS at the *Escherichia coli* *dps* promoter. *Mol. Microbiol* 68:1366–77 [PubMed: 18452510]
49. Groat RG, Schultz JE, Zychlinsky E, Bockman A, Matin A. 1986 Starvation proteins in *Escherichia coli*: kinetics of synthesis and role in starvation survival. *J. Bacteriol* 168:486–93 [PubMed: 3536847]
50. Gualdi L, Tagliabue L, Landini P. 2007 Biofilm formation-gene expression relay system in *Escherichia coli*: modulation of σ^S -dependent gene expression by the CsgD regulatory protein via σ^S protein stabilization. *J. Bacteriol* 189:8034–43 [PubMed: 17873038]
51. Hengge R 2009 Proteolysis of σ^S (RpoS) and the general stress response in *Escherichia coli*. *Res. Microbiol* 160:667–76 [PubMed: 19765651]
52. Hengge R 2011 The general stress response in gram-negative bacteria. See Ref 135, pp. 251–89
53. Hengge-Aronis R 2002 Signal transduction and regulatory mechanisms involved in control of the σ^S (RpoS) subunit of RNA polymerase. *Microbiol. Mol. Biol. Rev* 66:373–95 [PubMed: 12208995]
54. Hengge-Aronis R, Fischer D. 1992 Identification and molecular analysis of *glgS*, a novel growth-phase-regulated and *rpoS*-dependent gene involved in glycogen synthesis in *Escherichia coli*. *Mol. Microbiol* 6:1877–86 [PubMed: 1324388]
55. Hengge-Aronis R, Klein W, Lange R, Rimmele M, Boos W. 1991 Trehalose synthesis genes are controlled by the putative sigma factor encoded by *rpoS* and are involved in stationary-phase thermotolerance in *Escherichia coli*. *J. Bacteriol* 173:7918–24 [PubMed: 1744047]
56. Hirsch M, Elliott T. 2002 Role of ppGpp in *rpoS* stationary-phase regulation in *Escherichia coli*. *J. Bacteriol* 184:5077–87 [PubMed: 12193624]
57. Hirsch M, Elliott T. 2005 Fis regulates transcriptional induction of RpoS in *Salmonella enterica*. *J. Bacteriol* 187:1568–80 [PubMed: 15716427]
58. Holmqvist E, Reimegard J, Sterk M, Grantcharova N, Romling U, Wagner EG. 2010 Two antisense RNAs target the transcriptional regulator CsgD to inhibit curli synthesis. *EMBO J* 29:1840–50 [PubMed: 20407422]
59. Hussein R, Lim HL. 2010 Disruption of small RNA signaling caused by competition for Hfq. *Proc. Natl. Acad. Sci. USA* 108:1110–15 [PubMed: 21189298]

60. Ito A, May T, Kawata K, Okabe S. 2008 Significance of *rpoS* during maturation of *Escherichia coli* biofilms. *Biotechnol. Bioeng* 99:1462–71 [PubMed: 17979199]
61. Jin Y, Watt RM, Danchin A, Huang J-D. 2009 Small noncoding RNA GcvB is a novel regulator of acid resistance in *Escherichia coli*. *BMC Genomics* 10:165 [PubMed: 19379489]
62. Jishage M, Ishihama A. 1995 Regulation of RNA polymerase sigma subunit synthesis in *Escherichia coli*: intracellular levels of σ^{70} and σ^{38} . *J. Bacteriol* 177:6832–5 [PubMed: 7592475]
63. Jishage M, Ishihama A. 1997 Variation in RNA polymerase sigma subunit composition within different stocks of *Escherichia coli* W3110. *J. Bacteriol* 179:959–63 [PubMed: 9006056]
64. Jishage M, Iwata A, Ueda S, Ishihama A. 1996 Regulation of RNA polymerase sigma subunit synthesis in *Escherichia coli*: intracellular levels of four species of sigma subunit under various growth conditions. *J. Bacteriol* 178:5447–51 [PubMed: 8808934]
65. Kabir MS, Sagara T, Oshima T, Kawagoe Y, Mori H, et al. 2004 Effects of mutations in the *rpoS* gene on cell viability and global gene expression under nitrogen starvation in *Escherichia coli*. *Microbiology* 150:2543–53 [PubMed: 15289551]
66. Kandror O, DeLeon A, Goldberg AL. 2002 Trehalose synthesis is induced upon exposure of *Escherichia coli* to cold and is essential for viability at low temperatures. *Proc. Natl. Acad. Sci. USA* 99:9727–32 [PubMed: 12105274]
67. Kenyon CJ, Walker GC. 1980 DNA-damaging agents stimulate gene expression at specific loci in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 77:2819–23 [PubMed: 6771759]
68. Keseler IM, Bonavides-Martinez C, Collado-Vides J, Gama-Castro S, Gunsalus RP, et al. 2009 EcoCyc: a comprehensive view of *Escherichia coli* biology. *Nucleic Acids Res* 37:D464–70 [PubMed: 18974181]
69. Kiley PJ, Donohue TJ. 2011 Global responses of bacteria to oxygen deprivation. See Ref 135, pp. 175–89
70. King T, Ishihama A, Kori A, Ferenci T. 2004 A regulatory trade-off as a source of strain variation in the species *Escherichia coli*. *J. Bacteriol* 186:5614–20 [PubMed: 15317765]
71. Klauck E, Lingnau M, Hengge-Aronis R. 2001 Role of the response regulator RssB in σ^S recognition and initiation of σ^S proteolysis in *Escherichia coli*. *Mol. Microbiol* 40:1381–90 [PubMed: 11442836]
72. Kvint K, Farewell A, Nystrom T. 2000 RpoS-dependent promoters require guanosine tetraphosphate for induction even in the presence of high levels of σ^S . *J. Biol. Chem* 275:14795–98 [PubMed: 10747855]
73. Lacour S, Kolb A, Boris Zehnder AJ, Landini P. 2002 Mechanism of specific recognition of the *aidB* promoter by σ^S -RNA polymerase. *Biochem. Biophys. Res. Commun* 292:922–30 [PubMed: 11944903]
74. Lange R, Fischer D, Hengge-Aronis R. 1995 Identification of transcriptional start sites and the role of ppGpp in the expression of *rpoS*, the structural gene for the σ^S subunit of RNA polymerase in *Escherichia coli*. *J. Bacteriol* 177:4676–80 [PubMed: 7642494]
75. Lange R, Hengge-Aronis R. 1991 Identification of a central regulator of stationary-phase gene expression in *Escherichia coli*. *Mol. Microbiol* 5:49–59 [PubMed: 1849609]
76. Lange R, Hengge-Aronis R. 1994 The cellular concentration of the σ^S subunit of RNA polymerase in *Escherichia coli* is controlled at the levels of transcription, translation, and protein stability. *Genes Dev* 8:1600–12 [PubMed: 7525405]
77. Lease RA, Belfort M. 2000 Riboregulation by DsrA RNA: transactions for global economy. *Mol. Microbiol* 38:667–72 [PubMed: 11115103]
78. Lee IS, Lin J, Hall HK, Bearson B, Foster JW. 1995 The stationary-phase sigma factor σ^S (RpoS) is required for a sustained acid tolerance response in virulent *Salmonella typhimurium*. *Mol. Microbiol* 17:155–67 [PubMed: 7476202]
79. Lemaux PG, Herendeen SL, Bloch PL, Neidhardt FC. 1978 Transient rates of synthesis of individual polypeptides in *E. coli* following temperature shifts. *Cell* 13:427–34 [PubMed: 350413]
80. Lucchini S, McDermott P, Thompson A, Hinton JCD. 2009 The H-NS-like protein StpA represses the RpoS (σ^{38}) regulon during exponential growth in *Salmonella typhimurium*. *Mol. Microbiol* 74:1169–86 [PubMed: 19843227]

81. Majdalani N, Chen S, Murrow J, St. John K, Gottesman S. 2001 Regulation of RpoS by a novel small RNA: the characterization of RprA. *Mol. Microbiol* 39:1382–94 [PubMed: 11251852]
82. Majdalani N, Cuning C, Sledjeski D, Elliott T, Gottesman S. 1998 DsrA RNA regulates translation of RpoS message by an anti-antisense mechanism, independent of its action as an antisilencer of transcription. *Proc. Natl. Acad. Sci. USA* 95:12462–67 [PubMed: 9770508]
83. Majdalani N, Gottesman S. 2005 The Rcs phosphorelay: a complex signal transduction system. *Annu. Rev. Microbiol* 59:379–405 [PubMed: 16153174]
84. Majdalani N, Hernandez D, Gottesman S. 2002 Regulation and mode of action of the second small RNA activator of RpoS translation, RprA. *Mol. Microbiol* 46:813–26 [PubMed: 12410838]
85. Mandel MJ, Silhavy TJ. 2005 Starvation for different nutrients in *Escherichia coli* results in differential modulation of RpoS levels and stability. *J. Bacteriol* 187:434–42 [PubMed: 15629914]
86. Mandin P, Gottesman S. 2010 Integrating anaerobic/aerobic sensing and the general stress response through the ArcZ small RNA. *EMBO J* 29:3094–107 [PubMed: 20683441]
87. Marschall C, Labrousse V, Kreimer M, Weichart D, Kolb A, Hengge-Aronis R. 1998 Molecular analysis of the regulation of *csiD*, a carbon starvation-inducible gene in *Escherichia coli* that is exclusively dependent on σ^S and requires activation by cAMP-CRP. *J. Mol. Biol* 276:339–53 [PubMed: 9512707]
88. Martinez A, Kolter R. 1997 Protection of DNA during oxidative stress by the nonspecific DNA-binding protein Dps. *J. Bacteriol* 179:5188–94 [PubMed: 9260963]
89. McCullen C, Benhammou JN, Majdalani N, Gottesman S. 2010 Mechanism of positive regulation by DsrA and RprA small noncoding RNAs: pairing increases translation and protects *rpoS* mRNA from degradation. *J. Bacteriol* 192:5559–71 [PubMed: 20802038]
90. Merrikh H, Bougdour A, Olivier-Mason A, Lovett ST. 2009 A DNA damage response in *Escherichia coli* involving the alternative sigma factor, RpoS. *Proc. Natl. Acad. Sci. USA* 106:611–16 [PubMed: 19124769]
91. Merrikh H, Ferrazzoli AE, Lovett ST. 2009 Growth phase and (p)ppGpp control of IraD, a regulator of RpoS stability, in *Escherichia coli*. *J. Bacteriol* 191:7436–46 [PubMed: 19820090]
92. Mika F, Hengge R. 2005 A two-component phosphotransfer network involving ArcB, ArcA, and RssB coordinates synthesis and proteolysis of σ^S (RpoS) in *E. coli*. *Genes Dev* 19:2770–81 [PubMed: 16291649]
93. Mitchell JE, Oshima T, Piper SE, Webster SL, Westblade LF, et al. 2007 The *Escherichia coli* regulator of sigma 70 protein, Rsd, can up-regulate some stress-dependent promoters by sequestering sigma 70. *J. Bacteriol* 189:3489–95 [PubMed: 17351046]
94. Moreno M, Audia JP, Bearson SM, Webb C, Foster JM. 2000 Regulation of sigma S degradation in *Salmonella enterica* var typhimurium: in vivo interactions between sigma S, the response regulator MviA (RssB) and ClpX. *J. Mol. Microbiol. Biotechnol* 2:245–54 [PubMed: 10939250]
95. Muffler A, Barth M, Marschall C, Hengge-Aronis R. 1997 Heat shock regulation of σ^S turnover: a role for DnaK and relationship between stress responses mediated by σ^S and σ^{32} in *Escherichia coli*. *J. Bacteriol* 179:445–52 [PubMed: 8990297]
96. Muffler A, Fischer D, Altuvia S, Storz G, Hengge-Aronis R. 1996 The response regulator RssB controls stability of the σ^S subunit of RNA-polymerase in *Escherichia coli*. *EMBO J* 15:1333–39 [PubMed: 8635466]
97. Muffler A, Fischer D, Hengge-Aronis R. 1996 The RNA-binding protein HF-I, known as a host factor for phage Q β RNA replication, is essential for *rpoS* translation in *Escherichia coli*. *Genes Dev* 10:1143–51 [PubMed: 8654929]
98. Muffler A, Traulsen DD, Fischer D, Lange R, Hengge-Aronis R. 1997 The RNA-binding protein HF-I plays a global regulatory role which is largely, but not exclusively, due to its role in expression of the σ^S subunit of RNA polymerase in *Escherichia coli*. *J. Bacteriol* 179:297–300 [PubMed: 8982015]
99. Muffler A, Traulsen DD, Lange R, Hengge-Aronis R. 1996 Posttranscriptional osmotic regulation of the σ^S subunit of RNA polymerase in *Escherichia coli*. *J. Bacteriol* 178:1607–13 [PubMed: 8626288]

100. Mukhopadhyay S, Audia JP, Roy RN, Schellhorn HE. 2000 Transcriptional induction of the conserved alternative sigma factor RpoS in *Escherichia coli* is dependent on BarA, a probable two-component regulator. *Mol. Microbiol* 37:371–81 [PubMed: 10931332]
101. Nair S, Finkel SE. 2004 Dps protects cells against multiple stresses during stationary phase. *J. Bacteriol* 186:4192–98 [PubMed: 15205421]
102. Ogasawara H, Yamada K, Kori A, Yamamoto K, Ishihama A. 2010 Regulation of the *Escherichia coli csgD* promoter: interplay between five transcription factors. *Microbiology* 156:2470–83 [PubMed: 20466769]
103. Opdyke JA, Kang J-G, Storz G. 2004 GadY, a small-RNA regulator of acid response genes in *Escherichia coli*. *J. Bacteriol* 186:6698–705 [PubMed: 15466020]
104. Paul BJ, Barker MM, Ross W, Schneider DA, Webb C, et al. 2004 DksA: a critical component of the transcription initiation machinery that potentiates the regulation of rRNA promoters by ppGpp and the initiating NTP. *Cell* 118:311–22 [PubMed: 15294157]
105. Paul BJ, Ross W, Gaal T, Gourse RL. 2004 rRNA transcription in *Escherichia coli*. *Annu. Rev. Genet* 38:749–70 [PubMed: 15568992]
106. Peterson CN, Carabetta VJ, Chowdhury T, Silhavy TJ. 2006 LrhA regulates *rpoS* translation in response to the Rcs phosphorelay system in *Escherichia coli*. *J. Bacteriol* 188:3175–81 [PubMed: 16621809]
107. Peterson CN, Mandel MJ, Silhavy TJ. 2005 *Escherichia coli* starvation diets: essential nutrients weigh in distinctly. *J. Bacteriol* 187:7549–53 [PubMed: 16267278]
108. Peterson CN, Ruiz N, Silhavy TJ. 2004 RpoS proteolysis is regulated by a mechanism that does not require the SprE (RssB) response regulator phosphorylation site. *J. Bacteriol* 186:7403–10 [PubMed: 15489452]
109. Phadtare S, Inouye M. 2001 Role of CspC and CspE in regulation of expression of RpoS and UspA, the stress response proteins in *Escherichia coli*. *J. Bacteriol* 183:1205–14 [PubMed: 11157932]
110. Piper SE, Mitchell JE, Lee DJ, Busby SJW. 2009 A global view of *Escherichia coli* Rsd protein and its interactions. *Mol. Biosyst* 5:1943–47 [PubMed: 19763331]
111. Potrykus K, Cashel M. 2008 (p)ppGpp: still magical? *Annu. Rev. Microbiol* 62:35–51 [PubMed: 18454629]
112. Pratt LA, Silhavy TJ. 1996 The response regulator SprE controls the stability of RpoS. *Proc. Natl. Acad. Sci. USA* 93:2488–92 [PubMed: 8637901]
113. Pratt LA, Silhavy TJ. 1998 Crl stimulates RpoS activity during stationary phase. *Mol. Microbiol* 29:1225–36 [PubMed: 9767590]
114. Pruteanu M, Hengge-Aronis R. 2002 The cellular level of the recognition factor RssB is rate-limiting for σ^S proteolysis: implications for RssB regulation and signal transduction in σ^S turnover in *Escherichia coli*. *Mol. Microbiol* 45:1701–13 [PubMed: 12354235]
115. Ranquet C, Gottesman S. 2007 Translational regulation of the *Escherichia coli* stress factor RpoS: a role for SsrA and Lon. *J. Bacteriol* 189:4872–79 [PubMed: 17449615]
116. Repoila F, Gottesman S. 2001 Signal transduction cascade for regulation of RpoS: temperature regulation of DsrA. *J. Bacteriol* 183:4012–23 [PubMed: 11395466]
117. Repoila F, Gottesman S. 2003 Temperature sensing by the *dsrA* promoter. *J. Bacteriol* 185:6609–14 [PubMed: 14594834]
118. Resch A, Afonyushkin T, Lombo TB, McDowall KJ, Blasi U, Kaberdin VR. 2008 Translational activation by the noncoding RNA DsrA involves alternative RNase III processing in the *rpoS* 5'-leader. *RNA* 14:454–59 [PubMed: 18192613]
119. Resch A, Vecerek B, Palavra K, Blasi U. 2010 Requirement of the CsdA DEAD-box helicase for low temperature riboregulation of *rpoS* mRNA. *RNA Biol* 7:1–7
120. Ruiz N, Peterson CN, Silhavy TJ. 2001 RpoS-dependent transcriptional control of *sprE*: Regulatory feedback loop. *J. Bacteriol* 183:5974–81 [PubMed: 11566997]
121. Ruiz N, Silhavy TJ. 2003 Constitutive activation of the *Escherichia coli* Pho regulon upregulates *rpoS* translation in an Hfq-dependent fashion. *J. Bacteriol* 185:5984–92 [PubMed: 14526009]

122. Sak BD, Eisenstark A, Touati D. 1989 Exonuclease III and the catalase hydroperoxidase II in *Escherichia coli* are both regulated by the *katF* gene product. *Proc. Natl. Acad. Sci. USA* 86:3271–75 [PubMed: 2541439]
123. Schellhorn HE, Hassan HM. 1988 Transcriptional regulation of *katE* in *Escherichia coli* K-12. *J. Bacteriol* 170:4286–92 [PubMed: 3045091]
124. Schurdell MS, Woodbury GM, McCleary WR. 2007 Genetic evidence suggests that the intergenic region between *pstA* and *pstB* plays a role in the regulation of *rpoS* translation during phosphate limitation. *J. Bacteriol* 189:1150–53 [PubMed: 17142395]
125. Schweder T, Lee K-H, Lomovskaya O, Martin A. 1996 Regulation of *Escherichia coli* starvation sigma factor (σ^S) by ClpXP protease. *J. Bacteriol* 178:470–76 [PubMed: 8550468]
126. Sharma AK, Payne SM. 2006 Induction of expression of *hfq* by DksA is essential for *Shigella flexneri* virulence. *Mol. Microbiol* 62:469–79 [PubMed: 17020583]
127. Shiba T, Tsutsumi K, Yano H, Ihara Y, Kameda A, et al. 1997 Inorganic polyphosphate and the induction of *rpoS* expression. *Proc. Natl. Acad. Sci. USA* 94:11210–15 [PubMed: 9326588]
128. Silhavy TJ, Beckwith JR. 1985 Uses of *lac* fusions for the study of biological problems. *Microbiol. Rev* 49:398–418 [PubMed: 3005818]
129. Sledjeski DD, Gottesman S. 1996 Osmotic shock induction of capsule synthesis in *Escherichia coli* K-12. *J. Bacteriol* 178:1204–6 [PubMed: 8576059]
130. Sledjeski DD, Gupta A, Gottesman S. 1996 The small RNA, DsrA, is essential for the low temperature expression of RpoS during exponential growth in *Escherichia coli*. *EMBO J* 15:3993–4000 [PubMed: 8670904]
131. Soper T, Mandin P, Majdalani N, Gottesman S, Woodson SA. 2010 Positive regulation by small RNAs and the role of Hfq. *Proc. Natl. Acad. Sci. USA* 107:9602–7 [PubMed: 20457943]
132. Soper TJ, Woodson SA. 2008 The *rpoS* mRNA leader recruits Hfq to facilitate annealing with DsrA sRNA. *RNA* 14:1907–17 [PubMed: 18658123]
133. Spira B, Silberstein N, Yagil E. 1995 Guanosine 3',5'-bispyrophosphate (ppGpp) synthesis in cells of *Escherichia coli* starved for Pi. *J. Bacteriol* 177:4053–38 [PubMed: 7608079]
134. Stoebel DM, Hokamp K, Last MS, Dorman CJ. 2009 Compensatory evolution of gene regulation in response to stress by *Escherichia coli* lacking rpoS. *PLoS Genet* 5:e1000671 [PubMed: 19798444]
135. Storz G, Hengge R, eds. 2011 *Bacterial Stress Responses*. Washington, DC: ASM 2nd ed.
136. Storz G, Spiro S. 2011 Sensing and responding to reactive oxygen and nitrogen species. See Ref 135, pp. 157–73
137. Sutton A, Buencamino R, Eisenstark A. 2000 *rpoS* mutants in archival cultures of *Salmonella enterica* serovar Typhimurium. *J. Bacteriol* 182:4375–79 [PubMed: 10913067]
138. Tanaka K, Kusano S, Fujita N, Ishihama A, Takahashi H. 1995 Promoter determinants for *Escherichia coli* RNA polymerase holoenzyme containing σ^{38} (the rpoS gene product). *Nucleic Acids Res* 23:827–34 [PubMed: 7708499]
139. Tanaka K, Takayanagi Y, Fujita N, Ishihama A, Takahashi H. 1993 Heterogeneity of the principal σ factor in *Escherichia coli*: the *rpoS* gene product, σ^{38} , is a second principal σ factor of RNA polymerase in stationary-phase *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 90:8303 [PubMed: 8367498]
140. Tapley TL, Franzmann TM, Chakraborty S, Jakob U, Bardwell JCA. 2010 Protein refolding by pH-triggered chaperone binding and release. *Proc. Natl. Acad. Sci. USA* 107:1071–76 [PubMed: 20080625]
141. Touloukhonov II, Shulgina I, Hernandez VJ. 2001 Binding of the transcription effector ppGpp to *Escherichia coli* RNA polymerase is allosteric, modular, and occurs near the N terminus of the β subunit. *J. Biol. Chem* 276:1220–25 [PubMed: 11035017]
142. Tramonti A, De Canio M, De Biase D. 2008 GadX/GadW-dependent regulation of the *Escherichia coli* acid fitness island: transcriptional control at the *gadY-gadW* divergent promoters and identification of four novel 42 bp GadX/GadW-specific binding sites. *Mol. Microbiol* 70:965–82 [PubMed: 18808381]
143. Travers AA. 1980 Promoter sequence for stringent control of bacterial ribonucleic acid synthesis. *J. Bacteriol* 141:973–76 [PubMed: 6154042]

144. Traxler MF, Chang D-E, Conway T. 2006 Guanosine 3',5'-bispyrophosphate coordinates global gene expression during glucose-lactose diauxie in *Escherichia coli*. *Proc. Natl. Acad. Sci. USA* 103:2374–79 [PubMed: 16467149]
145. Traxler MF, Summers SM, Nguyen H-T, Zacharia VM, Hightower GA, et al. 2008 The global, ppGpp-mediated stringent response to amino acid starvation in *Escherichia coli*. *Mol. Microbiol* 68:1128–48 [PubMed: 18430135]
146. Traxler MF, Zacharia VM, Marquardt S, Summers SM, Nguyen H-T, et al. 2011 Discretely calibrated regulatory loops controlled by ppGpp partition gene induction across the “feast to famine” gradient in *Escherichia coli*. *Mol. Microbiol* 79:830–45 [PubMed: 21299642]
147. Trotochaud AE, Wassarman KM. 2006 6S RNA regulation of *pspF* transcription leads to altered cell survival at high pH. *J. Bacteriol* 188:3936–43 [PubMed: 16707685]
148. Trotochaud AE, Wassarman KM. 2004 6S RNA function enhances long-term cell survival. *J. Bacteriol* 186:4978–85 [PubMed: 15262935]
149. Tu X, Latifi T, Bougdour A, Gottesman S, Groisman EA. 2006 The PhoP/PhoQ two-component system stabilizes the alternative sigma factor RpoS in *Salmonella enterica*. *Proc. Natl. Acad. Sci. USA* 103:13503–8 [PubMed: 16938894]
150. Typas A, Barembruch C, Possling A, Hengge R. 2007 Stationary phase reorganisation of the *Escherichia coli* transcription machinery by Crl protein, a fine-tuner of σ^S activity and levels. *EMBO J* 26:1569–78 [PubMed: 17332743]
151. Typas A, Becker G, Hengge R. 2007 The molecular basis of selective promoter activation by the σ^S subunit of RNA polymerase. *Mol. Microbiol* 63:1296–306 [PubMed: 17302812]
152. van Dyk T, Myers BL, Morgan RW, Larossa RA. 1998 Constricted flux through the branched-chain amino acid biosynthetic enzyme acetolactate synthase triggers elevated expression of genes regulated by *rpoS* and internal acidification. *J. Bacteriol* 180:785–92 [PubMed: 9473030]
153. Volkert MR, Hajec LI, Matijasevic Z, Fang FC, Prince R. 1994 Induction of the *Escherichia coli* *aidB* gene under oxygen-limiting conditions requires a functional *rpoS* (*katF*) gene. *J. Bacteriol* 176:7638–45 [PubMed: 8002588]
154. Wassarman KS. 2007 6S RNA: a regulator of transcription. *Mol. Microbiol* 65:1425–31 [PubMed: 17714443]
155. Waterman SR, Small PL. 1996 Identification of σ^S -dependent genes associated with the stationary-phase acid-resistance phenotype of *Shigella flexneri*. *Mol. Microbiol* 21:925–40 [PubMed: 8885264]
156. Webb C, Moreno M, Wilmes-Riesenberg M, Curtiss RI, Foster JW. 1999 Effects of DksA and ClpP protease on sigma S production and virulence in *Salmonella typhimurium*. *Mol. Microbiol* 34:112–23 [PubMed: 10540290]
157. Weber H, Polen T, Heuveling J, Wendisch VF, Hengge R. 2005 Genome-wide analysis of the general stress response network in *Escherichia coli*: σ^S -dependent genes, promoters, and sigma factor selectivity. *J. Bacteriol* 187:1591–603 [PubMed: 15716429]
158. Xiao H, Kalman M, Ikehara K, Zemel S, Glaser G, Cashel M. 1991 Residual guanosine 3',5'-bispyrophosphate synthetic activity of *relA* null mutants can be eliminated by *spoT* null mutations. *J. Biol. Chem* 266:5980–90 [PubMed: 2005134]
159. Xu J, Johnson RC. 1995 Fis activates the RpoS-dependent stationary-phase expression of *proP* in *Escherichia coli*. *J. Bacteriol* 177:5222–31 [PubMed: 7545153]
160. Yamashino T, Ueguchi C, Mizuno T. 1995 Quantitative control of the stationary phase-specific sigma factor, σ^S , in *Escherichia coli*: involvement of the nucleoid protein H-NS. *EMBO J* 14:594–602 [PubMed: 7859747]
161. Yoshida T, Ueguchi C, Mizuno T. 1993 Physical map location of a set of *Escherichia coli* genes (*hde*) whose expression is affected by the nucleoid protein H-NS. *J. Bacteriol* 175:7747–48 [PubMed: 8244952]
162. Yuan AH, Gregory BD, Sharp JS, McCleary KD, Dove SL, Hochschild A. 2008 Rsd family proteins make simultaneous interactions with regions 2 and 4 of the primary sigma factor. *Mol. Microbiol* 70:1136–51 [PubMed: 18826409]

163. Zambrano MM, Siegele DA, Almiron M, Tormo A, Kolter R. 1993 Microbial competition: *Escherichia coli* mutants that take over stationary phase cultures. *Science* 259:1757–60 [PubMed: 7681219]
164. Zgurskaya HI, Keyhan M, Matin A. 1997 The sigma S level in starving *Escherichia coli* cells increases solely as a result of its increased stability, despite decreased synthesis. *Mol. Microbiol* 24:643–51 [PubMed: 9179856]
165. Zhang A, Altuvia S, Tiwari A, Argaman L, Hengge-Aronis R, Storz G. 1998 The *oxyS* regulatory RNA represses *rpoS* translation by binding Hfq (HF-1) protein. *EMBO J* 17:6061–68 [PubMed: 9774349]
166. Zhao B, Houry WA. 2010 Acid stress response in enteropathogenic Gammaproteobacteria: an aptitude for survival. *Biochem. Cell Biol* 88:301–14 [PubMed: 20453931]
167. Zhou Y, Gottesman S. 2006 Modes of regulation of RpoS by H-NS. *J. Bacteriol* 188:7022–25 [PubMed: 16980505]
168. Zhou Y, Gottesman S, Hoskins JR, Maurizi MR, Wickner S. 2001 The RssB response regulator directly targets σ^S for degradation by ClpXP. *Genes Dev* 15:627–37 [PubMed: 11238382]
169. Zhou Y-N, Gottesman S. 1998 Regulation of proteolysis of the stationary-phase sigma factor RpoS. *J. Bacteriol* 180:1154–58 [PubMed: 9495753]
170. Zwir I, Shin D, Kato A, Nishino K, Latifi T, et al. 2005 Dissecting the PhoP regulatory network of *Escherichia coli* and *Salmonella enterica*. *Proc. Natl. Acad. Sci. USA* 102:2862–67 [PubMed: 15703297]

SUMMARY POINTS

1. The RpoS regulon is induced when RpoS levels rise, primarily by changes in translation of *rpoS*, mediated by sRNAs, and by changes in RpoS degradation, mediated by antiadaptors, each induced under different conditions.
2. There is major overlap between the (p)ppGpp-induced stringent response and the RpoS response.
3. Many RpoS-dependent genes can also be read by RpoD under specific circumstances; H-NS plays a significant role in repressing the basal expression of these genes by RpoD and also negatively affects RpoS. The combined effect is to limit expression of the RpoS regulon to conditions of stress.
4. Overlap between genes necessary for resistance to high osmolarity as well as low and high temperature suggests common intracellular stresses under these conditions.
5. RpoS and some of the downstream genes have key roles in biofilm formation; the inducing environment in this case may be contact with a surface.
6. The Rcs phosphorelay and the PhoPQ two-component system contribute to RpoS induction through downstream connectors (sRNAs and antiadaptors), whereas the Arc and OxyR systems, among others, negatively regulate RpoS.

FUTURE ISSUES

1. What are the “core” RpoS regulon genes, dependent only on RpoS? What stress do these genes deal with, and does this define the broadest role for RpoS?
2. What RpoS-dependent genes are necessary for surviving starvation for C, P, and N, and how do they contribute to survival?
3. A variety of experiments and isolation of *E. coli* from the environment suggest that there are many conditions under which the cell accumulates mutations in *rpoS*. Why then is RpoS retained, when is it critical, and when is it bad for the cell?
4. What other stresses feed into the RpoS system? Have all inducing signals been defined?
5. When the RpoS system is induced, the cell is resistant to a wide range of stress and starvation treatments. During exponential growth, when RpoS is not induced, the cell has specific responses to specific stresses. To what extent do the effector genes for specific and global stress responses overlap, and if they do, how is specificity achieved? If they do not, why is it preferable to have different responses?
6. To what extent do the lessons learned from the *E. coli* general stress response extrapolate to other bacteria, and what can be learned from other bacteria to help in understanding the *E. coli* response?

Regulon:

a set of genes induced together in response to a signal or stress, under the control of a specific regulatory protein

Sigma factor:

a protein that binds to the RNA polymerase core and allows it to initiate transcription at different classes of promoters

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

RNAP:

RNA polymerase

(p)ppGpp (guanosine 3', 5'-bispyrophosphate):

the nucleotides pppGpp and ppGpp, derived, respectively, from GTP and GDP

Rcs phosphorelay:

lipoprotein RcsF transduces signals to kinase RcsC; phosphate then moves from RcsC via phosphotransfer protein RcsD to response regulator RcsB

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Cyclic AMP (cAMP):

the small molecule cAMP accumulates when cells are growing on suboptimal carbon sources

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Catabolite response protein (CRP):


binds cAMP and binds to DNA sites, positively and negatively regulating transcription

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript



ArcB/ArcA two-component system:

under anaerobic conditions, the ArcB sensor kinase activates ArcA; ArcA activates some anaerobic genes and represses aerobically expressed genes

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Hfq:

an RNA chaperone necessary for the function of many small RNAs in *E. coli* and other bacteria

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

SpoT:

a bifunctional enzyme that synthesizes but mainly degrades (p)ppGpp; plays a role in response to many kinds of non-amino acid starvation

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1.

Regulation of RpoS expression, stability, and activity. See text for details and references relative to these regulatory mechanisms. Different levels of regulation are shown in boxes: (a) transcriptional regulation, (b) translational regulation, (c) regulation of RpoS activity, and (d) proteolytic regulation. The major transcript for *rpoS* transcription is the one initiating at *PrpoS* within the *nlpD* open reading frame. This transcript forms a repressive hairpin loop that prevents ribosome binding, but positively acting small RNAs, with the help of the Hfq chaperone, release this inhibition and promote *rpoS* translation. OxyS small RNA inhibits *rpoS* translation likely by titrating out Hfq. Other proteins also impinge positively (CsdA, CspC, CspE, HU, DksA) or negatively (RNase III, H-NS) on *rpoS* translation. Once the RpoS (shown here as σ^S) protein is synthesized, it can have two fates: In exponential phase, it is bound by the adaptor protein RssB and degraded by the ClpXP protease, unless the cell encounters some stress that leads to synthesis of one of the antiadaptors (IraP, M, or D) that then interferes with RssB and leads to stabilization of RpoS. In stationary phase and under stress conditions, stable RpoS can bind free core RNA polymerase; this protects RpoS from degradation and allows the transcription of RpoS-dependent genes. 6S RNA, Crl, Rsd, and (p)ppGpp favor RpoS in its competition with RpoD (σ^{70}) for core. Some regulators [for instance, (p)ppGpp] act at multiple levels.

Table 1

Stress induction of the RpoS regulon

| Stress | Upstream mediator of stress signal | Level of regulation | Regulator of RpoS | Phenotype of <i>rpoS</i> mutant | Effector genes ^a | References |
|-----------------------------|------------------------------------|----------------------------|-----------------------|---|-----------------------------|-----------------------|
| Late exponential phase | Multiple signals; high (p)ppGpp | Translation | Multiple; ND | ND ^b | ND | (76, 91) |
| | | Degradation | Multiple: IraD, IraP? | | | |
| Phosphate starvation | SpoT/(p)ppGpp | Degradation | IraP | ND | ND | (16, 91) |
| DNA damage/UV | ND | Degradation | IraD | Sensitivity to AZT, H ₂ O ₂ | <i>dps</i> | (15, 90, 101) |
| Magnesium starvation | PhoQ/PhoP | Degradation | E.c: IraM S.t.: IraP | ND | ND | (15, 149) |
| Carbon starvation | Competition for protease | Degradation | ClpXP titration? | Loses viability faster, long-term starvation | ND | (44, 76) |
| Low temperature | DsrA promoter | Translation | DsrA | Slow growth, loss of viability at low temperature | <i>otsAB</i> | (66, 130) |
| High temperature | Secondary structure of mRNA? | Translation/mRNA stability | RNase III resistance | Loss of viability at high temperature | <i>otsAB, dps</i> | (55, 89, 101) |
| | ND | Degradation | DnaJ, DnaK | | | (55, 95) |
| Surface | Rcs phosphorelay | Translation | RprA | Poor biofilm formation | ND | (29, 84) |
| Aerobic/energy status | ArcB/ArcA negative regulation | Transcription | ArcA repression | ND | ND | (92) |
| | | Translation | ArcZ | | | (86) |
| | | Degradation | RssB state | | | (92) |
| Low pH | ND | Degradation? | ND | Loss of viability at low pH | <i>hdeA, cfa, dps, gadC</i> | (11, 101, 155) |
| Constricted amino acid flux | ND | ND | ND | Acidification | ND | (152) |
| High pH | ND | ND | ND | Loss of viability at high pH | <i>dps</i> | (101) |
| High osmolarity | ND | Translation | RprA | Loss of viability at high osmolarity | <i>otsAB</i> | (55, 76, 81, 99, 134) |
| | ND | Degradation | ND | | | (99) |
| Oxidative stress | ND | ND | ND | H ₂ O ₂ sensitive | <i>xthA; dps</i> | (2, 35, 122) |
| N starvation | ND | Activity | Sigma competition? | ND | ND | (65, 85, 107) |

^a Genes included here have been shown, in experiments testing deletions or inactivation, to contribute to resistance to the listed stress.

^b ND, not determined. For some stresses, there is no specific evidence of RpoS induction, although resistance to the stress is seen, dependent on RpoS (i.e., oxidative stress).