patients with severe asthma is associated with a decline in pulmonary function (7).

Epithelial repair requires cell migration, proliferation, and differentiation to restore the structure after injury (8). As progenitor/stem cells, basal cells play a critical role in epithelial repair and undergo proliferation and differentiation in response to injury. Because they showed a persisting expansion after BT, we evaluated their proliferation. We observed that the number of proliferative (Ki-67⁺) p63⁺ cells decreased ≥ 12 months after BT, whereas the total number of p63⁺ cells remained high. These data suggest that the local proliferation of basal cells participated in the observed p63⁺ cell expansion shortly after BT, and that later other sources of progenitor cells likely sustained the observed higher number of p63⁺ cells (9).

This real-life study did not have a control group, and patients were asked to undergo bronchial biopsies at the time of the BT procedures. However, we believe that bronchial biopsies taken in the left lower lobe at the first bronchoscopy for right lower lobe BT treatment constitute an acceptable baseline.

In conclusion, in severe asthma, the bronchial epithelium could be considered as an important target of BT treatment. BT causes epithelial injury, induces proliferation of basal progenitor cells, reduces mucin expression, and improves epithelial integrity. These results support BT as an effective treatment option for patients with severe uncontrolled asthma, and suggest that the observed clinical benefits could be related to BT-induced changes of the bronchial epithelium structure in addition to airway mucosa smooth muscle mass and nerve fiber changes. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Acknowledgment: The authors thank the patients for their participation, Serge Simard for statistical analysis, Sophie Plante for technical help, Lyne Ringuette for patient recruitment, and Sabrina Biardel (Quebec Respiratory Health Network Tissue Bank) for providing tissue samples.

Ikhlass Haj Salem, Ph.D.*
Université Laval
Québec City, Québec, Canada

Delphine Gras, Ph.D.*
Université d'Aix-Marseille
Marseille, France
and
Assistance Publique Hôpitaux de Marseille
Marseille, France

Philippe Joubert, M.D., Ph.D.
Louis-Philippe Boulet, M.D.
Noel Lampron, M.D.
Simon Martel, M.D.
Krystelle Godbout, M.D.
Université Laval
Québec City, Québec, Canada

Pascal Chanez, M.D.
Université d'Aix-Marseille
Marseille, France
and
Assistance Publique Hôpitaux de Marseille
Marseille, France

Michel Laviolette, M.D.
Jamila Chakir, Ph.D.†
Université Laval
Québec City, Québec, Canada

*These authors contributed equally to this work and are joint first authors.
†Corresponding author (e-mail: jamila.chakir@med.ulaval.ca).

References

- Wechsler ME, Laviolette M, Rubin AS, Fiterman J, Lapa e Silva JR, Shah PL, *et al.*; Asthma Intervention Research 2 Trial Study Group. Bronchial thermoplasty: long-term safety and effectiveness in patients with severe persistent asthma. *J Allergy Clin Immunol* 2013;132:1295–1302.
- Salem IH, Boulet LP, Biardel S, Lampron N, Martel S, Laviolette M, *et al.* Long-term effects of bronchial thermoplasty on airway smooth muscle and reticular basement membrane thickness in severe asthma. *Ann Am Thorac Soc* 2016;13:1426–1428.
- Chakir J, Haj-Salem I, Gras D, Joubert P, Beaudoin EL, Biardel S, *et al.* Effects of bronchial thermoplasty on airway smooth muscle and collagen deposition in asthma. *Ann Am Thorac Soc* 2015;12:1612–1618.
- Pretolani M, Bergqvist A, Thabut G, Dombret MC, Knapp D, Hamidi F, *et al.* Effectiveness of bronchial thermoplasty in patients with severe refractory asthma: clinical and histopathologic correlations. *J Allergy Clin Immunol* 2017;139:1176–1185.
- Crosby LM, Waters CM. Epithelial repair mechanisms in the lung. *Am J Physiol Lung Cell Mol Physiol* 2010;298:L715–L731.
- Kanoh S, Tanabe T, Rubin BK. IL-13-induced MUC5AC production and goblet cell differentiation is steroid resistant in human airway cells. *Clin Exp Allergy* 2011;41:1747–1756.
- Loxham M, Davies DE, Blume C. Epithelial function and dysfunction in asthma. *Clin Exp Allergy* 2014;44:1299–1313.
- Brightling CE, Gupta S, Gonen S, Siddiqui S. Lung damage and airway remodelling in severe asthma. *Clin Exp Allergy* 2012;42:638–649.
- Nickerson DW, Presson AP, Weigt SS, Gregson AL, Belperio JA, Gomperts BN. Quantification of cytokeratin 5 mRNA expression in the circulation of healthy human subjects and after lung transplantation. *PLoS One* 2009;4:e5925.
- LeBlanc A, Robichaud P, Lacasse Y, Boulet LP. Quantification of asthma control: validation of the asthma control scoring system. *Allergy* 2007;62:120–125.

Copyright © 2019 by the American Thoracic Society

High Prevalence of Laryngeal Obstruction during Exercise in Severe Asthma

To the Editor:

Exertional dyspnea is a key and pervasive feature in severe asthma and often persists despite optimization of asthma treatment (1). Laryngeal or vocal cord dysfunction, an inappropriate and excessive closure of the glottis during the breathing cycle, is recognized to be present in up to half of patients with severe asthma (2, 3). This laryngeal closure can lead to transient airflow obstruction during exercise and

Supported by Asthma UK and the National Institute for Health Research (J.H.H.).

Author Contributions: J.H.H. and A.M.-G. conceived and designed this study; J.H.H. collected data; all authors provided analysis of results and preparation of manuscript.

Originally Published in Press as DOI: 10.1164/rccm.201809-1734LE on December 20, 2018

exertional dyspnea (4). Although the normal response of the laryngeal structures is to abduct during exercise, in some individuals, the glottic aperture can narrow inappropriately during exercise. It is currently unclear whether laryngeal closure is present and contributes to the etiology of exertional dyspnea in patients with severe asthma, yet such a finding may explain the “treatment refractory” nature of this symptom and have implications for management.

The aim of this study was to evaluate exercise-related laryngeal function in a cohort of patients with severe asthma and exertional dyspnea. We used the technique of continuous laryngoscopy during exercise and hypothesized that laryngeal obstruction would be prevalent and relate to lung function, dyspnea, and exercise intolerance.

Methodology

Study population. Adults with severe asthma and persistent exertional dyspnea, affecting activities of daily living, were recruited between 2011 and 2013 to complete a continuous laryngoscopy during exercise test, spirometry, and questionnaires of both asthma control (asthma control test) (5) and dyspnea (D12) (6). Participants had completed a severe asthma systematic assessment, and those with a contraindication to exercise testing and/or laryngoscopy and current or significant ex-smokers (>5 -pack-year history) were excluded. The study was approved by a research ethics committee (11/LO/1404–NRES–London East).

Study procedures. Continuous laryngoscopy during exercise testing was performed (7) with a high-definition fiberoptic nasendoscope (Olympus ENF-VQ) placed superior to the laryngeal aperture and fixed in headgear to ensure stable image acquisition during a maximal cycle-ergometer cardiopulmonary exercise test, with dyspnea characterized with the Borg scale (0–10). Images of laryngeal movement at rest and near-maximal exercise were captured and closure was assessed, using a closure score (8), by an experienced operator (V.B.) blinded to subject characteristics. The predominant location (i.e., glottic vs. supraglottic level) and breathing phase were recorded, and only a closure score ≥ 2 (i.e., at least moderate severity) was classified as abnormal. Laryngopharyngeal reflux (LPR) was scored by a laryngologist (G.S.) and defined by a reflux finding score >7 (9).

Statistical analysis. Data are expressed as mean (SD) for parametric variables and as median (interquartile range) otherwise. Differences in continuous variables were assessed using one-way ANOVA or a nonparametric equivalent (Kruskal-Wallis), and proportions assessed with chi-squared testing. All statistical calculations were made with SPSS v.21 (SPSS Inc.), and a P value < 0.05 was considered significant.

Results

Subject characteristics and pulmonary function. Thirty-two patients, aged 18–61 years (78% female), completed assessment (Table 1). Patients reported an asthma diagnosis for a median of 36 (range, 3–52) years; 75% were atopic and 66% had a blood eosinophil count $\geq 0.3 \times 10^9/L$. All patients were prescribed treatment $\geq$ GINA step 4 treatment, with 23 (75%) patients prescribed continuous or near-continuous ($\geq 50\%$ of the year) oral corticosteroids. Patients exhibited poor asthma control, with a median asthma control test score of 12 (range, 6–23).

Exercise and laryngoscopy data. Exercise was limited by dyspnea in 64% of subjects with peak exercise capacity reduced to lower limits of normal (Table 1). Exercise-induced bronchoconstriction (i.e., $>10\%$ reduction in FEV_1) was present in 16% of subjects.

There was no evidence of any structural laryngeal defect; however, LPR was present in the majority (88%). During exercise, excessive laryngeal closure was apparent in 15 (47%) patients, in the expiratory phase in six patients (40%), inspiratory phase in seven patients (47%), and in both phases in two patients (13%; Figure 1). Expiratory-phase closure was seen to be exclusively at the glottic level ($P = 0.001$), but inspiratory-phase closure occurred predominantly at the supraglottic ($P = 0.04$) level (Figure 2).

Patients with abnormal laryngeal movement were similar for age, body mass index, sex, atopic status, LPR score, and presence of eosinophilia to those without (all $P > 0.05$; Table 1). Patients with inspiratory phase closure had better lung function and peak oxygen uptake ($P < 0.05$) than those with expiratory closure, and yet were more likely to report that breathing limited their peak exercise capacity ($P < 0.01$; Table 1).

Discussion

Excessive laryngeal closure during exercise was apparent in approximately half the patients with severe asthma, and was associated with dyspnea and impaired pulmonary function. This laryngeal closure was heterogeneous: apparent at different phases of the respiratory cycle and at differing anatomical locations.

Several studies have highlighted an increased prevalence of excessive laryngeal movement in patients with severe asthma (2, 3). Low and colleagues (2) reported that approximately half of all patients with difficult-to-treat asthma had evidence of excessive laryngeal closure on computed tomography imaging. Laryngeal closure has most commonly been classified in the resting state with or without supine positioning, and to date, the functional implications of this excessive laryngeal movement have not been clearly described.

The heightened airflow mandated by exercise presents a direct challenge to the pulmonary system. Laryngeal closure likely acts to modulate intrathoracic pressure; however, excessive closure, especially during inspiration, may actually be maladaptive and increase work of breathing (4). The variable pattern of laryngeal closure we describe has been reported previously, with expiratory closure at the glottic level associated with poorer lung function. We reported a similar finding in chronic obstructive pulmonary disease, where expiratory-phase closure appeared to modulate lung emptying (10); that is, it may be adaptive. In the current study, a high proportion of patients had evidence of LPR. Gastrointestinal reflux is prevalent in asthma, and thus may be acting to sensitize the glottis. Alternatively, these changes may simply reflect the local mechanical consequence of excessive movement.

In the current study, the diagnosis of laryngeal dysfunction was based on endoscopic confirmation of abnormal or excessive laryngeal movement. We found that it was not possible to discern the exercise laryngeal response pattern from bedside clinical assessment, or from the questionnaires administered (i.e., the D12 or Nijmegen Questionnaire). It remains to be determined, however, whether more specific scoring laryngeal-focused systems (e.g., the Pittsburgh Vocal Cord Dysfunction Index [11]) may be diagnostically valuable in this setting. We also found that the resting and/or postexercise flow-volume loop did not permit diagnosis of the presence of in-exercise laryngeal closure. This finding is in keeping

Table 1. Subject Characteristics, Pulmonary Function, and Exercise Data

	All (N = 32)	Normal Laryngeal Function (n = 17)	Abnormal Laryngeal Function		
			All (n = 15)	Expiratory Only (n = 6)	Inspiratory Only (n = 7)
Age, yr	44 (12)	42 (13)	46 (12)	47 (14)	42 (15)
Sex, M/F, %	22/78	18/82	27/73	14/86	50/50
BMI, kg/m ²	29 (25–33)	33 (9–17)	28 (24–31)	28 (25–30)	29 (24–35)
Characteristics					
ACT, x/25	12 (9–16)	12 (9–17)	12 (7–15)	13 (9–25)	9 (6–15)
D12, x/36	18 (9–23)	13 (7–26)	19 (16–23)	17 (10–25)	20 (19–23)
HADS	7 (4–10)	5 (3–9)	8 (5–10)	7 (5–9)	9 (4–10)
Oral steroid use, %	50	41	60	67	57
Serum eosinophils	0.4 (0.1–0.7)	0.3 (0.2–0.6)	0.6 (0.1–0.8)	0.4 (0.0–0.9)	0.7 (0.1–0.8)
Pulmonary function					
FEV ₁ , L	2.1 (0.7)	2.2 (0.6)	1.9 (0.7)	1.7 (0.7)	2.2 (0.8)
FEV ₁ , % predicted	67.7 (18.0)	72.4 (15.0)	61.8 (20.7)	51.2 (15.0)*	73.8 (18.2)
FVC, L	3.0 (0.8)	3.3 (0.7)	3.0 (0.8)	3.1 (1.1)	3.1 (0.8)
FVC, % predicted	86.0 (12.8)	91.4 (10)	80.3 (13.9) [†]	71.8 (12.7)*	89.5 (12.3)
FER (FEV ₁ /FVC)	0.6 (0.1)	0.6 (0.1)	0.6 (0.2)	0.6 (0.1)*	0.7 (0.1)
PEF/PIF	1.3 (0.4)	1.3 (0.3)	1.3 (0.4)	1.3 (0.5)	1.5 (0.5)
Exercise data					
Reason for exercise cessation					
Legs, %	18.8	6.7	29.4	29.4	6.7
Breathing, %	62.5	86.7	35.3 [‡]	35.3 [§]	86.7
Other, %	6.3	0	17.6	17.6	0
Missing, %	12.5	6.7	17.6	17.6	6.7
Peak exercise BORG, x/10	6 (2)	5 (2)	6 (2) [†]	7 (3)	6 (2)
Peak oxygen uptake, L/min	1.6 (1.3–1.7)	1.7 (1.4–1.9)	1.3 (1.3–1.3)	1.4 (0.9–1.7)	1.6 (1.3–1.7)
Peak oxygen uptake, % predicted	81.5 (72.3–94.0)	87.0 (74.5–101.5)	80.0 (64.0–87.0)	68.5 (56.3–74.5)*	85 (80.0–93.0)

Definition of abbreviations: ACT = Asthma Control Test; BMI = body mass index; D12 = dyspnea 12 score (x-36); HADS = Hospital Anxiety and Depression Score; FER = forced expiratory ratio; PEF = peak expiratory flow; PIF = peak inspiratory flow.

Reason for stopping, lack of clear cause/missing data: n = 4. Reason for stopping "other" represents headache, dizziness and chest pain. Parametric data are presented as mean (SD), and nonparametric data are presented as median (interquartile range).

*P < 0.05, expiratory versus inspiratory.

[†]P < 0.05, abnormal versus normal laryngeal movement.

[‡]P < 0.01, abnormal versus normal laryngeal movement.

[§]P < 0.01, expiratory versus inspiratory.

Figure 1. Outcome and classification of continuous laryngoscopy during exercise assessment in severe asthma. CLE = continuous laryngoscopy during exercise; EILO = exercise-induced laryngeal obstruction.

Figure 2. Representative case examples of laryngoscopic images obtained during exercise, illustrating (A) expiratory closure pattern with excessive glottic-level closure, (B) normal response, and (C) inspiratory closure pattern with excessive supraglottic closure.

with previously published work (12), indicating the poor precision of perieercise flow-volume loops in the diagnosis of exercise-induced laryngeal obstruction.

In conclusion, exercise laryngeal closure appears to be a prevalent comorbid phenomenon in severe asthma and may contribute to the development of exertional dyspnea and exercise impairment. The continuous laryngoscopy during exercise test offers a robust means of characterizing laryngeal movement, and our findings highlight the need for future work to determine whether targeted laryngeal intervention, particularly for inspiratory phase laryngeal closure, can improve dyspnea and exercise capacity in severe asthma. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

James H. Hull, Ph.D., F.R.C.P.*
Royal Brompton Hospital
London, United Kingdom
and
Imperial College London
London, United Kingdom

Emil S. Walsted, M.D., Ph.D.
Royal Brompton Hospital
London, United Kingdom
and
Copenhagen University Hospital Bispebjerg
Copenhagen, Denmark

Matt J. Pavitt, M.R.C.P.
Royal Brompton Hospital
London, United Kingdom

Andrew Menzies-Gow, F.R.C.P., Ph.D.
Royal Brompton Hospital
London, United Kingdom
and
Imperial College London
London, United Kingdom

Vibeke Backer, M.D., D.M.Sc.
Copenhagen University Hospital Bispebjerg
Copenhagen, Denmark

Guri Sandhu, M.D., F.R.C.S.
Charing Cross Hospital
London, United Kingdom

ORCID IDs: 0000-0003-4697-1526 (J.H.H.); 0000-0002-6640-7175 (E.S.W.).

*Corresponding author (e-mail: j.hull@rbht.nhs.uk).

References

1. Cordova-Rivera L, Gibson PG, Gardiner PA, McDonald VM. A systematic review of associations of physical activity and sedentary time with asthma outcomes. *J Allergy Clin Immunol Pract* 2018;6:1968–1981.
2. Low K, Lau KK, Holmes P, Crossett M, Vallance N, Phyland D, et al. Abnormal vocal cord function in difficult-to-treat asthma. *Am J Respir Crit Care Med* 2011;184:50–56.

3. Tay TR, Radhakrishna N, Hore-Lacy F, Smith C, Hoy R, Dabscheck E, *et al.* Comorbidities in difficult asthma are independent risk factors for frequent exacerbations, poor control and diminished quality of life. *Respirology* 2016;21:1384–1390.
4. Walsted ES, Faisal A, Jolley CJ, Swanton LL, Pavitt MJ, Luo YM, *et al.* Increased respiratory neural drive and work of breathing in exercise-induced laryngeal obstruction. *J Appl Physiol (1985)* 2018;124:356–363.
5. Nathan RA, Sorkness CA, Kosinski M, Schatz M, Li JT, Marcus P, *et al.* Development of the asthma control test: a survey for assessing asthma control. *J Allergy Clin Immunol* 2004;113:59–65.
6. Yorke J, Moosavi SH, Shuldhham C, Jones PW. Quantification of dyspnoea using descriptors: development and initial testing of the Dyspnoea-12. *Thorax* 2010;65:21–26.
7. Heimdahl J-H, Roksund OD, Halvorsen T, Skadberg BT, Olofsson J. Continuous laryngoscopy exercise test: a method for visualizing laryngeal dysfunction during exercise. *Laryngoscope* 2006;116:52–57.
8. Maat RC, Roksund OD, Halvorsen T, Skadberg BT, Olofsson J, Ellingsen TA, *et al.* Audiovisual assessment of exercise-induced laryngeal obstruction: reliability and validity of observations. *Eur Arch Otorhinolaryngol* 2009;266:1929–1936.
9. Belafsky PC, Postma GN, Koufman JA. The validity and reliability of the reflux finding score (RFS). *Laryngoscope* 2001;111:1313–1317.
10. Baz M, Haji GS, Menzies-Gow A, Tanner RJ, Hopkinson NS, Polkey MI, *et al.* Dynamic laryngeal narrowing during exercise: a mechanism for generating intrinsic PEEP in COPD? *Thorax* 2015;70:251–257.
11. Traister RS, Fajt ML, Landsittel D, Petrov AA. A novel scoring system to distinguish vocal cord dysfunction from asthma. *J Allergy Clin Immunol Pract* 2014;2:65–69.
12. Christensen PM, Maltbæk N, Jørgensen IM, Nielsen KG. Can flow-volume loops be used to diagnose exercise induced laryngeal obstructions?: a comparison study examining the accuracy and inter-rater agreement of flow volume loops as a diagnostic tool. *Prim Care Respir J* 2013;22:306–311.

Copyright © 2019 by the American Thoracic Society

Diagnostic Failures in Asthma

To the Editor:

I commend Aaron and colleagues on their outstanding concise review on the contemporary state of asthma diagnosis (1). I agree that asthma is susceptible to both underrecognition and inappropriate diagnostic labels, and I applaud the authors for raising awareness around diagnostic failures. However, I would like to clarify the definitions of underdiagnosis and overdiagnosis, elaborate why precise use of these terms matters, and discuss inherent tensions in creating any universal disease definition.

In its narrowest sense, overdiagnosis refers to situations in which a diagnostic label is applied to a condition unlikely to cause symptoms or harm (2–4). True overdiagnosis occurs almost exclusively in the context of screening and represents an extreme form of lead-time bias. For example, indiscriminate prostate cancer screening in adult men increases the incidence of cancers detected, but many cancers are unlikely to progress to clinically relevant disease (2).

Similarly, when applied to large populations, spirometry may identify asymptomatic patients with reversible airflow obstruction (1). However, a subset of these patients will remain asymptomatic indefinitely, and labeling them as asthmatic represents overdiagnosis, not underdiagnosis. Even if we consider

underreporting of symptoms and lower quality of life for these patients on a population level, we must also acknowledge that some patients may suffer the consequences of overdiagnosis. Thus, clinicians must counterbalance intensive case-finding against unnecessary disease labeling and overtreatment (2).

Aaron and colleagues also describe situations in which another disease (e.g., allergic rhinitis) is incorrectly labeled as asthma. (1) This situation is not overdiagnosis but, rather, diagnostic error, which is a “failure to establish an accurate and timely explanation of the patient’s health problem(s) or communicate that explanation to the patient” (4). By calling out diagnostic error explicitly, we can target additional parts of the diagnostic process around asthma amenable to solutions, such as patient empowerment and access to appropriate diagnostic testing (4).

Aaron and colleagues also argue that the term asthma should not be used to describe patients with sustained clinical remission (1). I am less convinced of the need to call this example overdiagnosis. Fundamentally, if we conceive of diagnostic labeling as a form of meaning-making to guide prognostication, testing, and treatment (5), then continuing to use “asthma” to describe such patients has continued relevance for patients and clinicians alike. The patients’ underlying genotype and environmental susceptibilities greatly increase the probability that they will exhibit the asthma phenotype in the future.

Ultimately, the semantics matter less than the goal: to provide outstanding evidence-based care to our patients. In caring for patients with asthma, we need diagnostic definitions that suit patients, clinicians, scientists, and epidemiologists. Diagnosis is an uncertain science, and binary classifications of patients as asthmatic versus nonasthmatic may have unintended consequences. Complex diagnostic definitions that suit researchers and specialists may not always translate to better recognition of disease at the “front line” of medicine. In its quest for diagnostic excellence, I hope the asthma community can continue to refine diagnostic criteria that are simple to apply at the bedside and provide the right balance of overlabeling and underrecognition. ■

Author disclosures are available with the text of this letter at www.atsjournals.org.

Paul A. Bergl, M.D.*
Medical College of Wisconsin
Milwaukee, Wisconsin

ORCID ID: 0000-0002-9406-2792 (P.A.B.).

*Corresponding author (e-mail: pbergl@mcw.edu).

References

1. Aaron SD, Boulet LP, Reddel HK, Gershon AS. Underdiagnosis and overdiagnosis of asthma. *Am J Respir Crit Care Med* 2018;198:1012–1020.
2. Kale MS, Korenstein D. Overdiagnosis in primary care: framing the problem and finding solutions. *BMJ* 2018;362:k2820.
3. Carter SM, Rogers W, Heath I, Degeling C, Doust J, Barratt A. The challenge of overdiagnosis begins with its definition. *BMJ* 2015;350:h869.
4. National Academies of Sciences, Engineering, and Medicine. Improving diagnosis in health care. Washington, DC: The National Academies Press; 2015.