Research Digest

Magnesium sulfate is effective for severe acute asthma treated in the emergency department

BH Rowe
JA Bretzlaff
C Bourdon
GW Bota
CA Camargo, Jr

Cochrane Library. Issue
1. Oxford (England):
Update software, 1999.

Correspondence to:
Dr Rowe

Division of Emergency
Medicine, University of
Alberta, 1G1 Walter
Mackenzie Centre, 8440
112th St, Edmonton,
Alberta T6G 2B7,
Canada

Funding: Canadian
Association of
Emergency Physicians
and National Institutes
of Health, Bethesda,
MD

This paper was originally
published in *ACP
Journal Club*
1999;131:36

QUESTION

What is the effectiveness of intravenous magnesium sulfate in patients with acute asthma who were treated in the emergency department?

DATA SOURCES

Studies were identified from the Cochrane Airways Review Group Register, MEDLINE, EMBASE/Excerpta Medica, CINAHL (Cumulative Index to Nursing and Allied Health), and the *Cochrane Library*. We also manually searched 20 high-quality respiratory journals and bibliographies of relevant articles and contacted authors and content experts.

STUDY SELECTION

Studies were selected if they were randomized controlled trials or quasi-randomized trials, included adults or children seen at an emergency department with acute asthma, and compared the use of intravenous magnesium sulfate with placebo.

DATA EXTRACTION

Data on study and patient characteristics, hospital admissions, pulmonary function, vital signs, and adverse events were extracted.

MAIN RESULTS

We reviewed 27 studies for inclusion, and 7 trials (665 patients; 5 trials of adult and 2 of pediatric patients, with 6 from the United States and 1 from India) met the inclusion criteria. Analyses of all patients and of patients with severe asthma showed a reduction in hospital admission rates (table), whereas no difference was shown for patients with mild to moderate asthma. Studies that included all patients showed no differences in measures of

pulmonary function (peak expiratory flow rate or forced expiratory volume in 1 second) or vital signs (heart rate, respiratory rate, or blood pressure). In studies of patients with severe asthma, the peak expiratory flow rate improved by 52 L/min (95% confidence interval 27-78 L/min; 3 studies) and the forced expiratory volume in 1 second by 8% of the predicted value (95% confidence interval 5%-12%; 3 studies). Data were insufficient to assess adverse events.

CONCLUSION

The use of intravenous magnesium sulfate reduces the rate of hospital admissions and improves pulmonary function in patients with severe acute asthma treated in the emergency department.

COMMENTARY

J Mark FitzGerald, Vancouver General Hospital, Vancouver, BC, Canada

The systematic review by Rowe and colleagues is an important addition to the management of acute asthma. Optimal therapy currently consists of bronchodilation with short-acting β_2 agonists, adjunct therapy with ipratropium bromide (in patients with more severe disease), oxygen, and corticosteroids.¹ The results of this review indicate that magnesium sulfate use reduces hospital admission in patients with a forced expiratory volume in 1 second of less than 30% at presentation or those who do not respond to initial treatment or improve beyond a value of 60% after 1 hour.

It is tempting to extrapolate from these data that all patients with moderately severe acute asthma should receive magnesium sulfate. An important consideration, however, is the discretionary use of adjunct therapies, such as aminophylline (2 studies) and ipratropium bromide (1 study), included in the meta-analysis. Both of these interventions have been shown to reduce hospital admission rates without improving lung function.²⁻⁴ Until patients with less severe obstruction treated with these therapies show greater benefit with the incremental addition of magnesium sulfate than with placebo, magnesium sulfate should be reserved for patients who fulfill the criteria given above.

- 1 FitzGerald JM, Grunfeld A. Status asthmaticus. In: Lichtenstein LM, Fauci AS, eds. *Current therapy in allergy, immunology, and rheumatology*. 5th ed. Philadelphia (Pa): Mosby; 1996.
- 2 Wrenn K, Slovis CM, Murphy F, et al. Aminophylline therapy for acute bronchospastic disease in the emergency room. *Ann Intern Med* 1991;115:241-247.
- 3 Lanes SF, Garrett JE, Wentworth CE III, et al. The effect of adding ipratropium bromide to salbutamol in the treatment of acute asthma: a pooled analysis of three trials. *Chest* 1998;114:365-372.
- 4 Qureshi F, Pestian J, Davis P, et al. Effect of nebulized ipratropium on the hospitalization rates of children with asthma. *N Engl J Med* 1998;339:1030-1035.

*Magnesium sulfate vs placebo for hospital admissions among patients with acute asthma at end of study**

Patients	Weighted event rates		Relative risk reduction (95% CI)	Number needed to treat (CI)
	Magnesium sulfate	Placebo		
All	23%	45%	30% (2 to 49)	5 (3 to 63)
Severe asthma	51%	91%	44% (29 to 56)	3 (2 to 4)

*Relative risk reduction, 95% confidence interval, and number needed to treat calculated from data in article