pneumonia and allergic bronchopulmonary aspergillosis: in particular Churg–Strauss manifests, in up to 75% of patients, with mononeuritis multiplex.⁷ At twenty-four months our patient remained free from neurological symptoms, and had no evidence of cardiac, skin, joint or renal involvement. Hypereosinophilic syndrome may represent a preleukaemic state, or may herald the onset of Hodgkin's disease or T-cell lymphoblastic lymphoma.⁸ An underlying lymphoproliferative or myeloproliferative disorder is unlikely in the present case in view of the normal bone marrow, rapid response to hydrocortisone and lack of recurrence of the peripheral eosinophilia. Mild eosinophilia associated with adrenocortical insufficiency is well recognized in critically ill patients. However, the pulmonary infiltrates, mediastinal lymphadenopathy and marked eosinophilia seen in our patient cannot be explained by hypocorticism. Furthermore, the strong IgE response suggests a hypersensitivity phenomenon such as a drug reaction.

Acknowledgments We thank Dr Jeremy Berger for his assistance with diagnostic radiology, Dr Andrew Gallimore for providing histopathology images and Dr Adrian Barnes for his helpful comments.

REFERENCES

- 1 Lombardi C, Passalacqua G. Eosinophilia and diseases: clinical revision of 1862 cases. *Arch Intern Med* 2003;**163**:1371–3
- 2 Schatz PL, Mesologites D, Hyun J, Smith GK, Lahiri B. Captopril-induced hypersensitivity lung disease. An immune-complex-mediated phenomenon. *Chest* 1989;**95**:685–7
- 3 Benzaquen-Forner H, Dournovo P, Tandjaori-Lambiotte H, *et al.* Hypoxic eosinophilic pneumonia in two patients treated with ACE inhibitors. *Rev Mal Respir* 1998;**15**:804–10
- 4 Camus PH, Foucher P, Bonniaud PH, Ask K. Drug-induced infiltrative lung disease. *Eur Respir J* 2001;**18**(suppl. 32):93–1005
- 5 Terzano C, Petroianni A. Clarithromycin and pulmonary infiltration with eosinophilia. *BMJ* 2003;**326**:1377–8
- 6 White A, Woodmansee DP. Adrenal insufficiency from inhaled corticosteroids. *Ann Intern Med* 2004;**140**:E497
- 7 Conron M, Beynon HLC. Churg–Strauss syndrome. *Thorax* 2000;**55**:870–7
- 8 Weller PF. The idiopathic hypereosinophilic syndrome. *Blood* 1994;**83**:2759–79

Renal bone disease

Jeffrey H M Lee MRCP Malcolm Stodell FRCP
J C Fowler FRCP

J R Soc Med 2005;**98**:165–166

In rare instances of renal failure a clue to the underlying disease can be had from plain skeletal radiographs.

CASE HISTORY

A man of 65 was referred after five months of lethargy and poor appetite, with loss of 12 kg in weight. He had a history of hypertension and glaucoma and for several years had been taking quinapril and atenolol and using latanoprost eye drops. On clinical examination the only noteworthy finding was pallor. Haemoglobin was 8.7 g/dL and MCV 72.8 fL, erythrocyte sedimentation rate 93 mm/h, creatinine 205 µmol/L (having been 117 µmol/L 2 years previously).

On the chest X-ray the mediastinal contour was normal and the lungs were clear but unusual symmetrical sclerotic changes were noted in the humeral metaphyses (Figure 1). Bone scintigraphy revealed symmetrically increased uptake predominantly in the long bones, affecting the metaphyses and adjacent diaphyses, with sparing of the epiphyses especially within the distal femora and proximal humeri (Figure 2). On plain films a coarsening of bone texture correlated with the scintigraphic uptake. This strongly suggested a diagnosis of Erdheim–Chester disease, and CT of the abdomen reinforced this impression, revealing stranding of retroperitoneal soft tissues, especially in the perinephric fat, causing mild hydronephrosis (Figure 3). Because this combination of findings was so typical we did not seek biopsy confirmation. He was started on prednisolone and nine months later, taking 5 mg daily, he was feeling well, with serum creatinine 167 µmol/L and creatinine clearance 29 mL/min.

COMMENT

Erdheim–Chester disease is a histiocytosis characterized by diffuse infiltration of the affected organs by lipid-laden histiocytes and Touton-type giant cells, immuno-histochemically different from Langerhans-cell histiocytosis.¹ Radiologically, the most specific finding is bilateral and symmetrical cortical osteosclerosis affecting mainly the metaphyseal and diaphyseal regions of the long bones with sparing of the epiphyses.²

Departments of ¹Medicine and ²Radiology, Luton and Dunstable NHS Trust, Lewsey Road, Luton LU4 0DZ, UK

Correspondence to: Dr Malcolm Stodell
E-mail: malcolm.stodell@ldh-tr.anglox.nhs.uk

Figure 1 Detail from chest radiograph showing upper humerus

Fewer than a hundred cases of Erdheim–Chester disease have been reported.³ Average age at presentation is around 53 years but the youngest patient was 7. The condition can affect not only the bones and retroperitoneum, as in this case, but also the hypothalamic/pituitary axis (leading to diabetes insipidus), the pericardium, the lungs and the skin (xanthomas).^{3–5} The most common presenting complaint is bone pain along with general symptoms such as fever,

weight loss and weakness.⁴ Hydronephrosis is due to infiltration of retroperitoneal fat.⁷

The prognosis depends largely on the extent and distribution of extraosseous disease. Systemic corticosteroids, chemotherapy and radiation treatment have all been used but formal trials are lacking.

REFERENCES

- 1 Pertuiset E, Laredo JD, Liote F, Wassef M, Jagueux M, Kuntz D. Erdheim–Chester disease: report of a case, review of the literature, and discussion of relationships with Langerhans cell histiocytosis. *Rev Rhum Engl Ed* 1993;**60**:504–11
- 2 Breuil V, Brocq O, Pellegrino C, Grimaud A, Euler-Ziegler L. Erdheim–Chester disease: typical radiological bone features for a rare xanthogranulomatosis. *Ann Rheum Dis* 2002;**61**:199–200
- 3 Veyssier-Belot C, Cacoub P, Caparros-Lefebvre D, *et al.* Erdheim–Chester: clinical and radiological characteristics of 59 cases. *Medicine (Baltimore)* 1996;**75**:157–69
- 4 Eble J, Rosenberg A, Young R. Retroperitoneal xanthogranuloma in a patient with Erdheim–Chester disease. *Am J Surg Pathol* 1994;**18**:843–8
- 5 Johnson MD, Aulino JP, Jagasia M, Mawn LA. Erdheim–Chester disease mimicking multiple meningiomas syndrome. *Am J Neuroradiol* 2004;**25**:134–7
- 6 Shamburek RD, Brewer HB, Gochuico BR. Erdheim–Chester disease: a rare multisystem histiocytic disorder associated with interstitial lung disease. *Am J Med Sci* 2001;**321**:66–75
- 7 André M, Delevaux I, de Fraissinette B, *et al.* Two enlarged kidneys: a manifestation of Erdheim–Chester disease. *Am J Nephrol* 2001;**21**:315–17

Fibrinous peritonitis in a patient with rheumatoid arthritis

R Kapoor BMSci BMBS¹ E Dean MBBS¹
T Palferman MB FRCP² Z Khan MRCP³

J R Soc Med 2005;**98**:166–167

Ascites in rheumatoid arthritis has been linked to drug-induced liver damage but not previously to peritoneal disease.

CASE HISTORY

A man of 76, diagnosed 15 years ago with seropositive rheumatoid arthritis, had for 5 years been taking methotrexate as a disease-modifying agent (currently 7.5 mg per week). When he sought advice because of

Figure 2 Scintigrams

Departments of ¹General Medicine, ²Rheumatology and ³Gastroenterology, Yeovil District Hospital, Higher Kingston, Yeovil BA21 4AT, UK

Correspondence to: Dr R Kapoor

E-mail: rajkapoor@doctors.org.uk