Figure 1 Peritoneal biopsy specimen

shortness of breath and abdominal distension the methotrexate was immediately stopped. On examination he had gross ascites. Constrictive pericarditis secondary to rheumatoid arthritis was excluded by cardiac MRI. Diagnostic peritoneal tap revealed an exudate and so the possibility of peritoneal disease was further investigated by laparoscopy, at which peritoneal and liver biopsies were obtained. The liver biopsy was normal but the peritoneal biopsy showed a fibrinous peritonitis with a mild chronic inflammatory infiltrate (Figure 1) similar to that seen in fibrinous pericarditis associated with rheumatoid arthritis.

Repeated drainage of the ascites was necessary after the patient's discharge from hospital, but after institution of prednisolone 15 mg daily there was no recurrence. The methotrexate was not reintroduced because the rheumatoid arthritis was well controlled.

COMMENT

An extensive search of the published work has yielded two reported cases of ascites related to rheumatoid arthritis.^{1,2} In these instances the aetiology was judged to be methotrexate-induced liver damage and the ascites resolved on withdrawal of the drug. The only other documented cause of fibrinous peritonitis is practolol, an extinct beta-blocker that our patient had never received.³⁻⁶ The fibrinous peritonitis in this patient was histologically very reminiscent of the constrictive pericarditis seen in rheumatological arthritis.

Acknowledgments We thank Mrs Andrea Bradshaw for the digital imaging and photography and Dr J Sheffield for his guidance.

REFERENCES

- McRorie ER, Wright RA, Errington ML, Luqmani RA. Rheumatoid constrictive pericarditis. *Br Rheumatol* 1997;**36**:100-3
- Clegg DO, Furst DE, Tolman KG, Pogue R. Acute, reversible hepatic failure associated with methotrexate treatment of rheumatoid arthritis. *J Rheumatol* 1989;**16**:1123-6

- Kujala GA, Shamma'a JM, Chang WL, Brick JE. Hepatitis with bridging fibrosis and reversible hepatic insufficiency in a female with rheumatoid arthritis taking methotrexate. *Arthritis Rheum* 1990;**33**:1037-41
- Gurry JF, Cunningham IG, Brooke BN. Beta-blockers and fibrinous peritonitis. *BMJ* 1975;**ii**:498
- Thompson RP, Jackson BT. Beta-blockers and fibrinous peritonitis. *BMJ* 1975;**ii**:747
- Allan D, Cade D. Delayed fibrinous peritonitis after practolol treatment. *BMJ* 1975;**iv**:40
- Windsor WO, Durrein F, Dyer NH. Fibrinous peritonitis: a complication of practolol therapy. *BMJ* 1975;**ii**:68

Two cases of hypothalamic-pituitary sarcoidosis

D Smith MD MRCPi M J Cullen FRCPi

J R Soc Med 2005;**98**:167-169

Sarcoidosis is a multisystem granulomatous disorder that can affect the hypothalamus and pituitary glands.¹ Three decades ago the mean survival in hypothalamic-pituitary sarcoidosis, in a review of 19 cases, was 45 months.² We report on two patients still alive after diagnosis 20 years ago.

CASE HISTORIES

Case 1

A man of 34 sought advice in 1981 after 10 months of nausea, weight loss, depression and reduced libido. Severe depression was diagnosed and he was admitted to a psychiatric hospital. On examination, he was noted to have a nasal erythematous rash, sausage shaped fingers, reduced secondary sexual hair and small testes. Plasma sodium was 112 mmol/L. Anterior pituitary hormone stimulation tests (Table 1) and a water deprivation test demonstrated panhypopituitarism. A biopsy specimen of the nasal rash was reported as showing granulomatous inflammation consistent with lupus pernio, and in radiographs of the hands there were translucent cystic lesions likewise consistent with sarcoidosis. On CT the pituitary gland appeared normal. A Kveim test was positive. Cerebrospinal fluid had a raised protein (0.48 g/L) and white cell count (12/ μ L, all lymphocytes) but angiotensin converting enzyme (ACE) activity was normal. A whole-body gallium scan was normal.

Department of Endocrinology, St James's Hospital, James's Street, Dublin 2, Ireland

Correspondence to: Diarmuid Smith, Department of Endocrinology, St Vincent's Hospital, Elm Park, Dublin 4, Ireland

E-mail: d.smith@st-vincents.ie

Table 1 Results of insulin tolerance test, GnRH and TRH stimulation tests

Time (min)	Glucose (mmol/L)	Cortisol (nmol/L)	GH (mU/L)	IGF-1 (μ g/L)	TSH (mU/L)	T4 (mU/L)	FSH (U/L)	LH (U/L)	Testosterone (nmol/L)
0	4.9	158	0.1	84	<0.1	58.4	0.5	<0.1	4.6
20	2.4	180					0.5	<0.1	
30	1.8	210	0.1		<0.1				
45	3.8	240	0.2						
60	4.4	230	0.4		<0.1		0.8	<0.1	
90	4.8	267	0.1						
120	5.6	258	0.4						

After treatment with prednisolone 60 mg daily and pituitary hormone replacement his symptoms and depression resolved. After 6 months the prednisolone dose was reduced by 5 mg every 2–3 months until stabilized at 7.5 mg daily. In 1995, however, he developed acute ataxia. MRI of the brain showed hydrocephalus and basal meningeal enhancement of the hemisphere and pituitary fossa consistent with leptomeningitis (Figure 1). The prednisolone was increased to 60 mg per day and within three months his ataxia had completely resolved. In 2004, 23 years after diagnosis, he was still well, on a regimen of full pituitary hormone replacement and prednisolone 10 mg daily.

Figure 1 MRI of the brain showing hydrocephalus

Case 2

A 59-year-old male farmer was seen in 1988 with a 6-month history of polyuria, polydipsia, dry mouth and weight loss. An oral glucose tolerance test was normal but dynamic pituitary testing revealed anterior and posterior pituitary dysfunction. Bilateral hilar lymphadenopathy was seen on chest radiography and transbronchial biopsy of the nodes showed non-necrotizing granulomas consistent with sarcoidosis. A Kveim test was positive. As in case 1 the cerebrospinal fluid protein (0.51 g/L) and white cell count (34/ μ L) were raised but ACE was normal. CT of the pituitary showed an infiltrate within the gland. A whole-body gallium scan was normal. He was started on pituitary hormone replacement, together with prednisolone 60 mg daily, tapered to 7.5 mg daily. When last seen, 14 years after diagnosis, he was fit and active, still working on the farm.

COMMENT

The nervous system is affected in 5–16% of patients with sarcoidosis, and presentations range from peripheral or cranial neuropathy to granulomatous meningitis and, as in our two cases, hypothalamic-pituitary dysfunction. Any or all of these can occur without evidence of pulmonary or other systemic features of sarcoidosis.³

Hypothalamic-pituitary sarcoidosis, which occurs in less than 10% of patients with neurosarcoidosis,¹ continues to present challenges both in diagnosis and in treatment. Originally the neuroendocrine dysfunction was attributed to a destructive process affecting the pituitary, but pituitary responsiveness to synthetic hypothalamic releasing factors indicates that hypothalamic insufficiency is the major cause for the hypopituitarism.⁴ The infiltrative process often involves a patchy destruction of the hypothalamic-pituitary tract, resulting in various combinations of anterior and posterior pituitary hormone dysfunction and sometimes isolated dysfunction. Diagnosis is difficult because of the

variety of resultant symptoms. Contrast enhanced MRI is the imaging procedure of choice for diagnosis and follow-up. Gallium scanning and ACE measurement in serum and cerebrospinal fluid may be helpful but lack sensitivity and specificity. The same is true of increased protein and mild pleocytosis (usually lymphocytosis) in the cerebrospinal fluid. Definitive diagnosis of sarcoidosis requires biopsy evidence of non-caseating necrotizing granulomas.⁵

Hypothalamic-pituitary sarcoidosis is treated with corticosteroids together with anterior and posterior pituitary hormone replacement. Corticosteroids lower the CD4/CD8 lymphocyte ratio, decrease interleukin-2 production, and inhibit collagen synthesis, all of which are abnormal at the sites of active disease within the brain. When the dose of prednisolone is lowered to 20–25 mg daily the symptoms commonly recur. There are few data on other immunomodulatory therapies, but ciclosporin, radiotherapy, azathioprine, chlorambucil and cyclophosphamide have all been used with disappointing results.¹ Surgical intervention is indicated in cases of hydrocephalus or when an expanding mass lesion causes an increase in intracranial pressure.

Clearly, treatment of hypothalamic-pituitary sarcoidosis has made great advances since 1971, when the median survival was only 45 months.² The patients reported here were two of only three diagnosed in our unit over that period. The third, a 26-year-old man, was seen in 1986 with panhypopituitarism. On replacement therapy he remains well 19 years later. Unfortunately the condition can still cause sudden unexpected death.⁶

REFERENCES

- 1 Sharma OP. Neurosarcoidosis. *Chest* 1997;**112**:220–8
- 2 Heffernan A, Cullen M, Towers R, O'Donovan DK. Sarcoidosis of the hypothalamus: a case report with a review of the literature. *Hormones* 1971;**2**:1–12
- 3 Lawton FG, Shalet SM, Beardwell CG, Daws RA. Hypothalamic-pituitary disease as the sole manifestation of sarcoidosis. *Postgrad Med J* 1982;**58**:771–2
- 4 Stuart C, Neelson F, Lebovitz E. Hypothalamic insufficiency: the cause of hypopituitarism in sarcoidosis. *Ann Intern Med* 1978;**88**:589–94
- 5 Zajicek JP, Scolding NJ, Foster O, *et al.* Central nervous system sarcoidosis—diagnosis and management. *Quart J Med* 1999;**92**:103–1177
- 6 Gleckman AM, Patalas ED, Joseph JT. Sudden unexpected death resulting from hypothalamic sarcoidosis. *Am J Forensic Med Pathol* 2002;**23**:48–51

Giant colonic diverticulum

Nishath Altaf MRCSEd¹ Sue Geary FRCR²
Irfan Ahmed FCPS FRCS¹

J R Soc Med 2005;**98**:169–170

A large air cyst in the peritoneal cavity can present diagnostic and therapeutic dilemmas.

CASE HISTORY

A man aged 82 was seen after a week of gradually increasing dull left upper quadrant abdominal pain, worsened by movement and associated with abdominal distension and constipation. There was no history of nausea, vomiting, change in bowel habit or weight loss. The medical history included rheumatoid arthritis and peptic ulcer disease. On examination he had a non-tender central abdominal mass, resonant on percussion. A plain abdominal X-ray showed a large central air-filled cyst, and air was seen throughout the colon and rectum. On contrast-enhanced CT (Figure 1) the cyst was 15 cm in diameter and did not seem to communicate with the bowel (no entry of contrast material taken orally).

At laparotomy the cyst proved to be adherent to the bladder and sigmoid colon. It contained only gas—no urine, fluid or faeces—and no communication with the bowel wall or bladder wall was seen. The large bowel was free from diverticular disease but a few diverticula were noted along the mesenteric border of the jejunum.

The cyst was excised. Postoperatively, the obstructive symptoms completely resolved but the patient later developed pneumonia and died. On histological examination, faecal and food material was identified in granulation tissue within the lining of the cyst wall. In addition there was a small amount of colonic muscularis mucosa. The lesion was therefore categorized as a type I giant colonic diverticulum (Box 1).

COMMENT

Preoperative diagnosis of a giant colonic diverticulum can be difficult even with the aid of contrast-enhanced CT and barium enemas. Most patients are in their 60s or over, and the presentation can be acute or chronic. In some the

¹Professorial Unit of Surgery, Queen's Medical Centre, Nottingham NG7 2UH;

²Department of Academic Radiology, University of Nottingham, Nottingham NG7 2UH, UK

Correspondence to: Mr Nishath Altaf, Research Fellow, Department of Surgery, Queen's Medical Centre, Nottingham NG7 2UH, UK

E-mail: altafnishath@yahoo.co.uk