Embryo emergent: elucidating the cell biology of development

The Santa Cruz Conference on Developmental Biology 2000

The meeting was organized by Cori Bargmann and Richard Harland and held at The University of California, Santa Cruz on July 21–25, 2000.

Introduction

With the rapid rise in genomic information, increasing knowledge about the intricacies of cell signaling pathways, and a greater understanding of conserved gene functions, the field of developmental biology has reached a new level, where researchers face a different set of challenges. Some of these challenges and the exciting approaches being taken to surmount them were described at the Santa Cruz Conference in Developmental Biology. In this review, we highlight methodological trends and new areas of focus that emerged from the meeting.

Converging pathways, molecular complexity

At the time of the previous Santa Cruz conference in 1996, considerable progress had been made in unraveling signaling and transcriptional regulatory pathways that underlie cell fate choices. A recurring theme of the 2000 meeting was that linear pathways are an oversimplification, and that future research must take into account the many feedback loops and cross-signaling networks that link regulatory pathways throughout the cell. The complexity of cross-talk between pathways was ably demonstrated in the yeast by Ira Herskowitz (San Francisco, CA), who described the interplay between the cellular response to

high osmolarity, the pheromone response and pseudohyphal growth. These seemingly distinct pathways are mediated by partially overlapping MAP kinase cascades that negatively regulate one another. For example, during the high osmolarity response, the HOG1 MAP kinase is both an activator of this pathway and an inhibitor of FUS1, a target in the pheromone pathway (O'Rourke and Herskowitz, 1998). Yeast researchers are now exploiting DNA microarray technology to assay genome-wide responses to extrinsic signals, to identify pathway components and to define epistatic relationships among key regulators. The hope is that the technology will become adaptable to more complex systems such as developing *Xenopus*, where evidence is also accumulating for signaling cross-talk in pattern formation of the early embryo. Janet Heasman (Cincinnati, OH) reported that *chordin* expression in the *Xenopus* organizer requires both the VegT/mesoderm-inducing pathway and the β -catenin/dorsalizing pathway. At what level the pathways converge to specify dorsal mesoderm and promote the differentiation of mesodermal cell types is still not resolved.

The ability of one signaling pathway to inhibit another may also determine how neurons respond to chemoattractant or repulsive cues. Marc Tessier-Lavigne (San Francisco, CA) described commissural axons that are initially attracted by netrin to grow towards the neural tube midline but become repelled by it once they cross. As in other systems, this response appears to be mediated in part by the midline repellent Slit, although in mammals Semaphorins appear to contribute as well (Zou *et al.*, 2000). Activation of the Slit receptor Robo causes it to bind and inactivate the netrin receptor DCC, effectively causing navigating axons to turn a deaf ear to the attractive call of netrins.

Another revolution in the field of signal transduction comes from the realization that the 'lock and key' analogy for receptor/signal interactions no longer suffices to describe what actually happens *in vivo*. Norbert Perrimon (Boston, MA) discussed how enzymes involved in glycosaminoglycan biosynthesis can affect different signaling pathways in surprisingly different ways (Perrimon and Bernfield, 2000). In the FGF pathway the UDP-glucose dehydrogenase encoded by *sugarless* and the sulfotransferase encoded by *sulfateless* are required to stabilize the receptor/signal interaction (co-receptor role), whereas in the Hedgehog (Hh) pathway, the heparan sulfate co-polymerase encoded by *tout velu* is needed to facilitate the movement of

cholesterol modified Hh between cells. The distribution of extracellular Wingless is also under complex control, which as shown by Amy Bejsovec (Durham, NC) involves feedback from APC, a negative regulator of Wingless signaling (McCartney *et al.*, 1999). By trapping extracellular Wingless protein on cell surfaces, the proteoglycans Dally and Dally-like also affect signal delivery (Perrimon). Other ligands, such as TGF- β family members, form complexes with proteins that modulate their activity. Work described by Michael Oelgeschlager from the DeRobertis laboratory (Los Angeles, CA) suggests that *Xenopus* Twisted gastrulation is an agonist that dislodges latent BMPs in BMP-chordin complexes to activate BMP signaling (Oelgeschlager *et al.*, 2000). Ali Hemmati-Brivanlou (New York, NY) reported on another TGF- β binding protein, LTBP (latent TGF- β binding protein). Originally described as a TGF- β -masking protein, LTBP can synergize with Activin to promote mesoderm induction and thereby act as a positive regulator.

Modification of receptors is yet another strategy to modulate signaling. Vladimir Panin from Tom Vogt's group (Princeton, NJ) reported that Fringe, a known modulator of Notch activity, catalyses the addition of *N*-acetylglucosamine to Notch's extracellular domain (Moloney *et al.*, 2000). As a result, the relative affinity of Notch for its ligands and/or the proteolytic events that occur downstream of the ligand/receptor interaction could be affected. Trudi Schupbach (Princeton, NJ) showed that fine tuning of signaling can also be achieved through receptor turnover (Pai *et al.*, 2000). During dorsal-ventral patterning of the *Drosophila* embryo, EGF-Receptor (EGFR) activity must be turned off ventrally to ensure specification of the most ventral cell types. This downregulation involves Cbl, a tyrosine kinase binding protein shown in other systems to promote degradation of EGFR by recruiting ubiquitin-activating and -conjugating enzymes. Clearly, how ligands and receptors are presented, selectively modified, stored and degraded have become paramount issues in the field.

Nucleocentric no more!

A significant trend is that developmental biologists' interests are shifting away from gene regulation toward the cellular machinery powering developmental change. As pointed out by speakers working in invertebrate and vertebrate model systems, and articulated by Mike Levine (Berkeley, CA), 'the challenge is to fit complex transcriptional regulatory networks to the nuts and bolts of cellular morphogenesis.' Toward this goal, Levine undertook a molecular screen to identify genes controlling cell movements that shape the *Ciona intestinalis* notochord. Bill Smith (Santa Barbara, CA) is using a mutagenesis screen cleverly adapted to another ascidian species to recover mutations that disrupt notochord extension (Moody *et al.*, 1999). Richard Harland (Berkeley, CA) discussed related cell movements in the frog embryo. Labeling membranes with green fluorescent protein (GFP) revealed that cells undergo polarized behaviors to perform the movements of convergence and extension. This polarization depends on *Xenopus* Dishevelled, implicating a pathway equivalent to the *Drosophila* planar cell polarity signaling cascade in vertebrate gastrulation (Wallingford *et al.*, 2000).

Several other groups reported on the utility of GFP to characterize the behavior of subcellular structures during development. For example, using a tubulin-GFP fusion, Allan Spradling (Baltimore, MD) described how the fusome, an asymmetric membrane-rich organelle essential for oogenesis in *Drosophila*, influences the polarization of the microtubule network in early germ cells (Grieder *et al.*, 2000). Complementary experiments are also revealing a role for the fusome in organizing mitochondrial inheritance during oogenesis. Adapting similar visual strategies to mouse embryology is the goal of Scott Fraser (Pasadena, CA), who is devising several non-invasive imaging techniques suitable for vertebrate systems.

Departure from a nucleocentric view was further evidenced by the closer focus on activities occurring at the edges of cells. Chris Doe (Eugene, OR) working in *Drosophila* and Michel Labouesse (Illkirch, France) in *Caenorhabditis elegans* described a group of cortically-enriched proteins (Lethal Giant Larvae, Disc Large and Scribble) that regulate asymmetric protein localization in cells with distinct apical/basal polarity (Legouis *et al.*, 2000). Mutations in Lethal giant larvae can be suppressed by reducing the level of myosin II, revealing a negative interaction between these cortical proteins and the actomyosin network (Peng *et al.*, 2000). Eric Wieschaus (Princeton, NJ) reported on another protein on the edge, Nullo, which is required for the formation of specialized basal junctions during cellularization of the *Drosophila* blastoderm. Nullo must be degraded before gastrulation, as its prolonged expression blocks the formation of mature apical adherens junctions between cells (Hunter and Wieschaus, 2000). Thus, developmental biologists are characterizing molecules involved in basic cellular processes as they probe how cells modify their shape and contacts with neighbors in the context of the growing embryo.

Genetics gets green and goes genomic

Genetic screens are still the mainstay for developmental analyses, as mutations continue to implicate unanticipated proteins and biochemical processes. Arnon Rosenthal (South San Francisco, CA) reported that the zebrafish *foggy* locus, first identified because mutants produce extra serotonergic neurons and fewer dopaminergic neurons in the hypothalamus, encodes a highly conserved regulator of transcriptional elongation. How exactly this protein works to determine classes of neurons in the brain is unclear, although the mutation was found to disrupt a repressive activity within the transcriptional elongation machinery. Didier Stainier (San Francisco, CA) showed that the zebrafish mutation, *miles apart*, which produces cardia bifida due to migratory defects of the bilateral heart primordia, encodes a receptor for the lipid sphingosine-1-phosphate (Kupperman *et al.*, 2000). Whether lipid-mediated signals are required for cell polarity or for migration of the myocardial epithelium remains to be determined. Another heart mutant, *heart and soul*, is defective for an atypical protein kinase that was previously implicated in the formation of tight junctions in mammalian cells and in asymmetric cell division in *C. elegans*. New players acting in well-studied pathways are also being discovered. Using a forward genetic approach in the mouse (Anderson, 2000), Kathryn Anderson (New York, NY) identified

several mutations that interfere with Hedgehog signaling, two of which likely define new modifiers of the Hh pathway.

While the power of the classical genetic approach remains undisputed, the criteria for mutagenesis screens are changing. In particular, an increased understanding of the underlying cell biology is allowing investigators to design focused screens that target the mechanism of interest more directly. Genetic dissection of intricate cellular processes in *Drosophila*, such as the formation and branching of tracheal tubes (Mark Krasnow, Stanford, CA) or of neuromuscular synapses (Graeme Davis, San Francisco, CA) is expected to bring us closer to the basis of lung branching structure and synaptic physiology in humans. As is the case in studies of subcellular organization, GFP labeling of proteins is proving to be an important breakthrough for discovering subtle mutant phenotypes at a high resolution. For instance, a synaptobrevin–GFP fusion enabled Yishi Jin (Santa Cruz, CA) to paint nerve endings green and recover *C. elegans* mutants with abnormal synapses (Zhen *et al.*, 2000).

Innovative tools are also being developed for systematic genome-wide analysis of gene function. Proof of the old adage ‘you are what you eat’ comes from *C. elegans* laboratories capitalizing on the ‘feeding’ method devised by Lisa Timmons and Andy Fire (Timmons *et al.*, 2000). Specific gene function is obliterated after worms dine on bacteria engineered to produce the appropriate double-stranded RNA (RNAi or RNA-mediated interference). Julie Ahringer (Cambridge, UK) is creating a veritable feeding frenzy in efforts to survey the loss-of-function phenotypes of all ~19 000 ORFs predicted by the *C. elegans* sequencing project. To gain more insight into the RNAi phenomenon itself, Craig Mello (Worcester, MA) has isolated mutants resistant to ingested dsRNA. Intriguingly, some of the same mutants remain sensitive to dsRNA introduced by injection.

Bill Skarnes (Berkeley, CA) reported that the ‘secretory trap’ insertional mutagenesis scheme, designed to target all membrane or secreted proteins in the mouse, has paid off with the recovery of a number of interesting loci. One encodes a potential new Wnt receptor that is structurally related to the receptor for low-density lipoprotein (Pinson *et al.*, 2000). Technology for single gene inactivation has also increased in sophistication, with the combined use of Flp and Cre recombinases to create hypomorphic or tissue-specific null alleles. Such chromosomal constructs allow the functional dissection of signals that are involved in multiple processes, such as FGF8 (Gail Martin, San Francisco, CA), or that have complicated spatial and temporal modes of action in the early embryo, such as Nodal (Elizabeth Robertson, Boston, MA).

Development diversifies

Another advantage of the new molecular genetic tools is that developmental biologists are no longer confined to the realm of the embryo but are reaching out to tackle some of the ‘big questions’ in biology. Issues such as speciation and the basis for morphological change, sexual orientation, neuronal discrimination, synaptic plasticity and the regulation of life stages are but a few being actively explored. In some cases, this may require a bold move away from the usual model systems, as illustrated by David Kingsley (Stanford, CA), studying the remarkable diversity of the bony elements of stickleback fishes. Isolated fish popula-

tions are well-suited for the study of the genetic and environmental forces underlying morphological variation, which, ideally, will reveal general principles that have been at work in vertebrate skeletal evolution. With the framework of a stickleback genetic map in hand, the identification of polymorphisms linked to a wide array of physical and behavioral traits holds promise for future gene identification.

For other uncharted areas, however, model systems continue to deliver results and an entry to related processes in other species. Genetic analysis of the internal clock that controls the timing of developmental stages has progressed in *C. elegans*. Gary Ruvkun (Boston, MA) described recent work on *let-7*, a small RNA that regulates the transition from larva to adult, in part, by inhibiting translation of proteins that promote larval cell fates (Reinhart *et al.*, 2000). The discovery of temporally regulated fly and human *let-7* orthologues leads to the tantalizing idea that conserved mechanisms may very well coordinate the timing of developmental stages as diverse as larval instars and human puberty. Comparative approaches have also begun to pay off in the sex determination field, where genes orthologous to the regulators of male development *mab-3* (*C. elegans*) and *double sex* (*Drosophila*) are turning up in birds, turtles and mice. David Zarkower (Minneapolis, MN) indicated that *mab-3* is required in *C. elegans* for males to remain attracted to the opposite sex (Yi *et al.*, 2000), suggesting that insights into the regulation of sexual behavior may soon be an added bonus. Control of life span, another big question well-served by model systems, was discussed by Cynthia Kenyon (San Francisco, CA), who proposed that the integration of sensory inputs from the environment and internal inputs from the germline modulates the rate of aging in *C. elegans* (Hsin and Kenyon, 1999; Apfeld and Kenyon, 1999). Finally, Cori Bargmann (San Francisco, CA) addressed the topic of neuronal diversification in olfaction. She showed how a novel signaling pathway akin to lateral inhibition (but involving calcium signaling rather than Notch) leads to the differential expression of odorant receptors (Troemel *et al.*, 1999), potentially allowing bilaterally symmetric neurons to smell differently.

Conclusion

A strong message from the year 2000 Santa Cruz meeting is that, while developmental biologists are still entranced by the exquisite pattern of the embryo, new directions are emerging for attacking both smaller and bigger issues. In the millennial crystal ball, one foresees that by the next Santa Cruz meeting the divide between the cell biologist and the embryologist will dissolve, as we gain a greater appreciation of the roles of subcellular trafficking, cyto-architectural remodeling and organelle function in shaping developmental events. As anticipated by Eric Wieschaus (Princeton, NJ), ‘if we could understand what genes do in a cell biological context, we will come a long way in understanding development.’ To fulfill the mandate, directed genetic screens will take on even higher levels of sophistication with tools for visualizing cellular detail, as well as for monitoring tissue interactions and cell physiology *in situ*, leading to surprising discoveries about organ form and function. Finally, inroads will be made into biological problems previously thought to be too complex or even intractable, from the growing

interchange between genetic/genomic strategies and molecular lessons first learned from the developing embryo.

References

- Anderson, K.V. (2000) Finding the genes that direct mammalian development ENU mutagenesis in the mouse. *Trends Genet.*, **16**, 99–102.
- Apfeld, J. and Kenyon, C. (1999) Regulation of lifespan by sensory perception in *Caenorhabditis elegans*. *Nature*, **402**, 804–809.
- Grieder, N.C., de Cuevas, M. and Spradling, A.C. (2000). The fusome organizes the microtubule network during oocyte differentiation in *Drosophila*. *Development* **127**, 4253–4264.
- Hsin, H. and Kenyon, C. (1999) Signals from the reproductive system regulate the lifespan of *C. elegans*. *Nature*, **399**, 362–366.
- Hunter, C. and Wieschaus, E. (2000) Regulated expression of *nullo* is required for the formation of distinct apical and basal adherens junctions in the *Drosophila* blastoderm. *J. Cell Biol.*, **150**, 391–401.
- Kupferman, E., An, S., Osborne, N., Waldron, S. and Stainier D.Y. (2000) A sphingosine-1-phosphate receptor regulates cell migration during vertebrate heart development. *Nature*, **406**, 192–195.
- Legouis, R., Gansmuller, A., Sookhareea, S., Boshier, J.M., Baillie, D.L. and Labouesse M. (2000) LET-413 is a basolateral protein required for the assembly of adherens junctions in *Caenorhabditis elegans*. *Nature Cell Biol.*, **2**, 415–422.
- McCartney, B.M., Dierick, H.A., Kirkpatrick, C., Moline, M.M., Baas, A., Peifer, M. and Bejsovec A. (1999) *Drosophila* APC2 is a cytoskeletally-associated protein that regulates wingless signaling in the embryonic epidermis. *J. Cell Biol.*, **146**, 1303–1318.
- Moloney, D.J. *et al.* (2000) Fringe is a glycosyltransferase that modifies Notch. *Nature*, **406**, 369–375.
- Moody, R., Davis, S.W., Cubas, F. and Smith, W.C. (1999) Isolation of developmental mutants of the ascidian *Ciona savignyi*. *Mol. Gen. Genet.*, **262**, 199–206.
- Oelgeschlager, M., Larraín, J., Geissert, D. and De Robertis, E.M. (2000) The evolutionarily conserved BMP-binding protein Twisted gastrulation promotes BMP signalling. *Nature*, **405**, 757–763.
- O'Rourke, S.M. and Herskowitz, I. (1998) The Hog1 MAPK prevents cross talk between the HOG and pheromone response MAPK pathways in *Saccharomyces cerevisiae*. *Genes Dev.*, **12**, 2874–2886.
- Pai, L.-M., Barcelo, G. and Schüpbach, T. (2000) D-cbl, a negative regulator of the Egfr pathway, is required for dorsoventral patterning in *Drosophila* oogenesis. *Cell*, **103**, 51–61.
- Peng, C.-Y., Albertson, R., Manning, L. and Doe, C.Q. (2000). The tumor suppressor genes *lgl* and *dlg* regulate basal protein targeting in *Drosophila* neuroblasts. *Nature*, in press.
- Perrimon, N. and Bernfield, M. (2000) Specificities of heparan sulphate proteoglycans in developmental processes. *Nature*, **404**, 725–728.
- Pinson, K.I., Brennan, J., Monkley, S., Avery, B.J. and Skarnes, W.C. (2000) An LDL-receptor-related protein mediates Wnt signalling in mice. *Nature*, **407**, 535–538.
- Reinhart, B.J., Slack, F.J., Basson, M., Pasquinelli, A.E., Bettinger, J.C., Rougvie, A.E., Horvitz, H.R. and Ruvkun, G. (2000) The 21-nucleotide let-7 RNA regulates developmental timing in *Caenorhabditis elegans*. *Nature*, **403**, 901–906.
- Timmons, L., Court, D.L., II and Fire, A. (2000) Ingestion of bacterially expressed dsRNAs can produce specific and potent genetic interference in *Caenorhabditis elegans*. *Gene*, in press.

- Troemel, E.R., Sagasti, A. and Bargmann, C.I. (1999) Lateral signaling mediated by axon contact and calcium entry regulates asymmetric odorant receptor expression in *C. elegans*. *Cell*, **99**, 387–398.
- Wallingford, J.B., Rowning, B.A., Vogeli, K.M., Rothbacher, U., Fraser, S.E. and Harland, R.M. (2000) Dishevelled controls cell polarity during *Xenopus* gastrulation. *Nature*, **405**, 81–85.
- Yi, W., Ross, J.M. and Zarkower, D. (2000) *mab-3* is a direct *tra-1* target gene regulating diverse aspects of *C. elegans* male sexual development and behavior. *Development*, **127**, 4469–4480.
- Zhen, M., Huang, X., Bamber, B. and Jin, Y. (2000) Regulation of presynaptic terminal organization by *C. elegans* RPM-1, a putative guanine nucleotide exchanger with a RING-H2 finger domain. *Neuron*, **26**, 331–343.
- Zou, Y., Stoeckli, E., Chen, H. and Tessier-Lavigne, M. (2000) Squeezing axons out of the gray matter: a role for Slit and Semaphorin proteins from

The authors* and organizers⁺ from left to right are: Geraldine Seydoux*, Richard Harland⁺, Cori Bargmann⁺ and Marnie E. Halpern*

midline and ventral spinal cord. *Cell*, **102**, 363–375.

Marnie Halpern¹ and Geraldine Seydoux²

¹Carnegie Institution of Washington, Department of Embryology, Baltimore, MD and ²Johns Hopkins School of Medicine, Department of Molecular Biology and Genetics Baltimore, MD, USA

Received October 11, 2000; revised October 24, 2000; accepted October 27, 2000

+Corresponding author. Tel: +1 410 554 1218; Fax: +1 410 243 6311; E-mail: halpern@ciwemb.edu

DOI: 10.1093/embo-reports/kvd115