Htm1p, a mannosidase-like protein, is involved in glycoprotein degradation in yeast

Claude A. Jakob^{1,*}, Daniel Bodmer^{1,‡}, Urs Spirig¹, Patrick Bättig^{1,§}, Anne Marcil², Daniel Dignard², John J.M. Bergeron³, David Y. Thomas^{2,¶} & Markus Aebi¹

¹Institute of Microbiology, ETH Zurich, Schmelzbergstrasse 7, CH-8092 Zurich, Switzerland, ²Genetics Group, Biotechnology Research Institute, National Research Council of Canada, Montreal, PQ, H4P 2R2 and ³Department for Anatomy and Cell Biology, McGill University, Montreal, H3A 2B2, Canada

Received October 25, 2000; revised and accepted March 1, 2001

Misfolded proteins are recognized in the endoplasmic reticulum (ER), transported back to the cytoplasm and degraded by the proteasome. Processing intermediates of N-linked oligosaccharides on incompletely folded glycoproteins have an important role in their folding/refolding, and also in their targeting to proteolytic degradation. In *Saccharomyces cerevisiae*, we have identified a gene coding for a non-essential protein that is homologous to mannosidase I (*HTM1*) and that is required for degradation of glycoproteins. Deletion of the *HTM1* gene does not affect oligosaccharide trimming. However, deletion of *HTM1* does reduce the rate of degradation of the mutant glycoproteins such as carboxypeptidase Y, ABC-transporter Pdr5-26p and oligosaccharyltransferase subunit Stt3-7p, but not of mutant Sec61-2p, a non-glycoprotein. Our results indicate that although Htm1p is not involved in processing of N-linked oligosaccharides, it is required for their proteolytic degradation. We propose that this mannosidase homolog is a lectin that recognizes Man₈GlcNAc₂ oligosaccharides that serve as signals in the degradation pathway.

INTRODUCTION

Most secretory and membrane proteins are co-translationally translocated into the lumen of the endoplasmic reticulum (ER) (Rapoport *et al.*, 1996) where they acquire N-linked oligosaccharides, catalyzed by oligosaccharyltransferase. Also in the ER, secretory and membrane proteins attain their correct three-dimensional structure by the concerted action of chaperones, isomerases and folding mediators (Ellgaard *et al.*, 1999). The

N-linked oligosaccharides are processed by glucosidase I, glucosidase II and mannosidase I (Herscovics, 1999a). It has been shown that oligosaccharide processing and protein folding are linked processes coordinating protein folding, ER retention/retrieval and protein degradation (Ellgaard *et al.*, 1999; Herscovics, 1999b; Jakob and Burda, 1999).

Mechanisms that discern misfolded secretory and membrane proteins have been discovered in the past few years (Bonifacino and Weissman, 1998). Misfolded proteins in the ER lumen are specifically recognized by as yet unknown mechanisms and subjected to a cytosolically located proteolytic process termed ER-associated degradation (ERAD). The targeted proteins are transported back into the cytoplasm by a mechanism that is formed, at least partially, by constituents of the ER import machinery (Römisch, 1999). Many of the retrotranslocated proteins are polyubiquitin modified by the action of the cytosolically located enzymes Ubc6p, Ubc7p and Hrd1/Der3p, followed by degradation by the proteasome (Brodsky and McCracken, 1997; Kopito, 1997; Sommer and Wolf, 1997; Bays *et al.*, 2001; Deak and Wolf, 2001).

The degradation of misfolded mutant carboxypeptidase Y [CPY* (*prc1-1*); Finger *et al.*, 1993] has been shown to be N-glycan dependent (Knop *et al.*, 1996) and, moreover, the structure of this oligosaccharide plays an important role in rapid protein degradation (Jakob *et al.*, 1998). Specifically, glycoproteins carrying Man₈GlcNAc₂ sugars are more rapidly degraded than others. This finding supported the 'mannose timer hypothesis' (Su *et al.*, 1993; Helenius *et al.*, 1997) and has led to the proposal that the ER lumen contains a Man₈GlcNAc₂-binding

*Corresponding author. Tel: +41 1 632 3327; Fax: +41 1 632 1148; E-mail: jakob@micro.biol.ethz.ch

‡Present address: Clinic for Neurosurgery, University Hospital Zurich, CH-8091 Zurich, Switzerland

§Present address: Cytos Biotechnology AG, CH-8952 Schlieren-Zurich, Switzerland

¶Present address: Biochemistry Department, McGill University, Montreal, PQ, H3G 1Y6, Canada
C.A. Jakob and D. Bodmer contributed equally to this work

C.A. Jakob *et al.*

protein that initiates glycoprotein ERAD (Liu *et al.*, 1997; Jakob *et al.*, 1998; Yang *et al.*, 1998).

In the yeast genome we have identified a good candidate for this $\text{Man}_8\text{GlcNAc}_2$ -specific lectin (Htm1p) that is homologous to mannosidase I (Camirand *et al.*, 1991). We show that strains deleted for *HTM1* are defective in the degradation of a mutant soluble protein (CPY*), and also two mutant membrane proteins (Pdr5-26p, Stt3-7p). Our data strongly suggest that the Htm1 protein is the putative $\text{Man}_8\text{GlcNAc}_2$ -binding lectin.

RESULTS AND DISCUSSION

HTM1 codes for a gene homologous to mannosidase and is involved in protein degradation

We searched the *Saccharomyces cerevisiae* genome for open reading frames with homology to ER mannosidase since we hypothesized that a 'degradation lectin' might contain similar domains. The reading frame *YHR204w* (termed gene *homologous to mannosidase*; *HTM1*) was identified, showing a 40% amino acid sequence similarity to yeast mannosidase.

To test whether *HTM1* was involved in protein degradation, we deleted the gene in a yeast strain containing CPY* (Wolf and Fink, 1975) and Pdr5* [mutant Pdr5 protein (*pdr5-26*); Egner *et al.*, 1998], two substrates that have previously been shown to undergo ubiquitin-dependent proteasomal degradation. As controls, we used mutant strains that were deficient in the biosynthesis of complete oligosaccharide chains (Δalg12) or in trimming of the oligosaccharides (Δmns1). These mutations prevent the biosynthesis of $\text{Man}_8\text{GlcNAc}_2$ oligosaccharide or its formation by processing, respectively, and stabilize misfolded CPY* (Jakob *et al.*, 1998). Equal numbers of cells were inoculated and *de novo* protein synthesis was stopped by the addition of cycloheximide; the levels of CPY* and Pdr5* were determined by immunoblotting and quantified at different time points. The CPY* degradation rate in Δalg12 and Δmns1 cells was reduced in comparison to 'wild-type' cells, but also in the *HTM1*-deficient cells (Figure 1). In addition, we also found a higher steady-state level of CPY* in Δalg12 , Δmns1 and Δhtm1 cells as compared with wild-type cells (time point 0; Figure 1A, compare lanes 1, 4, 7 and 10). The mutant membrane protein Pdr5* showed a similar change in its degradation: steady-state levels of Pdr5* were increased in Δalg12 and Δmns1 , and also Δhtm1 cells (Figure 1), and degradation rates were significantly reduced (Figure 1).

To test whether *HTM1* was involved in general protein degradation, we studied the mutant Sec61-2 protein, a membrane protein with no *N*-glycosylation sites. The *sec61-2* mutant cells display a temperature-sensitive growth phenotype. Biederer *et al.* (1996) showed that Sec61-2 protein was degraded at restrictive temperature. Impairing protein degradation by deleting the *UBC6* and *UBC7* genes allowed the cells to survive restrictive growth conditions (Figure 2A). We deleted the *MNS1* and *HTM1* genes in the *sec61-2* mutants and tested whether these two genes influenced degradation of the Sec61-2 protein by cell growth at selective temperature and by western blot analysis. Neither of the two mutant cells (*sec61-2* Δmns1 and *sec61-2* Δhtm1) improved growth on plates at restrictive temperature

Fig. 1. Oligosaccharide-dependent degradation of the mutant proteins CPY* and Pdr5*. (A) Isogenic cells were grown to mid-log phase. Equal cell numbers were then incubated with 100 $\mu\text{g/ml}$ cycloheximide and aliquots removed at given time points. Crude protein extract was separated by SDS-PAGE and analyzed by immunoblotting using anti-CPY (CPY*) and anti-HA (Pdr5*) antisera. (B) Protein amounts from at least two independent experiments were quantified and plotted against time, setting the quantity of time point 0 (steady-state level) to 100%.

(Figure 2A) nor reduced the degradation rate of Sec61-2 protein (Figure 2B).

Fig. 2. The *HTM1* gene does not influence the degradation of mutant Sec61-2 protein. (A) *Δmns1* and *Δhtm1* do not suppress temperature-sensitive growth of *sec61-2* mutant cells. Isogenic *sec61-2* single- and double-mutant cells were plated in serial dilutions on rich medium and grown at 30 and 37°C for 3 days. (B) Degradation of Sec61-2 protein is not affected in *mns1*- and *htm1*-deficient cells. Cells were grown at 30°C to mid-log phase and shifted to 37°C for 2 h. Equal cell numbers were incubated with 100 μg/ml cycloheximide, aliquots were removed and cells broken at given time points. Extracted proteins were separated by SDS-PAGE, and analyzed and quantified by immunoblotting using anti-Sec61 antiserum. The protein quantity of time point 0 (steady-state level) was set to 100%.

These results support the hypothesis that the *HTM1* gene is required for efficient secretory and membrane glycoprotein degradation. Degradation of proteins lacking *N*-glycans was not affected. Glycoprotein degradation might require a specific lectin that recognizes *N*-linked $\text{Man}_8\text{GlcNAc}_2$ oligosaccharides (Jakob *et al.*, 1998). For this we reasoned that such a lectin might be similar to ER mannosidase I. The *HTM1* gene codes for a polypeptide of 796 amino acids with significant sequence similarity to yeast mannosidase I (Figure 3A), and amino acid

sequences similar to Htm1 protein are found in fungal, plant, insect, worm and mammal genomes. Phylogenetic analysis of these sequences revealed two distinct sequence groups, of which one contained all class I $\alpha 1,2$ -mannosidases and the other proteins homologous to Htm1p (data not shown). Many amino acids reported to be involved in oligosaccharide binding, catalysis and Ca^{2+} coordination in Mns1p (Vallee *et al.*, 2000) are conserved in Htm1 protein and its homologs. One of the characteristic features of the Htm1 protein family group was the

C.A. Jakob et al.

Fig. 3. Similarity of Mns1p and Htm1p proteins. **(A)** Areas of homology of Mns1p and Htm1p. Areas of high amino acid similarity are drawn as boxes. **(B)** Htm1 family proteins lack a highly conserved cysteine residue. Amino acid sequences of α -mannosidases and Htm1 family proteins of plant, mammalian, insect, worm and yeast cells were aligned with Clustal_W. The sequences were grouped by sequence similarity. The arrowhead indicates the position of an alanine residue in Htm1 proteins at the position of Cys340 (in *S. cerevisiae*) conserved in and most likely essential for maintaining the correct structure of α 1,2-mannosidases. The first two letters indicate the species: at, *Arabidopsis thaliana*; ce, *Caenorhabditis elegans*; dm, *Drosophila melanogaster*; hs, *Homo sapiens*; sc, *S. cerevisiae*; sp, *Schizosaccharomyces pombe*. The following numbers refer to the enzyme name or DDBJ/EMBL/GenBank accession No.

absence of the conserved cysteine residues 340 and 385 (position given for Mns1p of *S. cerevisiae*), which form a disulfide bond. These two cysteine residues are conserved in α 1,2-mannosidase family members and are most likely to be important for maintaining the correct structure of processing mannosidases (Lipari and Herscovics, 1996) (Figure 3B).

HTM1 is not involved in processing of N-linked oligosaccharides

Processing of N-glycans to create the $\text{Man}_8\text{GlcNAc}_2$ glycan is required for efficient degradation in yeast (Jakob et al., 1998 and see above). We determined whether altered processing of oligosaccharides was the reason for reduced glycoprotein degradation in Δhtm1 cells. For this, oligosaccharides were metabolically labeled *in vivo*. N-linked sugars were extracted and released from the protein by peptide N-glycosidase F (PNGase F) treatment and analyzed by HPLC. Attention was focused on sugars that were not processed by Golgi mannosyltransferases (Figure 4) and, as expected, wild-type cells contained $\text{Man}_8\text{GlcNAc}_2$ oligosaccharides, and Δmns1 cells contained $\text{Man}_9\text{GlcNAc}_2$ (Figure 4). However, oligosaccharides

Fig. 4. Normal N-linked oligosaccharide processing in *htm1*-deficient yeast cells. N-linked oligosaccharides of isogenic strains were metabolically labeled with [^3H]mannose, extracted, released with PNGase F and separated by HPLC.

obtained from cells deleted for *HTM1* showed $\text{Man}_8\text{GlcNAc}_2$ but no $\text{Man}_9\text{GlcNAc}_2$ oligosaccharides. These results confirm that normal processing of N-glycans occurs in Δhtm1 cells and, therefore, the Htm1 protein is not a processing mannosidase. Although Htm1p displays sequence similarity to class I mannosidases, it is not involved in the specific trimming of N-glycans in the ER, which is the only process that requires mannosidase activity in yeast (Dean, 1999; Gemmill and Trimble, 1999; Herscovics, 1999a). This is consistent with our conclusion that Htm1p is not a mannosidase, and with our inference that it may be a lectin.

There have been several reports that indicate a distinct function of N-glycan trimming by mannosidase I in glycoprotein degradation (Liu et al., 1997, 1999; Yang et al., 1998; Cabral et al., 2000; Mancini et al., 2000; Wang and Androlewicz, 2000; Wang and White, 2000). These reports are in good agreement with results found in *S. cerevisiae*, where a specific $\text{Man}_8\text{GlcNAc}_2$ glycan is required for efficient protein degradation in the ER (Knop et al., 1996; Jakob et al., 1998). This specific oligosaccharide on degradation substrates might be recognized by the Htm1 protein and by this facilitate glycoprotein degradation.

Degradation of the Stt3-7 protein is oligosaccharide specific and mediated by the ubiquitin–proteasomal pathway

In yeast, oligosaccharyltransferase (OST), the enzyme complex that transfers oligosaccharides from lipid-linked precursors to

Fig. 5. *UBC6*-dependent degradation of *stt3-7* protein. **(A)** Overexpression or deletion of the *UBC6* gene suppressed the temperature-sensitive growth phenotype of *stt3-7* mutant cells. Isogenic *stt3-7* single- and double-mutant cells were plated in serial dilutions on rich medium and grown at 23 and 30°C for 3 days. **(B)** Overexpression or deletion of the *UBC6* gene stabilized the mutated *Stt3-7* protein. Cells were shifted to 37°C and protein was extracted. Steady-state protein levels and CPY glycosylation were determined by western analysis using the indicated antisera. Hexokinase was used as the control for equal protein loading.

nascent polypeptides at N-X-S/T, is composed of nine membrane proteins (Knauer and Lehle, 1999), of which *Stt3p* is the largest subunit and is also essential for the N-linked glycosylation process *in vivo* (Zufferey *et al.*, 1995). Assembly of the OST complex is a multistep process, and integration of either the *Ost3p* or *Ost6p* subunit is dependent upon the presence of *Stt3p* (U. Spirig, D. Bodmer, M. Wacker and M. Aebi, in preparation). The *stt3-7* conditional mutant has been shown to severely impair

Fig. 6. Oligosaccharide-dependent suppression of conditional growth of *stt3-7* by deletion of *MNS1* and *HTM1* genes. Isogenic *stt3-7* single- and double-mutant cells were plated in serial dilutions on rich medium and grown at 23 and 30°C for 3 days.

OST activity, leading to incomplete N-glycosylation of nascent glycoproteins at restrictive temperature (Spirig *et al.*, 1997). We observed that overexpression or deletion of the *UBC6* gene could suppress the temperature-sensitive phenotype of mutant *stt3-7* strains (Figure 5A).

UBC6 encodes the E2 ubiquitin-conjugating enzyme situated on the cytosolic face of the ER membrane and plays a pivotal role in the degradation of misfolded proteins in the ER (Sommer and Wolf, 1997). Overexpression or deletion of the *UBC6* gene increased the steady-state level of the mutant *Stt3-7* protein as well as levels of the OST subunits *Ost3p* and *Ost6p*, as shown by western blot analysis (Figure 5B). N-linked glycosylation was strongly affected in *stt3-7* mutant cells (visualized by the presence of hypoglycosylated CPY) and was only slightly improved by reducing protein degradation using *UBC6* mutants (Figure 5B). Our finding shows that mutant *Stt3-7* protein is degraded by the ubiquitin-proteasome pathway at restrictive temperature. We speculated that reducing protein degradation by deleting the *UBC6* gene increased the steady-state level of *Ost3p* and *Ost6p* as well as mutant *Stt3-7* protein, permitting the integration of *Ost3p*, *Ost6p* and *Stt3-7p* into a partially functional oligosaccharyltransferase complex. This partially restored enzymatic activity then increases N-linked glycosylation to allow mutant cells to survive at restrictive temperature.

Deletion of a gene (*HTM1*) homologous to mannosidases suppresses the *stt3-7* mutation and stabilizes misfolded protein in the ER

UBC6-dependent growth of *stt3-7* strains is a convenient method to monitor the effect of mutant genes in this process. If reduced protein degradation can rescue these mutant cells, we reasoned that we might find similar results upon creating the *stt3-7* $\Delta mns1$ or *stt3-7* $\Delta htm1$ double mutants since *Stt3* protein is N-glycosylated. In validation of this assay, $\Delta mns1$ suppressed

Table I. Yeast strains used

Strain	Genotype	Reference
SS328	<i>MATα ade2-101 ura3-52 his3Δ200 lys2-801</i>	Vijayraghavan <i>et al.</i> (1989)
YG543	<i>MATa ade2-101 ura3-52 his3Δ200 lys2-801 leu2 stt3-7</i>	Spirig <i>et al.</i> (1997)
YG885	<i>MATα ade2-101 ura3-52 his3Δ200 lys2-801 Δubc6::HIS3</i>	this study
YG950	<i>MATα ade2-101 ura3-52 his3Δ200 lys2-801 stt3-7 Δmns1::KanMX</i>	this study
YG894	<i>MATα ade2-101 ura3-52 his3Δ200 lys2-801 leu2 stt3-7 Dubc6::HIS3</i>	this study
YG618	<i>MATα ade2-101 ura3-52 his3Δ200 lys2-801 prc1-1</i>	Jakob <i>et al.</i> (1998)
YG777	<i>MATa ade2-101 ura3-52 his3Δ200 tyr1 Δmns1::KanMX prc1-1</i>	Jakob <i>et al.</i> (1998)
YG807	<i>MATα ade2-101 ura3-52 his3Δ200 lys2-801 Δalg12::KanMX prc1-1</i>	Jakob <i>et al.</i> (1998)
YG1211	<i>MATα ade2-101 ura3-52 his3Δ200 lys2-801 Δmns1::KanMX prc1-1</i>	this study
YCJ1	<i>MATα ade2-101 ura3-52 his3Δ200 lys2-801 Δhtm1::KanMX prc1-1</i>	this study
YCJ82	<i>MATa ade2-101 ura3-52 his3Δ200 lys2-801 stt3-7 Dhtm1::KanMX</i>	this study
YCJ88	<i>MATa ade2-101 ura3-52 his3Δ200 lys2-801 stt3-7 Dhtm1::KanMX</i>	this study
RSY533	<i>MATα sec61-2 ade2 leu2-3 leu2-112 pep4-3 ura3-52</i>	R. Schekman
YG1350	<i>MATα sec61-2 ade2 leu2-3 leu2-112 pep4-3 ura3-52 Δubc6::KanMX</i>	this study
YG1351	<i>MATα sec61-2 ade2 leu2-3 leu2-112 pep4-3 ura3-52 Δubc7::KanMX</i>	this study
YG1352	<i>MATα sec61-2 ade2 leu2-3 leu2-112 pep4-3 ura3-52 Δmns1::KanMX</i>	this study
YG1353	<i>MATα sec61-2 ade2 leu2-3 leu2-112 pep4-3 ura3-52 Δhtm1::KanMX</i>	this study

Table II. Primers for gene deletions

Deletion	Name	Sequence 5' → 3' ^a
<i>HTM1</i>	HR204forKan	tcgatcatgtagcatgcgaagacgatcgctactcattcacttctat cgatgaattcgagctc
	HR204revKan	gccattggaagtgagcacaggactatgtttcttgattgtacac cgtagctgcaggtcgac
	HR204–90u	gcaataaagaaggacgcg
	HR204+350L	ccggtcttcagtatagc
<i>UBC6</i>	UBC6forKan	gatgttggaacccctccaccatattcttgcctcgcccaacgaagat cgatgaattcgagctc
	UBC6revKan	tgaagaactatcattaggttcttgccatcttcttgaattatttct cgtagctgcaggtcgac
	UBC6–180u	actaccatcgcatatcgc
	UBC6+130L	gttatcattctgatagccg
<i>UBC7</i>	UBC7forKan	ctcctcaaggagcttcaacagttaattaagattctccacctggtatata cgatgaattcgagctc
	UBC7revKan	ctcactcaacatgctcataacttaataagaattttctacatttgc cgtagctgcaggtcgac
	UBC7–70u	ggcgtttagcgtagcgaag
	UBC7+130L	agcgtagcggtatctgg
All	KanMXu	gtattgatgttggacgag

^aBold letters represent sequences specific to plasmid pFA6aKanMX4 for the amplification of the KanMX resistance gene (Wach *et al.*, 1994).

the temperature sensitivity of *stt3-7* cells at 30°C, indicating a direct role of *N*-glycan trimming in the degradation of Stt3-7 protein (Figure 6). Moreover, the *stt3-7 Δhtm1* double mutant could grow at elevated temperature, similar to the *stt3-7 Δmns1* strain (Figure 6). This result further supported the interpretation that Htm1 protein plays an important role in efficient degradation of the mutant glycoprotein Stt3-7p.

Our phenotypic analysis of *Δhtm1* mutant strains is compatible with the idea that Htm1 protein is in fact the lectin involved in

this process. Detailed biochemical studies will be required to confirm this hypothesis.

METHODS

Materials. Yeast strains used are detailed in Table I. The strains harboring the *prc1-1* allele were described previously (Jakob *et al.*, 1998). The pRS313 HA-*pdr5-26* plasmid was kindly provided by Dr D.H. Wolf (University of Stuttgart, Germany).

For western blot analysis, antisera raised against CPY, Sec61p, Ost3p, Ost6p (Dr R. Knauer, University of Regensburg, Germany), hexokinase (Dr S. Schröder, Göttingen, Germany) and anti-hemagglutinin (anti-HA) (Santa Cruz, CA) were used.

Yeast manipulations. Standard protocols were followed for yeast manipulation (Guthrie and Fink, 1991). Cells were grown at permissive temperature in either YPD (2% bactopectone, 1% yeast extract, 2% glucose) or, for maintaining plasmids, in minimal medium (0.67% yeast nitrogen base, 2% glucose, with the appropriate supplements to allow growth). Whole-cell protein was extracted as previously described (Jakob *et al.*, 1998). Extracts containing the HA-tagged Pdr5* gene were incubated at 50°C for 15 min instead of boiled. The cycloheximide chase was performed as described by Plemper *et al.* (1998). Serial dilutions of cells for growth assays, starting at 5×10^5 cells, were spotted on YPD plates and incubated at the given temperatures for 3 days.

Construction of yeast strains. The loci *YHR204w/HTM1*, *UBC6* and *UBC7* were deleted by replacing the major part of the gene with the KanMX cassette (Wach *et al.*, 1994) using the primers given in Table II. The correct integration was verified by PCR. Deletion of the *MNS1* gene was described previously (Jakob *et al.*, 1998).

Labeling of N-linked oligosaccharides. The procedures for the metabolic labeling, recovery of glycoproteins and release of N-glycans by treatment with PNGase F (Roche Diagnostics, Rotkreuz, Switzerland) were as described previously (Jakob *et al.*, 1998).

ACKNOWLEDGEMENTS

We are greatly indebted to Dr D.H. Wolf (pRS313-HA-*pdr5-26* plasmid), Dr R. Knauer (anti-Ost6p) and Dr S. Schröder (anti-hexokinase). The authors would like to thank Dr Charles J. Waechter for critically reading the manuscript. This work has been supported by the Swiss National Science Foundation (fellowship 81ZH-054340 to C.A.J.; grants 31-57082.99 to M.A.), and grants from the Canadian Institutes for Health Research (to D.Y.T. and J.J.M.B.).

REFERENCES

- Bays, N.W., Gardner, R.G., Seelig, L.P., Joazeiro, C.A. and Hampton, R.Y. (2001) Hrd1p/Der3p is a membrane-anchored ubiquitin ligase required for ER-associated degradation. *Nature Cell Biol.*, **3**, 24–29.
- Biederer, T., Volkwein, C. and Sommer, T. (1996) Degradation of subunits of the Sec61p complex, an integral component of the ER membrane, by the ubiquitin-proteasome pathway. *EMBO J.*, **15**, 2069–2076.
- Bonifacino, J.S. and Weissman, A.M. (1998) Ubiquitin and the control of protein fate in the secretory and endocytic pathways. *Annu. Rev. Cell Dev. Biol.*, **14**, 19–57.
- Brodsky, J.L. and McCracken, A.A. (1997) ER-associated and proteasome-mediated degradation: how two topologically restricted events came together. *Trends Cell Biol.*, **7**, 151–155.
- Cabral, C.M., Choudhury, P., Liu, Y. and Sifers, R.N. (2000) Processing by endoplasmic reticulum mannosidases partitions a secretion-impaired glycoprotein into distinct disposal pathways. *J. Biol. Chem.*, **275**, 25015–25022.
- Camirand, A., Heysen, A., Grondin, B. and Herscovics, A. (1991) Glycoprotein biosynthesis in *Saccharomyces cerevisiae*. Isolation and characterization of the gene encoding a specific processing α -mannosidase. *J. Biol. Chem.*, **266**, 15120–15127.
- Deak, P.M. and Wolf, D.H. (2001) Membrane topology and function of Der3/Hrd1p as an ubiquitin-protein ligase (E3) involved in endoplasmic reticulum degradation. *J. Biol. Chem.*, **276**, 10663–10669.
- Dean, N. (1999) Asparagine-linked glycosylation in the yeast Golgi. *Biochim. Biophys. Acta*, **1426**, 309–322.
- Egner, R., Rosenthal, F.E., Kralli, A., Sanglard, D. and Kuchler, K. (1998) Genetic separation of FK506 susceptibility and drug transport in the yeast Pdr5 ATP-binding cassette multidrug resistance transporter. *Mol. Biol. Cell*, **9**, 523–543.
- Ellgaard, L., Molinari, M. and Helenius, A. (1999) Setting the standards: quality control in the secretory pathway. *Science*, **286**, 1882–1888.
- Finger, A., Knop, M. and Wolf, D.H. (1993) Analysis of two mutated vacuolar proteins reveals a degradation pathway in the endoplasmic reticulum or a related compartment in yeast. *Eur. J. Biochem.*, **218**, 565–574.
- Gemmill, T.R. and Trimble, R.B. (1999) Overview of N- and O-linked oligosaccharide structures found in various yeast species. *Biochim. Biophys. Acta*, **1426**, 227–237.
- Guthrie, C. and Fink, G.R. (1991) Guide to yeast genetics and molecular biology. *Methods Enzymol.*, **194**, 1–270.
- Helenius, A., Trombetta, E.S., Hebert, D.N. and Simons, J.F. (1997) Calnexin, calreticulin and the folding of glycoproteins. *Trends Cell Biol.*, **7**, 193–200.
- Herscovics, A. (1999a) Processing glycosidases of *Saccharomyces cerevisiae*. *Biochim. Biophys. Acta*, **1426**, 275–285.
- Herscovics, A. (1999b) Importance of glycosidases in mammalian glycoprotein biosynthesis. *Biochim. Biophys. Acta*, **1473**, 96–107.
- Jakob, C.A. and Burda, P. (1999) Quality control in biosynthetic pathways of N-linked glycoproteins in the yeast endoplasmic reticulum. *Protoplasma*, **207**, 1–7.
- Jakob, C.A., Burda, P., Roth, J. and Aeby, M. (1998) Degradation of misfolded endoplasmic reticulum glycoproteins in *Saccharomyces cerevisiae* is determined by a specific oligosaccharide structure. *J. Cell Biol.*, **142**, 1223–1233.
- Knauer, R. and Lehle, L. (1999) The oligosaccharyltransferase complex from yeast. *Biochim. Biophys. Acta*, **1426**, 259–273.
- Knop, M., Hauser, N. and Wolf, D. (1996) N-glycosylation affects ER degradation of a mutated derivative of carboxypeptidase yscY in yeast. *Yeast*, **12**, 1229–1238.
- Kopito, R.R. (1997) ER quality control: the cytoplasmic connection. *Cell*, **88**, 427–430.
- Lipari, F. and Herscovics, A. (1996) Role of the cysteine residues in the α 1,2-mannosidase involved in N-glycan biosynthesis in *Saccharomyces cerevisiae*—the conserved Cys(340) and Cys(385) residues form an essential disulfide bond. *J. Biol. Chem.*, **271**, 27615–27622.
- Liu, Y., Choudhury, P., Cabral, C.M. and Sifers, R.N. (1997) Intracellular disposal of incompletely folded human α 1-antitrypsin involves release from calnexin and post-translational trimming of asparagine-linked oligosaccharides. *J. Biol. Chem.*, **272**, 7946–7951.
- Liu, Y., Choudhury, P., Cabral, C.M. and Sifers, R.N. (1999) Oligosaccharide modification in the early secretory pathway directs the selection of a misfolded glycoprotein for degradation by the proteasome. *J. Biol. Chem.*, **274**, 5861–5867.
- Mancini, R., Fagioli, C., Fra, A.M., Maggioni, C. and Sitia, R. (2000) Degradation of unassembled soluble Ig subunits by cytosolic proteasomes: evidence that retrotranslocation and degradation are coupled events. *FASEB J.*, **14**, 769–778.
- Plemper, R.K., Egner, R., Kuchler, K. and Wolf, D.H. (1998) Endoplasmic reticulum degradation of a mutated ATP-binding cassette transporter Pdr5 proceeds in a concerted action of Sec61 and the proteasome. *J. Biol. Chem.*, **273**, 32848–32856.
- Rapoport, T.A., Jungnickel, B. and Kutay, U. (1996) Protein transport across the eukaryotic endoplasmic reticulum and bacterial inner membranes. *Annu. Rev. Biochem.*, **65**, 271–303.
- Römisch, K. (1999) Surfing the Sec61 channel: bidirectional protein translocation across the ER membrane. *J. Cell Sci.*, **112**, 4185–4191.

C.A. Jakob et al.

- Sommer, T. and Wolf, D.H. (1997) Endoplasmic reticulum degradation: reverse protein flow of no return. *FASEB J.*, **11**, 1227–1233.
- Spirig, U., Glavas, M., Bodmer, D., Reiss, G., Burda, P., Lippuner, V., te Heesen, S. and Aebi, M. (1997) The STT3 protein is a component of the yeast oligosaccharyltransferase complex. *Mol. Gen. Genet.*, **256**, 628–637.
- Su, K., Stoller, T., Rocco, J., Zemsky, J. and Green, R. (1993) Pre-Golgi degradation of yeast prepro- α -factor expressed in a mammalian cell. Influence of cell type-specific oligosaccharide processing on intracellular fate. *J. Biol. Chem.*, **268**, 14301–14309.
- Vallee, F., Lipari, F., Yip, P., Sleno, B., Herscovics, A. and Howell, P.L. (2000) Crystal structure of a class I α 1,2-mannosidase involved in *N*-glycan processing and endoplasmic reticulum quality control. *EMBO J.*, **19**, 581–588.
- Vijayraghavan, U., Company, M. and Abelson, J. (1989) Isolation and characterization of pre-mRNA splicing mutants of *Saccharomyces cerevisiae*. *Genes Dev.*, **3**, 1206–1216.
- Wach, A., Brachat, A., Pöhlmann, R. and Philippsen, P. (1994) New heterologous modules for classical or PCR-based gene disruptions in *Saccharomyces cerevisiae*. *Yeast*, **10**, 1793–1808.
- Wang, Y. and Androlewicz, M.J. (2000) Oligosaccharide trimming plays a role in the endoplasmic reticulum-associated degradation of tyrosinase. *Biochem. Biophys. Res. Commun.*, **271**, 22–27.
- Wang, J. and White, A.L. (2000) Role of calnexin, calreticulin, and endoplasmic reticulum mannosidase I in apolipoprotein(a) intracellular targeting. *Biochemistry*, **39**, 8993–9000.
- Wolf, D.H. and Fink, G.R. (1975) Proteinase C (carboxypeptidase Y) mutant of yeast. *J. Bacteriol.*, **123**, 1150–1156.
- Yang, M., Omura, S., Bonifacino, J.S. and Weissman, A.M. (1998) Novel aspects of degradation of T cell receptor subunits from the endoplasmic reticulum (ER) in T cells: importance of oligosaccharide processing, ubiquitination, and proteasome-dependent removal from ER membranes. *J. Exp. Med.*, **187**, 835–846.
- Zufferey, R., Knauer, R., Burda, P., Stagljär, I., te Heesen, S., Lehle, L. and Aebi, M. (1995) *STT3*, a highly conserved protein required for yeast oligosaccharyltransferase activity *in vitro*. *EMBO J.*, **14**, 4949–4960.

DOI: 10.1093/embo-reports/kve089