Is there a rationale for rationing chronic dialysis?

Two points need clarification

EDITOR—I agree with Chandna et al that late referral is an important factor in the survival of patients receiving dialysis.¹ I am unaware, however, of the source of their information that an NHS consensus statement recommends that nephrology referrals be made at a serum creatinine concentration $>135 \mu\text{mol/l}$ in women and $>180 \mu\text{mol/l}$ in men. The workload generated by this practice would be so enormous that British renal units under present conditions would be unable to cope.

I am also concerned that of the authors' 292 patients receiving dialysis, 26 (admittedly high risk) patients spent 44% of their lives in hospital. The costs of inpatient treatment for these high risk patients would be substantially higher than the average cost of £250 a day that the authors quote.

Rasheed Ahmad *Consultant renal physician*
Royal Liverpool University Hospital, Liverpool
L7 8XP
TheAhmads@aol.com

1 Chandna SM, Schulz J, Lawrence C, Greenwood RN, Farrington K. Is there a rationale for rationing chronic dialysis? A hospital based cohort study of factors affecting survival and morbidity. *BMJ* 1999;318:217-23. (23 January.)

Study's objectives were not achieved

EDITOR—Chandna et al's study asking whether there is a rationale for rationing chronic dialysis does not fulfil its stated purpose.¹ The study successfully shows the groups of patients already receiving dialysis who could be expected to do poorly, and this might be used as a rationale for withdrawing treatment in this small group. The cost savings that their unit would accrue are nevertheless small (just 3.2% of the cost of their chronic dialysis programme).

Even in the authors' high risk group five of the 26 patients were long term survivors. This is similar to the effect of interferon treatment in hepatitis C in terms of long term response and better than results of in vitro fertilisation programmes. One could argue, therefore, that these patients should not be refused treatment either. Quality of life considerations would be important in this group since these five patients may have had an excellent quality of life, as indeed may some of the other patients.

This study cannot be considered a basis for rationing access to renal replacement. The authors do not report the characteristics or reasons for refusal of patients

referred to them for renal replacement. Their assertion that Wiltshire Health Authority's recommendations may be used as a starting point in deciding prioritisation is also not valid. They agree that these recommendations are couched in general terms but do not specify them in their paper. Furthermore, they state that patients were assessed on an individual basis, with no formal criteria being used.

The study shows that the criteria used to decide acceptance on to the authors' renal replacement programme work for the patients who are accepted. What it does not tell us is whether the correct criteria are being used for those refused renal replacement. It would be interesting to know what happened to those refused replacement and on what grounds this decision was made.

T Farrant *Specialist registrar in gastroenterology*
Jersey General Hospital, St Helier, Jersey, Channel Islands
106712.1276@compuserve.com

1 Chandna SM, Schulz J, Lawrence C, Greenwood RN, Farrington K. Is there a rationale for rationing chronic dialysis? A hospital based cohort study of factors affecting survival and morbidity. *BMJ* 1999;318:217-23. (23 January.)

Question is difficult to address on basis of retrospective studies

EDITOR—Chandna et al's paper confirms the influence of comorbidity on survival of patients receiving renal replacement that has been shown in previous retrospective studies.¹ The question of rationing dialysis, however, while pertinent, is difficult to address on the basis of such studies.

Chandna et al costed the potential savings from rationing chronic dialysis on the basis of comorbidity. As no source or methods were given it was not clear whether the costs were from the perspective of the institution, the healthcare system, or society. Nor was it clear whether these costs related to haemodialysis or peritoneal dialysis. To be included in any rationing debate, clinical and cost data must be robust.

An obvious limitation of any retrospective study is that some of the information needed may not have been recorded in the case notes. This makes the assessment of functional data (Karnofsky scores) and comorbidity unreliable and difficult to reproduce. A subjective comorbidity scoring system, completed in many cases after death, is a potential source of bias, and we are currently studying prospectively the influence

of comorbidity and functional status on outcomes of renal replacement in various parts of Europe.

Chandna et al compared their comorbidity severity score with that used by Khan et al in a retrospective survival analysis² and found considerable differences in the proportion of patients in each risk group (for example, high risk group 51% and 23% respectively). Although this may reflect higher comorbidity in Chandna et al's patients, their interpretation of the risk stratification system may have differed. Furthermore, patients receiving a transplant or recovering renal function (18%) were left in Chandna et al's survival analysis whereas Khan et al censored patients at such events. The better prognosis of this subgroup would certainly influence the overall survival rates.

The purpose of risk stratification to study outcomes should be twofold: to enable valid comparison of outcomes between centres, and to identify groups of patients for whom efforts should be directed at improving outcomes rather than to deny such patients treatment. The risk stratification system used by Khan et al and Wright was developed to correct for case mix and allow comparison of survival between centres.^{2,3} Recognising that a patient is at high risk also allows us to be more informed when

Advice to authors

We prefer to receive all responses electronically, sent either directly to our website or to the editorial office as email or on a disk. Processing your letter will be delayed unless it arrives in an electronic form.

We are now posting all direct submissions to our website within 72 hours of receipt and our intention is to post all other electronic submissions there as well. All responses will be eligible for publication in the paper journal.

Responses should be under 400 words and relate to articles published in the preceding month. They should include ≤ 5 references, in the Vancouver style, including one to the BMJ article to which they relate. We welcome illustrations.

Please supply each author's current appointment and full address, and a phone or fax number or email address for the corresponding author. We ask authors to declare any competing interest. Please send a stamped addressed envelope if you would like to know whether your letter has been accepted or rejected.

Letters will be edited and may be shortened.

www.bmj.com
letters@bmj.com

discussing prognosis and outlook with patients receiving dialysis and their families.

Fergus J Caskey *Clinical research fellow*
f.caskey@abdn.ac.uk

Wendy Metcalfe *Clinical research fellow*
Alison M MacLeod *Reader*

Department of Medicine and Therapeutics,
University of Aberdeen, Aberdeen AB25 2ZD

Izhar H Khan *Consultant physician*
Renal Unit, Aberdeen Royal Infirmary, Aberdeen
AB9 2ZB

1 Chandna SM, Schulz J, Lawrence C, Greenwood RN, Farrington K. Is there a rationale for rationing chronic dialysis? A hospital based cohort study of factors affecting survival and morbidity. *BMJ* 1999;318:217-23. (23 January.)

2 Khan IK, Catto GRD, Edward N, Fleming LW, Henderson IS, MacLeod AM. Influence of coexisting disease on survival on renal replacement therapy. *Lancet* 1993;341:415-8.

3 Wright LE. Survival in patients with end-stage renal disease. *Am J Kidney Dis* 1991;17:25-8.

Authors' reply

EDITOR—Ahmad is right that there is no NHS consensus recommending nephrology referrals. This was a typographical error and should have read NIH (National Institutes of Health).¹ Encouraging early referral undoubtedly means more work in nephrology outpatient departments, but this may partially be offset by reduced work and costs owing to fewer unplanned presentations for dialysis; at present these account for 44% of all patients entering the long term dialysis programme.

Ahmad may be right in expecting higher than average inpatient costs for the 26 high risk patients. This does not, however, alter our conclusions, since many of these patients did not live long. In total they had 428 inpatient days during the first three months, and even if these are costed at an average of £350 a day, the figure in our table 3 only changes from 3.2% to 3.5%.

We do not agree with Farrant that our study's objectives were not achieved. We identified several factors affecting survival and hospitalisation in patients receiving dialysis and, using these, defined a small high risk group with a poor one year survival. We suggested that the methods used to define this group might be tested prospectively. We did not advocate using them as a basis for formal rationing, but they might contribute to the understanding of a complex decision making process. Farrant poses the same question that we did: is a one year survival of 19% low enough to justify withholding dialysis? This is a value judgment that a study such as ours cannot make.

We agree with Caskey et al that the question of rationing is difficult to address, but it is important to make a start. The costs we quoted were those of dialysis (both haemodialysis and peritoneal dialysis) and hospitalisation. Clearly, other costs also arise, but these are difficult to quantify retrospectively. We look forward to the results of Caskey et al's prospective work.

The original purpose of risk stratification was to compare outcomes between centres.^{2,3} We hypothesised that a more tightly defined high risk group could be identified in which renal replacement was inappropriate.

ate.⁴ Our analysis did not substantiate this but may have resulted in an improved definition of high risk in this setting. This new definition may indeed be better geared for counselling patients with potentially poor prognosis. Wright's and Khan et al's classification, with a two year survival of 55% in high risk group in our population, is clearly not suited for this purpose.

Shahid M Chandna *Associate specialist*
shahid.chandna@lister.org.uk

Roger N Greenwood *Consultant*

Ken Farrington *Consultant*
Department of Nephrology, Lister Hospital,
Stevenage SG1 4AB

1 National Institutes of Health. Consensus conference statement. Morbidity and mortality of renal dialysis. *Ann Intern Med* 1994;121:62-70.

2 Wright LE. Survival in patients with end-stage renal disease. *Am J Kidney Dis* 1991;17:25-8.

3 Khan IK, Catto GRD, Edward N, Fleming LW, Henderson IS, MacLeod AM. Influence of coexisting disease on survival on renal-replacement therapy. *Lancet* 1993;341:415-8.

4 Farmery E, Milner P. *Renal replacement therapy: purchasing review and recommendations*. Devizes: Wiltshire Health Authority, 1997.

Rationing of sildenafil

Nobody needs an erection at public expense

EDITOR—I am a BMA member, but Drs John Chisholm and Ian Bogle do not speak for me on sildenafil (Viagra).^{1,2} I understood that we had been asking successive governments for years to accept the need for explicit rationing decisions. As soon as this government makes one—a sensible one that the general public accepts—we kick it in the teeth.

The grounds for differentiating between different causes of erectile dysfunction outlined by the health secretary, Frank Dobson, were imperfect but not unreasonable³; for the first time there was an overt admission that totally comprehensive demand-led care had to be limited. This was not the time to pick a fight. The media comment suggests that the public accepts that when resources are limited conditions such as cataract and stroke should be given higher priorities than erectile dysfunction.

I have been telling my patients who request sildenafil that the NHS cannot afford it. I still take that view, but the BMA's reported comments will make my position harder to sustain. Sildenafil should be available at cost without prescription; it will cost about as much as an acceptable bottle of table wine.

Stephen Hayes *General practitioner*
Southampton SO30 2AA
stephen.hayes1@virgin.net

1 Chisholm J. Viagra: a botched case for rationing. *BMJ* 1999;318:273-4. (30 January.)

2 British Medical Association. *GPC Viagra statement*. London: BMA, 1999. (Press release 21 January.)

3 Department of Health. *Viagra: NHS prescription proposals announced*. London: DoH, 1999. (Press release 21 January.)

True debate is needed

EDITOR—The proposals for rationing of sildenafil (Viagra) put forward by the secretary of state are unethical,¹ and the General Practitioners Committee is correct

to advise general practitioners to prescribe the drug on prescription to all patients who require it.²

Many doctors would welcome a mature debate regarding rationing in the NHS, recognising that in an underfunded service where rationing has always existed such a discussion should be held openly and honestly. Few, however, can have expected that the first politically instigated, clearly financial, limitation of treatment would be decided on the basis of aetiology and not clinical need. To implement such a decision would be to break the fundamental principle of medicine that patients should be treated according to need, regardless of the chain of events that placed them in that position of need.

Impotence has many similarities with menopausal hot flushes. Neither are life threatening, both cause distress, and both are common conditions often but not always associated with a particular age and viewed by some as natural. A general practitioner willing to give hormone replacement treatment only to those women whose symptoms were due to an oophorectomy or radiotherapy and not those whose symptoms were the result of natural ovarian failure, and who justified the decision on grounds of cost, would not only be acting unethically but also be in breach of his or her terms and conditions of service.

The proposed loophole whereby patients with impotence caused by other processes can have NHS treatment only if judged by a urologist to be in extreme distress places a burden of psychological assessment on this clinician; the clinician will have little access to detailed psychiatric history, poor knowledge of the patient's social situation, little time for a mental state examination, and no training or experience in psychiatry. It would be surprising if decisions made by such a professional were any more valid than those made by the patient's general practitioner. This measure represents only a crude attempt to appease opponents of the rationing decision and does nothing to solve logically the problems that such new treatments produce.

Andrew Green *General practitioner*
Hedon Group Practice, Hedon, East Yorkshire
HU12 8JD
orchardh@globalnet.co.uk

1 Department of Health. *Viagra: NHS prescription proposals announced*. London: DoH, 1999. (Press release 21 January.)

2 British Medical Association. *GPC Viagra statement*. London: BMA, 1999. (Press release 21 January.)

Older doctor derived great benefit from sildenafil

EDITOR—A retired doctor in my 60s, I had come to accept impotence as part of normal ageing. But taunts in an unhappy marriage led me to explore the problem with a specialist colleague. Soon afterwards widowed, and eventually enjoying a new relationship with a much younger widow, I was confronted again with the bogey of erectile dysfunction. After further psychiatric counselling a trial of the latest new remedy proved unhelpful, so I reluctantly accepted my limitations once again.

Later, a consultant urologist consulted for nocturnal frequency advised review of the erectile dysfunction. Onward referral to a uroandrogologist established organic causation amenable to treatment. He recommended androgen replacement and was reassuring about its risks. Unfortunately, despite improved hormonal concentrations, testosterone alone failed to restore potency. The alternative oral regimen proved hard to sustain without any discernible benefit. Injections of another remedy into the penis proved too repugnant to tolerate.

The latest reputedly magic cure—sildenafil (Viagra)—became available eventually to nominated specialists. Before prescribing it my uroandrogologist sought psychiatric reassurance about my relationship with my new wife, holding a prevalent belief that such problems usually had a psychological component. Before our joint psychiatric appointment, sildenafil became freely available abroad. Tablets bought over the counter on holiday in Switzerland and taken in a low dose restored normal potency immediately, after 15 years' impotence. The benefits to a retired doctor in his early 70s and his premenopausal wife do not need to be spelt out.

No untoward side effects occurred, so the uroandrogologist recommended continuation of hormone replacement plus sildenafil (by then licensed in the United Kingdom), and the joint prescription was implemented by my general practitioner. Later, having studied official advice, she stopped the sildenafil; she would continue to prescribe the androgen, but I should seek sildenafil privately.

Accustomed to my health needs being met by the NHS, to which I had given a lifetime of service, I asked her to reconsider her decision. After reviewing my extensive specialist investigations she exercised her clinical judgment and resumed NHS prescription of testosterone with sildenafil, pending the health minister's advice and the BMA's response to it.

It is anomalous that a joint NHS prescription, shown as curative for a seemingly irremediable organic condition, is apparently to become impermissible again unless the proposed list of "acceptable" reasons for impotence is reviewed sensibly and promoted on an advisory basis. Even if a more logical list was devised it should not deny doctors the right to exercise clinical discretion individually.

Name supplied *Consultant psychiatrist*

Summary of electronic responses

The Viagra debate attracted about 30 responses.¹ The following quotes cover the broad range of opinions.

- Nobody seems to be against rationing:
"Until we have all the surgeons and community nurses who could help some patients, there is no case for providing drugs for all" (P West).

"We should welcome and applaud the bravery of the secretary of state for health in

opening up the whole issue of rationing" (L Miller).

"The core services provided by the NHS need to be redefined ... the less money spent on Viagra the better. Impotence is hardly life threatening" (M Sparks).

"The real issue is that we will soon no longer be able to operate by using the prescription pad as a blank cheque" (T Hyde).

"Guidelines seem to be necessary for rationing. If no guideline is enforced, leaving it to the individual GP will lead to enormous variation in prescribing, as shown for many other drugs. Fairness requires a consistent response" (P West).

- Guidelines and the tricky business of private prescription, however, are of concern:

"The secretary of state, no doubt in a state of panic, may have initially come up with a rather shoddy list of indications for Viagra, but hopefully this will be modified when the dust has settled and the spot-light is off" (J Stapleton).

"If the drug is to be rationed, then consider all the possibilities" (S Shaffi).

"Supposedly there are many acceptable indications for eligibility. Patients will be eligible to obtain Viagra on the NHS if they have had their 'prostate removed.' Does this include all forms of prostate surgery, or does it only include patients who have had a radical prostatectomy for cancer? ... Further, there is the question who is going to assess if a patient is suffering from 'severe distress?'" (R Beck), and this is one urologist who certainly would not be making such an assessment.

"Maybe a blanket ban would be fairer than one with more holes than a Swiss cheese" (D Hopkins).

"GPs prescribing sildenafil would be in breach of guidance if they use an FP10, and in breach of their terms of service if they issue a private prescription to their own patients" (P English).

- Specialist consultants are better off:

"I cannot write a private prescription on NHS premises. Thus the patients end up having to make a private appointment and pay for the drug! Crazy, but good for my income" (G Caldwell).

"Why not make sildenafil (Viagra) available over the counter from pharmacists?" (C West).

¹ Electronic responses, Viagra debate. *BMJ* 1999; 318. (www.bmj.com/cgi/content/full/318/7178/DC1). (30 January)

Patient centred care of diabetes in general practice

Doctors and nurses must understand meaning of "communication"

EDITOR—In Kinmonth et al's randomised controlled trial of patient centred care of diabetes one group of health professionals (the intervention group) received training in patient centred care while the other (the comparison group) did not.¹ Subsequently, scores on quality of communication with

patients favoured the intervention group, but scores on patients' knowledge about their disease favoured the comparison group—to put it crudely, patients assigned to the intervention group were relatively happy but ignorant.

Sadly, the straightforward paradox of saying, in effect, "he communicates well but tells you nothing" is all but lost in the way that the word communication is used in medical education these days. Indeed, in a manner that many medical educators would find unexceptionable, Kinmonth et al gloss this term as meaning "the ability to tell the doctor or nurse personal or troubling things and feeling understood"—as if the transfer of information from professional to patient was something else. Where does this content free definition come from?

Perhaps the problem is that the whole communication skills movement arose from a perception that medicine (if not nursing) had insufficient empathy; the result—the common, lazy identification of empathy with patient centredness, and of both with communication—therefore has its own historical logic. But we cannot propagate a view of communication that suggests that "just to be there is enough."

We need a more grown up and integrated view of what is at stake. Put simply, health professionals have skill: if they didn't, patients wouldn't visit them. It is hardly patient centred to act as if things were otherwise, or rational to deny that it is knowledge that people seek to communicate.

John Skelton *Senior lecturer in communication skills*
Department of Primary Care and General Practice,
Medical School, University of Birmingham,
Birmingham B15 2TT
j.r.skelton@bham.ac.uk

¹ Kinmonth AL, Woodcock A, Griffin S, Spiegel N, Campbell MJ on behalf of the Diabetes Care from Diagnosis Research Team. Randomised controlled trial of patient centred care of diabetes in general practice: impact on current wellbeing and future disease risk. *BMJ* 1998;317:1202-8. (31 October.)

Study failed to measure patient centredness of GPs' consulting behaviour

EDITOR—Kinmonth et al should not claim that their study "shows the power of a consultation to affect patients' health and wellbeing."¹ They evaluated some of the outcomes of training health professionals in patient centred clinical method but failed to measure the patient centredness of general practitioners' or nurses' consulting behaviour. Consequently, we have no idea whether the training actually resulted in clinicians providing more patient centred care; the study provides circumstantial evidence about the influence of patient centred clinical method on clinical outcomes.

Before we can accurately investigate the effects of patient centred care we need reliable instruments for measuring the relevant aspects of doctor-patient communication.² Although several communication coding systems exist, few have been validated adequately for their intended uses.³ One of these research tools was designed to measure clinicians' patient centredness,⁴ and there is some evidence of its reliability⁴ and validity.³

Further development of this instrument will clarify its strengths and weaknesses for measuring clinicians' patient centredness in routine consultations in primary care. This would enable those investigating the effects of patient centred care to use it to monitor doctor-patient communication. Although this is methodologically challenging and likely to be costly, it could ultimately lead to trials providing more relevant information. If Kinmonth et al had used this approach they would perhaps have produced hard evidence to relate their unexpected findings to clinicians' consulting behaviours.

Tim Coleman *Clinical lecturer*
Department of General Practice and Primary Health Care, University of Leicester, Leicester General Infirmary, Leicester LE5 4PW

- 1 Kinmonth AL, Woodcock A, Griffin S, Spiegel N, Campbell MJ on behalf of the Diabetes Care from Diagnosis Research Team. Randomised controlled trial of patient centred care of diabetes in general practice: impact on current wellbeing and future disease risk. *BMJ* 1998;317:1202-8. (31 October.)
- 2 Inui TS, Carter WB. Problems and prospects for health services research on provider-patient communication. *Med Care* 1985;23:521-38.
- 3 Ong LM, de Haes JC, Hoos AM, Lammes FB. Doctor-patient communication: a review of the literature. *Soc Sci Med* 1995;40:903-18.
- 4 Henbest RJ, Stewart MA. Patient-centredness in the consultation. 1. A method for measurement. *Fam Pract* 1989;6:249-53.
- 5 Henbest RJ, Stewart M. Patient-centredness in the consultation. 2. Does it really make a difference? *Fam Pract* 1990;7:28-33.

Might difference in prescribing explain some of the findings?

EDITOR—Kinmonth et al's study has important implications for general practice as the authors suggest that practitioners who concentrate on being patient centred might be at risk of losing the focus on disease management.¹ One of the findings that supports this conclusion is that patients in the intervention group gained weight.

In the discussion, the authors suggest that this difference between the two groups could not be accounted for by small differences in prescribing rates of hypoglycaemic agents (50% in the intervention group and 46% in the comparison group). There were, however, larger differences between the groups in the type of hypoglycaemic agent prescribed. In the intervention group 47% of the patients were prescribed a sulphonylurea and 11% a biguanide. In the comparison group the figures were 40% and 15% respectively. Given that sulphonylureas tend to cause weight gain and biguanides promote weight loss, might not some of the difference in weight be accounted for by this difference in prescribing?

Christopher Hand *Adviser in postgraduate general practice, Anglia and Oxford Region*
School of Health Policy and Practice, University of East Anglia, Norwich NR4 7TJ
c.hand@uea.ac.uk

- 1 Kinmonth AL, Woodcock A, Griffin S, Spiegel N, Campbell MJ on behalf of the Diabetes Care from Diagnosis Research Team. Randomised controlled trial of patient centred care of diabetes in general practice: impact on current wellbeing and future disease risk. *BMJ* 1998;317:1202-8. (31 October.)

Authors' reply

EDITOR—Defining and measuring patient centred consulting and its outcomes are

problematic, as our study shows. We defined patient centred consulting as an integration of the individual patient's perspective with the practitioner's general knowledge of disease, recognising that doctors and nurses do not deal with diseases alone; they deal also with individuals who are ill or concerned about their health.

Skelton argues for the primacy of communication of medical knowledge in the consultation, but our assertion is that empathy is the means by which the knowledge possessed by the practitioner becomes applicable by the patient. The prevailing approach to the management of patients with newly diagnosed type 2 diabetes remains one of "downloading information," despite evidence of its relative ineffectiveness in enabling self care.¹ Approaches taking account of the particular situation and feelings of the individual patient seem more likely to support the integration of information into perception and action.² But teaching this integrated approach did not prove easy and deserves further effort.

Coleman suggests that we should have directly observed the patient centredness of practitioners' consulting behaviour. Evidence is emerging, however, that it is patients' views of the process of the consultation, rather than researchers' perceptions of it, that are key predictors of health outcomes.^{3,4} We observed consultations during pilot work⁴ and subsequently measured the effect of training on practitioners' attitudes, behaviours, and style of care through patients' and practitioners' reports.⁵

Weight was higher in the intervention than the comparison group, but only among the 112 (48%) patients prescribed hypoglycaemic drugs. Hand wonders if this is because of a 7% excess of sulphonylurea prescribing (which might promote weight gain) in the intervention group and a 4% excess of metformin prescribing (which might promote weight loss) in the comparison group. We also wondered about this, but adjustment for category of drug in a regression analysis made no impact: the difference between groups in body mass index at one year was 2.00 kg/m² after adjustment for sulphonylurea and 2.12 kg/m² after adjustment for metformin compared with 2.04 kg/m² before adjustment. No measure was made of adherence to the prescribed

drug in the study. Without such a measure we cannot tell whether the training given to practitioners produced the reported difference in outcomes through greater adherence to the drug in the intervention group.

Ann Louise Kinmonth *Professor of general practice*
alk25@medschl.cam.ac.uk

Simon Griffin *University lecturer in general practice*
General Practice and Primary Care Research Unit, Institute of Public Health, Cambridge CB2 2SR

Alison Woodcock *Lecturer*
Department of Psychology, Royal Holloway, University of London, Egham, Surrey TW20 0EX

Mike Campbell *Professor of medical statistics*
School of Health and Related Research, Northern General Hospital, Sheffield S5 7AU
On behalf of the Diabetes Care from Diagnosis Research Team

- 1 Griffin SJ, Kinmonth AL, Skinner C, Kelly J. *Educational and psychosocial interventions for adults with diabetes*. London: British Diabetic Association, 1999.
- 2 Stewart M, Brown JB, Weston WW, McWhinney IR, McWilliam CL, Freeman TR. *Patient-centred medicine: transforming the clinical method*. Thousand Oaks, CA: Sage, 1995.
- 3 Little P, Gould C, Williamson I, Warner G, Gantley M, Kinmonth AL. Reattendance and complications in a randomised trial of prescribing strategies for sore throat: the medicalising effect of prescribing antibiotics. *BMJ* 1997;315:350-2.
- 4 Kinmonth A, Spiegel N, Woodcock A. Developing a training programme in patient-centred consulting for evaluation in a randomised controlled trial: diabetes care from diagnosis in British primary care. *Patient Educ Couns* 1996;29:75-86.
- 5 Woodcock AJW, Kinmonth AL, Campbell MJ, Griffin SJ, Spiegel NM. Diabetes care from diagnosis: effects of training in patient-centred care on beliefs, attitudes and behaviour of primary care professionals. *Patient Educ Couns* 1999;37:65-79.

Prenatal and perinatal risk factors for psychiatric diseases of early onset

Results are different if seasons are categorised differently

EDITOR—Hultman et al found that late winter births are associated with increased risks of schizophrenia and affective psychosis, the odds ratios being 1.4 and 1.5 respectively.¹ The authors presented the excess of late winter births as one of their key messages, although the finding reached significance only among male subjects with affective psychosis.

Using the northern Finland 1966 birth cohort (5636 male and 5381 female subjects), we partially replicated Hultman et al's study. The cohort is a genetically homogeneous population, and data are available

Odds ratios (with 95% confidence intervals) for incidence of schizophrenia and numbers of births in relation to season of birth in northern Finland 1966 birth cohort study

	Male subjects		Female subjects	
	Odds ratio (95% CI)	No of cases/controls	Odds ratio (95% CI)	No of cases/controls
Season of birth according to study by Hultman et al¹				
January to April	0.99 (0.59 to 1.66)	22/1861	1.21 (0.64 to 2.29)	15/1699
May to December (reference category)	1.0	42/3510	1.0	26/3560
Season of birth defined by solstices and equinoxes				
February to April	2.01 (0.91 to 4.44)	20/1397	1.65 (0.65 to 4.20)	12/1267
May to July	1.62 (0.55 to 1.65)	16/1386	1.69 (0.68 to 4.21)	14/1439
August to October	2.02 (0.91 to 4.49)	19/1320	1.04 (0.38 to 2.88)	8/1335
November to January (reference category)	1.0	9/1268	1.0	7/1218

on biological, socioeconomic, environmental, and health conditions for every individual from mid-pregnancy of their mother to the age of 31. Psychiatric diagnoses were made on the basis of extracts from the Finnish national hospital discharge register and a detailed diagnostic validation process.² Birth seasonality was examined separately in those with schizophrenia (*Diagnostic and Statistical Manual of Mental Disorders*, third edition, revised, code 295; n=105) and those with no mental disorder (controls; n = 10 630). Our study (table) did not show a significant association between late winter births and schizophrenia in either male or female subjects when seasons were defined according to the study by Hultman et al.

The method that Hultman et al used gives us cause for concern on three counts.

Firstly, the authors should have taken into account the birth seasonality in the general population, because in Western countries most births occur during the first half of the year.^{3 4}

Secondly, because their study population was young, the "age incidence effect" as well as the "age prevalence effect" may explain their finding and should at least have been discussed.⁵⁻⁵

Thirdly, the authors' categorisation of seasons should have taken account of natural photoperiods. Although the mechanisms of "the season of birth phenomenon" in psychiatric disorders remain obscure and may be related to seasonality of environmental factors or circadian rhythms of internal biological processes,^{3 4} the observed seasonal effect is mainly based on the seasonal variations in climatic factors (light, temperature, humidity, etc). In Sweden as well as Finland, weather conditions vary considerably between seasons. Hultman et al used May to December as their reference season, but that period includes months with extreme climatic factors, and its use in the statistical analyses is therefore questionable. The use of monthly, bimonthly, or quarterly periods around the solstices and equinoxes in the authors' analyses would have been preferable, allowing comparisons with previous studies.⁵⁻⁵ As our table shows, the results are different when the categorisation of seasons is changed.

Pirkko Räsänen Senior research fellow
pirkko.rasanen@oulu.fi

Helinä Hakko Biostatistician
Department of Psychiatry, University of Oulu,
FIN-90210 Oulu, Finland

Marjo-Riitta Järvelin Professor
Department of Public Health Science and General
Practice, FIN-90220 Oulu

1 Hultman CM, Sørensen P, Takei N, Murray RM, Cnattingius S. Prenatal and perinatal risk factors for schizophrenia, affective psychosis, and reactive psychosis of early onset: case-control study. *BMJ* 1999;318:421-6. (13 February.)

2 Isohanni M, Mäkiyö T, Moring J, Räsänen P, Hakko H, Partanen U, et al. A comparison of clinical and research DSM-III-R diagnoses of schizophrenia in a Finnish national birth cohort. *Soc Psychiatry Psychiatr Epidemiol* 1997;32:303-8.

3 Fossey E, Shapiro CM. Seasonality in psychiatry—a review. *Can J Psychiatry* 1992;37:299-308.

4 Torrey EF, Miller J, Rawlings R, Yolken RH. Seasonality of births in schizophrenia and bipolar disorder: a review of the literature. *Sch Res* 1997;28:1-38.

Monthly distribution of births of subjects who developed schizophrenia and affective psychosis, their controls, and study base of Swedish children born in 1973-9 who survived until 1 January 1987

	Schizophrenia		Affective psychosis		Study base (n=693 312)
	Cases (n=167)	Controls (n=835)	Cases (n=198)	Controls (n=990)	
January	9.6	8.4	11.1	9.8	8.4
February	8.4	10.2	8.6	7.6	8.2
March	15.6	8.4	10.1	9.4	9.7
April	10.2	9.3	14.1	9.4	9.6
May	7.2	10.4	9.1	9.4	9.3
June	7.2	7.9	5.1	9.0	8.4
July	4.8	9.6	9.1	7.8	8.3
August	6.6	8.4	9.1	6.7	8.1
September	10.8	7.3	6.1	7.4	8.1
October	5.4	7.1	6.6	8.8	7.8
November	5.4	5.5	3.5	7.1	7.1
December	9.0	7.5	7.6	7.8	7.1

5 Lewis MS, Griffin PA. An explanation for the season of birth effect in schizophrenia and certain other diseases. *Psychol Bull* 1981;89:589-96.

Author's reply

EDITOR—Season of birth is one of the most robust epidemiological findings in research into schizophrenia, with a 5-15% excess of births in the winter and spring. As an increased rate of perinatal obstetric difficulties among patients with schizophrenia born in the winter has been shown,¹ we estimated the independent effect of season of birth after adjusting for other possible risk factors.

In the multivariate analyses birth in the late winter was associated with an increased risk of schizophrenia (odds ratio 1.4) and affective psychosis (odds ratio 1.5). The risk effect that we found is in line with those in other large studies.² For schizophrenia, the point estimates were similar in the sex specific analyses; the differences between the total sample and separate analyses for sex (with fewer patients) were a matter of power. As Räsänen et al point out, the season of birth effect was more prominent among male than female subjects with affective psychosis, but these differences should be interpreted with caution, as the numbers of cases in the analyses stratified for sex were low.

Season of birth was categorised into the periods January-April and May-December according to work by Dalén.³ Our control sample was randomly selected, and we had no reason to believe that these subjects did not follow the birth seasonality of the general population. The table shows the monthly distribution of births in the patients, their controls, and the study base of 693 312 Swedish children.

If births were analysed bimonthly, as Räsänen et al suggest, March-April had a maximum excess relative to the control sample and the general population for both schizophrenia and affective psychosis. For schizophrenia the shift from maximum deficit to excess seems to lie earlier than January, and we consequently argue that the effect is genuine and not an age incidence effect.

We agree with Räsänen et al that season of birth examined by quarterly periods around the solstices and equinoxes is of interest, with seasonal effects (harmful or protective) of light, temperature, and

weather conditions possibly having a role. Other possible explanations are seasonal effects of nutrition and infectious agents.² When we repeated our original multivariate analysis with the suggested categorisation of season around natural photoperiods, schizophrenia was associated with birth in February to April (odds ratio 1.91 (95% confidence interval 1.18 to 3.09); May to July as the reference category), with maternal and perinatal and perinatal factors being controlled for.

Much larger sample sizes than ours are required to explain the season of birth effects, which also occur in other neurodevelopmental disorders.

Christina M Hultman Research fellow
Department of Neuroscience, Psychiatry, Ulleråker,
University of Uppsala, S-750 17 Uppsala 17,
Sweden
Christina.Hardman@ulpsyk.uu.se

1 Wrede G, Mednick SA, Huttunen MO, Nilsson CG. Pregnancy and delivery complications in the births of an unselected series of Finnish children with schizophrenic mothers. *Acta Psychiatr Scand* 1980;163:522-34.

2 Cotter D, Larkin C, Waddington J, O'Callaghan E. Season of birth in schizophrenia; clue or cul-de-sac. In: Waddington J, Buckley P, eds. *The neurodevelopmental bases of schizophrenia*. Heidelberg: Springer-Verlag, 1996.

3 Dalén P. Season of birth: a study of schizophrenia and other mental disorders. Amsterdam: Elsevier/North Holland, 1975.

Antibiotics for acute sinusitis in general practice

Entry criteria were too dissimilar for studies to be combined for meta-analysis

EDITOR—De Ferranti et al's meta-analysis on antibiotics in acute sinusitis could be criticised in many ways.¹ Published in the General Practice section of the *BMJ*, it is presumably aimed at general practitioners.

The authors first looked at randomised controlled trials on the effectiveness of antibiotic against non-antibiotic management and then carried out a meta-analysis of the six placebo controlled trials they identified. It is implied that the patients being randomised were similar in each of the studies, so that a meta-analysis could be performed. This is not the case. One of the papers (that by Wald et al, in 1986) concerned the treatment of children and

cannot be combined with the other five papers, which were adult studies.

Table 1 in the authors' paper shows the clinical settings and the entry criteria for the other five studies. In looking at this, readers have to identify which study is applicable to their own clinical setting. The first study required a positive result of diagnostic irrigation for a patient to be included in the study, the second required a positive bacteriological culture, the third required a positive computed tomogram, and the fourth required a positive plain x ray film. The entry criteria in these four studies are thus grossly different and invalidate their being combined. The fifth study was the only one that had clearly defined symptomatic entry criteria that readers could apply to their own practice and did not rely on investigation—likely to be the case in most British primary care centres.

The title of Stalman et al's paper embodies the usual situation that general practitioners face—"adults with acute sinusitis-like complaints."² Their study showed no effect of antibiotic (doxycycline) versus placebo on the mean number of days until resolution of facial pain and resumption of daily activities in 192 randomised patients. Though a power analysis is not reported to have been performed to exclude a type I error, the number of patients is sufficiently large to exclude a clinically material difference. Hence this trial should be the main evidence on which general practitioners should base their practice—that antibiotics have no effect in adults with acute sinusitis-like complaints—until other studies in similar situations show otherwise. A meta-analysis of which antibiotic is best in acute sinusitis is thus made redundant and the rest of the paper irrelevant.

G G Browning *Professor of otolaryngology and head and neck surgery*
University of Glasgow, Royal Infirmary, Glasgow
G31 2ER
sec@ihr.gla.ac.uk

1 de Ferranti SD, Ioannidis JPA, Lau J, Anninger WV, Barza M. Are amoxycillin and folate inhibitors as effective as other antibiotics for acute sinusitis? A meta-analysis. *BMJ* 1998;317:632-7. (5 September.)

2 Stalman W, van Essen GA, van der Graaf Y, de Melker RA. The end of antibiotic treatment in adults with acute sinusitis-like complaints in general practice? A placebo-controlled double-blind randomized doxycycline trial. *Br J Gen Pract* 1997;47:794-9.

Authors' reply

EDITOR—We strongly disagree with Browning's approach to our meta-analysis and to meta-analysis in general. The notion that patients in randomised trials must be closely similar for the trials to be included in a meta-analysis is unrealistic.^{1,2} No two patients are identical or even closely alike. The six trials in our meta-analysis show no statistical evidence of heterogeneity, but this does not mean that biological heterogeneity does not exist.

In our paper we pointed out that the trial that included patients simply on the basis of sinusitis-like symptoms had higher spontaneous cure rates in the placebo arm

and failed to show a beneficial effect of antibiotics.³ Nevertheless, the 95% confidence intervals in this moderate-size trial extend to a more than 50% reduction in the risk of clinical failures. We strongly disagree that a study of 176 patients with a confidence interval that includes such a large effect is "sufficiently large to exclude a clinically material difference."

Browning has misinterpreted our meta-analysis as a recommendation to use antibiotics in all patients with sinusitis-like symptoms. We clearly state that, on average, over two thirds of patients improve spontaneously without antibiotics. The percentage might be higher in patients with predominantly viral sinusitis. Our results show that antibiotics reduce the risk of clinical failure by half. For patients with a low risk of bacterial sinusitis, antibiotics would be unnecessary.

Better evidence is needed on how to differentiate bacterial from non-bacterial acute sinusitis. Cost effective approaches should be defined for not only the treatment but also the diagnosis of this condition. We have performed a decision analysis to address these issues, using data from diagnostic and therapeutic studies.⁴ This analysis suggests that when the prevalence of bacterial sinusitis is expected to be <25%, symptomatic management is the most cost effective option, with antibiotics being reserved for patients whose condition has not improved after a week. For a prevalence of 25-83%, applying clinical criteria (a history of purulent rhinorrhoea, facial pain, presence of pus⁵) to decide whether to use antibiotics becomes the most cost effective approach. For a higher prevalence, immediate treatment with antibiotics is best.

We do not agree that excluding five of the six available studies provides most insight into this common condition. On the contrary, we need more data on different clinical settings and different patient populations to determine which approaches serve patients best.

John P A Ioannidis *Research fellow*
Sarah D de Ferranti *Research associate*
Michael Barza *Professor*
Joseph Lau *Associate professor*
New England Medical Center, Boston, MA 02111, USA
joseph.lau@es.nemc.org

1 Lau J, Ioannidis JPA, Schmid CH. Summing-up evidence: one answer is not always enough. *Lancet* 1998;351:123-7.

2 Ioannidis JPA, Lau J. Heterogeneity of the baseline risk within clinical trial populations: a proposed evaluation algorithm. *Am J Epidemiol* 1998;148:1117-26.

3 Stalman W, van Essen GA, van der Graaf Y, de Melker RA. The end of antibiotic treatment in adults with acute sinusitis-like complaints in general practice? A placebo-controlled double-blind randomized doxycycline trial. *Br J Gen Pract* 1997;47:794-9.

4 *Diagnosis and treatment of acute bacterial sinusitis. Evidence report/technology assessment No 17.* Rockville, MD: Agency for Health Care Policy and Research, 1998. (Prepared by New England Medical Center under contract No 290-97-0019. AHCPR publication No 99-E0017.)

5 Berg O, Carenfelt C. Analysis of symptoms and clinical signs in the maxillary sinus empyema. *Acta Otolaryngol* 1988;105:343-9.

Breast feeding and working mothers

Summary of electronic responses

Mary Black's personal view on the problems she encountered because she chose to work full time while breast feeding¹ caused much correspondence. We received about 25 letters, and most are published on our website.² Most of the respondents were mothers who had breast fed or were currently breast feeding their baby.

"Breast feeding is more than just a healthy way of getting food and drink into babies. It is a social choice, not just a nutritional one, which mothers make in the teeth of social pressures" (H Welford). "Modern Western society does not make it easy for mothers to integrate their public and private lives" (K A Dettwyler), and "young mothers find society poorly supportive in many ways" (B Symon).

However, mothers who breast fed at work "may improve their world by 'normalising' breast feeding for the people around them" (G S Hertz). "By doing this, and insisting on getting the appropriate conditions to do so, breast feeding mothers may change the working environment slowly but surely" (M Leibovich). And "by the time these breast fed children reach parenthood things will be different" (G S Hertz).

But for now breast feeding mothers returning to work "need more than sympathetic rhetoric. The minimum requirements are a private area for pumping milk close to handwashing facilities and refrigerator space to store milk" (K Farrell). "Legislation is presently being considered in the US House of Representatives to ensure a woman has the time and place to express her milk. I say this is silly, to have to pass a law for ensuring a woman can do this. But, unfortunately, it is needed" (M Griesse).

1 Black ME. Breast feeding and my brain. *BMJ* 1999;318:545. (20 February.)

2 Electronic responses. Breast feeding and my brain. *eBMJ* 1996;318. (www.bmj.com/cgi/content/full/318/7182/545#responses)

Britain's Academy of Medical Sciences has been busy in recent months

EDITOR—I was pleased to see Minerva's concern about the reality of the Academy of Medical Sciences.¹ By her own count, her mention of it raises its media score from 0 to 1. However, she can be reassured that the academy is not a dream.

The academy has had a busy time since its launch last November. It has recruited Dr Jolyon Oxley as its executive secretary and has moved into offices in 10 Carlton House Terrace, London. It is setting up its network of links with the government, with medical and academic bodies, and with sister academies overseas. A handbook should be ready by September, and a website (www.acmedsci.ac.uk) was established this

month. The 100 successful candidates in the academy's first annual election of fellows is available on the website.

Four working parties have already been set up and are at various stages of their work. Reports on "Career structure and prospects for clinical scientists in the UK"; "The interaction of the medical sciences with the media"; "The state of academic bacteriology in the UK"; and, as part of National Academies Policy Advisory Group, "Intellectual fraud and misconduct" should appear at the end of the year.

Several national and regional scientific meetings are at the planning stage. One that should certainly go into Minerva's diary is on 9 July, when the admission of the new fellows will be marked by a high profile scientific and social meeting. And of vital importance is the academy's brochure (written, designed, and produced by Burnett Publications, London EC2A 3AY), which will form the core of its fundraising and profile building activities.

Minerva might do well to contemplate these lines from A H Clough:

Say not the struggle naught availeth,
The labour and the wounds are vain,
The enemy faints not, nor faileth.
And as things have been they remain
For while the tired waves, vainly breaking,
Seem here no painful inch to gain,
Far back, through creeks and inlets making,
Comes silent, flooding in, the main.

Peter Lachmann *President*
Academy of Medical Sciences, London SW1Y 5AH
pjl1000@cam.ac.uk

1 Minerva. *BMJ* 1999;318:884. (27 March.)

Early diet in preterm babies and later intelligence quotient

Surely study showed that breast milk is feed of choice for premature babies

EDITOR—Lucas et al's paper on early diet in preterm babies and their later intelligence quotient (IQ)¹ sparked intense media interest: "Premature babies need enriched diets," said the BBC.² The authors didn't point out any of the known advantages of breast milk for premature babies.

The paper compared babies who received only infant formula with babies who received breast milk and infant formula. Babies received either standard term formula or preterm formula. The main outcome, IQ at age 7-8, showed no advantage for preterm formula. Subgroup analyses looked at results for boys and girls. Few details were given about how successful the randomisation was in terms of the sex of the baby in relation to weight. Is it then valid to draw separate conclusions for boys and girls?

We suggest that the following conclusion would be more accurate: "Breast milk continues to be the feed of choice for premature babies. The results did not show an advantage for preterm formula in terms of cognitive performance at 7½-8 years.

Further work with a cohort randomised by sex would be of interest."

Publication of this study seems to have been delayed. The last data must have been collected in 1992—and indeed were used in a paper published in that year.³ If the study had contained anything to change practice it should have been published as soon as the results were known.

Breast feeding is the physiological norm for infants. Any other method of feeding babies must be assumed to be risky until proved otherwise. The onus is on researchers to prove that substitutes are both safe and beneficial. Indeed, it is recommended that all research studies on infant feeding should include a breast feeding reference group to put the findings into context.⁴ When research compares two breast milk substitutes, neither one can be concluded to be safer or better than breast milk. When the sponsors of research stand to gain from increased sales of infant formula, readers must be aware of the conflict of interest.

Researchers and reporters have a responsibility to parents and babies. Headlines such as "Premature babies need enriched diets" can undermine women. It would confirm the work of health professionals and lay workers, and support parents providing breast milk for their babies, if press releases concerning infant feeding reiterated the simple message that "breast is best."

Deborah Behrman *Breastfeeding supporter*

Mary Broadfoot *Breastfeeding supporter*
broadfoot@btinternet.com

Phyll Buchanan *Breastfeeding supporter*

Carolanne Lamont *Breastfeeding supporter*

Magda Sachs *Breastfeeding supporter*
Breastfeeding Network PO Box 11126,
Paisley PA2 8YB

1 Lucas A, Morley R, Cole TJ. Randomised trial of early diet in preterm babies and later intelligence quotient. *BMJ* 1998;317:1481-7. (28 November.)

2 BBC News, 26 November 1998, Health. Premature babies need enriched diets. <http://news.bbc.co.uk/1/health/health/newsid%5F222000/222453.stm>

3 Lucas A, Morley R, Cole TJ, Lister G, Leeson-Payne C. Breast milk and subsequent intelligence quotient in children born preterm. *Lancet* 1992;339:261-4.

4 Committee on Medical Aspects of Food and Nutrition Policy. *Guidelines on the nutritional assessment of infant formulas* 47. London: Stationery Office, 1996.

Authors' reply

EDITOR—Behrman et al disappoint us, since we have provided as much evidence as any other authors, internationally, supporting breast milk for neonatal care. Our preterm studies, suggesting that breast milk reduces necrotising enterocolitis, systemic sepsis, and feed intolerance and promotes neurodevelopment, have strongly influenced mothers to provide their milk.¹⁻⁴ We recommend that mothers of preterm infants be counselled to provide breast milk on health grounds. Against this, any implication that our study was biased towards promoting sales of infant formula is misplaced.

Behrman et al miss the point. They say that the claim that premature babies need enriched diets undermines women. What mothers readily appreciate, given proper explanation, is that it is undermining to the health of premature babies not to receive

their special nutritional requirements. We cannot suppose that human milk evolved to feed babies born up to four months too soon, too immature to feed, who would have died before modern care. Had these babies remained in utero they would have been fed via the placenta; thus no enteral feed—breast milk or formula—is "physiological" for them.

As it happens, objective research strongly favours human milk, but not as a sole source of nutrition. Substantial work over 50 years shows that preterm infants have special nutrient needs—for instance, the brain at 28 weeks' gestation is growing 2.5 times faster than the brain at term.⁵ Their rapid growth, bone mineralisation, and need to establish tissue nutrient stores often require fuelling by the additional use of specialised nutritional products.⁵ Moreover, special formulas are essential when mothers express insufficient milk.

Behrman et al should re-examine our paper to resolve their misunderstanding of the randomised groups and reappraise our justification for subgroup analyses. They infer incorrectly that our database was complete by 1992. Our paper described a randomised study providing evidence of serious consequences of feeding preterm infants suboptimally. These data should not, however, undermine the use of expressed breast milk for premature babies or the establishment of breast feeding, both of which occurred in our study population.

Supporters of breast feeding must appreciate that standard public health advice and campaigns relevant to healthy full term infants cannot be applied to the complex hospital care of premature babies. Any advocacy role undertaken by this or any similar organisation must be backed by a medical understanding of this special circumstance.

Alan Lucas *Clinical research professor, MRC Childhood Nutrition Research Centre*
A.Lucas@ich.ucl.ac.uk

Tim Cole *MRC senior scientist, Department of Epidemiology and Public Health*
Institute of Child Health, London WC1N 1EH

Ruth Morley *Senior research fellow*
Menzies Centre for Population Health Research, Hobart, and Clinical Epidemiology and Biostatistics Unit, Department of Paediatrics, University of Melbourne, Parkville, Victoria 3052, Australia

1 Lucas A, Cole TJ. Breast milk and neonatal necrotising enterocolitis. *Lancet* 1990;336:1519-23.

2 Lucas A. Feeding low birthweight infants. In: Robertson NRC, ed. *Textbook of neonatology*, 2nd ed. London: Churchill Livingstone, 1992:279-88.

3 Lucas A, Morley RM, Cole TJ, Lister G, Leeson-Payne C. Breast milk and subsequent intelligence quotient in children born preterm. *Lancet* 1992;339:261-4.

4 Lucas A, Fewtrell M, Morley R, Lucas P, Baker B, Lister G, et al. Randomized outcome trial of human milk fortification and developmental outcome in preterm infants. *Am J Clin Nutr* 1996;2:142-51.

5 Tsang R, Lucas A, Uauy R, Zlotkin S, eds. *Nutritional needs of the preterm infant*. Pawling, USA: Caduceus Medical, 1993.

Rapid responses

Rapid responses submitted directly to our website are available on www.bmj.com