Website: www.bmj.com
Email: letters@bmj.com

Children born after intracytoplasmic sperm injection

What do authors mean by control infants?

EDITOR—Sutcliffe et al compare babies born through intracytoplasmic sperm injection with control babies (matched for social class, maternal education, region, and sex) born after unassisted conception.¹ In their table they give means, standard deviations, and 95% confidence intervals for the two groups separately. These confidence intervals tell us about the estimation of the mean in the population of which the babies are a random sample, which they are not. They tell us nothing about the comparison between the cases and controls.

These controls have been chosen to be similar to the study babies in many ways; they are not a representative sample of babies. It is the difference between the study babies and the control babies that is of interest, and the confidence interval we should have is that for this difference in mean. We cannot work this out from the data given, because this is a paired study and we have unpaired standard deviations. All we have for the comparison is the implication, not actually stated, that the differences are not significant. What we should be given are

confidence intervals that tell us how different the cases and controls are.

J M Bland *Professor of medical statistics*
St George's Hospital Medical School, London
SW17 0RE
mbland@sghms.ac.uk

¹ Sutcliffe AG, Taylor B, Li J, Thornton S, Grudzinskas JG, Lieberman BA. Children born after intracytoplasmic sperm injection: population control study. *BMJ* 1999;318:704-5. (13 March.)

Authors' reply

EDITOR—We do not think that a recalculation is necessary or valid for the conclusions that we made. Bland supposes that our study was a 1:1 paired case-control study. This was not so, but at the time of our report the groups were coincidentally the same size. The study (which is ongoing) now has 208 study children and 221 control children.

The controls were not selected from a general population but from a population likely to match the study infants. There was close similarity between the control and study groups in terms of factors known to affect developmental outcome. We believe that our results are valid, as is our conclusion that intracytoplasmic sperm injection seems safe in terms of children's development to age 2.

Alastair G Sutcliffe *Lecturer in child health*

Brent Taylor *Professor of child health*

Jun Li *Epidemiologist*

Royal Free and University College Medical School,
Department of Community Child Health, Royal
Free Campus, London NW3 2PF
icsi@rfhsm.ac.uk

Simon Thornton *Clinical director*

Centre for Assisted Reproduction, Park Hospital,
Nottingham NG5 8RX

Gedis Grudzinskas *Professor of obstetrics and gynaecology*

Department of Obstetrics and Gynaecology, St
Bartholomew's and the Royal London School of
Medicine and Dentistry, Royal London Hospital,
London E1 1BB

Brian Lieberman *Consultant in obstetrics and gynaecology*

Department of Reproductive Medicine, St Mary's
Hospital, Manchester M13 0JH

UK inquiry should establish why contaminated blood products were given to people with haemophilia

EDITOR—A government inquiry has been announced into the infection by contaminated blood products of people with haemophilia in the Republic of Ireland; this follows a

public inquiry into these issues in Canada.^{1 2} A similar inquiry should be conducted into the plight of people with haemophilia in the United Kingdom who have been infected in the same way.

Some 4800 people with haemophilia in the United Kingdom are believed to have been infected with hepatitis C virus through the use of contaminated blood products in their treatment by the NHS before 1986. Of these, more than 1200 were also infected with HIV. Nearly every patient with haemophilia who was treated before 1986 was infected; many have died as a result.

People with haemophilia throughout the world have been affected, but the response from various governments has differed. The Irish government, in addition to setting up a tribunal of inquiry, has set up a compensation scheme to provide financial assistance to all those who contracted hepatitis C or HIV infection, or both, from blood or blood products. Similarly, after a four year inquiry the Canadian government has set aside C\$1.1bn (£458m) to compensate an estimated 20 000 people. In France, Switzerland, and Japan government ministers and officials have faced charges relating to blood safety and infection. In the United Kingdom, however, the government has failed even to hold an inquiry into how and why patients with haemophilia came to be infected with bloodborne viruses before 1986.

The previous government ordered a look back study among patients infected with hepatitis C virus through blood transfusions, but this did not include patients with haemophilia who were infected by blood products. This omission means that there are still no official figures on hepatitis C in haemophiliac patients, despite these patients being one of the most widely monitored sectors of the UK population for bloodborne viruses.

The Haemophilia Society believes that it is important to establish how what has been described as one of the greatest treatment disasters in the history of the NHS took place. It has called on the health secretary for a full inquiry and also contacted the chairman of the select health committee. Why is an inquiry not appropriate in the United Kingdom? Firm information about something that has wrecked so many lives would help in the targeting of NHS services to those affected. It would also ease the anguish of those who have suffered, by giving them the answers they need and recognition of their plight.

Karin Pappenheim *Chief executive*
Haemophilia Society, London NW1 3AU
sue@haemophilia.org.uk

Advice to authors

We prefer to receive all responses electronically, sent either directly to our website or to the editorial office as email or on a disk. Processing your letter will be delayed unless it arrives in an electronic form.

We are now posting all direct submissions to our website within 72 hours of receipt and our intention is to post all other electronic submissions there as well. All responses will be eligible for publication in the paper journal.

Responses should be under 400 words and relate to articles published in the preceding month. They should include ≤ 5 references, in the Vancouver style, including one to the *BMJ* article to which they relate. We welcome illustrations.

Please supply each author's current appointment and full address, and a phone or fax number or email address for the corresponding author. We ask authors to declare any competing interest. Please send a stamped addressed envelope if you would like to know whether your letter has been accepted or rejected.

Letters will be edited and may be shortened.

www.bmj.com
letters@bmj.com

- 1 Spurgeon D. Canada creates new national blood service. *BMJ* 1997;315:1564.
- 2 Kondro W. Final Krever report paints picture of regulatory dysfunction. *Lancet* 1997;350:1688.

Proposed appraisal system and political correctness

Consultant appraisals will be useful

EDITOR—Kerr's negative attitude to the idea of consultant appraisals¹ reminded me of that of surgeons in the 19th century, who were said to sharpen their scalpels on their boots to show their contempt for the microbe theory. He also shows a common misunderstanding of the difference between appraisal and assessment. Appraisal is for the benefit of the one being appraised—akin to formative assessment in educational jargon.

Speaking for myself, I look forward to my first appraisal. If it provides the opportunity to draw the trust's attention to the ever rising pressures on us and the resulting threats to safe patient care then this much political correctness I will happily accept.

Andrew Whitehouse *Consultant physician*
16 Coleshill Road, Atherstone, Warwickshire
CV9 1BW
106354.574@compuserve.com

- 1 Kerr D. Political correctness is behind proposed appraisal system. *BMJ* 1999;318:1013. (10 April.)

Correspondent rather than the royal college may have lost touch with reality

EDITOR—We thank Kerr for his suggestion that the Royal College of Physicians may be suffering from political correctness and is disconnected from the real world.¹ It may be, however, that Kerr rather than the royal college has lost touch with reality if he has not noted the almost daily accounts in the media of doctors, some of them physicians, who have failed to achieve appropriate standards. This is why government and the profession are seeking ways to ensure continuing competence.

No medical or clinical director should have been appointed without having had proper training in appraisal, and the directors will themselves be subject to appraisal by trust managers. Appraisals are already routine in universities and many other professions, being an ideal vehicle for both career development and detecting problems early. Appraisals themselves will not affect the care of patients, but physicians whose weaknesses or gaps in knowledge are pointed out early, or who are able to discuss constructively their difficulties, are more likely to maintain high standards than are those who remain unaware of or hide such problems.

The college's document recognises the resource consequences of the several elements of clinical governance,² and it regularly points this out to the government. If the public truly wants assurance of continuing high standards of medical care it must pay for it.

K G M M Alberti *President*
John R Bennett *Treasurer*
Royal College of Physicians, London NW1 4LE.

- 1 Kerr D. Political correctness is behind proposed appraisal system. *BMJ* 1999;318:1013. (10 April.)
- 2 Royal College of Physicians. *Physicians maintaining good medical practice. Clinical governance and self regulation*. London: RCP, 1999.

Combining the two neonatal examinations

In primary care, second examination is useful

EDITOR—The paper by Glazener et al defines neonatal screening in narrow terms.¹ In primary care the second neonatal examination is usually performed about 6 weeks after birth. The main purpose of this check is not just screening for medical conditions such as hip and heart anomalies. It is also a good opportunity to screen for postnatal depression; Hearn et al found the incidence of this to be 15% when they used the Edinburgh depression score in primary care.²

The six week check also provides an excellent insight into the way that the mother interacts with her baby as well as providing an excellent opportunity to give advice about vaccinations and prevention of cot death and accidents.

Chris Harnden *General practitioner*
Prescott Surgery, Baschurch, Shropshire SY4 2EU
char992120@aol.com

- 1 Glazener CMA, Ramsay CR, Campbell MK, Booth P, Duffy P, Lloyd DJ, et al. Neonatal examination and screening trial (NEST): a randomised, controlled, switchback trial of alternative policies for low risk infants. *BMJ* 1999;318:627-31. (6 March.)
- 2 Hearn G, Iliff A, Kirby A, Ormiston P, Parr P, Rout J, et al. Postnatal depression in the community. *Br J Gen Pract* 1998;48:1064-6.

Midwives perform a neonatal examination, so was this counted?

EDITOR—Glazener et al described a study of two or one neonatal examinations done by medical staff.¹ In our hospital all newborn babies are examined by the midwife soon after birth for gross anomalies, which include cleft palate, spinal deformities, skull and facial deformities, limb abnormalities such as syndactyly, and number of vessels in the umbilical cord. The abnormalities are recorded on the back page of the mother's partogram and the doctor notified. As an example of this, I was recently asked to see a baby born the previous night, the attending midwife having noted a tooth in the baby's lower gum when she examined the baby after birth. I remember that this was the practice during my 10 years of service in NHS hospitals in England and would think that it is still so today.

If that is so the babies in the study would have been examined three times instead of two or twice instead of once. It would be useful to know how many of the gross superficial abnormalities in Glazener et al's study were detected soon after birth by the midwives.

T H H G Koh *Senior specialist in neonatology*
Kirwan Hospital for Women, Townsville, Great Barrier Reef, QLD 4817, Australia
KohT@health.qld.gov.au
Competing interests: None declared

- 1 Glazener CMA, Ramsay CR, Campbell MK, Booth P, Duffy P, Lloyd DJ, et al. Neonatal examination and screening trial (NEST): a randomised, controlled, switchback trial of alternative policies for low risk infants. (With commentary by JJ Deeks) *BMJ* 1999;318:627-32. (6 March.)

Authors' reply

EDITOR—Our study was confined to those neonatal screening examinations occurring in hospital and performed by paediatric staff. General practitioners who provide obstetric services have a statutory obligation to see a new mother and baby after delivery. This, as Harnden says, provides an opportunity to identify postnatal problems that develop at home in both the mother and baby and for health promotion advice.

Koh reminds us that all newborn babies are examined immediately after birth by the delivery suite midwife. Thus, in our study, babies allocated to the "one screen" policy on day 3 after birth were actually examined twice, and those allocated to the "two screen" policy were actually examined three times. We excluded from the trial babies who were admitted to the neonatal unit; these would have included those found in the labour ward by the midwife's examination to have a major abnormality that required treatment. We cannot tell how many of the remaining congenital abnormalities were detected before the paediatrician's formal screening examination(s), only the total number diagnosed in each arm of the trial.

As Deeks stated in his commentary on our paper, there was no reason to suspect non-random clustering of cases within wards. Allocation bias could have arisen only if mothers of high risk babies were systematically assigned to postnatal care on a ward practising one of the policies. As this changed every month, and assignment to a particular ward was done by the consultant at the booking clinic in early pregnancy, it is difficult to see how bias could have occurred. Furthermore, if we had randomised by individual baby, mothers in adjacent beds would have had different policies, which would have caused confusion in maintaining treatment allocation.

Cathryn M A Glazener *Clinical research fellow*
Health Services Research Unit, Aberdeen
AB25 2ZD
c.glazener@abdn.ac.uk

GMC's current proposals for revalidation are flawed

EDITOR—According to the General Medical Council, the object of revalidation is to establish the principle that "to maintain their registration, all doctors must be able to demonstrate regularly that they continue to be fit to practise in their chosen field."¹ In theory, revalidation has much to commend it, but the GMC's proposals for its implementation are flawed. They state: "local profiling of performance: evidence about the doctor's performance and continuing professional development" will be "gathered and reviewed locally" and that "doctors' pro-

files are likely to take into account the views of patients, colleagues and employers."

Local evaluation of a doctor's performance can seldom be completely unbiased, since pre-existing personal and social relationships between the doctor and his or her appraisers may falsely sway evaluations. There is a further risk of bias on the part of all three categories of appraiser. Patients may misjudge the calibre of their doctors (a pleasant and plausible but incompetent doctor may attract a favourable verdict, whereas a brusque but efficient doctor may be unjustly condemned). Colleagues may be reluctant to record an adverse verdict on less than competent doctors (who may be personal friends), while highly efficient but cantankerous doctors may not receive the favourable appraisal they deserve. There is an even greater risk of misappraisal by doctors' employers. Doctors who comply diligently with the directives of administrators will earn top marks, whereas those who constantly and openly inveigh against the bureaucratic folly of some employers may be blacklisted.

Revalidation, certainly when operated locally, is a misconceived and unworkable concept, which the GMC would be well advised to abandon. A more effective approach would be for the council, with government approval, to establish a statutory, easily understood, and confidential procedure for the submission of complaints against doctors from any source. It would investigate such complaints fairly and thoroughly, and, if they are proved, take much more rapid and decisive action against the offenders than it is usually prepared to do at present. Doctors found guilty of serious blunders or shortcomings would be punished by summary suspension of their registration.

A sanction such as summary suspension of doctors' registration, if rigorously enforced but safeguarded by a rapid and effective appeals procedure, is much more likely to contribute to an improvement in the standard of medical care than the cumbersome, expensive, and inequitable process of revalidation advocated by the GMC.

Ian W B Grant Retired consultant physician
Nether Balchandy, By Pitlochry, Perthshire
PH16 5JT

1 General Medical Council. Revalidation—the profession moves forward. *GMC News* 1999;5:1-4.

Cyclosporin neurotoxicity after chemotherapy

Case had features of thrombotic thrombocytopenic purpura and haemolytic uraemic syndrome

EDITOR—Tweddle et al's drug point described neurotoxicity due to cyclosporin after chemotherapy for lymphoproliferative disease after cardiac transplantation; the authors commented that it had not been reported in this setting previously.¹ The arti-

cle failed to mention thrombotic microangiopathy, which it is essential to consider in such patients.

Thrombotic thrombocytopenic purpura and the haemolytic uraemic syndrome are closely related conditions; they are associated with fever, thrombocytopenia, red cell fragmentation, and thrombotic lesions in various organs and result in neurotoxicity and renal impairment. The aetiology of thrombotic thrombocytopenic purpura and haemolytic uraemic syndrome is poorly understood, but it has been well described after treatment with immunosuppressive drugs including cyclosporin, chemotherapy, cancer, and bone marrow transplantation.²⁻⁵

The case described had features of thrombotic thrombocytopenic purpura and the haemolytic uraemic syndrome—namely, neurotoxicity, fever, and renal impairment with hypertension. No comment is made on the patient's full blood count, reticulocyte count, and lactate dehydrogenase concentration or whether fragmented cells were present in the blood film. Thrombocytopenia is likely, given that the events occurred six days after chemotherapy.

It is important to recognise thrombotic microangiopathy as plasma exchange is an effective treatment, although less so in patients who have undergone bone marrow transplantation. It is essential to consider the diagnosis in any patient with unexplained neurotoxicity who is taking cyclosporin.

Guy Pratt Research registrar in haematology
Ben Kennedy Specialist registrar in haematology
Graeme Smith Consultant haematologist
Department of Haematology, General Infirmary at Leeds, Leeds LS1 3EX

1 Tweddle DA, Windebank KP, Hewson QC, Yule SM. Cyclosporin neurotoxicity after chemotherapy. *BMJ* 1999;318:1113. (24 April.)

2 Bonser AS, Adu D, Franklin I, McMaster P. Cyclosporin induced haemolytic uraemic syndrome in liver allograft recipient. *Lancet* 1984;ii:1337.

3 Beaufils H, de Groc F, Gubler MC, Wechsler B, Le Hoang P, Baumelou A, et al. Hemolytic uraemic syndrome in patients with Behcet's disease treated with cyclosporin A: report of 2 cases. *Clin Nephrol* 1990;34:157-62.

4 Lesesne JB, Rothschild N, Erickson B, Korec S, Sisk R, Keller J, et al. Cancer-associated hemolytic-uraemic syndrome: an analysis of 85 cases from a national registry. *J Clin Oncol* 1989;7:781-9.

5 Pettitt AR, Clark RE. Thrombotic microangiopathy following bone marrow transplantation. *Bone Marrow Transplant* 1994;14:495-504.

Cyclosporin causes reversible posterior leukoencephalopathy syndrome

EDITOR—Tweddle et al describe a syndrome that is well recognised¹ and has been associated with several conditions, including hypertensive encephalopathy, pre-eclampsia, immunosuppression, rapid blood transfusion, and use of cyclosporin.² The common feature behind all of these conditions is usually a rapid and severe rise in blood pressure, although posterior leukoencephalopathy has been described with normotension.³ Hypertension is not mentioned in Tweddle et al's description of the initial event but occurred when they rechallenged the patient with cyclosporin.

The reversible posterior leukoencephalopathy syndrome is associated with headaches, vomiting, confusion, seizures, visual disturbance, and motor signs, in various

degrees. Neuroimaging typically shows bilateral, often strikingly symmetrical, white matter changes in the posterior regions of the cerebral hemispheres, particularly the occipital lobes and the posterior parietal lobes. The clinical and radiological findings usually resolve within two to three weeks if the underlying cause is removed.^{2,3}

I suspect that the case reported represents an expression of this syndrome through the mechanism of hypertension rather than a direct toxic effect of the interaction between cyclosporin and cyclophosphamide on neural tissue.

M B Lewis Specialist registrar in neurology
St James's University Hospital, Leeds LS9 7TF
m-k-lewis@email.msn.com

1 Tweddle DA, Windebank KP, Hewson QC, Yule SM. Cyclosporin neurotoxicity after chemotherapy. *BMJ* 1999;318:1113. (24 April.)

2 Hinchey J, Chaves C, Appignani B, Breen J, Pao L, Wang A, et al. A reversible posterior leukoencephalopathy syndrome. *N Engl J Med* 1996;334:494-500.

3 Ay H, Buonanno FS, Schaefer PW, Le DA, Wang B, Gonzalez RG, et al. Posterior leukoencephalopathy without severe hypertension. *Neurology* 1998;51:1369-76.

Authors' reply

EDITOR—Pratt et al suggest a possible diagnosis of thrombotic microangiopathy in relation to our recent Drug Point. The full blood count at the time of the seizures was haemoglobin concentration 93 g/l, white cell count $2.5 \times 10^9/l$ (neutrophils $1.9 \times 10^9/l$), and platelets $288 \times 10^9/l$; a blood film performed the next day was normal. In view of the normal platelet count and the absence of evidence of haemolysis in the blood film we did not request any biochemical investigations of haemolysis. We acknowledge that thrombotic microangiopathy can occur with cyclosporin neurotoxicity and after chemotherapy, but it was not apparent in this case.

The clinicoradiological features we described are consistent with the posterior leukoencephalopathy syndrome, a syndrome that is also associated with hypertensive encephalopathy. Lewis suggests that hypertension was the underlying mechanism for neurotoxicity in our case. Our patient did have a moderately raised systolic blood pressure of 130 mm Hg (previous baseline systolic blood pressure 110 mm Hg), both on admission with seizures and when rechallenged with cyclosporin. Cyclosporin neurotoxicity is often associated with mild to moderate hypertension, but this is unlikely to be the sole mechanism of neurotoxicity as cyclosporin itself has direct toxic effects on vascular endothelium.¹

We recently identified one previous report of cyclosporin neurotoxicity after doxorubicin chemotherapy for lymphoproliferative disease after transplantation.² In this case the authors proposed that an impaired blood-brain barrier from long term use of cyclosporin enabled doxorubicin to produce neurotoxicity. There are no previous reports of cyclosporin neurotoxicity with the chemotherapy regimen that we used.³ In our case we suggested that, despite normal concentrations of cyclosporin, its metabolism may have been affected by concomitant high dose chemotherapy (as evidenced by a raised

alanine aminotransferase concentration), leading to the accumulation of toxic metabolites; these could damage the blood-brain barrier and cause neurotoxicity.⁴

In view of the increasing incidence of malignancy related to immunosuppression, we reiterate the risk of continuing treatment with cyclosporin during chemotherapy.

Deborah A Tweddle *Research registrar in paediatric oncology*

D.A.Tweddle@newcastle.ac.uk

Kevin P Windebank *Senior lecturer*

Department of Paediatric Oncology, Institute of Child Health, Royal Victoria Infirmary NHS Trust and University of Newcastle, Newcastle upon Tyne NE1 4LP

- 1 Benigni A, Morigi M, Perico N, Zoja C, Amuchastegui CS, Piccinelli A, et al. The acute effect of FK506 and cyclosporine on endothelial cell function and renal vascular resistance. *Transplantation* 1992;54:775-80.
- 2 Barbui T, Rambaldi A, Parenzan L, Zucchielli M, Perico N, Remuzzi G. Neurological symptoms and coma associated with doxorubicin administration during chronic cyclosporin therapy. *Lancet* 1992;339:1421.
- 3 Tweddle DA, Gennery AR, Reid MM, Thomas JA, Burke M, Hamilton JRL, et al. Posttransplantation B lymphoblastic leukemia with Burkitt-like features. *Transplantation* 1999;67:1379-80.
- 4 Lucey MR, Kolars JC, Merion RM, Campbell DA, Aldrich M, Watkins PB. Cyclosporin toxicity at therapeutic blood levels and cytochrome P-450IIIa. *Lancet* 1990;335:11-5.

Devolution in Latin America has had poor effects on health care

EDITOR—The Editor's choice of 1 May expressed "excitement" at the possible effects of devolution on health care in the United Kingdom.¹ Had the experiences of some Latin American countries been examined in addition to those of Canada, Spain, and Finland, apprehension might have been a more appropriate sentiment.

Within Latin America, Chile, Brazil, Mexico, and Bolivia have all experimented with devolution of publicly funded health services to subnational government bodies. In these countries, as in the United Kingdom, the motives for decentralisation were political and so little consideration was given to the possible impact on the performance of health systems. In Mexico, Bolivia, and Brazil popular demands for greater democracy at all levels were instrumental in bringing about the changes. In Chile, where municipalities were given control of primary health care in 1981, a desire to elude the constraints of public service employment regulations seems to have been the main motivation.

Despite an almost universal feeling that decentralisation has to be a good thing, there is little evidence in Latin America that it leads to more locally appropriate services, innovation, greater accountability, community participation, or better management. Devolution in Bolivia, Mexico, and Brazil may have yielded greater equity in resource allocation through the introduction of various forms of capitation based funding, but this has not necessarily produced greater equity in use of health services or in health outcomes.

The richest states, provinces, and municipalities are able to top up central

government funding with their own revenues or from user charges. In the poorer regions, people are generally less healthy and less likely to have health insurance, so publicly funded services are more heavily burdened. The best administrative and technical expertise is available in the wealthiest areas and will serve deprived areas only if given attractive incentive packages.

Some other problems have been encountered. In Bolivia, devolution led to neglect of nationwide vertical disease control programmes with serious consequences, as exemplified by the world's first ever outbreak of urban yellow fever.² In Brazil, as in Finland,³ when responsibility for health care was devolved to small municipalities, economies of scale and risk pooling were lost. Though a logical solution would be to establish consortiums of municipalities, local politicians have been slow to forge any such accords.

Many of the problems associated with devolution can be avoided, but as the experience in Latin America shows, what it does to the health system is not a major concern for the politicians.

Peter Sandiford *Consultant*

Institute for Health Sector Development, London EC1V 9HL

- 1 Editor's choice. The excitement of devolution. *BMJ* 1999;318 (7192). (1 May.)
- 2 Van der Stuyf P, Gianella A, Pirard M, Cespedes J, Lora J, Peredo C, et al. Urbanisation of yellow fever in Santa Cruz, Bolivia. *Lancet* 1999;353:1558-62.
- 3 Koivusalo M. Decentralisation and equity of healthcare provision in Finland. *BMJ* 1999;318:1198-200. (1 May.)

Scientific foundation of mammographic screening is based on inconclusive research in Sweden

EDITOR—In a letter Rosén and Rehnqvist criticised our analysis of mammographic screening in Sweden.¹ We respond here to their criticism. The Swedish Board of Health and Welfare claims that mammographic screening has been shown to be effective in Swedish high quality clinical trials. A re-evaluation of these trials casts serious doubt on that conclusion.

A meta-analysis of five Swedish clinical trials showed a significant 24% reduction in mortality from breast cancer, but the control groups in four of the trials were invited to screening 4-5 years after randomisation.² The fifth trial showed an excess mortality in the first six years of mammographic screening.³ This finding could be of great importance as screening in the control group would lead to false "excess mortality" and explain the significant decrease in breast cancer in the study group in the meta-analysis.²

Rosén and Rehnqvist consider the two county trial to be one of the best mammographic screening trials. In both the 1985 and 1989 papers, however, it is repeatedly stated that "the control group (passive study population) was not invited to screening."^{4,5} In fact, screening started in the control group in 1982-3.² With this in mind,

can we stay assured that the positive findings in the statistical analysis of the trial are not the result of a "fishing expedition" whereby different ways of using the controls and of subgrouping the data were the means by which the magic $P < 0.05$ finally appeared? Interestingly, the design of the two county trial was not presented in the papers in 1985 or 1989. The original protocol stated that the trial should be a five year, controlled, and cluster randomised trial with two rounds of mammographic screening in the study group and no screening in the control group.

Because several assumptions critical to the reliability of a controlled clinical trial are not fulfilled, estimates of the efficacy of mammographic screening can therefore be based on the results of only one of the five Swedish trials—the Malmö trial. Those results are compatible with our analysis of the effects of mammographic screening in daily practice in 17 Swedish counties (that is, no reduction in mortality from breast cancer). As the meta-analysis of 1993 includes the two county trial, thereby accepting intervention in the control groups in violation of the original protocol, its conclusions are invalid and not reliable.²

Göran Sjönell *Family physician*

Kvartersakuten Matteus, Surbrunnsgatan, 66 SE 11327 Stockholm, Sweden
wonca@sjonell.se

Lars Ståhle *Assistant professor*

Department of Clinical Pharmacology, Karolinska Institute, Huddinge University Hospital, SE-14186 Huddinge, Sweden
Competing interests: None declared.

- 1 Rosén M, Rehnqvist N. Breast screening. *BMJ* 1999;318:809-10. (20 March.)
- 2 Nyström L, Rutqvist LE, Wall S, Lundgren A, Lindqvist M, Rydén S, et al. Breast cancer screening with mammography: overview of Swedish randomised trials. *Lancet* 1993;341:973-8.
- 3 Andersson I, Aspegren U, Janson L, Landberg T, Lindholm K, Linell F, et al. Mammographic screening and mortality from breast cancer: the Malmö mammographic screening trial. *BMJ* 1988;297:943-48.
- 4 Tabár L, Fagerberg CJG, Gad A, Baldertorp L, Holmberg LH, Grönroth O, et al. Reduction in mortality from breast cancer after mass screening with mammography. *Lancet* 1985;i:829-32.
- 5 Tabár L, Fagerberg G, Duffy SW, Day NE. The Swedish two-county trial of mammographic screening for breast cancer: recent results and calculation of benefit. *J Epidemiol Community Health* 1989;43:107-17.

Funding long term care for older people

Heirs will have to forgo their inheritance

EDITOR—Murphy makes several points about who the true beneficiaries of the suggested reforms of funding long term care would be.¹ They would be not disabled older people of modest means (who might benefit if more money were put into service improvements), poorly paid care workers in homes, or unpaid carers at home but the already well off heirs of older people with assets. Media attention has been focused on old people forced into selling their homes to pay for care. But sensible older people (or their representative) will sell their home on permanently entering care whatever the

financial arrangements for paying for the care; the alternative, often, is leaving it empty until their death (though when there is a co-resident spouse or other carer then his or her interests should be safeguarded).

As Murphy points out, the preserved capital is unlikely to be spent by the older person but will pass to heirs—who are likely to be well off already. Results from the 1994 round of the retirement survey (of people then aged 60-74) showed that 19% had inherited property.² Among those from social classes IV and V this proportion was only 7%, compared with 30% among professionals. Couples who had inherited property had average incomes of £306 a week, while those who had not had average incomes of £226 a week (some of this difference may of course be an effect of the inheritance).

In many cases heirs will not be direct descendants of the old person, as single and childless older people are strongly overrepresented among those in institutional care.³ It is worth noting that under the proposals of the Royal Commission on the Funding of Long Term Care a daughter or son who gives up a career to provide care for a disabled parent at home will not gain appreciably compared with a daughter or son whose parent is cared for in a nursing home.

What we need is a system that ensures that the minority of old people with disabilities that preclude independent living receive good quality care, either at home or in a communal setting. If funding such a system requires a generally affluent section of society to forgo some of their inheritances then so be it.

Emily Grundy *Reader in social gerontology*
Centre for Population Studies, London School of Hygiene and Tropical Medicine, London WC1E 7HT
e.grundy@lshtm.ac.uk

1 Murphy E. With respect to old age. *BMJ* 1999;318:681-2. (13 March.)

2 Disney R, Johnson P, Stears G. Coverage, levels and changes in financial assets. In: Disney R, Grundy E, Johnson P, eds. *The dynamics of retirement: analyses of the retirement surveys*. London: Stationery Office, 1997. (DSS research report No 72.)

3 Grundy E. Population review: the population aged 60 and over. *Popul Trends* 1996;84:14-20.

Taxes ensure equal quality care for all at time it is needed

EDITOR—There's something wrong and perplexing with the view that "Universal state funding ... would depress standards ... and impede the improvement of wages for poorly paid staff."¹ Visiting students from the United States tell me how much they admire the quality of care offered to people in local authority homes in the United Kingdom; they contrast the pride and dignity of the care staff with their own experiences in American nursing homes—the staff there are despised and suspected, the lowest of the low.² In our local homes, conditions for residents and certainly staff are usually better in the public than the private sector. Is there a parallel universe: "State bad; private good"?

Murphy's review of history is instructive, but her comprehension of current and future aspirations of the population in general, and

those who have entered old age in particular, is flawed. Robbing the richer who are in difficulty to pay for the poorer who are in similar difficulty is not an elegant model. It is not necessary and will not be accepted. We are a cohesive society and have more wealth available to us than did previous generations. We can make corporate provision for our predictable future needs, including nursing care. It is illogical to declare costs to be alarmingly expensive when sponsored through taxes but affordable if raised through personal insurance or loan schemes.

Distribution via taxes is unobtrusive, life-long, and to an agreed formula and delivers equal quality care for all at the time it is needed. Distribution by other systems is less assured, may be irregular, and is likely to skew higher quality of care toward subscribers to particular schemes (leaving the poor to make do with what they can get). Many good things began in 1948. Those whose vision gave us these starts would applaud the main recommendations of the Royal Commission on the Funding of Long Term Care³ and be pleased that we are, thanks to them, able to act on them.

I think that the royal commission has taken the visionary leap that Murphy calls for, but she hasn't seen it. Perhaps she was looking in the wrong direction, back to the 1980s.

David Jolley *Professor of old age psychiatry*
Penn Hospital, Wolverhampton WV4 5HN
DESSJOL@aol.com

1 Murphy E. With respect to old age. *BMJ* 1999;318:681-2. (13 March.)

2 Butler RN. *Why survive? Being old in America*. New York: Harper and Rowe, 1975.

3 Royal Commission on the Funding of Long Term Care. *With respect to old age: long term care—rights and responsibilities*. London: Stationery Office, 1999. (Cmnd 4192-1.)

Redistribution of wealth from rich to poor is not function of a care service

EDITOR—One of the purposes of a royal commission is to listen to what people have to say. In thousands of letters received by the Royal Commission on Long Term Care for the Elderly¹ older people expressed their sense of having been betrayed by society and by the government. Responding to the commission's report, Murphy fails to acknowledge this sense of betrayal.²

Most older people are fit and well, their health only minimally compromised. An important minority, however, are rendered frail and debilitated by chronic illness. Only these people must depend on others for personal care: being helped to the toilet, to bathe, and to dress. The need for this care is a direct consequence of illness and disease. It is not medical care but is undoubtedly health care.

For too long the medical profession has responded poorly to the needs of those for whom no effective treatment exists. This situation has now been compounded by the systematic exclusion of the healthcare needs of frail elderly people from health service provision. Most people on the royal commission were unable to find any moral justi-

fication for the continuation of this exclusion and recommended that intimate personal care of frail elderly people should be provided free on the basis of assessed need. The commission was clear that costs associated with daily living should remain the responsibility of the individual.

Like the dissenting minority of the royal commission, Murphy deplores the recommendation of free personal care on the basis that it will favour those who are better off and can afford to pay for it. But the redistribution of wealth from rich to poor is not the function of a care service; it is properly the function of the taxation system. The financial responsibility of a care service funded by the state is to provide universal inclusive liability for the potentially catastrophic costs that the misfortune of illness can incur. At present, frail older people suffer twice—through the effects of illness and disease and through being forced to pay for the care that their ill health makes necessary. This is rightly perceived as unjust by older people and those who care for them.

If those who assume positions of responsibility and influence in the NHS consider that universal inclusive liability "smacks of outdated paternalism" then what hope is there for meaningful social inclusion and the rebuilding of trust?

Iona Heath *General practitioner and member of Royal Commission on Long Term Care for the Elderly*
Caversham Group Practice, London NW5 2UP
pe31@dial.pipex.com

1 Royal Commission on the Funding of Long Term Care. *With respect to old age: long term care—rights and responsibilities*. London: Stationery Office, 1999. (Cm 4192-1.)

2 Murphy E. With respect to old age. *BMJ* 1999;318:681-2. (13 March.)

Magnetic resonance necropsy is offered routinely in University College London Hospitals

EDITOR—Bisset suggests that magnetic resonance imaging might be an alternative to necropsy.¹ The first public presentation of work specifically assessing the use of magnetic resonance imaging as an alternative to necropsy was given by two of us (JASB, WRL) in 1995 (abstract presented at Radiological Society of North America's scientific programme, Chicago, 1995). The first peer reviewed publication of this work appeared in 1996.² Further developmental work continues at University College London Hospital, London,³ and has been reported at several meetings.

Magnetic resonance necropsy is now offered as an established service in the imaging department of this hospital. At a time when rates of necropsy not requested by a coroner have fallen to less than 10% for adults and about 50% for infants who die in the perinatal period (the most prognostically important age group for necropsy) magnetic resonance imaging offers an acceptable alternative.⁴ Imaging diagnosis has become the established norm for the living (for example,

abdominal computed tomographic scanning has almost entirely replaced diagnostic laparoscopy). Surely it can no longer be denied to the dead, most of whose relatives find surgical necropsy unacceptable. By increasing the rate of uptake of postmortem examination in this way we may once again be in a position to derive postmortem information of epidemiological importance relevant to today's population.

We applaud Bisset's efforts and look forward to the formal publication of his results.

J A S Brookes *Consultant radiologist*
joebrookes@msn.com

M Hall-Craggs *Consultant radiologist*
W R Lees *Professor*
Department of Medical Imaging, University
College London Hospitals, Middlesex Hospital,
London W1N 8AA

- 1 Bisset R. Magnetic resonance imaging may be alternative to necropsy. *BMJ* 1998;317:145. (21 November.)
- 2 Brookes JAS, Hall-Craggs MA, Sams VR, Lees WR. Non-invasive perinatal necropsy by MRI. *Lancet* 1996;348:1139-41.
- 3 Deng J, Brookes JAS, Gardener JE, Rodeck CH, Lees WR. 3D MR imaging of the post-mortem fetal heart. *Fetal Diagn Ther* 1996;11:147-21.
- 4 Brookes JS, Hall-Craggs MA. Postmortem perinatal examination: the role of MRI. *Ultrasound Obstet Gynaecol* 1997;9:145-7.

Arguments in editorial were not "biologically implausible"

EDITOR—In a news article Woodman reports that "controversy about the contraceptive pill and its effect on the risk of breast cancer was rekindled" in an editorial that I wrote in the *Journal of Epidemiology and Community Health*.¹ The editorial was intended in part to reduce undue complacency about the safety of the pill. Ex cathedra statements by Professor John Guillebaud in Woodman's article—that my argument "is biologically implausible"—need to be well justified, since many people rely on him for advice on matters of contraception. Biological plausibility—an elusive concept, which keeps epidemiologists in business, of course—is never well served by unjustified assertions.

In this case, use of oral contraception could be an important part of a breast cancer initiation process without being an initiator itself, simply by speeding up cell proliferation rates at vulnerable times, whatever they may be.² This could happen with a wholly different potency before first term pregnancy than afterwards, and this is well supported by biological and epidemiological data. Firstly, the epidemiology of the effect of menarche and first pregnancy on the risk of breast cancer is already well understood.³ Secondly, the only large study to investigate the complex determinants of mitotic activity in resting breast tissue samples found a dramatic difference in the adjusted proliferation rate attributable to the oestrogen dose of current oral contraception, but only among nulliparous women.⁴ Among parous women no effect whatsoever was observed, and this difference was highly significant.

Cell proliferation rates are widely thought to be a strong determinant of the

risk of breast cancer.⁵ If they are then the latency problem is indeed only going to exist among users before first pregnancy and could be as long as 30 years. Such a postulate is not biologically implausible, and to say that it is misleading, particularly without evidence, and is likely to induce more undue complacency.

An important issue, of course, is that such "effect modifications" are usually difficult to detect because most studies lack sufficient power. When they are detected, as in this case—rather starkly—they should be heeded, not dismissed. But as Gillian Reeves says in Woodman's article, we certainly do need more data; meanwhile I repeat that we also need less illegitimate certainty.

Klim McPherson *Professor of public health epidemiology*
Cancer and Public Health Unit, London School of Hygiene and Tropical Medicine, London WC1E 7HT
k.mcpherson@lshtm.ac.uk

- 1 Woodman R. Controversy rekindled over pill and risk of breast cancer. *BMJ* 1999;318:1095. (24 April.)
- 2 Kelsey JL. Breast cancer epidemiology: summary and future directions. *Epidemiol Rev* 1993;15:256-65.
- 3 McPherson K, Steel CM, Dixon JM. Breast cancer—epidemiology, risk factors and genetics. In: Dixon JM, ed. *ABC of breast diseases*. London: BMJ, 1995:18-21.
- 4 Anderson TJ, Battersby S, King RJ, McPherson K, Going JJ. Oral contraception use influences resting breast proliferation. *Hum Pathol* 1989; 20:1139-44.
- 5 Preston-Martin S, Pike MC, Ross RK, Jones PA, Henderson BE. Increased cell division as a cause of human cancer. *Cancer Res* 1990;50:7415-21.

Surely doctors should follow the spirit, rather than the letter, of the law

EDITOR—Am I the only one who thinks, after reading the article and comments on retrieving semen from a dead patient,¹ that the world is going mad? We seem to be worshipping a goddess called Idealised Consent. The central theme seemed to be not the advisability of posthumous conception (which could well be debated) but the validity of the consent.

In the case quoted the husband had signed an authorisation that his sperm might be used after death. Play was made of the fact that this authorisation had not been witnessed, but was there any doubt as to the authenticity of the signature? The fact that the authorisation did not apply exactly to the circumstances of death seems to me to testify to its validity.

The dead patient had no objection to his corpse being mutilated in a good cause. He had already consented to his organs being given to strangers. It seems clear too that he both wanted to father children posthumously and was happy for his body to be operated on after his death. Nothing in the articles seems to question these statements. The argument that he might have been swayed by his cerebrovascular malformation seems weak—we may all have biological causes for some of the decisions we make.

The argument seems only to be about whether doctors should follow the spirit of the law or the letter. If, like lawyers, we

become trapped by the letter of the law, and not the spirit, it will be a sad day for our profession, and for our patients.

R Hugh James *Consultant anaesthetist*
Department of Anaesthesia, Leicester Royal Infirmary, Leicester LE1 5WW

- 1 Swinn M, Emberton M, Ralph D, Smith M, Serhal P, Bewley S, et al. Retrieving semen from a dead patient. *BMJ* 1998;317:1583-5. (5 December.)

Patients were more satisfied with chiropractic than other treatments for low back pain

EDITOR—Patient satisfaction is increasingly recognised as an important outcome in trials. Whatever interpretation Ernst and Assendelft place on changes in the Oswestry score, they did not mention in their letter¹ that significantly more of those patients in our trial who were treated by chiropractic expressed satisfaction with their outcome at three years than those treated in hospital²—84.7% (127/150) v 65.5% (76/116) for those referred by chiropractors ($P < 0.0001$) and 79.2% (103/130) v 60.2% (71/118) for those recruited from hospitals ($P = 0.001$). This aspect was assessed three years after entry to the trial.

The main stimuli to the Medical Research Council's current trial were the suggestive results and further questions raised by our initial observations. Assendelft et al's critique can hardly be considered to have been one of its starting points.³

T W Meade *Professor*
MRC Epidemiology and Medical Care Unit,
Wolfson Institute of Preventive Medicine, London EC1M 6BQ

- 1 Ernst E, Assendelft WJJ. Chiropractic for low back pain. *BMJ* 1999;318:261. (23 January.)
- 2 Meade TW, Dyer S, Browne W, Frank AO. Randomised comparison of chiropractic and hospital outpatient management for low back pain: results from extended follow up. *BMJ* 1995;311:349-51.
- 3 Assendelft WJJ, Bouter LM, Kessels AGH. Effectiveness of chiropractic and physiotherapy in the treatment of low back pain: a critical discussion of the British randomized clinical trial. *J Manipulative Physiol Ther* 1991;14:281-6.

Surveillance of congenital rubella in Great Britain

Rubella can be mistaken for parvovirus B19 infection

EDITOR—Tookey and Peckham are mistaken in stating that guidance says that all pregnant women with suspected symptoms of rubella or who have been exposed to the disease should be offered diagnostic investigation only in the first four months of pregnancy.¹ The Department of Health's current guidance places no such time limit on the need for appropriate investigation.²

The point is important because, without appropriate investigation, parvovirus B19 infection can be mistaken for rubella, and maternal parvovirus B19 infection is associated with fetal loss beyond the first four months of pregnancy.³

A J L Turner *Consultant virologist*
Newcastle Public Health Laboratory, Newcastle
General Hospital, Newcastle upon Tyne NE4 6BE
newaturn@newphls.demon.co.uk

- 1 Tookey PA, Peckham CS. Surveillance of congenital rubella in Great Britain, 1971-96. *BMJ* 1999;318:769-70. (20 March.)
- 2 United Kingdom Health Departments. *Immunisation against infectious disease*. London: HMSO, 1996.
- 3 Public Health Laboratory Service Working Party on Fifth Disease. Prospective study of human parvovirus B19 infection in pregnancy. *BMJ* 1990;300:1166-70.

Authors' reply

EDITOR—We acknowledge that rubella infection and parvovirus infection can be difficult to distinguish clinically. When there is little rubella about, suspected rubella does often turn out to be parvovirus infection, and this should certainly be considered if a woman or her close contact has a rubella-like illness at any stage in pregnancy.

Pat A Tookey *Coordinator of national congenital rubella surveillance programme*
Catherine S Peckham *Professor*
Department of Epidemiology and Public Health,
Institute of Child Health, London WC1N 1EH
P.Tooley@ich.ucl.ac.uk

Withdrawing low risk women from cervical screening programmes

Conclusions cannot yet be drawn

EDITOR—Sherlaw-Johnson et al evaluated policies for withdrawing women from the cervical cancer screening programmes before the recommended age of 64, using a mathematical model.¹ Their results were obtained with specific and uncertain model assumptions, which were insufficiently subjected to validation and sensitivity analysis.

From the description of the model in cited earlier papers, most new cases of cervical intraepithelial neoplasia seem to originate at younger ages. The duration is assumed to be independent of age and very long on average (50 years for cervical intraepithelial neoplasia grade III). This implies that most invasive cancers occurring over age 50 started as cervical intraepithelial neoplasia before age 50, which could thus be detected by screening before age 50. Hence this model is bound to predict only small increases in incidence when women are withdrawn from screening before the recommended age of 64.

The sensitivity analysis considers only small adaptations of this basic assumption. Other models, for which detailed analysis of screening data and data on the incidence of cancer was used, resulted in much lower estimates of the mean duration of cervical intraepithelial neoplasia.²⁻⁴ These models would predict less favourable effects of withdrawal policies.

Present data on human papillomavirus allow for widely different models, some of which are and some of which are not favourable for use in screening.⁵ Given this uncertainty, it is not yet possible to come to conclusions about the impact of withdraw-

ing women from cervical screening programmes if results of their smear test and a simultaneous test for high risk types of human papillomavirus are negative.

M E van den Akker-van Marle *Mathematician*
vanmarle@mgz.fgg.eur.nl

M van Ballegooijen *Epidemiologist*
R Boer *Statistician*

G J van Oortmarssen *Mathematician*

J D F Habbema *Professor of medical decision science*
Department of Public Health, Erasmus University,
PO Box 1738, 3000 DR Rotterdam, Netherlands

- 1 Sherlaw-Johnson C, Gallivan S, Jenkins D. Withdrawing low risk women from cervical screening programmes: mathematical modelling study. *BMJ* 1999;318:356-61. (6 February.)
- 2 Brookmeyer R, Day NE. Two-stage models for the analysis of cancer screening data. *Biometrics* 1987;43:657-69.
- 3 Gustafsson L, Adami HO. Natural history of cervical neoplasia: consistent results obtained by an identification technique. *Br J Cancer* 1989;60:132-41.
- 4 van Oortmarssen GJ, Habbema JD. Epidemiological evidence for age-dependent regression of pre-invasive cervical cancer. *Br J Cancer* 1991;64:559-65.
- 5 van Ballegooijen M, van den Akker-van Marle ME, Warmerdam PG, Meijer CJ, Walboomers JM, Habbema JD. Present evidence on the value of HPV testing for cervical cancer screening: a model-based exploration of the (cost-) effectiveness. *Br J Cancer* 1997;76:651-7.

Authors' reply

EDITOR—The authors of this letter criticise our assumptions about mean durations of cervical intraepithelial neoplasia, yet our paper makes no mention of such quantities; indeed, there is mathematical ambiguity about what this "mean" represents in the context of disease progression, which is why we have found it wise to avoid the term. Certainly, readers should not be given the impression that our paper is based on the nonsensical assumption that those who develop malignancy have had cervical intraepithelial neoplasia grade III for an average of 50 years. It is not. Our analysis uses assumptions about rates of progression of premalignancy. We have tried to tease out fact from opinion in the authors' letter, and the core criticism seems to be that the progression rates that we assumed are too low.

Ethics preclude direct observation of the rate at which cervical intraepithelial neoplasia grade III progresses when left untreated, and thus inference about transition rates relies on indirect methods. Notwithstanding the authors' high opinion of work investigating such rates, the fact is that little is known with precision, as we point out.

Given this, as reported, we carried out extensive sensitivity analysis to judge the robustness of the conclusions, and assumptions about progression rates were tested over a wide range of values. We have even considered the possibility of regression from cervical intraepithelial neoplasia grade III—an assumption some may find barely credible from a clinical viewpoint. This allowed us to investigate the effects of assuming increased rates of progression from cervical intraepithelial neoplasia grade III (with compensating increases in regression rates to maintain compatibility with United Kingdom data). The effects of this sensitivity analysis on the results reported in our paper are small.

The authors imply that our results favour a policy of early withdrawal. This is an inappropriate value judgment. We predict an increased risk of invasive cancer among women who are withdrawn early. As we have pointed out, the size of the predicted increase depends on several factors, such as the clinical course of disease in older women and the natural course of human papillomavirus infection, for which complete information is not available. The important question is whether these increases are justified by the resource savings and any other favourable effects of early withdrawal. This touches on ethical and pragmatic issues that are beyond the scope of our study.

Chris Sherlaw-Johnson *Senior research fellow*
c.sherlaw-johnson@ucl.ac.uk

Steve Gallivan *Professor of operational research*
Clinical Operational Research Unit, Department of
Mathematics, University College London, London
WC1E 6BT

David Jenkins *Reader in pathology*
Division of Pathology, University of Nottingham,
Queen's Medical Centre, Nottingham NG7 2UH

Frequent changes in policy risk confusion among health workers

EDITOR—The study by Usen et al¹ adds weight to their previous work,² with a larger sample and better blinding of investigators considering a wide range of clinical predictors of hypoxaemia. We question the practical importance of this work in terms of the World Health Organisation's current guidelines for community health workers.³ Their simple algorithm improves specificity, which would allow more efficient use of resources, albeit at the expense of sensitivity, the importance of which is debatable.

If the authors are suggesting that their chosen predictors should be incorporated into, or even replace, the World Health Organisation's guidelines for community health workers then we would question whether a change in these guidelines is justified. Retraining has substantial financial and practical costs. Furthermore, frequent changes in policy carry risks of confusion and disagreement among community health workers and may decrease their credibility in the community.

P Brett *Medical student*
L Guinness *Medical student*
L Lim *Medical student*
S Robb *Medical student*
Department of Epidemiology and Public Health,
University of Newcastle upon Tyne, Newcastle upon
Tyne NE2 4HH
Elizabeth.lim@ncl.ac.uk

- 1 Usen S, Weber MW, Mulholland EK, Jaffar S, Oparaugo A, Omosigbo C, et al. Clinical predictors of hypoxaemia in Gambian children with acute lower respiratory tract infection: prospective cohort study. *BMJ* 1999;318:86-91. (9 January.)
- 2 Weber MW, Usen S, Palmer A, Jaffar S, Mulholland EK. Predictors of hypoxaemia in hospital admissions with acute lower respiratory tract infection in a developing country. *Arch Dis Child* 1997;76:1-5.
- 3 World Health Organisation Programme for the Control of Acute Respiratory Tract Infections. *Acute respiratory tract infection in children: case management in small hospitals in developing countries*. Geneva: WHO, 1990. (WHO/ARI/90.5.)

High security facilities must not be closed until viable alternatives are in place

EDITOR—The future of special hospitals, including the possibility of closure, has been much debated.^{1,2} Although it is assumed that up to one third of the current 1400 patients could be managed in long term medium security,³ most of the research underpinning this assumption may be out of date.⁴ Special hospitals have tightened their admission criteria and have fewer elderly patients than they used to; these patients have been replaced by younger and more dangerous patients.

We conducted an interview and case note study of all 119 patients from two health authorities (Lambeth, Southwark and Lewisham, and Bexley and Greenwich) who were currently at special hospitals. We noted the views of their clinical team and their catchment area medium secure unit and made our own estimate based on Eastman's criteria (N Eastman, unpublished report to South Thames Executive Region).

The estimates of the research and clinical teams showed that between 30 and 39 patients did not require maximum security. Of these patients, 19 were already on trial leave or on a waiting list for an identified placement. The number of misplaced patients, for whom the system is not working, therefore ranged between 11 and 20.

Although delays inevitably occur, lack of a placement was the barrier to transfer for about half of all patients judged suitable for less secure care. The development of new places in long term medium security may allow an additional 9-17% of patients to move out of high security. This is a worthwhile objective but suggests that previous assumptions are overoptimistic and the potential for reducing the number of high security beds may be less than anticipated. It must also be remembered that the special hospitals often have long waiting lists. It is naive to expect that moving patients out would leave empty beds.

Patients have a right to be treated in the least secure conditions consistent with public safety. We should move out those who can be managed elsewhere, but closure is unrealistic except in the long term and should be driven by the provision of alternative facilities. Until viable alternatives are in place we cannot afford to lose the limited high security facilities we have. A pilot project is urgently needed in one or two areas to measure the reduction in the use of high security beds that follows the development of a long term medium secure unit.

Kapil Sayal *Specialist registrar*
Maudsley Hospital, London SE5 8AF
k.sayal@iop.kcl.ac.uk

Anthony Maden *Senior lecturer*
Department of Forensic Psychiatry, Institute of Psychiatry, London SE5 8AZ

1 Basson JV. Fate of special hospitals in England has not yet been decided. *BMJ* 1998;316:1243.

- 2 Murphy E. The future of Britain's high security hospitals. *BMJ* 1997;314:1292.
3 Thake T, Jobbins C, Jones D. *High Security Psychiatric Services Commissioning Board needs assessment report phase 1*. London: HSPSCB, 1998.
4 Maden A, Curle C, Meux C, Burrow S, Gunn J. *Treatment and security needs of special hospital patients*. London: Whurr, 1995.

Annual general meeting of the BMA

BMA or UKMA?

EDITOR—Unless second thoughts prevail the BMA's annual general meeting on 7 July will be asked to make a grievous and unjust error. It will be proposed that our 4000-odd overseas members (some of those in the armed forces) be disenfranchised and deprived of their entitlement to membership benefits. Why and how did this come about?

Apart from four overseas branches our members are scattered worldwide, from China to Peru. Some of them have sought help from BMA House on "terms of service" and the like. While this is run of the mill stuff for our staff when sent in by members from the United Kingdom they have neither the expertise, nor time, to cope with queries from Argentina or Zimbabwe. What could be done? Someone last summer thought up the special resolution as an answer: reduce them all to second class citizens without a vote, and then they can importune in vain.

Having decided this, our council did not notify or consult the overseas branches. It also failed to instruct our lawyers to look for the knock on effects of the changes on other articles and byelaws. As a result, if the resolution is passed, for 12 months at least anomalies and doubts will give BMA House more trouble than the original problem. Just two examples will do: no overseas member will be obliged to pay a subscription, and none who is now a fellow may remain so.

The council did not think the proposal merited a mention in its annual report, so it remained a secret until the AGM notice in the *BMJ* of 5 June—four weeks before the decision was to be taken.¹ What a way to treat the membership: a total denial of natural justice—think of the reaction in our Hong Kong branch, already under pressure to disband.

If the resolution is put and passed, then surely our overseas membership will wither away. Why remain as a second class member when one can subscribe to the *BMJ* at a lower rate as a non-member?

The resolution is neither urgent nor necessary. There are alternative ways of amending the articles to cope with the problem; a nut requires no sledgehammer. I have suggested such alternatives to BMA House but so far have had no reply. So, unless there is a change of heart by then, I urge all present at the AGM to refer the resolution back to the council for reconsideration.

David Gullick *Former deputy secretary of the BMA*
1 Heathbrow Road, Welwyn, Herts AL6 0GQ

1 BMA's annual general meeting. *BMJ* (clinical research) 1999;318:1523.

Reply from the BMA

EDITOR—I welcome the opportunity to reply to Gullick's letter. I have believed for a long time that overseas members of the BMA were an undervalued resource for the association. We need to find out from them what range of services is appropriate and hope that this will lead to a substantial increase in their numbers. The council is proposing to the annual general meeting to put in place the foundation on which a new relationship between the BMA and its overseas members may be built. The BMA is not in a position to provide trade union services for its overseas members—at most it can point them in the direction of organisations which may be able to assist them. In that respect Gullick is in error in suggesting that the amendments to the articles will create a hiatus.

E M Armstrong *Secretary of the BMA*
BMA House, London WC1H 9JP

Iceland's medical database is insecure

EDITOR—Ragnarsson states that personally identifiable data cannot be linked to medical data held in the proposed Icelandic central database, as "it is made impossible through encryption."¹ This is not so. I evaluated the design of the database on behalf of the Icelandic Medical Association, and it became clear that the proposal to encrypt personal identity numbers into pseudonyms was inadequate.²

Longitudinal record linkage means that the encryption function cannot change over time, so if someone can enter a record into the computer system of an Icelandic hospital or general practice and then observe its "de-identified" version appearing in the central database, they can deduce the linkage between the patient's identity and the pseudonym. Even if pseudonyms could not be deduced directly, de-identified databases suffer from the intrinsic limitation that many patients can be identified from partial information about their circumstances, and this may be particularly problematic in a database that will contain genealogical data as well as medical records.

Doctors in Britain should take note of Iceland's problems when considering, for example, whether to construct central databases of episodes of illness in patients with HIV infection or AIDS that are identified with encrypted NHS numbers.

Ross Anderson *Lecturer*
Cambridge University Computer Laboratory,
Cambridge CB2 3QG
rja14@cl.cam.ac.uk

1 Ragnarsson G. Opposition to the Icelandic database is based on false information *BMJ* 1999;318:1354. (15 May.)
2 Anderson R. The DeCODE proposal for an Icelandic health database. *Icelandic Medical Journal* 1998;84:874-5.

Rapid responses

Correspondence submitted electronically
is available on our website www.bmj.com