Understanding how the brain reorganises itself may eventually facilitate the treatment of patients with intractable phantom pain. Ramachandran, for example, has proposed a novel technique for the relief of the cramp-like pain caused by a clenched phantom fist. By observing the reflection of their free moving normal hand in a mirror, so that it appears in the position of the phantom, some patients report an immediate and compelling illusion whereby they are able to generate voluntary movements in the phantom hand. In some cases, the sensation of unclenching the phantom hand relieves the pain.⁵ The efficacy of this method will

require a prospective controlled trial, but it offers the prospect of a real clinical benefit from an exciting insight into how the brain organises itself.

Peter W Halligan *MRC senior research fellow*

Department of Experimental Psychology, University of Oxford, Oxford OX1 3UD

Adam Zeman *consultant neurologist*

Department of Clinical Neurosciences, University of Edinburgh, Western General Hospital, Edinburgh EH4 2XU

Abi Berger *science editor, BMJ*

- 1 Mitchell SW. *Injuries of nerves and their consequences*. Philadelphia: Lippincott, 1872:348.
- 2 Aglioti S, Cortese F, Franchini C. Rapid sensory remapping in the adult brain as inferred from phantom breast perception. *Neuroreport* 1994;5:473-6.
- 3 Halligan PW, Marshall JC, Wade DT. Three arms: a case study of supernumerary phantom limb after right hemisphere stroke. *J Neurol Neurosurg Psych* 1993;56:159-66.
- 4 Weinstein S. Neuropsychological studies of the phantom. In: Benton AL, ed. *Contributions to clinical neuropsychology* Chicago: Aldine, 1969:73-107.
- 5 Ramachandran VS, Hirstein W. The perception of phantom limbs. *Brain* 1998;121:1603-30.
- 6 Ramachandran VS, Stewart M, Rogers-Ramachandran DC. Perceptual correlates of massive cortical reorganization. *Neuroreport* 1992;3:583-6.

- 7 Halligan PM, Marshall LC, Wade DT, Davey J, Morrison D. Thumb in cheek? Sensory reorganization and perceptual plasticity after limb amputation. *Neuroreport* 1993;4: 233-6.
- 8 Kew JJM, Halligan PW, Marshall JC, Rothwell JC, Ridding MC, Sooriakumaran S, et al. Abnormal access of axial vibrotactile input to deafferented somatosensory cortex in human upper limb amputees. *J Neurophysiol* 1997;77:2753-64.
- 9 Jones EG, Pons TP. Thalamic and brainstem contributions to large scale plasticity of primate somatosensory cortex. *Science* 1998;282:1121-5.
- 10 Sadato N, Pascual-Leone A, Grafman J, Ibanez V, Deiber MP, Dold G, et al. Activation of the primary visual cortex by Braille reading in blind subjects. *Nature* 1996;380:526-8.

Pituitary surgery for acromegaly

Should be done by specialists

The objective of treatment in acromegaly is not only to alleviate symptoms, improve quality of life, and prevent complications but also to reverse the increased (two to threefold) mortality from vascular and respiratory disease and colon cancer. That this can be achieved if "target" basal growth hormone concentrations of <5 mU/l (2.5 ng/ml) are reached has been shown,^{1,2} with the proviso that serum insulin-like growth factor 1 (IGF-1) values may need to be reduced to the normal age-related reference range.³ Transsphenoidal adenomectomy is the most rapid means of reducing serum growth hormone and IGF-1 in acromegaly. The question then arises of how effective surgery is at achieving the target growth hormone and IGF-1 values. Subsidiary issues are: (a) what determines the outcome of surgery? (b) what is the relapse rate? (c) what is the complication rate? and (d) what is the most cost effective service provision?

Surgical outcomes in terms of safe target growth hormone concentrations have recently been reported in over 1000 patients.³⁻¹¹ All these series show that large tumour size, together with the degree of extrasellar extension (particularly lateral), and high preoperative serum growth hormone concentrations are major determinants of initial surgical failure. Thus, in 1999 a patient with a microadenoma (<1 cm in diameter) or mesoadenoma (>1 cm but confined to the sella turcica) should have upwards of 80% chance of successful adenomectomy. However, the surgical outcome for those with a macroadenoma (>1 cm with extension outside the sella) is by no means so successful, with at best 60% achieving the safe target growth hormone concentration. However, even for macro-

adenomas it is important to reduce the serum growth hormone level as much as possible, since the outcome of adjuvant treatment with somatostatin analogues, dopamine agonists, or radiotherapy is directly related to the pretreatment growth hormone concentration.⁸

What is new in the recent reports is the demonstration that the outcome of surgical treatment for acromegaly is related to the experience of the operator. In the study from Manchester,⁸ where 73 patients were operated on by any of nine surgeons, fewer than 20% of patients achieved a growth hormone <5 mU/l, an outcome the authors attribute to the few cases operated on by any one surgeon. Likewise, the success rate in Birmingham was also low at 33%,⁵ where eight surgeons were involved. However, in Oxford, Newcastle, or St Bartholomew's Hospital, London (each of which had only one surgeon performing pituitary surgery), a patient had a chance of a successful adenomectomy of 42-56%.^{4,6,10} Interestingly, Yamada et al in Japan report a doubling of the success rate when a single surgeon started operating on their patients with acromegaly (from 37% to 81%).⁷ Similarly, the proportion of patients achieving the target basal growth hormone concentration of <5 mU/l in Birmingham doubled (from 33% to 64%) when a single surgeon took over all operations.¹¹ The Oxford authors also report improvement in results of a single surgeon as his experience grew over 20 years.¹⁰

The relapse rate 5-10 years after successful initial surgery is low, at 0-7%, and major complication rates (leaks of cerebrospinal fluid, meningitis, permanent diabetes insipidus, new pituitary hypofunction) are also low, at less than 10%. Thus, in experienced centres

BMJ 1999;319:588-9

adenomectomy for small adenomas secreting growth hormone is effective and safe and has low relapse and complication rates. However, even in the best hands some 15-20% of microadenomas and 40-50% of macroadenomas will not achieve safe growth hormone concentrations after initial surgery and will require additional long term adjunctive treatment (radiotherapy or medical treatment). Since adjunctive treatment is expensive, and may be required for many years, it is clearly in everyone's interests for people with acromegaly to be operated on by experienced pituitary surgeons. It is difficult to measure cost of treatment, but the cost of the most effective medical treatment, somatostatin analogues, alone is around £12 000 per patient per year.

Endocrinologists are normally responsible for advising patients with acromegaly. Since this is usually pituitary surgery first they should choose the "best" surgeon, even if this means referring outside their immediate local service. Not only will this be best for the patient but it will also result in the most effective use of NHS resources since fewer patients will require adjuvant treatment. Pressure must be exerted to put in

place a system whereby many fewer centres and surgeons in the United Kingdom operate on patients with acromegaly and other pituitary tumours. Where this has happened, in Japan⁷ and in Birmingham,¹¹ the effect has been dramatic.

How many centres should there be? Our own view is that the minimum population that should be served is 5 000 000, which would produce 20-30 new cases of acromegaly a year. With two surgeons per centre this would mean no more than 10 centres in the UK.

R N Clayton *professor of medicine*

Keele University and North Staffordshire Hospital, Stoke on Trent ST4 7QB

P M Stewart *professor of medicine*

Queen Elizabeth Hospital, Birmingham B15 2TH

S M Shalet *professor of endocrinology*

Christie Hospital, Manchester M20 4BX

J A H Wass *professor of endocrinology*

Radcliffe Infirmary, Oxford OX2 6HE

- 1 Bates AS, van't Hoff W, Jones JM, Clayton RN. An audit of outcome of treatment in acromegaly. *Quart J Med* 1993;86:293-9.
- 2 Orme SM, McNally RJQ, Cartwright RA, Belchetz PE. Mortality and cancer incidence in acromegaly: a retrospective cohort study. *J Clin Endocrinol Metab* 1998;83:2730-4.
- 3 Swearingen B, Barker FG, Katznelson L, Biller BMK, Grinspoon S, Klibanski A, et al. Long-term mortality after transphenoidal surgery and adjunctive therapy for acromegaly. *J Clin Endocrinol Metab* 1998;83:3419-26.
- 4 Osman IA, James RA, Chatterjee S, Mathias D, Kendall-Taylor P. Factors determining the long-term outcome of surgery for acromegaly. *Quart J Med* 1994;87:617-23.
- 5 Jenkins S, O'Brien I, Johnson A, Shakespear R, Sheppard MC, Stewart PM. The Birmingham pituitary database: auditing the outcome of the treatment of acromegaly. *Clin Endocrinol* 1995;43:517-22.
- 6 Sheaves R, Jenkins P, Blackburn P, Huneidi AH, Afshar F, Medbak S, et al. Outcome of transphenoidal surgery for acromegaly using strict criteria for surgical cure. *Clin Endocrinol* 1996;45:407-13.
- 7 Yamada S, Aiba T, Takada K, Ozawa Y, Shimizu T, Sawano S, Shishiba Y, Sano T. Retrospective analysis of long-term surgical results in acromegaly: preoperative and postoperative factors predicting outcome. *Clin Endocrinol* 1996;45:291-8.
- 8 Lissett CA, Peacey SR, Laing I, Tetlow L, Davis JRE, Shalet SM. The outcome of surgery for acromegaly: the need for a specialist pituitary surgeon for all types of growth hormone secreting adenomas. *Clin Endocrinol* 1998;49:653-7.
- 9 Abosch A, Tyrell JB, Lamborn KR, Hannegan LT, Applebury CB, Wilson CB. Trans sphenoidal microsurgery for growth hormone secreting pituitary adenomas: initial outcome and long-term results. *J Clin Endocrinol Metab* 1998;83:3411-18.
- 10 Ahmed S, Elsheikh M, Page RCL, Adams CBT, Wass JAH. Outcome of transphenoidal surgery for acromegaly and its relationship to surgical experience. *Clin Endocrinol* 1999;50:561-7.
- 11 Gittoes NJL, Johnson AP, Sheppard MC, Stewart PM. Outcome of surgery for acromegaly: the experience of a single surgeon. *Quart J Med* (in press).

British guidelines on managing hypertension

Provide evidence, progress, and an occasional missed opportunity

Education and debate
p 630

This issue of the *BMJ* includes a summary of the new British Hypertension Society guidelines for managing hypertension.¹ Recently, one of the authors of these guidelines commented on the strengths of the fifth and sixth reports of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure (JNC): "The JNC reports score highly on two counts—they took note of important new evidence, and unlike the others, delivered a clear message."² The same can be said of this version of the British guidelines, which represent an outstanding summary of the existing evidence about the diagnosis and treatment of hypertension. In contrast to the World Health Organisation-International Society of Hypertension³ and the Joint National Committee VI guidelines,⁴ the brevity of this version will make the British Hypertension Society guidelines especially useful to clinicians.

The authors aimed to produce an evidence based document, and in the full version (reference 1 in ¹) they applied the North of England Group criteria to grade

the evidence for most recommendations. For instance, a low dose thiazide diuretic is listed as the first line treatment for patients with uncomplicated hypertension. This evidence based recommendation, appropriately, received a grade A rating, which signifies evidence from at least one randomised trial or from a meta-analysis of randomised trials.

Implicit in the recommendations about drug choice is a distinction not often made in the traditional evidence based grading systems. For the main drug treatment recommendation the authors have relied on the results of randomised clinical trials that assess major disease end points rather than surrogate end points such as level of blood pressure.⁵ All drugs approved for treating hypertension lower blood pressure. But the purpose of treatment is to reduce the risk of devastating hypertensive complications such as myocardial infarction, stroke, and heart failure. Few antihypertensive drugs, only low dose diuretics and β blockers, have consistently passed this higher standard of evidence—one which is appropriate for public

BMJ 1999;319:589-90