Website: bmj.com
Email: letters@bmj.com

Authors' reply to Camelford letters

EDITOR—Many of the comments of David, Esmonde, McMillan, and Murray et al are incorrect and overlap.¹ Owing to space constraints we have addressed points of fact below and opinion based comments in bmj.com (bmj.com/cgi/eleletters/320/7245/1337#EL1) or in the printed² and longer web versions of our paper (bmj.com/cgi/content/full/319/7213/807/DC1).

Few normative data are published on large series of flash or pattern stimulated visual evoked potentials or the difference in timing between them. In a given subject the flash-pattern difference might be large because of a quicker (shorter) pattern latency than normal, but, to our knowledge, pathologically short pattern evoked responses have not been reported in the world literature. As in our studies, most of the many studies on dementia report that only flash-pattern difference, rather than absolute values, correlates with severity.

We acknowledged and addressed the problem of self selection bias. David's belief that our data "are totally inadequate" is unfounded. Our methods were clear and have been published with illustrative waveforms.³ We never used the term flash minus pattern evoked latency, we did not give absolute latencies for the 55 Camelford subjects, and the 28 cases quoted in his table were part of a separate methodological quality control study.

David and Murray et al incorrectly refer to the results of Sloan and Fenton⁴ and imply that depression may have led to our observations on visual evoked potentials. The flash-pattern difference increases with age, and Sloan and Fenton's patients were all over 65 (mean 70), considerably older than our subjects. David has not transcribed Sloan and Fenton's results correctly in his table.

Sloan and Fenton found that mean flash-pattern differences at the start of the longitudinal study were 37.7 (SD 23.3) ms in normal controls, 33.7 (21.1) ms in depressed patients, and 45.0 (25.4) ms in those with Alzheimer's disease. The only significant results were that the Alzheimer's group had larger flash-pattern differences than controls and that regression coefficients with time of flash latency and flash-pattern difference were significant in Alzheimer's disease. We could find only two other English language references to visual evoked potentials in depression, and both showed no significant change.^{5,6}

Esmonde is wrong to deny the many published reports of visual evoked potential abnormalities in Alzheimer's disease, some of which were referred to in our paper. Controversy exists about aluminium and Alzheimer's disease but not about the neurotoxicity of aluminium. How it leads to these effects is not clear, but it may not be solely through neuronal death. We chose unexposed siblings to provide genetic (phenotypic) controls, a standard way of establishing acquired rather than inherited abnormalities. Only 15 such siblings were available—it would have been wrong not to proceed, and the paper was refereed by a medical statistician.

We were concerned about the possible effects that publication might have on people involved in the Camelford incident, but it would have been wrong to withhold it. Had our studies not been designed to provide scientifically useful information (despite the inevitable limitations) and been used only to satisfy the legal process, the clinical and scientific communities would have been deprived of interesting results that are valid and worthy of debate.

Paul Altmann consultant nephrologist
Oxford Kidney Unit, Oxford Radcliffe Hospital,
Oxford OX3 7LJ

John Cunningham consultant nephrologist and
physician

Frank Marsh consultant nephrologist and physician
The Royal London Hospital, London E1 1BB

Usha Dhanesha principal optometrist
Paybody Eye Unit, Coventry and Warwickshire
Hospital, Coventry CV1 4FH

Margaret Ballard consultant clinical psychologist
Priory Hospital, London SW15 5JJ

James Thompson senior lecturer in psychology
University College London Medical School,
London W1N 8AA

1 Correspondence. Cerebral dysfunction after water pollution incident in Camelford. *BMJ* 2000;320:1337-8. (13 May.)

2 Altmann P, Cunningham J, Dhanesha U, Ballard M, Thompson J, Marsh F. Disturbance of cerebral function in people exposed to drinking water contaminated with aluminium sulphate: retrospective study of the Camelford water incident. *BMJ* 1999;319:807-11. (25 September.)

3 Altmann P, Dhanesha U, Hamon C, Cunningham J, Blair J, Marsh F. Disturbance of cerebral function by aluminium in haemodialysis patients without overt aluminium toxicity. *Lancet* 1989;ii:7-12.

4 Sloan EP, Fenton GW. Serial visual evoked potential recordings in geriatric psychiatry. *Electroenceph Clin Neurophys* 1992;84:325-31.

5 Jordan SE, Nowacki R, Nuwer M. Computerized electroencephalography in the evaluation of early dementia. *Brain Topogr* 1989;1:271-82.

6 Swanwick GR, Rowan M, Coen RF, O'Mahony D, Lee H, Lawlor BA, et al. Clinical application of electrophysiological markers in the differential diagnosis of depression and very mild Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 1996;60:82-6.

Drug use has declined among teenagers in United Kingdom

EDITOR—During 1995 a survey of illicit drug use was conducted among teenagers in Europe. The United Kingdom and 22 other countries participated.^{1,2} British teenagers reported the highest rates of drug use. The survey was repeated in 1999.

The population sampled in 1999 consisted of 15 and 16 year old students attending state and private schools in the United Kingdom. Information was elicited from 1280 boys and 1361 girls in 223 schools.

A third of girls and 39.5% of boys had used illicit drugs. A total of 36.5% had used cannabis. Glues and solvents had been used by 18.6%, lysergide by 5.2%, amphetamines by 8.9%, and ecstasy (methylenedioxymethamphetamine) by 5.0% (tables 1 and 2).

There were several regional variations. Students in Scotland were more likely than others to have used cannabis, any illicit drug, and amphetamines. Respondents in Scotland and Northern Ireland were more likely than others to have used ecstasy and heroin. Those in Northern Ireland reported the greatest use of glues and solvents.

Most forms of drug use had declined since 1995. Girls showed significant reductions in their use of cannabis, any illicit drug, solvents, lysergide, amphetamines, "pills" combined with alcohol, ecstasy, and tranquillisers. Among boys the significant falls were in cannabis, any illicit drug, solvents, lysergide, amphetamines, "pills" combined with alcohol, ecstasy, and crack cocaine. The only exception to this trend was heroin, whose use, though still rare, had risen in both sexes.

These reductions may be temporary, but they are striking.

Martin Plant director
mplant@ahrc.onet.co.uk

Patrick Miller senior research fellow
Alcohol and Health Research Centre, City Hospital,
Edinburgh EH10 5SB

We thank Steve Chalmers, Janet Docherty, Björn Hibell, Barbro Andersson, and the Pompidou Group for their help. This study was mainly supported by the Alcohol Education and Research Council, the Department of Health and Social Security, Belfast, and the Health Education Authority. Support was also provided by Allied Domecq, the PF Charitable Trust, the North British Distillery Company, the Sir James Miller (Edinburgh) Trust, and the Drapers' Fund.

1 Miller P, Plant MA. Drinking, smoking, and illicit drug use among 15 and 16 year olds in the United Kingdom. *BMJ* 1996;313:394-7.

2 Hibell B, Andersson B, Bjarnason T, Kokkevi A, Morgan M, Narusk A. The 1995 ESPAD report: alcohol and other drug use among students in 26 European countries. Stockholm, Swedish Council for Information on Alcohol and Other Drugs, 1997.

Girls

Table 1 Proportions (percentages; 95% confidence intervals) of girls aged 15 and 16 using illicit drugs. Percentage for all regions are weighted

	England	Northern Ireland	Scotland	Wales	All regions in 1999	All regions in 1995
Cannabis (No of times):						
1-39	87/319 (27.3; 25.1 to 29.5)	115/410 (28.0; 23.1 to 33.3)	152/514 (29.6; 26.3 to 33.9)	30/108 (27.8; 18.2 to 37.4)	384/1351 (27.7; 24.2 to 31.2)	1525/4009 (38.0; 35.1 to 40.9)*
≥40	13/319 (4.1; 2.2 to 6.3)	7/410 (1.7; 0.0 to 6.8)	36/514 (7.0; 4.6 to 9.2)	7/108 (6.5; 1.8 to 11.2)	63/1351 (4.4; 2.7 to 6.1)	
Type of illicit drug ever used:						
Any	103/319 (32.3; 26.9 to 37.7)	126/412 (30.6; 25.3 to 35.9)	198/522 (37.9; 31.5 to 44.4)	38/108 (35.2; 24.2 to 46.2)	465/1361 (33.0; 28.9 to 37.2)	1628/4092 (39.8; 37.1 to 42.5)
Any by injection	2/318 (0.6; 0.0 to 1.5)	3/411 (0.7; 0.0 to 1.7)	3/520 (0.6; 0.0 to 1.4)	0/107	8/1356 (0.6; 0.0 to 1.3)	25/3987 (0.6; 0.2 to 1.0)
Glues and solvents	49/318 (15.4; 11.4 to 19.4)	96/410 (23.4; 18.9 to 28.2)	19.5 (102/522; 14.4 to 24.7)	24/105 (22.8; 17.0 to 31.2)	271/1355 (16.9; 13.8 to 20.0)	847/4031 (21.0; 18.9 to 23.1)
Lysergide or other hallucinogens	9/318 (2.8; 1.0 to 4.6)	17/412 (4.1; 1.6 to 6.7)	28/521 (5.4; 2.7 to 8.0)	6/107 (5.6; 1.3 to 9.8)	60/1358 (3.4; 1.9 to 4.8)	489/4017 (12.2; 10.2 to 14.2)
Amphetamines	22/319 (6.9; 3.8 to 10.0)	18/412 (4.4; 1.7 to 7.1)	59/522 (11.3; 7.0 to 15.6)	10/107 (9.3; 4.0 to 14.5)	109/1360 (7.4; 5.0 to 9.8)	493/4015 (12.3; 10.5 to 14.1)
"Pills" combined with alcohol	37/313 (11.8; 7.9 to 15.7)	65/407 (16.0; 11.3 to 20.6)	64/513 (12.5; 8.9 to 16.0)	19/107 (17.8; 8.3 to 27.2)	185/1340 (12.5; 9.4 to 15.5)	1014/4016 (25.3; 22.9 to 27.7)
Ecstasy†	8/319 (2.5; 0.8 to 4.2)	26/410 (6.3; 3.4 to 9.2)	27/520 (5.2; 2.7 to 7.7)	5/107 (4.7; 0.7 to 8.6)	66/1356 (3.2; 1.9 to 4.5)	293/3999 (7.3; 5.9 to 8.7)
Tranquillisers or sedatives	8/319 (2.5; 0.5 to 4.5)	15/412 (3.6; 1.8 to 5.5)	25/522 (4.8; 2.1 to 7.4)	8/107 (7.5; 2.4 to 12.6)	56/1360 (3.1; 1.7 to 4.5)	380/4010 (9.5; 7.9 to 11.1)
Cocaine	13/319 (4.1; 2.0 to 6.2)	8/411 (1.9; 0.5 to 3.4)	11/521 (2.1; 0.5 to 3.7)	5/107 (4.7; 0.3 to 9.0)	37/1358 (3.7; 2.1 to 5.3)	95/4016 (2.4; 1.9 to 2.9)
Crack cocaine	8/318 (2.5; 0.9 to 4.1)	2/411 (0.5; 0.0 to 1.2)	6/521 (1.1; 0.1 to 2.2)	2/107 (1.9; 0.0 to 4.1)	18/1357 (2.2; 0.9 to 3.4)	89/4013 (2.2; 1.7 to 2.7)
Steroids	6/316 (1.9; 0.4 to 3.4)	4/411 (1.0; 0.0 to 2.0)	4/521 (0.8; 0.0 to 1.7)	2/106 (1.9; 0.0 to 4.2)	16/1354 (1.7; 0.5 to 2.8)	40/3987 (1.0; 0.6 to 1.4)
Heroin	10/319 (3.1; 1.3 to 5.0)	28/411 (6.8; 3.8 to 9.8)	28/521 (5.4; 2.8 to 7.9)	5/107 (4.7; 0.7 to 8.6)	71/1358 (3.7; 2.3 to 5.2)	61/4007 (1.5; 1.0 to 2.0)
"Relevin" (dummy drug)	0/319	1/409 (0.2; 0.0 to 0.8)	0/518	0/105	1/1351	12/4004 (0.3; 0.1 to 0.5)

*Cannabis ever.

†Methylenedioxymethamphetamine.

Boys

Table 2 Proportions (percentages; 95% confidence intervals) of boys aged 15 and 16 using illicit drugs. Percentages for all regions are weighted

	England	Northern Ireland	Scotland	Wales	All regions in 1999	All regions in 1995
Cannabis (No of times):						
1-39	100/331 (30.2; 24.6 to 34.9)	92/311 (29.6; 21.1 to 38.0)	159/510 (31.1; 27.4 to 34.8)	21/122 (17.2; 9.5 to 24.9)	372/1274 (29.3; 25.3 to 33.3)	1546/3546 (43.6; 41.0 to 46.2)*
≥40	27/331 (8.2; 4.8 to 11.8)	27/311 (8.7; 0.0 to 17.8)	72/510 (14.1; 11.1 to 17.1)	12/122 (9.8; 3.7 to 16.0)	138/1274 (9.1; 6.4 to 11.9)	
Type of illicit drug ever used:						
Any	132/336 (39.3; 33.7 to 44.8)	123/311 (39.5; 30.0 to 49.1)	234/511 (45.8; 40.0 to 51.6)	33/122 (27.1; 16.0 to 38.1)	522/1280 (39.5; 35.1 to 43.9)	1636/3630 (45.0; 42.6 to 47.4)
Any by injection	3/332 (0.9; 0.0 to 1.9)	2/308 (0.6; 0.0 to 1.6)	2/509 (0.4; 0.0 to 1.1)	0/120	7/1269 (0.8; 0.0 to 1.5)	26/3555 (0.7; 0.4 to 1.0)
Glues and solvents	43/335 (12.8; 9.3 to 16.3)	86/311 (27.7; 21.1 to 34.2)	70/511 (13.7; 9.1 to 18.3)	20/121 (16.5; 9.2 to 23.6)	219/1278 (13.8; 11.0 to 16.6)	705/3587 (19.7; 17.6 to 21.8)
Lysergide or other hallucinogens	16/334 (4.8; 2.0 to 7.5)	22/309 (7.1; 3.6 to 10.5)	32/511 (6.3; 3.2 to 9.3)	7/121 (5.8; 1.9 to 9.5)	77/1275 (5.1; 2.9 to 7.3)	606/3574 (17.0; 14.6 to 19.4)
Amphetamines	23/333 (6.9; 4.0 to 9.7)	19/308 (6.1; 2.5 to 9.7)	71/509 (13.9; 9.0 to 18.8)	12/121 (9.9; 3.6 to 16.1)	125/1271 (7.8; 5.5 to 10.2)	517/3566 (14.5; 12.5 to 16.5)
"Pills" combined with alcohol	30/326 (9.2; 5.4 to 13.0)	27/306 (8.8; 4.9 to 12.7)	40/500 (8.0; 5.1 to 10.9)	8/119 (6.7; 2.9 to 10.6)	105/1251 (8.9; 5.9 to 11.9)	499/3567 (14.0; 12.4 to 15.6)
Ecstasy†	9/333 (2.7; 0.9 to 4.5)	20/306 (6.5; 3.4 to 9.4)	30/506 (5.9; 3.2 to 8.9)	5/121 (4.1; 0.7 to 7.5)	64/1266 (3.4; 1.9 to 4.8)	326/3555 (9.2; 8.1 to 10.3)
Tranquillisers or sedatives	18/334 (5.4; 2.6 to 8.2)	20/308 (6.5; 3.6 to 9.3)	33/509 (6.5; 3.4 to 9.5)	4/121 (3.3; 0.0 to 6.7)	75/1272 (5.5; 3.2 to 7.7)	244/3556 (6.9; 5.8 to 8.0)
Cocaine	9/333 (2.7; 1.1 to 4.3)	6/309 (1.9; 0.0 to 3.9)	14/509 (2.7; 0.9 to 4.6)	2/121 (1.6; 0.0 to 3.7)	31/1272 (2.6; 1.3 to 3.9)	101/3571 (2.8; 2.2 to 3.4)
Crack cocaine	3/332 (0.9; 0.0 to 1.9)	4/309 (1.3; 0.0 to 3.1)	7/508 (1.4; 0.2 to 2.5)	3/121 (2.5; 0.0 to 4.9)	17/1270 (1.1; 0.3 to 1.8)	98/3569 (2.7; 2.1 to 3.3)
Steroids	8/332 (2.4; 0.7 to 4.1)	7/307 (2.3; 0.5 to 4.0)	5/508 (1.0; 0.0 to 2.2)	3/121 (2.5; 0.0 to 6.7)	23/1268 (2.3; 0.9 to 3.6)	80/3546 (2.2; 1.5 to 2.9)
Heroin	9/334 (2.7; 0.9 to 4.5)	24/309 (7.8; 4.5 to 11.0)	31/509 (6.1; 3.3 to 8.8)	5/121 (4.1; 0.7 to 7.5)	69/1273 (3.4; 2.0 to 4.9)	62/356 (1.7; 1.2 to 2.2)
"Relevin" (dummy drug)	1/331 (0.3; 0.0 to 0.9)	1/309 (0.3; 0.0 to 1.0)	1/508 (0.2; 0.0 to 0.7)	1/121 (0.8; 0.0 to 2.3)	4/1269 (0.3; 0.0 to 0.8)	8/3560 (0.2; 0.1 to 0.3)

*Cannabis ever.

†Methylenedioxymethamphetamine.

Unconventional approaches to nutritional medicine

Conventional doctors need more insight into nutritional medicine ...

EDITOR—Nutrition has become a popular topic of public discussion, but the conventional medical community has largely been caught wearing no clothes. In his electronic response to Vickers and Zollman's article on nutritional medicine in the ABC of complementary medicine,¹ Stargrove points out that somehow nutrition fell "out of the medical bag."² Commercial health care simply grew with the industrial economy. Now we are experiencing waves of more loudly expressed alternative medicine, which recognises a fundamental centrality in nutrition.

It seems to me that this complex discussion is in many ways a cover story for an enormous, often unspoken, public discourse on religion, politics, personal identity, group process, and public and private angst or joy. The many issues addressed by the authors¹ unwrap briefly some of the pieces. But the authors do not seem to do more than read the names of the cities from the map. They do not address where we are trying to go or what we might achieve by trying to do there.

Nutritional insights, therapeutics, and protocols offer genuine healthcare benefits for literally millions of people who presently lack at least some portion of the wherewithal to achieve them. Vickers and Zollman's article seems unconcerned with this and is instead drawn to the subject as if it were reviewing arcane fashions.

Ned Hoke private practitioner in ecological and oriental medicine
158 West Napa Street, Sonoma, CA, USA 94924
drhoke000@aol.com

1 Vickers A, Zollman C. ABC of complementary medicine: Unconventional approaches to nutritional medicine. *BMJ* 1999;319:1419-22. (27 November.)

2 Stargrove M. Nutrition—part of every medicine. *eBMJ* 1999;319 (www.bmj.com/cgi/letters/319/7222/1419#EL1).

... and can now get training in it

EDITOR—Vickers and Zollman's article on nutrition as complementary medicine¹ did not recognise the increasingly important contribution that nutrition is making to conventional clinical practice, creating a demand for training and education in nutrition. This has long been a neglected aspect of the undergraduate curriculum: many of today's doctors were not taught the scientific principles on which nutritional therapy is based. This deficiency is now being addressed by a joint intercollegiate initiative: representatives of 11 medical royal colleges have been working together since 1996 to improve postgraduate education and training in nutrition for medical graduates.

A programme of four-day intercollegiate courses has now been developed.² These courses may well be the first postgraduate exposure to nutrition that many medical graduates have, assuming that their only formal training has been at an undergraduate level (based on the principles set out in the

core curriculum).³ The intercollegiate course therefore represents a link in the vertical thread of training, as set out in *Tomorrow's Doctors*.⁴ It clarifies the general principles of human nutrition, including the range of nutritional requirements in health and disease throughout the life cycle, and the assessment and management of nutritional abnormalities, both undernutrition and overnutrition. This provides a systematic approach to the solution of practical problems. Particular attention is paid to the evidence base for applied nutrition.

There are special challenges in carrying out any intercollegiate activity, where the point of reference is those considerations that are common and draw specialties together. Our first experience has shown that nutritional science forms a suitable focus for enhancing cross disciplinary understanding. The Intercollegiate Group on Nutrition plans to hold the course three times a year at centres in England and Scotland. This should ultimately ensure that trainees, general practitioners, and consultants will have access to a suitable local course.

In contrast to the fairly narrow registration requirements and training suggested in complementary medicine,¹ the committee believes that nutritional considerations must be part of the overall specialty training of doctors and should be based on a strong scientific foundation of nutritional principles. Ultimately, clinical nutrition may have to be recognised as a subspecialty with its own pathway of postgraduate training.

A Shenkin chairman, Intercollegiate Group on Nutrition
Department of Clinical Chemistry, University of Liverpool, Liverpool L69 3GA
shenkin@liverpool.ac.uk

The intercollegiate group includes representatives of the Royal Colleges of Anaesthetists, General Practitioners, Obstetricians and Gynaecologists, Paediatrics and Child Health, Pathologists, Physicians (London, Edinburgh), Physicians and Surgeons (Glasgow), Psychiatrists, and Surgeons (England, Edinburgh), and British Dietetic Association.

1 Vickers A, Zollman C. ABC of complementary medicine: Unconventional approaches to nutritional medicine. *BMJ* 1999;319:1419-22. (27 November.)

2 Shenkin A. The Intercollegiate Group on Nutrition progress report. *Bull R Coll Pathol* 2000;109:39. (www.rcpath.org/activities/icgnutrition.html)

3 Department of Health. *The health of the nation. Nutrition for medical students; nutrition in the undergraduate curriculum*. London: DoH, 1996.

4 General Medical Council. *Tomorrow's doctors*. London: GMC, 1993.

Summary of rapid responses

Vickers and Zollman's article on unconventional approaches to nutritional medicine aroused much interest, and we received 12 responses on our website in addition to the letters published here.¹ They were from both doctors working in hospital and alternative practitioners, and five were an online argument between J Wedderburn and Vickers over whether veganism should be classified as restrictive. The general question debated in the responses was whether doctors should know more about nutrition than they do; the general view was that they should.

bmj.com

M Stargrove, a naturopathic physician, says that he looks "toward the day when every physician sees foods, herbs, and supplements as valuable tools in their therapeutic repertoire. The notion that such nutritive and therapeutic interventions are somehow 'alternative' strikes me as a historical accident that needs to be rectified."

K M Jacobie, who is in private practice providing massage therapy, says that "it seems to me that the most popular medical belief is that just about every disease is caused by a drug deficiency, and ultimately to whom is this belief most 'valid and useful'? Follow the profit for the answer. The facts suggest that there is just no 'money' in telling people that a lifetime of excellent nutrition could mean a lifetime of better health, so why learn about it?"

A Sali and L Vitetta, doctors from a university in Australia, say that the article "further illustrates that the 'regular' doctor is incomplete in medical training. This has significant implications for health and the future of conventional medicine." They point out that "the trend of available funds has been toward patented pharmaceutical products, ... the net result being a deficiency in the promotion ... of nutritional medicine products whose natural occurrence deems them impossible to patent." They quote Hippocrates: "Let your food be your medicine and let your medicine be your food."

H Farley, a patient who has rheumatoid arthritis, writes that an elimination diet showed her which foods she should avoid and that when she avoided them the disease was greatly improved. This was despite her rheumatologist having said that "diet played no role in the disease."

E C G Grant, a physician, says that "it is to be deplored that modern biochemical nutritional testing is not generally available to all doctors. Accurate analyses clearly demonstrate that deficiencies of essential nutrients ... are widespread in the general population. ... Following an exclusion diet is a rapid, drug-free way of lowering blood pressure and preventing headaches and migraine attacks. ... Until accurate biochemical nutritional testing becomes generally available, the medical profession will remain in the dark. Meanwhile dissatisfied patients will continue to seek out possibly dubious alternative practitioners."

1 Electronic responses. Unconventional approaches to nutritional medicine. *bmj.com* 1999;319 (www.bmj.com/cgi/content/full/319/7222/1419). (Accessed 25 February)

Child discipline

Weak evidence for a smacking ban

EDITOR—A ban of a medical intervention would never be supported on the basis of such meagre evidence as used by Waterston to support a ban of the parental intervention of smacking.¹ "Significant adverse effects" and a failure to "learn the desired behaviour" were based on a literature review that is unpublished² and that includes studies that included severe types of

corporal punishment such as "beating with a stick," "still hurt the next day," "burning," and "using a knife or gun." Most studies that were reviewed were cross sectional, which cannot disentangle the causal direction between smacking and child misbehaviour.²

In the only published review (in 1996) of child outcomes of non-abusive or customary physical punishment, only eight studies could disentangle the causal effects of smacking.³ All eight studies, including four randomised clinical trials, found that non-abusive smacking benefited children when it backed up milder disciplinary tactics with children aged 2 to 6 years.

Smacking, then, makes milder tactics more effective, not "harder to use" as concluded by Waterston.¹

Another study was cited to conclude that Swedish "public opinion on the need for physical punishment changed dramatically after a public education campaign" following the 1979 smacking ban.⁴ The so called dramatic change was artificially created because survey questions from before 1982 and from 1994 were compared. The 1994 survey question that was most similar to the previous question showed an increased endorsement of mild or moderate physical punishment as sometimes necessary—from 26% in both 1978 and 1981 to 34% in 1994.⁵ The 1994 Swedish survey also found that corporal punishment of teenagers was as prevalent after the 1979 ban as in previous generations and that, overall, the incidence of corporal punishment had decreased little.⁵

Consequently, the British proposal for a middle ground between the status quo and a 100% smacking ban is reassuring. As Waterston noted, parents are already motivated to find alternatives to smacking, and positive interaction between parents and children and enhancing appropriate child behaviours are good places to start. The most difficult puzzle for parents and professionals concerns effective methods for decreasing misbehaviour.

Eighteen studies in the 1996 review investigated alternative disciplinary tactics as well as smacking.³ Only grounding was more effective than smacking, in two studies of older children. In contrast, nine alternatives were associated with more detrimental outcomes in children than was smacking.

Parents need to be empowered with more effective alternatives, not disempowered by premature bans on traditional disciplinary tactics.

Robert E Larzelere *psychologist*
Psychology Department, Munroe-Meyer Institute,
University of Nebraska Medical Center, Omaha, NE
68198-5450, USA
rlarzelere@unmc.edu

1 Waterston T. Giving guidance on child discipline. *BMJ* 2000;320:261-2. (29 January.)

2 Gershoff ET. *The effects of parental corporal punishment on children: a process model and meta-analytic review*. Tempe, AZ: Arizona State University, 1999.

3 Larzelere RE. A review of the outcomes of parental use of nonabusive or customary physical punishment. *Pediatrics* 1996;98:824-8.

4 Durrant JE. *The status of Swedish children and youth since the passage of the 1979 corporal punishment ban*. London: Save the Children, 1997.

5 Sanden A. *Spanking and other forms of physical punishment: a study of adults' and middle school students' opinions, experience, and knowledge*. Stockholm: Statistics Sweden, 1996.

Parents need techniques for behavioural control

EDITOR—The art of child rearing is a complex process in which the outcome of a parent's efforts is influenced by many factors unique to the child, the parent, the environment, and the context. Waterston's editorial promoting a ban on all disciplinary physical punishment does not respect this complexity and oversimplifies the debate over a parent's use of spanking (smacking).¹

As a participant in the American Academy of Pediatrics consensus conference, I would like to clarify some of its findings. The group's goal was to develop consensus statements regarding the scientific evidence on the long term and short term effects of corporal punishment on children. Definitions were the first order of business for the group: corporal punishment was defined as "bodily punishment of any kind"; spanking was defined as "physically non-injurious, intended to modify behavior, and administered with the open hand to the buttocks or the extremities." Using strict definitions prevented the common mistake of mixing abusive physical punishment with non-injurious spanking. With these definitions, however, the committee could not reach any strong conclusions favouring or opposing a parent's use of disciplinary spanking for children aged 2-11 years.

Central to the conference was the exhaustive review of the literature on corporal punishment presented by clinical psychologist Robert Larzelere. He found stronger evidence of beneficial than detrimental effects of non-abusive spanking by parents with preschool children (aged 2 to 6 years). The conference chairpersons concluded: "Given a relatively 'healthy' family life in a supportive environment, spanking in and of itself is not detrimental to a child or predictive of later problems ... there is a lack of research related to the use of corporal punishment."²

Developmental research indicates that optimal outcomes in children result from an authoritative style of parenting that combines positive encouragement with consistent behavioural control of the young child.³ Waterston describes the process of encouragement well, but leaves parents short-handed on techniques for behavioural control. Young children need correction and punishment, but this is often ignored by the opponents of physical punishment. Time out and disapproval are effective tools but are not sufficient to control all problem behaviour with all children. Disciplinary spanking, when properly applied, can augment nonphysical measures and optimise the process of behavioural control. To remove spanking from the repertoire of parents of young children could promote child abuse and lead to increased violence among older, unruly children. This seems, from statistics, to be an effect that the Swedish ban has had on that society.

I urge the makers of public policy in the United Kingdom to move slowly and scientifically in analysing this issue. An in

depth analysis of this subject can be found at www.frc.org/fampol/fp96jpa.html.

Den A Trumbull *paediatrician*
4700 Woodmere Blvd, Montgomery, AL 36106, USA
trumbulld3@aol.com

1 Waterston T. Giving guidance on child discipline. *BMJ* 2000;320:261-2. (29 January.)

2 Friedman SB, Schonberg SK, Sharkey M, eds. The short and long term consequences of corporal punishment. *Pediatrics* 1996;98(suppl):857-8.

3 Baumrind D. The development of instrumental competence through socialization. *Minnesota Symp Child Psych* 1973;7:3-46.

Occasional smacking does no harm

EDITOR—In his article saying that smacking children is wrong, Waterston shows himself to be a caring paediatrician who wishes to speak up for children in what he sees as an important ethical issue.¹ However, his article shows the pitfalls of representing a viewpoint that is based on emotion as a considered, evidence based approach.

The evidence cited by Waterston has shown that smacking has adverse effects only when it is excessive. There is no evidence that occasional smacking is harmful, and indeed it would be astonishing if there were, given the trivial nature of the physical and psychological event. There are even theoretical reasons to suppose that smacking may be less harmful than some alternative strategies. It is, after all, quickly over and avoids protracted emotional withdrawal ("I won't love you if you're naughty") which, for many parents, is the alternative.

It can be inferred from Waterston's article that he acknowledges that a consistent and measured strategy of discipline that happens to include smacking is likely to benefit rather than to harm a child. So why does he oppose it so strongly? Perhaps because his experience is that for many parents smacking is not a measured or consistent strategy. Instead, it is a last resort when control is lost. It is this element of unpredictable, irrational, and potentially uncontrolled violence that is dangerous in smacking, rather than the smack itself.

By definition, this element will not be influenced by changing the law. In recommending legislation against smacking, Waterston ensures that chaotic, uncontrolled smacking will continue, and that only measured smacking, which does no harm and may even help a child understand discipline, will stop. In other words, he will have achieved the reverse of his intention.

There is an urgent need to support parents in developing parenting skills. Agitating, with little evidence and less logic, for the criminalisation of smacking does little to help in this endeavour. It does, however, do much to undermine the credibility of our profession in trying to advocate for children in this area.

Richard D W Hain *senior lecturer in paediatric palliative care*
Llandough Hospital, Penlan Road, Penarth
CF64 2XX

1 Waterston T. Giving guidance on child discipline. *BMJ* 2000;320:261-2. (29 January.)

Parents must be in charge of their children

EDITOR—Waterston articulates the arguments for restraint and possible legislation on the physical punishment of children.¹ Why then, as a child psychiatrist and family therapist who espouses and promotes these ideas daily in my professional practice, and as a father with two children under 5 against whom my wife and I have never raised a hand (though sorely tempted), do I still demur? Because I do not wish the state to be able to intervene where it should not. Similar arguments apply to child sexual abuse and seatbelt legislation. Both are now “state-intervenable” matters. Why is smacking different?

The best I can come up with (and it may not be intellectually rigorous) is that I also believe, professionally and personally, that it is important for parents to be in charge of their children. In this sense the use of physical punishment is qualitatively different from sexual abuse. There is a benign aim in the use of punishment that cannot be said to exist in child sexual abuse, even though all the evidence shows that there are better ways to achieve the aim. We must let parents parent in their own individual ways, even if that may sometimes involve physical chastisement. Even if children are given the power to be able to sue their parents for such chastisement, I am not encouraged to believe that this would produce a more benign environment for that child or, more importantly for public policy, for all children. Indeed this prospect may only lead to the inversion of the perhaps politically incorrect but bald fact that parents need to be more powerful than children in the hierarchical structure that we call family. At the very least parental power represents the best we have yet come up with for successful child rearing, in the early years at least. Public education (Waterston cites the example of Sweden) may be a more effective route to change.

Let us hope that the debate and consequent changing of minds continues, but let us not try to impose values that may not lead where we want to go. Perhaps the government has got it right.

Richard Fry consultant child and family psychiatrist
Child Family and Adolescent Consultation Service,
Uxbridge UB8 1BN

1 Waterston T. Giving guidance on child discipline. *BMJ* 2000;320:261-2. (29 January)

Author's reply

EDITOR—A key part of the case against a ban on corporal punishment is the desire to distinguish between injurious and non-injurious smacking. In legal terms this is not possible, as a smack on any part of the body is potentially injurious, whatever the intention.

Larzelere states that smacking has beneficial outcomes, on the evidence of his review published in 1996.¹ These outcomes were short term only and in artificial situations. Of the eight studies he mentions that showed benefit of non-abusive smack-

ing, five were laboratory studies by the same team,² the children were in an extreme group for disordered behaviour, and only immediate outcomes were assessed. Parents were trained to give two spansks while being observed by a therapist, and it would not be wise to draw general conclusions in relation to the population of smacking parents. The authors state that spanking was not a superior method of discipline. A sixth study was on one extremely disordered child with no control subject,³ and in the remaining studies in which parents used structured diaries to observe the effects of their own management of their children within specific parameters, the authors stated that “to use this research as general evidence supporting punitive parenting or corporal punishment outside these parameters would be totally inappropriate.”

In relation to the survey of the effects on public opinion of the ban on smacking in Sweden, the question used was identical in a series of four surveys conducted between 1965 and 1981.⁵ The data are as follows: in 1965, 53% of respondents to the survey believed that corporal punishment is necessary in child rearing; in 1968 the percentage was 42%; in 1971 it was 35%; and in 1981 it was 26%.

In 1994, a different question was used: Are you positively inclined toward physical punishment, even in its mildest forms? Eleven per cent of Swedes answered this affirmatively.

Sweden's rate of fatal child abuse is low. Between 1976 and 1990, no child in Sweden died as a result of abuse. Between 1990 and 1996, four children died from the effects of physical abuse; only one of these children was killed by a parent.

Trumbull attempts to distinguish between abusive physical punishment and non-injurious spanking. I do not believe that this is practicable in law. The increasing trend in reports of youth assaults in Sweden is at least partly attributable to increased enforcement.⁵

I agree with Hain that there is no evidence that occasional smacking is harmful, and that it provides a model to the child. Is the practice of violence by parents against a small child a good model to learn from?

Fry believes that parents need to be “more powerful than children.” Surely this should be through intellectual rather than muscular strength. The Swedish experience does not lead us to expect that children will start suing their parents.⁵ A change in the law following an educational campaign would set a marker that violence in the family is not to be condoned.

Tony Waterston consultant paediatrician
Community Paediatric Department, Newcastle
General Hospital, Newcastle upon Tyne NE4 6BE

- 1 Larzelere RE. A review of the outcomes of parental use of non-abusive or customary physical punishment. *Pediatrics* 1996;98:824-8.
- 2 Roberts MW, Powers SW. Adjusting chair timeout reinforcement procedures for oppositional children. *Behav Ther* 1990;21:257-71.
- 3 Bernal ME, Duryee JS, Pruett HL, Burns BJ. Behaviour modification and the brat syndrome. *J Consult Clin Psychol* 1968;32:447-55.

4 Larzelere RE, Schneider WN, Larson DB, Pike PL. The effects of discipline responses in delaying toddler misbehaviour recurrences. *Child Fam Behav Ther* 1996;18:35-57.

5 Durrant JE. Evaluating the success of Sweden's corporal punishment ban. *Child Abuse Negl* 1999;23:435-48.

Treatment of *Helicobacter pylori* infection

Development of resistance to antibiotics used must be avoided

EDITOR—We agree with de Boer and Tytgat that treatment to eradicate *Helicobacter pylori* in patients with proved peptic ulcer is cost effective and benefits the patient and society.¹ Treatment to eradicate the organism in patients with non-ulcer dyspepsia, however, is contentious. The efficacy, safety, and cost effectiveness are far from proved. Several recent trials, including a multicentre double blind placebo controlled trial, failed to show any significant long term benefit of *H pylori* eradication treatment in non-ulcer dyspepsia.²⁻⁵

In our hospital 12 patients with dyspepsia who had previously received treatment from their general practitioners to eradicate *H pylori* on the basis of positive serological results were found to have gastric carcinoma. In several cases the delay in referring the patients to a gastroenterologist was considerable (unpublished observations).

Perhaps the most compelling reason to desist from empirical or indiscriminate antibiotic treatment of non-ulcer dyspepsia is the recent observation that commensal oral and bowel flora rapidly develop resistance to antibiotics (clarithromycin, amoxycillin, and metronidazole) used in *H pylori* eradication treatment. The emergence and persistence of resistant strains was most pronounced in patients treated with clarithromycin.⁴ If this resistance then spreads to pathogenic bacteria these antibiotics will cease to be useful in a variety of infections. The development of such resistance will far outweigh any perceived benefits of antibiotic treatment of non-ulcer dyspepsia.

G Gopal Rao consultant microbiologist
gopal.rao@uhl.nhs.uk

J O'Donohue consultant gastroenterologist
C S Mahankali Rao research coordinator, microbiology department

H Fidler consultant gastroenterologist
University Hospital Lewisham, London SE13 6LH

Competing interests: None declared.

- 1 De Boer WA, Tytgat GNJ. Treatment of *Helicobacter pylori* infection. *BMJ* 2000;320:31-4. (1 January)
- 2 Blum AL, Talley NJ, O'Morain C, van-Zanten SV, Labenz J, Stolte M, et al. Lack of effect of treating *Helicobacter pylori* infection in patients with nonulcer dyspepsia. Omeprazole plus Clarithromycin and Amoxycillin Effect One Year after Treatment (OCAY) Study Group. *N Engl J Med* 1998;339:1875-81.
- 3 Talley NJ, Janssens J, Lauritse K, Racz I, Bolling-Sternevald E. Eradication of *Helicobacter pylori* in functional dyspepsia: randomised double blind placebo controlled trial with 12 months' follow up. *BMJ* 1999;318:833-7.
- 4 Adamsson I, Nord CE, Lundquist P, Sjöstedt S, Edlund C. Comparative effects of omeprazole, amoxycillin plus metronidazole versus omeprazole, clarithromycin plus metronidazole on oral, gastric and intestinal microflora of *Helicobacter pylori* infected patients. *J Antimicrob Chemother* 1999;44:629-40.

Quadruple treatment may be solution

EDITOR—De Boer and Tytgat review the regimens used for *Helicobacter pylori* infection and comment correctly that we lack large randomised controlled trials to compare different treatments.¹ A general finding, however, has been that earlier enthusiasm is not borne out by later experience; probably the old two week triple regimen of bismuth, tetracycline, and metronidazole should join dual treatment and be regarded as obsolete. One week's quadruple treatment with bismuth, tetracycline, and metronidazole plus proton pump inhibitor is more effective and halves the load of the rather toxic bismuth, tetracycline, and metronidazole component which makes it so hard for patients to take.

We should aim to use treatments with a 90% or better success rate per protocol. Even a proton pump inhibitor plus amoxycillin plus metronidazole does not always reach this in modern studies, which report 85-90% cures, and perhaps this regimen should be reserved for second line treatment.

One alternative that has been effective locally is quadruple treatment with lansoprazole 30 mg daily, tetracycline 500 mg twice daily, clarithromycin 250 mg twice daily, and metronidazole 400 mg twice daily for one week.² This has achieved a 95.3% success rate (negative result of a urea breath test one to two months after treatment) in 213 consecutive patients with duodenal ulcers at endoscopy and a positive result of a direct urease test immediately before treatment. Interestingly, all of the first 66 patients had negative results of breath tests, which emphasises the need for larger studies to assess effectiveness accurately. This regimen has been used locally for four years, and a recent audit has shown that the antibiotic sensitivity of *H pylori* to the drugs used was exactly the same in 1999 as it was in 1992-6. Results were a little better (and early duodenal ulcer healing rates may be expected to be superior) when lansoprazole was used for a month rather than a week. I commend this scheme for general use in treatment.

M C Bateson consultant physician
South Durham Health Care NHS Trust, Bishop Auckland General Hospital, Bishop Auckland, County Durham DL14 6AD

Competing interests: None declared.

1 De Boer WA, Tytgat GNJ. Treatment of *Helicobacter pylori* infection. *BMJ* 2000;320:31-4. (1 January.)

2 Bateson MC, Diffey BL. Improving management of duodenal ulcer disease. *Postgrad Med J* 1997;73:717-9.

Triple regimen based on ranitidine bismuth citrate could be solution

EDITOR—One of the key messages of de Boer and Tytgat's paper on the treatment of *Helicobacter pylori* infection is that triple regimens combining clarithromycin and metronidazole should not be used as there is no valid empirical back up regimen after failure.¹ This recommendation is based on the increasing evidence that antibiotic resistance has become the main factor affecting negatively the efficacy of treatment to eradicate

H pylori. De Boer and Tytgat's suggestion, however, is in contrast with the guidelines produced by the Maastricht consensus meeting and similar conferences worldwide. The Maastricht meeting recommended the regimens based on the two antibiotics as the best available first line treatments for eradicating *H pylori*.²

De Boer and Tytgat suggest adopting a triple treatment combining a proton pump inhibitor or ranitidine bismuth citrate with amoxycillin and clarithromycin as first line treatment in areas with low primary prevalence of resistance to clarithromycin. But the association of these two antibiotics showed on average a lower eradication rate than the combination of clarithromycin and metronidazole.³ The statement that the quadruple regimen containing metronidazole performs well in almost all populations should be more cautious, because a significant negative impact of resistance to metronidazole on the efficacy of this complex scheme has been found.⁴

In recent years a new triple treatment based on the combination of ranitidine bismuth citrate plus clarithromycin and metronidazole has been developed. This provides a high cure rate even in patients with resistant strains⁵ and prevents the induction of secondary resistance when treatment fails, as reported by de Boer and Tytgat themselves in their review. So this triple regimen based on ranitidine bismuth citrate may be considered not only a first choice treatment in areas with a high prevalence of resistance to metronidazole or clarithromycin (thus allowing us to reduce the costs and the frequent adverse effects of two subsequent courses of treatment if eradication is not obtained with the first cycle) but also an effective second line treatment.

Vincenzo Savarino associate professor of gastroenterology
Dipartimento di Medicina Interna e Specialità Mediche, Università di Genova, n6-16132 Genova, Italy
vsavarino@unige.it

Matteo Neri associate professor of internal medicine
Dipartimento di Medicina e Scienze dell'Invecchiamento, Università di Chieti, Policlinico Colle dell'Ara, 66100 Chieti, Italy

Sergio Vigneri associate professor of gastroenterology
Istituto di Medicina Interna e Geriatria, Università di Palermo, 90127 Palermo, Italy

Competing interests: Each author has received funding for speaking or research projects and reimbursement for attending a symposium from several companies producing some of the drugs mentioned in this letter.

- De Boer WA, Tytgat GNJ. Treatment of *Helicobacter pylori* infection. *BMJ* 2000;320:31-4. (1 January.)
- European *Helicobacter Pylori* Study Group. Current European concepts in the management of *Helicobacter pylori* infection: The Maastricht consensus report. *Gut* 1997;41:8-13.
- Pipkin GA, Williamson R, Wood JR. Review article: one-week clarithromycin triple therapy regimens for eradication of *Helicobacter pylori*. *Aliment Pharmacol Ther* 1998;12:823-37.
- Van der Hulst RWM, van der Ende A, Homan A, Roorda P, Dankert J, Tytgat GNJ. Influence of metronidazole resistance on efficacy of quadruple therapy for *Helicobacter pylori* eradication. *Gut* 1998;42:166-9.
- Savarino V, Zentilin P, Pivari M, Bisso G, Mele MR, Bilardi C, et al. The impact of antibiotic resistance on the efficacy of three 7-day regimens against *H pylori* infection. *Gastroenterology* 1999;116:G1326.

Stool immunoassay for *Helicobacter pylori* is not specific enough

EDITOR—The paper by Lehmann et al¹ confirms the view that stool immunoassay for *Helicobacter pylori* is a non-invasive, cheap, and reliable test for assessing *H pylori* infection in unselected patients. However, Lehmann et al overlook the major issue concerning this diagnostic tool—namely, whether it is suitable for evaluating the outcome of treatment.

Although some authors, such as Vaira et al, maintain that the test has a high specificity only four weeks after the end of the treatment,² others report a number of false positive results after eradication. In particular, when the test is performed even two months after the eradication of *H pylori*, it still shows an unsatisfactory specificity, with 18% false positive results.³

In defending their findings, Vaira et al seek support from recent studies, which are unpublished and therefore not available for evaluation.² Thus, for the time being, we must draw on the published information. Data presented at the recent international workshop of the European *Helicobacter pylori* Study Group held in Helsinki confirm that in treated patients the specificity of stool immunoassay for *Helicobacter pylori* is lower than that of the C13 urea breath test, indicating that even six months after eradication the number of false positives is high.^{4 5}

Mario Guslandi gastroenterologist
Gastroenterology Unit, S Raffaele Hospital, Milan 20132, Italy
guslandi.mario@hsr.it

- Lehmann F, Drewe J, Terracciano L, Stuber R, Frei R, Beglinger C. Comparison of stool immunoassay with standard methods for detecting *Helicobacter pylori* infection. *BMJ* 1999;319:1409-10. (27 November.)
- Vaira D, Malfentheiner P, Megraud P, Axon TRA. Diagnosis of *Helicobacter pylori* infection by HpSA test. *Lancet* 1999;354:1732.
- Trevisani L, Sartori S, Galvani F, Rossi MR, Ruina M, Chiamenti C, et al. Evaluation of a new enzyme immunoassay for detecting *Helicobacter pylori* in feces: a prospective study. *Am J Gastroenterol* 1999;94:1830-3.
- Masoero G, Lombardo L, Della Monica P, Andriani L, Vicari S, Sallio F, et al. Discrepancy between *Helicobacter pylori* stool antigen assay and urea breath test in the detection of *Helicobacter pylori* infection [abstract]. *Gut* 1999;45(suppl 11):A131.
- Forné M, Lite J, Domínguez J, Esteve M, Fernández-Bañares F, Galí N, et al. Accuracy of a non-invasive immunoassay for detection of *H pylori* antigen in stools in the diagnosis of infection and follow-up after eradication. *Gut* 1999;45(suppl 11):A124(abstract).

Health effects of landfill sites

Whether results are assertions or evidence is unclear

EDITOR—Fielder et al reported increased rates of congenital malformations in the area surrounding the Nant-y-Gwyddon landfill site, having examined five "exposed" wards and 22 comparison wards.¹ They imply that their findings are relevant to landfill sites receiving domestic, commercial, and industrial waste in general rather than being specific to the Nant-y-Gwyddon site.

Studies such as this are subject to several scientific constraints.² Paramount among

these are that comparison populations should not be exposed to potential risks from landfill sites and that exposed areas should be relatively unaffected by alternative potential pollutant sources.

Examination of site licence records suggests that there are up to seven other operating or closed landfill sites whose zones of potential influence include the comparison wards, including sites receiving similar wastes to the Nant-y-Gwyddon site. Given the absence of comment from the authors, we conclude that these sites are not having any measurable effects. Hence the study seems to show that one landfill site may have certain health effects while seven others do not.

Health concerns have been expressed about incinerators,³ particularly the older generation of municipal waste incinerators, which operated without modern emissions controls. A former municipal waste incinerator was located about 2 km to the southeast of the exposed area. We understand that it operated from 1974 and was decommissioned in 1987. Fielder et al's study shows a peak in rates of congenital anomaly in 1988 and 1989 in the exposed wards. We further understand that an older combustion plant had operated at roughly the same location since 1916. The more recent incinerator handled the area's municipal waste before the opening of the Nant-y-Gwyddon landfill site and was apparently closed because of local complaints, poor performance, and air pollution.

The incinerator was located in a steep northwesterly valley running through the exposed wards, and atmospheric emissions probably extended into the study area. As Fielder et al provide no direct evidence that the Nant-y-Gwyddon landfill site is the cause of the raised rates of congenital malformations, it could be equally asserted that the incinerator is a material factor, although there may be other causal factors.

We would suggest a refinement of Fielder et al's final recommendation: that protocols be developed to measure community exposures systematically in areas where raised rates of congenital anomalies have been identified. To pre-empt the findings of such exposure studies with an assumption that the cause is landfill seems to miss an important opportunity to answer the question of what causes the health effects and hence to resolve the legitimate concerns of local residents.

Dave Roberts *chairman*

Andy Redfearn *senior adviser*
aerc@aerc.co.uk

Jenny Dockerty *advisory officer*

Applied Environmental Research Centre, Feering,
Colchester, Essex CO5 9ES

1 Fielder HMP, Poon-King CM, Palmer SR, Moss N, Coleman G. Assessment of impact on health of residents living near the Nant-y-Gwyddon landfill site: retrospective analysis. [Commentary by Dolk H.] *BMJ* 2000;320:19-22. (The page range includes article and commentary.) (1 January.)

2 Redfearn A, Dockerty JC, Roberts RD. Assessment of health effects associated with landfills. *Wastes Management, the Scientific and Technical Review* (in press).

3 Friends of the Earth. *Incineration campaign guide*. London: FoE, 1997.

Analyses require high quality data

EDITOR—We are founders of the Welsh Congenital Anomaly Register and Information Service (CARIS) and are well aware of the difficulties with congenital anomaly data mentioned by Fielder et al and by Dolk in her commentary on their paper.¹ Analysis of our first year's data (1998) has confirmed the suspicion of previous substantial under-reporting to the national congenital anomaly monitoring system.²

As the Welsh service has data only from 1998 and numbers are small, we are unable to reproduce Fielder et al's study concerning midline abdominal wall defects. We can, however, report the numbers of cases of all anomalies reported in 1998 for the wards surrounding the Nant-y-Gwyddon landfill site, the 22 control wards used in the study, Rhondda Cynon Taf Unitary Authority (in which the landfill site is situated), and Wales as a whole (table). We excluded data on spontaneous fetal losses from the analysis as miscarriages were not included in Fielder et al's study.

Our data suggest a slightly higher overall prevalence of cases of congenital anomaly among babies and fetuses of women living in wards surrounding the landfill site than in both the control wards and Rhondda Cynon Taf Unitary Authority. Nevertheless, the percentage of babies and fetuses affected by congenital anomaly in the wards surrounding the Nant-y-Gwyddon landfill site is no higher than that reported overall for Wales. None of the differences found reached significance with χ^2 testing. The situation should become clearer as further years of data are collected.

Fielder et al's study refers to both actual cases of congenital anomaly and anomaly rates without clearly differentiating between these terms. The Welsh service defines a case as a baby or fetus and an anomaly as the defects detected in the case. Thus the numbers of cases and numbers of anomalies in a given population may be different. In Fielder et al's study, based on data from the Office for National Statistics, this difference may not be significant as the Office for National Statistics reports 1.2 anomalies per case.³ In 1998, however, the Welsh service reported an average of 2.5 anomalies per case for Wales.² In the future it will be

important to distinguish between numbers of cases and numbers of anomalies so that equivalent comparisons can be ensured.

Judith Greenacre *consultant in public health medicine*

Margery Morgan *consultant obstetrician and gynaecologist*

David Tucker *CARIS project manager*
CARIS Office, Singleton Hospital, Swansea
SA2 8QA
dave.tucker@swansea-tr.wales.nhs.uk

1 Fielder HMP, Poon-King CM, Palmer SR, Moss N, Coleman G. Assessment of impact on health of residents living near the Nant-y-Gwyddon landfill site: retrospective analysis [with commentary by H Dolk]. *BMJ* 2000;320:19-22. (1 January.)

2 Congenital Anomaly Register and Information Service for Wales. *Annual report 1998*. Swansea: CARIS, 1998.

3 Office for National Statistics. *Congenital anomaly statistics 1997*. London: ONS, 1997. (MB3 No 12.)

Haemoglobinopathy screening can be carried out in general practice

EDITOR—Modell et al found that only half the couples at risk of having a child with thalassaemia received a service that allowed them an informed choice in genetic screening for the condition, with wide variation by region and ethnic group.¹ For informed choice to be translated into the practical reality of a prenatal diagnostic procedure, haemoglobinopathy screening needs to be undertaken in the first trimester.

In general practice the window of opportunity is small, as many mothers do not present for confirmation of pregnancy until after their second missed period. Even with early electrophoresis, if the woman is found to be a carrier it is necessary then to find her partner and persuade him to come for testing.

To overcome these time constraints, the practice where I work offers all men and women haemoglobinopathy screening based on their self perception of risk. Screening is undertaken at a registration check and opportunistically within the practice, with counselling both before and after the test. In this work we use a CD Rom developed by the department of primary care and population sciences at the Royal Free Hospital and University College London School of Medicine; this CD Rom provides accessible genetic information to

Total births and numbers of babies and fetuses reported to be affected by congenital anomalies in Wales, Rhondda Cynon Taf Unitary Authority, wards surrounding Nant-y-Gwyddon landfill site, and 22 control wards in 1998. Figures are numbers except where stated otherwise

Area	Babies and fetuses considered		Cases of congenital anomaly reported to CARIS (excluding spontaneous fetal losses)				% of babies and fetuses affected
	All births reported by ONS	Births and terminations with congenital anomaly	Terminations of pregnancy	Live births	Stillbirths	All cases	
Wales	33 610	33 798	188	609	24	821	2.43
Rhondda Cynon Taf	2 824	2 840	16	39	1	56	1.97
Nant-y-Gwyddon wards	253	256	3	3	0	6	2.34
Control wards	1 349	1 356	7	17	0	24	1.77

ONS=Office for National Statistics.

CARIS=Welsh Congenital Anomaly Register and Information Service.

both patient and clinician and also publicity materials.

This service is acceptable to patients, with high uptake in appropriate ethnic groups and a high attendance at follow up visits, when they are given results and a haemoglobinopathy card. The results are readily available on the computer screen and in the general practice record. Any patients found to carry a significant trait are advised to present early for antenatal care should they or their partner become pregnant. Preconceptional haemoglobinopathy testing removes some of the additional anxiety for the patient in early pregnancy by identifying risk earlier and allows more time to organise prenatal diagnosis should it be necessary.

The practice where I work is recognised by the NHS Executive as a beacon practice for this service. We would be happy to share our experience with other professionals and provide them with free CD Roms to facilitate this.

Jane Logan *general practitioner*
Mawbey Brough Health Centre, London SW8 2UD
Mawbeygp@dircon.co.uk

1 Modell B, Harris R, Lane B, Khan M, Darlison M, Petrou M, et al. Informed choice in genetic screening for thalassaemia during pregnancy: audit from a national confidential inquiry. *BMJ* 2000;320:337-41. (5 February.)

Working in other countries

Work opportunities in developing countries broaden the mind

EDITOR—Working in other countries can enhance knowledge and skills and broaden one's perspective considerably, as Smith emphasises in her personal view about working for Voluntary Service Overseas.¹ Having

Advice to authors

We prefer to receive all responses electronically, sent either directly to our website or to the editorial office as email or on a disk. Processing your letter will be delayed unless it arrives in an electronic form.

We are now posting all direct submissions to our website within 24 hours of receipt and our intention is to post all other electronic submissions there as well. All responses will be eligible for publication in the paper journal.

Responses should be under 400 words and relate to articles published in the preceding month. They should include ≤ 5 references, in the Vancouver style, including one to the BMJ article to which they relate. We welcome illustrations.

Please supply each author's current appointment and full address, and a phone or fax number or email address for the corresponding author. We ask authors to declare any competing interest. Please send a stamped addressed envelope if you would like to know whether your letter has been accepted or rejected.

Letters will be edited and may be shortened.

bmj.com
letters@bmj.com

begun life in India, I qualified in medicine in the United Kingdom and then spent several years preparing to return to rural India as an ophthalmologist. Most people regarded this choice as quaint, but there were colleagues who showed genuine interest.

I worked for five years as an ophthalmologist in a leprosy hospital (also providing eye services for the general population) in eastern India. These were the most challenging, interesting, and exhausting years of my career. There were times when I felt ill equipped, ill informed, and totally inadequate for my responsibilities. I quickly had to learn how to budget, to buy equipment, to write project proposals, to train and manage staff, and to organise and teach on training courses and workshops. My training in the United Kingdom had not afforded me these skills.

Clinically, I did my best to put into practice what I had learnt in the United Kingdom and to give equivalent standards of care. This was despite caring for more patients, having less equipment, and there being no senior colleagues on hand to give advice.

I am now working in Glasgow, and enjoying it. I have plenty of colleagues with whom to discuss topics of interest, plentiful equipment for every procedure, and a wide range of treatments available to prescribe. When I am on call it is for ophthalmology alone. I don't have to set up drips on dehydrated infants, perform lumbar punctures, manage pancreatitis, or deliver babies.

Is such experience in India and elsewhere valued appropriately? I have worked with scores of dedicated Indian colleagues who practise as I did, with access to a tiny fraction of the budget we have in the United Kingdom. Few have had European or American postgraduate training, but most have a wide range of skills and abilities and a comprehensive "world view" rarely found in developed countries.

My experience has shown me that my training in the United Kingdom (albeit something I appreciate) has given me a fraction of the knowledge and skills that I regard as valuable, wherever I am working. Work opportunities in developing countries broaden the mind and make us realise how much we can learn from each other.

Kirsteen J Thompson *specialist registrar*
Tennent Institute of Ophthalmology, Gartnavel General Hospital, Glasgow G12 0YN
kirsteenjt@doctors.org.uk

1 Smith L. So who's teaching whom? *BMJ* 2000;320:323. (29 January.)

People may make choices at odds with their cultural norms

EDITOR—Like Smith,¹ I have worked in Africa. Like her, I am the coauthor of a training pack in race and cultural awareness.² But my interest was triggered not by Africa but by being a patient in a European country with an excellent and well resourced health service whose staff knew a great deal about the British way of life. I still, however, found myself a square peg squeezing into round Swedish holes, cared for by kind,

highly competent staff who made some very incorrect assumptions about me.

I do not agree that information booklets about "the different needs of Muslims, Sikhs, and Hindus" are what's needed. There are already far too many resource packs, textbooks, and training packs with this cookery book approach, with menus of cultural norms—what A eats, how B does her hair, how C dies. Such an approach, in which the special features of "the Asian family" or "the Afro-Caribbean child" are explored, with rigid and static notions of culture, can worsen the care that black people receive as it leads to pigeon holing, victim blaming, and stereotyping. In reality, people make choices that may be totally at odds with their cultural norms. Indeed, the whole notion of cultural norms in a multicultural society is questionable.

The idea of cultural norms arises from a simplistic and unidimensional view of people's lives which "ignores the effects of host cultures on minority ethnic communities, both collectively and at the level of the individual. Members of minority ethnic communities live at the intersection of two or even more cultures. At these intersections reactions take place that no textbook can predict."³

Of course professionals need access to information about, for example, Sikh naming systems and Hindu dietary norms. But to put such information in context and to apply it intelligently to their everyday practice, professionals need a structured opportunity to explore their own attitudes; recognise that they are not culturally neutral but a product of their own cultural conditioning; and understand how their own and others' attitudes towards race, colour, and religion interact with those towards class, age, sex, and disability.

Cultural competence is not about generalising on the contraceptive needs of Roman Catholics or about learning what Somali refugees eat for breakfast on Thursdays. Becoming culturally competent requires first learning a great deal about oneself.

Elspeth Webb *senior lecturer*
Department of Child Health, University of Wales College of Medicine, Cardiff CF4 4XN
john.clark@whicb.net

1 Smith L. So who's teaching whom? *BMJ* 2000;320:323. (29 January.)

2 Shah L, Thomas-Ramuset A, Webb E. *Equal rights, equal access—improving the health care of disabled and chronically ill children from minority ethnic communities: a training pack*. Cardiff: Department of Child Health, University of Wales College of Medicine, 1996.

3 Ingleby D. Psychosocial problems among children of migrants. In: *Child health care for migrants and refugees. Proceedings of the annual meeting of the European Society of Social Paediatrics*. Rotterdam: Stichting Ouder en Kind-zorg, 1995: 40-6.

Rapid responses

Correspondence submitted electronically is available on our website