Letters

Website: bmj.com
Email: letters@bmj.com

Prostatic symptoms

Essential simple investigations were not mentioned

EDITOR—Urologists have spent over a decade trying to get across the message that lower urinary tract symptoms in middle aged and elderly men are not synonymous with prostatic problems. A major attempt at getting this message across was an editorial by Abrams in 1994.¹ How disappointing, then, to see nocturia in a man of 68 automatically assumed to be a manifestation of prostatic disease.²

Almost the first action recommended to the general practitioner in Farmer's article is to "Show [the patient] a picture of ... the prostate ... and how it causes problems." The symptom described may well be due to age related changes in the detrusor, perhaps with some diminished renal concentrating power, reduced nocturnal secretion of antidiuretic hormone, and even subclinical accumulation of oedema by day and its reabsorption at night. A combination of these is at least as likely a cause of nocturia as is the prostate.

No mention is made of the cheapest, simplest, and safest investigation of all—completion of a frequency-volume chart, which often shows mild nocturnal polyuria. Nor is the measurement of urine flow rate

and residual volume mentioned; this is essential and is now widely available to general practitioners through open access or nurse led clinics. No man should be given treatment until these investigations have been done.

The last third of Farmer's article implies that management is mainly by α blocking drugs. We have moved from an era of inappropriate prostatectomy for nocturia and other irritative symptoms by urologists to one of inappropriate pharmacotherapy by general practitioners, at massive cost. This 10-minute consultation would have been much better used to explain to the man that his presenting symptom had no more than an even chance of being due to a prostate problem and that no treatment would be considered without the essential investigations mentioned above having been done.

K Baxby consultant urological surgeon
Department of Urology, Ninewells Hospital,
Dundee DD1 9SY
kbaxby@aol.com

1 Abrams P. New words for old: lower urinary tract symptoms for "prostatism." *BMJ* 1994;308:929-30.

2 Farmer A. 10-minute consultation: Prostatic symptoms. *BMJ* 2001;322:1468. (16 June.)

All possible causes of lower urinary tract symptoms must be investigated

EDITOR—The assessment of a man with urinary symptoms is hard to summarise comprehensively in a single page, but Farmer's 10-minute consultation has important errors and omissions.¹ I was surprised that an article in the *BMJ* should use the term "prostatic symptoms" as the journal was the first to use the more correct term "lower urinary tract symptoms."²

Studies of elderly men and women matched for age show that there is no such thing as prostatic symptoms. Abundant evidence indicates that at best there is only a weak association between individual urinary symptoms and any variable of lower urinary tract anatomy or physiology (prostate size, urinary flow rate, volume of residual urine, bladder outlet obstruction).³

For these reasons, the term lower urinary tract symptoms (which does not imply any particular pathophysiology) is more acceptable. Use of the term prostatic symptoms encourages us to forget the other pathological explanations that should be considered in the differential diagnosis of lower urinary tract symptoms, such as bladder cancer, prostate and other urinary tract infection, and

several causes of disturbed fluid balance. It also encourages us to treat the prostate when this may be unnecessary.

Watchful waiting was not mentioned as a treatment option. A randomised study has shown that men with moderate lower urinary tract symptoms and residual urine volumes of <350 ml do just as well with watchful waiting as with surgery.⁴

About 30% of men are not happy with the outcome after transurethral resection of the prostate, so this is hardly a "very effective treatment." It is incorrect to say that this treatment only "occasionally leads to complications." The national prostatectomy audit⁵ and other studies⁴ have shown that transurethral resection of the prostate for lower urinary tract symptoms is associated with cardiorespiratory problems in 3% of cases; the need for transfusion of >2 units of blood in 3%; return to theatre for, among other things, control of severe haemorrhage in 1.4%; septicaemia in 1%; transurethral resection of the prostate syndrome in 0.4%; and a 30 day mortality of 0.2%. In the long term there is a 3% rate of urethral stricture and a 3% chance of bladder neck stenosis. This is in the context of an elective procedure for symptoms that are not life threatening.

John Reynard consultant urologist
Department of Urology, Churchill Hospital, Oxford
OX3 7LJ
jmreynard@baus.org.uk

1 Farmer A. 10-minute consultation: Prostatic symptoms. *BMJ* 2001;322:1468. (16 June.)

2 Abrams P. New words for old: lower urinary tract symptoms for "prostatism." *BMJ* 1994;308:929-30.

3 Reynard JM, Yang Q, Donovan JL, Peters TJ, Schafer W, de la Rosette JJ, et al. The ICS-BPH study: uroflowmetry, lower urinary tract symptoms and bladder outlet obstruction. *Br J Urol* 1998;82:619-23.

4 Wasson JH, Reda DJ, Bruskewitz RC, Elinson J, Keller AM, Henderson WG for the Veterans Affairs Cooperative Study Group on Transurethral Resection of the Prostate. A comparison of transurethral surgery with watchful waiting for moderate symptoms of benign prostatic hyperplasia. *N Engl J Med* 1995;332:75-9.

5 Pickard R, Emberton M, Neal DE on behalf of the National Prostatectomy Audit Steering Group. The management of men with acute urinary retention. *Br J Urol* 1998;81:712-20.

Isn't use of prostate specific antigen tests just screening by the back door?

EDITOR—Farmer tackles a controversial subject in his article on managing prostatism.¹ He recommends prostate specific antigen testing for men with symptoms of prostatism, but we would question the value of this for two reasons.

Firstly, prostatism is caused by outflow obstruction of the bladder (generally due to benign prostatic hypertrophy) and occurs in the periurethral portion of the prostate gland, whereas cancers occur in the peripheral zone.² Symptoms of obstruction would

Advice to authors

We prefer to receive all responses electronically, sent directly to our website. Processing your letter will be delayed unless it arrives in an electronic form.

We are now posting all direct submissions to our website within 24 hours of receipt and our intention is to post all other electronic submissions there as well. All responses will be eligible for publication in the paper journal.

Responses should be under 400 words and relate to articles published in the preceding month. They should include ≤ 5 references, in the Vancouver style, including one to the *BMJ* article to which they relate. We welcome illustrations.

Please supply each author's current appointment and full address, and a phone or fax number or email address for the corresponding author. We ask authors to declare any competing interest. Please send a stamped addressed envelope if you would like to know whether your letter has been accepted or rejected.

Letters will be edited and may be shortened.

bmj.com
letters@bmj.com

therefore be expected to occur late, if at all, in prostate cancers. The relation between prostate cancer and benign prostatic hypertrophy is entirely coincidental because of the prevalence of the conditions in older men. Thus this test cannot be considered any more than screening.

Secondly, benign prostatic hypertrophy is one of the recognised causes of a raised level of prostate specific antigen in the absence of cancer. The positive predictive value of the prostate specific antigen test is about 30% in the general population and probably lower in men with prostatism.³ Yet all those with a raised result will require transrectal ultrasound guided prostatic biopsy to identify the 30% who have a cancer. The side effects of the biopsy are infection (sufficiently severe to require admission in 1% of cases), bleeding (33% of patients have some haematuria and haematospermia in the three months after the biopsy), and death (<0.01%).⁴

Because of the nature of the disease many of those found to have cancer will have indolent tumours that will not cause problems in their lifetime. There is currently no proved treatment for early (localised) prostate cancer⁵—hence the recommendation by the national screening committee not to offer prostate specific antigen screening as part of its national programme. Thus, in having this test, many men will have been subjected to an invasive investigation; in the 30% in whom early cancer is found many will have an otherwise “innocent” illness (except that now they have increased anxiety) and no proved effective treatment can be offered.

Men presenting with prostatism will probably know from the media about prostate specific antigen testing. They may develop prostate cancer in the future and wonder why it was not detected earlier. We would therefore recommend that testing is discussed and the advantages and disadvantages outlined. The patient would then be in a better position to make an informed decision about whether a test would help him.

Lucy Jenkins *general practitioner*
lucyjenkins@doctors.org.uk

Peter Rose *general practitioner*
Tim Wilson *general practitioner*
Mill Stream Surgery, Benson, Oxfordshire
OX10 6RL

1 Farmer A. 10-minute consultation: Prostatic symptoms. *BMJ* 2001;322:1468. (16 June.)

2 Selley S, Donovan J, Faulkner A, Coast J, Gillatt D. Diagnosis, management and screening of early localised prostate cancer. *Health Technol Assess* 1997;1(2):3.

3 Neal DE, Donovan JL. Prostate cancer: to screen or not to screen? *Lancet Oncology* 2000;1:17-24.

4 NHS Executive. *NHS prostate cancer programme*. London: Department of Health, 2000.

5 Wilt T, Brawer M. Prostate cancer: non-metastatic. In: Barton S, ed. *Clinical evidence* 4. London: BMJ Publishing Group, 2000.

Author's reply

EDITOR—The 10-minute consultation was intentionally titled to draw attention to the management of benign prostatic hypertrophy, one of the causes of lower urinary tract symptoms. Use of the international prostate

symptom score, as suggested in the article, is part of a systematic approach that will suggest the presence of obstruction or draw attention to the possibility of alternative explanations for the presenting symptoms.

The difficulties in interpreting results of the prostate specific antigen test raised by Jenkins et al are particularly relevant to decisions about population screening, as the level of evidence required to start a screening programme should be high. The general practitioner in the vignette is responding to a demand for care by a patient with symptoms—in this case an implicit request for reassurance about the possibility of prostate cancer. The prostate specific antigen test is probably the best available blood test for detecting cancer, and although it has limited specificity and sensitivity, it remains widely used for investigating lower urinary tract symptoms.¹

All three letters draw attention to the role of patient choice, whether in counselling about the advantages and disadvantages of the prostate specific antigen test or in discussions of the relative benefits of non-pharmacological management of symptoms, pharmacotherapy, and surgery. The degree to which these issues can be routinely explored in depth within the time constraints of a routine general practice consultation is limited. However, the provision of systematically developed educational resources² and the extension of open access nurse led assessment clinics for people with lower urinary tract symptoms may increase the range of management options available in primary care.

Andrew Farmer *honorary senior lecturer in general practice*
Institute of Health Sciences, Oxford OX3 7LF
andrew.farmer@public-health.oxford.ac.uk

1 Abrams P, Wein A. Recent advances: Urology. *BMJ* 2000;321:1393-6.

2 Murray E, Davis H, See Tai S, Coulter A, Gray A, Haines A. Randomised controlled trial of an interactive multimedia decision aid on benign prostatic hypertrophy in primary care. *BMJ* 2001;323:493-6. (1 September.)

Community pharmacy supply of emergency contraception

Impact of emergency contraception on women's and men's behaviour requires further exploration

EDITOR—Stammers in his letter expresses anxiety about the supply of emergency hormonal contraception in community pharmacies and the effect this may have on the spread of sexually transmitted infections.¹ He says that women in need of emergency hormonal contraception will be at risk of sexually transmitted infections. But many of these women are using the contraceptive pill and have forgotten to take it, or are in stable relationships. These women are not at increased risk of contracting infections, and pharmacy supply will have no impact on their spread. If a woman is attending for emergency contraception because of a burst condom or an episode of unprotected sex,

there is a risk of contracting sexually transmitted infections, but the level of risk is not directly affected by the wider availability of emergency contraception in the pharmacy. If women are at risk of sexually transmitted infections they do not need an emergency intrauterine device: the quick and accessible supply of levonorgestrel is even more pertinent here.

Stammers is wrong to imply that pharmacists provide suboptimal care because they do not make reference to sexually transmitted infections when supplying emergency contraception. The Lambeth, Southwark, and Lewisham health action zone, patient group direction for access to emergency hormonal contraception, makes specific reference to sexually transmitted infections, as does the patient group direction for Manchester, Salford, and Trafford health action zone. Furthermore, during our ongoing evaluation of both these schemes, such infections were a central concern of the participating pharmacists.

Stammers should bear in mind the views of women clients, who suggested that, when obtaining emergency hormonal contraception (from whatever source), they were most concerned about averting pregnancy rather than contracting sexually transmitted infections. We should not be complacent about the spread of these infections, but the consultation itself may not be the most appropriate moment for education. Leaflets and other means of providing individualised, tailored information also need to be considered.

Stammers is right to point to the possibility of risk displacement—the potential rise in the incidence of sexually transmitted infections as a result of pharmacy supply. This may, however, be conflating two quite dissimilar risks: that of an unplanned or unwanted pregnancy and that of sexually transmitted infections. Currently, there are few indications that women are changing their contraceptive behaviour because of wider availability of emergency hormonal contraception. The evidence from our evaluation is that women (largely) behave responsibly in relation to their use of contraception. None the less, the wider availability of emergency hormonal contraception and its impact on women's (and men's) sexual and contraceptive behaviour requires further empirical exploration.

Paul Bissell *lecturer in social pharmacy*
University of Nottingham, Nottingham NG7 2RD
Paul.Bissell@nottingham.ac.uk

Claire Anderson *director of pharmacy practice and social pharmacy*
University of Nottingham

Lesley Bacon *consultant in community reproductive health*
South London NHS Trust, London SE5 7RN

Beth Taylor *pharmacy manager, community health*
South London NHS Trust, London SE14 5ER

Karen O'Brien *project coordinator*
Manchester, Salford and Trafford Health Action Zone, Manchester M21 7RL

1 Stammers T. Emergency contraception from pharmacists misses opportunity. *BMJ* 2001;322:1245. (19 May.)

Collaboration is vital

EDITOR—The issue of pharmacists providing emergency contraception is controversial, as shown by the contribution by Stammers, and I am still uncertain about it.¹ Although the recent reclassification of emergency contraception can only broaden access to this useful product, I would have reservations about selling this item myself, not for any issue of conscience (which is causing great debate in the pharmacy literature in the United Kingdom) but because most British community pharmacies do not have the facilities to take a history in private. If pharmacists are going to provide this service then, as Stammers suggests, they should also be providing information about local sexual health services and advocating their use.

Contact your local sexual health centre and discuss this issue with them, and you will find that they are only too willing to help with the provision of relevant materials. This issue has the possibility to develop into yet another of those "them and us" scenarios between the medical and pharmacy professions. The time has come for everyone to work towards a common goal rather than everyone working in isolation.

Gary Ward *dispensary manager, Auckland Hospital*
3 Condor Place, Unsworth Heights, Auckland 1030, New Zealand
garyw@ahsl.co.nz

1 Stammers T. Emergency contraception from pharmacists misses opportunity. *BMJ* 2001;322:1245. (19 May.)

Family history and risk of venous thromboembolism with oral contraception

Family history is important tool

EDITOR—Cosmi et al in their article claim that family history has poor diagnostic test qualities to detect prothrombotic mutations.¹ They also indicate that overall population screening is not cost effective. Although it always was obvious that family history would never make a perfect test (of families with many cases of venous thrombosis, up to 40 or 50% have factor V Leiden²), the finding of an equal predictive value for a positive and a negative family history by Cosmi et al is surprising and might be due to the small numbers included or the type of history that was considered "positive."

Even if family history is far from perfect as a diagnostic test for one or two mutations, the question is whether one should refrain from using it. In our original publication, we tried to emphasise the meaning of family history for the prescription of oral contraceptives, and not really as a clue for the detection of mutations.³ Apart from any prothrombotic mutation, a strongly positive family history might point to a tendency for venous thrombosis that might be taken into account clinically in the decision whether or not to use oral contraceptives.

Moreover, even the meaning of a mutation is different in the context of a positive family history: the age at first venous thrombosis with factor V Leiden is about 10 years younger in persons with a positive family history than in consecutive patients.⁴ The most important information that one would like to obtain from a family history is not a proxy test for factor V Leiden or any other mutation, but the likelihood that a woman will develop thrombosis if she uses oral contraceptives. Whether or not family history would be valuable for that purpose is the ultimate test.

Jan P Vandenbroucke *professor of clinical epidemiology*

Department of Clinical Epidemiology
vdbroucke@mail.medfac.leidenuniv.nl

Felix J M van der Meer *physician*
Haemostasis and Thrombosis Centre

Frans M Helmerhorst *senior lecturer*
Department of Obstetrics and Gynaecology

Frits R Rosendaal *professor*
Department of Clinical Epidemiology and
Department of Obstetrics and Gynaecology, Leiden
University Medical Center, 2300 RC Leiden,
Netherlands

1 Cosmi B, Legnani C, Bernardi F, Coccheri S, Palareti G. Value of family history in identifying women at risk of venous thromboembolism during oral contraception: observational study. *BMJ* 2001;322:1024-5. (28 April.)

2 Rosendaal FR. Venous thrombosis: a multicausal disease. *Lancet* 1999;353:1167-73.

3 Vandenbroucke JP, van der Meer FJM, Helmerhorst FM, Rosendaal FR. Factor V Leiden: should we screen oral contraceptive users and pregnant women? *BMJ* 1996;313:1127-30.

4 Lensen RP, Rosendaal FR, Koster T, Allaart CF, de Ronde H, Vandenbroucke JP, et al. Apparent different thrombotic tendency in patients with factor V Leiden and protein C deficiency due to selection of patients. *Blood* 1996;88:4205-8.

Author's reply

EDITOR—The finding of an equal predictive value of a positive and negative family history of venous thromboembolism might be because of the type of family history that was considered to be positive—that is, one first or second degree relative. However, because of the demographic changes in Italy over the past three decades, modern families are now small and many siblings are extremely rare. As a result, it would be quite difficult to find young women with two or more relatives with a history of venous thromboembolism.

Our study was prompted by the question whether clinicians can identify women at risk of venous thromboembolism during contraception only on the basis of family history. A positive personal or family history, or both, can be considered in itself a contraindication to oral contraception regardless of any thrombophilic defect. In the case of a positive history, screening for thrombophilia may be irrelevant because, even if no thrombophilic defects are found, the decision may be based only on clinical grounds. The problem can arise when clinicians obtain a negative personal or family history. We agree that family history cannot be a proxy test for any thrombophilic mutation, but it indicates the likelihood that a woman will develop thrombosis during oral contraception.

We also know that oral contraceptives may trigger thromboembolic complications in women with unrecognised thrombophilic defects. Thus women with thrombophilic defects can be considered at higher risk of developing thrombotic complications, even though the absolute risk is low. Is it necessary to search for thrombophilic defects if personal or family history is negative? Possibly not if the problem is considered from an epidemiological point of view because universal screening is not cost effective. However, women cannot be denied the knowledge and the information about the possibility of screening. We do not yet have an understanding of the biological basis of thrombotic complications during oral contraception and we have no other means of identifying women at risk of such complications. Apart from the risk of fatal pulmonary embolism, the morbidity and cost and side effects of venous thromboembolism cannot be ignored by clinicians.

Moreover, clinicians constantly face litigation, which could arise from incomplete information and the finding of a thrombophilic defect after an episode of venous thromboembolism during oral contraception.

Benilde Cosmi *lecturer*

Cardiovascular Department, Division of Angiology,
Unità Ricerca Clinica sulla Trombofilia
"M Golinelli", University Hospital,
S Orsola-Malpighi, 40138, Bologna, Italy
bcosmi@med.unibo.it

Author's reply to criticism of study on benzodiazepines and risk of hip fracture

EDITOR—Sgadari et al's letter¹ commented on the study that I and several others carried out to see whether benzodiazepines are associated with an increased risk of hip fracture.² We found that they are not.

In their large case-control study of 9752 patients with hip fractures compared with 38 564 controls, Sgadari et al also found no association between the use of benzodiazepines and hip fracture.³ But like us, though in different subgroups, they did find an association between certain drugs and hip fracture. They looked at the metabolic pathways involved, and it seems that in a specific subgroup of the most elderly patients these drugs may confer more risk. If this is not the result of multiple post-hoc testing it is an interesting finding, and one we did not look at.

We stated that the results we found for individual drugs might be spurious, related to multiple testing and statistical variability. But the main result is still that overall there is no increased risk of fracture associated with benzodiazepines. The risks associated with the subgroups mentioned by Sgadari et al are minuscule and apparent only because of the size of their study. This is in clear contrast with the important and constantly found dose dependent, increased risk of falling, for all products, which is

consistent with their remarkably similar pharmacological properties.

It is time to try to understand why and how benzodiazepines increase the risk of falling but not of hip fracture, rather than concentrating on finding subgroups of benzodiazepines that do increase the risk of hip fracture.

Nicholas Moore *professor of clinical pharmacology*
Université Victor Segalen-Bordeaux2, Bordeaux,
France
nicholas.moore@pharmaco.u-bordeaux2.fr

- 1 Sgadari A, Gambassi G, Pedone C, Onder G. Study should have reported more data about associated diseases. *BMJ* 2001;322:112. (14 July.)
- 2 Pierfitte C, Macouillard G, Thicoïpe M, Chaslerie A, Pehourcq F, Aissou M, et al. Benzodiazepines and hip fractures in elderly people: case-control study. *BMJ* 2001;322:704-8. (24 March.)
- 3 Sgadari A, Lapane KL, Mor V, Landi F, Bernabei R, Gambassi G. Oxidative and nonoxidative benzodiazepines and the risk of femur fracture. *J Clin Psychopharmacol* 2000;20:234-9.

Journal should show balance, not bias, when reporting on PFI

EDITOR—Yet again the *BMJ* has compromised its integrity by uncritically yielding to its bias against hospital developments made under the private finance initiative (PFI). In reporting Dr Richard Taylor's election as a member of parliament, Gulland quotes from the Pollock report on Worcestershire's private finance initiative.¹ As Deakin and I have previously pointed out, this report is based on an incomplete and inaccurate analysis.²

Firstly, the cost of Worcestershire's private finance initiative has not increased from £49m to £108m. Pollock did not compare like with like. She compared the original building cost of the small South Worcestershire Hospital with the total cost of the larger pan-Worcestershire Hospital under the private finance initiative (building it, equipping it, and maintaining it for 30 years, including inflation costs). In practice the Worcestershire Hospital will be built and commissioned by March 2002 within budget and on time—an unusual occurrence for major capital projects in the NHS.

Secondly, the clinical review that triggered the changes at Kidderminster (still unchallenged by any of the independent reviews that have been commissioned in Kidderminster) preceded the private finance initiative business case by several years rather than being a consequence of the development.

Thirdly, Kidderminster Hospital is not closing. It is still open and has a bright future as an integral part of the NHS Plan in Worcestershire. It is just as busy as ever.

Brian McCloskey *director of public health*
Worcestershire Health Authority, Worcester
WR4 9RW
brian.mccloskey@wha.worcester-ha.wmids.nhs.uk

Professor McCloskey is employed by Worcestershire Health Authority and instigated the strategic clinical review that led to the changes at Kidderminster Hospital.

- 1 Gulland A. Doctor wins seat in fight to save hospital. *BMJ* 2001;322:1443. (16 June.)
- 2 McCloskey B, Deakin M. Private finance initiative. *BMJ* 2000;320:250.

Implementing clinical governance

Intellectual discussion would have been useful

EDITOR—What is upsetting about clinical governance is the way in which political zeal has eclipsed intellectual honesty. Halligan and Donaldson have listened to audiences composed of healthcare professionals who believe that clinical governance is the right idea.¹ From the response to my criticism of clinical governance,² and of audiences that I have addressed on the basis of that criticism, I know there are many healthcare professionals who are not so convinced. Halligan has himself heard important criticisms because he spoke at a conference convened at the Royal College of Physicians in March 2000 to discuss some of its difficulties.³ So why is my critical reference, or any other, not cited in the article by Halligan and Donaldson? The article is little more than a restatement of the rhetoric of the original article by Scally and Donaldson,⁴ plus some anecdotal examples of how they claim clinical governance has improved the delivery of service.

Anyone writing a review of a new medical treatment is expected to look for all the evidence. Halligan and Donaldson have ignored evidence that they knew was there—or does intellectual dissent not count as evidence? The *BMJ* has championed the rigorous searching for evidence that underlies evidence based medicine. Why did the assessors of this paper not point out Halligan and Donaldson's apparent blind spot? The paper appeared in the education and debate section of the journal, but should more rightly have been in a new section termed propaganda.

Neville W Goodman *consultant anaesthetist*
Southmead Hospital, North Bristol NHS Trust,
Bristol BS10 5NB
Nev.W.Goodman@bris.ac.uk

- 1 Halligan A, Donaldson L. Implementing clinical governance: turning vision into reality. *BMJ* 2001;322:1413-7. (9 June.)
- 2 Goodman NW. Clinical governance. *BMJ* 1998;317:1725-7.
- 3 Miles A, Hill A, Hurwitz B, eds. *Clinical governance in the NHS: enabling excellence or imposing control?* London: Aesculapius Medical Press (in press).
- 4 Scally G, Donaldson LJ. Clinical governance and the drive for quality improvement in the new NHS in England. *BMJ* 1998;317:61-5.

Medical training should include project management

EDITOR—As one of the first teams to graduate from the national development programme for clinical governance, and having successfully used techniques of clinical governance to produce important changes in our hospital, we endorse the paper by Halligan and Donaldson.¹

The establishment of a project culture, as discussed in the paper, was a key event in the implementation of clinical governance in our organisation. This was also the most invigorating element of our trust's response to the challenge set by our involvement with the development programme. Project cul-

ture enables effective change to take place in many different areas of a service without requiring the whole service itself to be reformulated. Project teams are especially effective because they can reach across traditional professional barriers while also addressing areas of service with which they are entirely familiar. As a result, hurdles that blocked the process of change previously are overcome.

It is also liberating to see how the quality of patient care can be improved substantially without resorting to grandiose schemes that seldom produce the desired result. Through project management, the ability to effect real change is placed within the grasp of those who see the need for it most.

The best way to engender a project culture throughout the NHS, however, requires consideration. Although the NHS support team for clinical governance is expanding and aiming to reach many professionals, we believe that ultimately the implementation of clinical governance would be well served if early medical training included instruction in project management. If the quality of the provision of medical care is a priority for all, then the ability to manage change is a desirable skill for all healthcare professionals to master. Improved quality does not arrive by chance but by well reasoned and well founded intervention.

The implementation of clinical governance is far too important a lesson to be taught at a relatively late stage in one's career. The NHS support team for clinical governance should consider visiting medical schools as well as trust boards.

Sean W O'Kelly *consultant anaesthetist*
swokelly@aol.com

Rachel Maxwell *senior sister*
Princess Margaret Hospital, Okus Road, Swindon
SN1 4UJ

- 1 Halligan A, Donaldson L. Implementing clinical governance: turning vision into reality. *BMJ* 2001;322:1413-7. (9 June.)

Perineal massage in pregnancy

Such massage significantly decreases perineal trauma at birth

EDITOR—Stamp et al's trial of the effect that perineal massage during labour has on perineal trauma adds to our knowledge about care of the perineum during childbirth,¹ but their report of the results of our trial is erroneous.²

We found an absolute increase of 9% in intact perineum in women who had not had a previous vaginal birth who were randomised to do antenatal perineal massage (24%, n=411) compared with the control group (15%, n=417). In the intention to treat analysis this difference was highly significant (95% confidence interval 4% to 15%; P=0.001).

The difference was also clinically important: one case of perineal trauma requiring suturing was avoided for every 11 women without a previous vaginal birth who were

assigned to the massage group. When the results were analysed by actual practice, women randomised to do massage who practised perineal massage on less than one third, one third to two thirds, and more than two thirds of the assigned days had an intact perineum in 20%, 23%, and 28% of cases, respectively (χ^2 for trend = 13.2; $P = 0.0003$).

Thus, in women approaching their first vaginal delivery, antenatal perineal massage significantly decreases the risk of perineal trauma.

Michel Labrecque *professor*
Department of Family Medicine, Laval University,
Quebec, Canada G1K 7P4
michel.labrecque@mfu.ulaval.ca

Erica Eason *associate professor*
Department of Obstetrics and Gynaecology,
University of Ottawa, Ottawa, Canada

Sylvie Marcoux *professor*
Department of Social and Preventive Medicine,
Laval University

- Stamp G, Kruzins G, Crowther C. Perineal massage in labour and prevention of perineal trauma: randomised controlled trial. *BMJ* 2001;322:1277-80. (26 May.)
- Labrecque M, Eason E, Marcoux S, Lemieux F, Pinault JJ, Feldman P, et al. Randomized controlled trial of prevention of perineal trauma by perineal massage during pregnancy. *Am J Obstet Gynecol* 1999;180:593-600.

True incidence of third degree tears should be ascertained

EDITOR—Stamp et al state that faecal incontinence after vaginal delivery is less likely in women whose perineum remains intact,¹ but it is well recognised that clinical examination of the perineum by a midwife or doctor in no way reflects the true damage done to the sphincter complex.² Anal endosonography, which images the sphincters in detail, shows a far higher incidence of sphincter disruption and laceration than does clinical examination.

A retrospective study of 8603 vaginal deliveries found that a third degree tear had been clinically diagnosed in only 50 women (0.6%),³ yet when the same authors used anal endosonography in a consecutive group of 202 deliveries there was evidence of third degree tears in 35% of primiparous and 44% of multiparous women—a startling incidence confirmed by other researchers.⁴ Furthermore, sonographically detected tears are strongly associated with physiological sphincter impairment and symptoms of anal incontinence, suggesting that they are functionally important.⁵

The message is clear: anal sphincter tears occur often during vaginal delivery, and clinical examination misses most of them. Inadequate training has a central role in this; a study of midwives and obstetricians found their anatomical knowledge to be generally inadequate—for example, only 7% of midwives and none of the doctors were aware that only the bulbospongiosus and superficial transverse perineal muscles were commonly divided during posterolateral episiotomy.⁵

The incidence of third degree tears in Stamp et al's study (1.7% in the massage group versus 3.8% in the control group) suggests that most tears went unrecognised. The authors state that their trial was underpowered to detect third degree tears and

that further research with a larger sample is warranted to determine if their trend towards reduced risk with perineal massage is real. Instead, it may be more scientifically valid to determine the true incidence by using endosonography; this would certainly be easier given the much smaller sample size needed. Alternatively, as third degree tears are so common, research should be directed towards assessing primary repair, who should perform it, and the training needed.

Steve Halligan *consultant radiologist*
St Mark's Hospital, Harrow, Middlesex HA1 3UJ
steve@halligan9.fsnet.co.uk

- Stamp G, Kruzins G, Crowther C. Perineal massage and prevention of perineal trauma: randomised controlled trial. *BMJ* 2001;322:1277-80. (26 May.)
- Sultan A, Kamm MA, Hudson CN, Thomas JM, Bartram CI. Anal sphincter disruption during vaginal delivery. *N Engl J Med* 1993;329:1905-11.
- Sultan AH, Kamm MA, Hudson CN, Bartram CI. Third degree obstetric anal sphincter tears: risk factors and outcome of primary repair. *BMJ* 1994;308:887-91.
- Fines M, Donnelly V, Behan M, O'Connell PR, O'Herlihy C. Effect of second vaginal delivery on anorectal physiology and faecal continence: a prospective study. *Lancet* 1999;354:983-6.
- Sultan AH, Hudson CN. Are junior doctors and midwives adequately prepared to repair the perineum? *J Obstet Gynaecol* 1993;13:484-5.

Removal from a GP's list: qualitative research is needed

EDITOR—Pickin et al offer a postal questionnaire survey of general practitioners' reasons for removing patients from their lists.¹ To date, published research on removals has relied on an analysis of routinely collected health authority data that do not allow the reasons for removal to be ascertained except for immediate removals on the grounds of violence.^{2,3}

We think, however, that there are important problems with the methods adopted by Pickin et al. They base their list of possible reasons for removal on two postal surveys of general practitioners, one of which is unpublished and has a response rate of less than 30%.⁴ These studies may have inappropriately limited the number, range, and type of reasons under investigation. Moreover, respondents may be especially prone to giving publicly acceptable reasons for removal in this kind of questionnaire survey, as Pickin et al themselves acknowledge.

We argue that removal and the reasons underlying it remain poorly understood phenomena which require good quality qualitative research to address them. We have begun this process by undertaking an in depth interview study of general practitioners' accounts of removing patients and patients' accounts of being removed from a general practitioner's list. Our preliminary findings support the conclusion by Pickin et al that there is little evidence that patients are removed on economic grounds.

We think, however, that Pickin et al have presented an overly simplified view of what is an extremely complex and stressful process, which has profound implications for our understanding of the relationship between doctor and patient and how it should be managed. For example, the issue of formal

and informal practice policies on removal are absent from the study by Pickin et al, as is the role of relationships between the patient and other practice staff such as receptionists. Perhaps, most importantly, Pickin et al do not consider the concept of the "difficult" patient in general practice and the important notion of a breakdown in the relationship between doctor and patient, which is found in published guidance on removal.⁵

Future studies of patient removal should be more soundly based on detailed qualitative investigations to inform the content and design of instruments so that they add to our understanding of this important primary care issue.

Tim Stokes *clinical lecturer in general practice*
Department of General Practice and Primary Health Care, University of Leicester, Leicester General Hospital, Leicester LE5 4PW
tms2@le.ac.uk

Robert K McKinley *senior lecturer in general practice*
Department of General Practice and Primary Health Care

Mary Dixon-Woods *lecturer in health policy*
Department of Epidemiology and Public Health, University of Leicester, Leicester LE1 6TP

- Pickin M, Sampson F, Munro J, Nicholl J. General practitioners' reasons for removing patients from their lists: postal survey in England and Wales. *BMJ* 2001;322:1158-9.
- O'Reilly D, Steele K, Gilliland A, Brown S. Patient removals from general practitioner lists in Northern Ireland: 1987-1996. *Br J Gen Pract* 1998;48:1669-73.
- Stokes T, McKinley RK, Dixon-Woods M, Braidwood C. Patient removals from GPs' lists. *Br J Gen Pract* 1999;49:144.
- Perry J. *Removed from care: a report of patients removed from GP lists at the doctor's request*. Maidstone: Kent Family Health Services Authority, 1995.
- General Practitioners Committee of the British Medical Association. *Removal of patients from GP lists*. London: BMA, 1999. (<http://web.bma.org.uk/gpc.nsf>)

Duodenal haematoma occurs in children with bicycle injuries

EDITOR—Closed duodenal injury is a good topic to highlight as a lesson of the week. Surgical injuries to the upper abdomen are a fairly rare childhood injury, and few doctors will diagnose them with the necessary speed.¹ Lam et al remind us to examine the paediatric abdomen carefully and frequently after blunt trauma and not to rely too heavily on imaging.

One other type of intra-abdominal damage caused by handlebar injury is not highlighted in this paper. The first and third parts of the duodenum may be crushed between the handlebar and the lumbar spine, causing rupture of the subserosal blood vessels and resulting in a slowly expanding intramural haematoma.² The child may be relatively free of symptoms until the haematoma causes complete obstruction of the duodenal lumen; as a result, the diagnosis is often delayed.

Important clues to the diagnosis include the history of upper abdominal trauma, persistent vomiting, and air-fluid levels in both the stomach and duodenum (the so called double bubble) in a plain erect abdominal x ray film. If doubt still exists, contrast study of the upper gastrointestinal tract will show partial to complete obstruction of the

duodenum, with a "stacked coins" or "coiled spring" appearance caused by oedematous mucosal folds proximal and distal to the point of maximum obstruction.

Perhaps the most important reason to confirm this diagnosis is that, as an isolated injury, duodenal haematoma does not require laparotomy. Patients can be successfully managed with free nasogastric drainage and parenteral nutrition, and duodenal patency should return within 10-14 days. Clinical deterioration at any stage, or failure of the condition to resolve with this conservative regimen, should raise suspicion of a misdiagnosis or other concomitant injury.

Road traffic injury was the commonest cause of duodenal haematoma in the small series of nine cases in South Africa that Voss and I reported,² but bicycle injury predominated in a larger series from North America.³ There, as in the United Kingdom, more children are likely to own and ride bicycles than in South Africa, and motor vehicles are driven with more restraint than they are in South Africa.

David Bass *head, medicolegal services*
Western Cape Provincial Government, Cape Town,
South Africa 8001
dbass@global.co.za

1 Lam JPH, Eunson GJ, Munro FD, Orr JD. Lesson of the Week: Delayed presentation of handlebar injuries in children. *BMJ* 2001;322:1288-9. (26 May.)

2 Voss M, Bass DH. Traumatic duodenal haematoma in children. *Injury* 1994;25:227-30.

3 Jewett TC Jr, Caldarola V, Karp MP, Allen JE, Cooney DR. Intramural haematoma of the duodenum. *Arch Surg* 1988;123:54-8.

Systematic reviews and Cochrane Collaboration

Cochrane Library is now available free to people working in NHS

EDITOR—It is encouraging to see Müllner publicising the Cochrane Library as it is an essential resource for the practice of evidence based care.¹ One point that he didn't mention is that people working in the NHS can now access the full content of this collection of databases without charge through the National Electronic Library for Health.²

Donald M Mackay *inquiry services manager*
Cairns Library, University of Oxford, John Radcliffe Hospital, Oxford OX3 9DU
donald.mackay@cairns.ox.ac.uk

1 Müllner M. Website of the week: Systematic reviews and the Cochrane Collaboration. *BMJ* 2001;323:116. (14 July.)

2 www.nelh.nhs.uk

Free full text access to Cochrane articles is available to all health professionals

EDITOR—Müllner's website of the week is relevant to a variety of health professionals.¹ I am a frequent user of Cochrane for both teaching and patient care. Müllner says that it is available only on subscription, but there is a site providing free full text access to Cochrane to health professionals.² They have to register for gold services offered by

the site, but the registration is free. Many of my students and residents have registered and enjoy free full text access.

P Badrinath *clinical assistant professor*
Faculty of Medicine and Health Sciences, United Arab Emirates University, Al-Ain, United Arab Emirates
badrishanthi@hotmail.com

1 Müllner M. Website of the week: Systematic reviews and the Cochrane Collaboration. *BMJ* 2001;323:116. (14 July.)

2 www.obgyn.net/cochrane.asp

Doctors should not participate in executions

EDITOR—At the time that Dunea's article on the execution of Timothy McVeigh (the Oklahoma bomber) was published,¹ I was in Strasbourg participating in the first world congress against the death penalty.² Although some opponents of the United States death penalty may have drawn the line at the McVeigh case, no such sentiments were detected in Strasbourg. The presence at the congress of former (and innocent) residents of the death rows of the United States (Kerry Max Cook, 22 years under sentence of death) and Japan (Sakae Menda, 34 years under sentence of death) underlined an important, but not the sole, reason for concern about the use of the death penalty.

Opponents of the death penalty argue that one innocent life is too great a price to pay for carrying out executions. In Illinois, Governor George Ryan agreed and in January 2000 imposed a moratorium after 13 men were freed from death row after exoneration during the previous 13 years.

Dunea notes the high cost of the McVeigh trial but does not acknowledge that the high cost of capital trials is due to the death penalty. Irreversible penalties require absolute certainty and ample appeal processes. As the Illinois example shows, sophisticated legal systems fail to deliver safe convictions but do ensure that it costs more to kill prisoners with due process than to imprison them for life. Other studies have shown racial bias and failure to take account of mental illness and learning disabilities in those on trial for their lives.³ Above all, preserving respect for human rights is not effectively accomplished by putting individuals to death.

The study cited by Dunea showing willingness on the part of doctors in the United States to assist in executions, by Farber et al,⁴ does indeed indicate possible gaps between those elaborating ethical codes and doctors who are supposed to work within the codes' framework. Such discrepancies do not mean, though, that we should throw out the existing ethics and replace them with more populist ones. Doctors should not participate in executions; the challenge is to persuade all doctors that the medical role does not stretch to assisting in the killing of prisoners on behalf of the state.

James Welsh *coordinator, medical programme*
Amnesty International, London WC1X 0DW
medical@amnesty.org

- 1 Dunea G. Executing Timothy McVeigh. *BMJ* 2001;322:1553. (23 June.)
- 2 First world congress against the death penalty, Strasbourg, 21-23 June 2001. http://press.coe.int/dossiers/108/E/ecart_program.htm (accessed 27 June 2001).
- 3 Human Rights Watch. *Beyond reason: the death penalty and offenders with mental retardation*. New York: HRW, 2001. www.hrw.org/reports/2001/ustat/ (accessed 27 June 2001).
- 4 Farber N, Davis EB, Widern J, Jordan J, Boyer EG, Ubel PA. Physicians' attitudes about involvement in lethal injection for capital punishment. *Arch Intern Med* 2000;160:2912-6.

Plight of Afghan people must not be forgotten

EDITOR—We are deeply concerned about the potential humanitarian consequences of war in Afghanistan.

Afghanistan is one of the poorest countries in the world with a life expectancy of only 43 years, and a death rate in those under 5 of 257 per 1000 live births.¹ Drought and conflict in the region and have already generated over four million displaced people. Since the attack on the World Trade Centre and Pentagon thousands more have left Afghanistan's cities, making the situation even worse. The withdrawal of all international aid, including essential food supplies provided by the World Food Programme, owing to the threat of military action, is most likely to result in famine and increased death rates in the camps for internally displaced people. The human impact, particularly on children, could dwarf even the death toll arising from the terrorist attacks.

We therefore urge the UK government to give priority to protecting the health of the Afghan population to prevent a further humanitarian disaster.

Ben Armstrong *head, environmental epidemiology unit*
Michel Coleman *head, cancer and public health unit*
Clive Davies *head, disease control and vector biology unit*
Diana Elbourne *head, medical statistics unit*
Astrid Fletcher *head, epidemiology unit*
Emily Grundy *head, centre for population studies*
Andy Haines *dean*
Andy Hall *head, infectious disease epidemiology unit*
Betty Kirkwood *head, public health intervention research unit*
Donna Lamping *head, health services research unit*
Michael Miles *head, pathogen molecular biology and biochemistry unit*
Ian Roberts *professor of epidemiology and public health, public health intervention research unit*
Egbert Sondorp *senior lecturer, public health and humanitarian aid*
London School of Hygiene and Tropical Medicine, London

The authors are writing in a personal capacity.

1 United Nations Development Programme. *Human development report 2001: making technologies work for human development*. Oxford: Oxford University Press, 2001.

Rapid responses

Correspondence submitted electronically is available on our website