Postoperative starvation after gastrointestinal surgery

Type of intravenous nutrition given in control groups is not indicated

EDITOR—We appreciated Lewis et al's wise conclusions in their study: "there seems to be no clear advantage to keeping patients nil by mouth after elective gastrointestinal resection."¹ The subtitle of the accompanying editorial sounded less prudent: "early feeding is beneficial."² This positive statement stimulates readers to raise several questions that cannot be answered from the study. Which diet is beneficial? By which route? And compared with which intravenous feeding?

From table 1 we note that four types of diet were used in the 11 studies (standard, elemental, oral, and immune enhancing), through four different routes (oral, nasoduodenal, nasojejunal, and jejunostomy), giving rise to seven different combinations. No indication is given of the quality and quantity of the intravenous nutrition in the control groups. This is because the original studies were more concerned with feasibility, safety, and tolerance of enteral feeding than with its effect on postoperative outcome. The control diet was consequently defined only as conventional or traditional treatment or routine intravenous crystalloid solutions or hypocaloric fluids. Moreover, in one study the control group was not nil by mouth, as a placebo was given through a nasoduodenal feeding tube.³

This issue is not an academic one. In a randomised clinical trial Lobo et al showed in postoperative patients that salt and water restriction (water ≤ 2 l and sodium ≤ 77 mmol/day) was significantly better than standard management (water roughly 3 l and sodium roughly 154 mmol/day) in reducing postoperative gastric emptying times, time to passage of flatus and stools, and postoperative stay.⁴

In the absence of clear information on the composition of the intravenous support in the control arm we can speculate that the advantage of early feeding, as reported in this meta-analysis, reflects not a true benefit but simply the lack of detriment due to the intravenous standard treatment.

Finally, we agree with Lewis et al that the role of early enteral feeding should be assessed in a randomised clinical trial. Such a trial, sponsored by the Italian Society of Parenteral and Enteral Nutrition, was recently done in 317 malnourished patients with cancer who were candidates for elective

gastrointestinal surgery. They were randomised to two postoperative isoenergetic and isoprotein nutritional regimens given enterally or parenterally; the results have just been reported.⁵

F Bozzetti surgeon
L Mariani biostatistician
mariani@istitutotumori.mi.it

Department of Surgery and Biometric Unit,
National Cancer Institute, 1 20133 Milan, Italy

- 1 Lewis SJ, Egger M, Sylvester PA, Thomas S. Early enteral feeding versus "nil by mouth" after gastrointestinal surgery: systematic review and meta-analysis of controlled trials. *BMJ* 2001;323:773-6. (6 October.)
- 2 Silk DBA, Menzies Gow N. Postoperative starvation after gastrointestinal surgery. *BMJ* 2001;323:761-2. (6 October.)
- 3 Beier-Holgersen R, Boesby S. Influence of postoperative enteral nutrition on postsurgical infections. *Gut* 1996;39:833-5.
- 4 Lobo DN, Bostock KA, Neal KR, Perkins AC, Rowlands BJ, Allison SP. Effect of salt and water balance on gastrointestinal function and outcome after abdominal surgery: a prospective randomised controlled study. *Clin Nutr* 2001;20 (suppl 3):35-6.
- 5 Bozzetti F, Braga M, Gianotti L, Gavazzi C, Mariani L. Post-operative enteral versus parenteral nutrition in malnourished patients with gastrointestinal cancer: a randomised multicentre trial. *Lancet* 2001;358:1487-92.

Meta-analysis was not appropriate

EDITOR—We agree with Lewis et al that an adequately powered clinical trial is needed to determine the benefit of early enteral feeding.¹ But subjecting the disparate papers in their study to meta-analysis to justify this was inappropriate.

The papers included four different types of feed and four different routes of administration. Only three papers used the same combination of feed and route. Feeding proximal to the anastomosis was analysed with feeding distal to the anastomosis, which, if the authors believe that food "stresses" an anastomosis, was particularly inappropriate. Enteral feeding was purported to start within 24 hours of surgery, but in four of the papers (and over half of all patients) only clear enteral fluids were given initially, with nutrition being withheld until these were tolerated.²⁻⁵ Moreover, it cannot be assumed that patients assigned to receive true nutrition actually received their metabolic requirements.

The authors considered anastomotic leakage a crucial end point. However, data regarding site of anastomosis were available for only eight of the 11 studies; in two of the six in which the anastomosis was distal to the site of feeding, anastomotic leakage was not recorded. The authors state that when combined "the effects of early enteral feeding failed to reach conventional levels of significance, but eight of nine studies ... reported

benefit." In so doing they have ignored their own statistics in favour of anecdote.

Hospital stay for postoperative patients is not normally distributed, and therefore the median and not the mean is the appropriate measure of spread. The significance of the small reduction in hospital stay might not be maintained if non-parametric analysis was performed. The definition of infection in the papers assessed was not clearly defined; it was therefore inappropriate to analyse a group presumed to comprise "all infections."

The authors themselves state that the trials included are "clinically heterogeneous," are "small and of doubtful methodological quality," and have "inadequate reporting," all of which make them unsuitable for meta-analysis. There is now the risk that this paper will be seen as an evidence based medicine resource, which it is not.

P B Goodfellow specialist registrar
Petegoodfellow@bigfoot.com

N J Everitt consultant surgeon
Chesterfield and North Derbyshire Royal Hospital,
Chesterfield S44 5BL

- 1 Lewis SJ, Egger M, Sylvester PA, Thomas S. Early enteral feeding versus "nil by mouth" after gastrointestinal surgery: systematic review and meta-analysis of controlled trials. *BMJ* 2001;323:773-6. (6 October.)
- 2 Ortiz H, Armendariz P, Yarnoz C. Is early postoperative feeding feasible in elective colon and rectal surgery? *Int J Colorectal Dis* 1996;11:119-21.
- 3 Reissman P, Teoh TA, Cohen SM, Weiss EG, Nogueras JJ, Wexner SD. Is early oral feeding safe after elective colorectal surgery? A prospective randomized trial. *Ann Surg* 1995;222:73-7.
- 4 Stewart BT, Woods RJ, Collopy BT, Fink RJ, Mackay JR, Keck JO. Early feeding after elective open colorectal resections: a prospective randomized trial. *Aust N Z J Surg* 1998;68:125-8.
- 5 Hartsell PA, Frazee RC, Harrison JB, Smith RW. Early postoperative feeding after elective colorectal surgery. *Arch Surg* 1997;132:518-20; discussion 520-1.

Authors' reply

EDITOR—We were interested in the pragmatic comparison of early compared with deferred feeding strategies after gastrointestinal surgery. Results from trials included in our review were fairly homogeneous despite clinically heterogeneous study populations and interventions. It therefore was appropriate to calculate average estimates of the effects of early feeding using meta-analysis. We modestly concluded that early feeding may be beneficial and that a large trial is required to answer this question.

The homogeneity between studies suggests that there may be a common underlying mechanism similar to the class effects observed in drug treatment. The mechanism underlying any beneficial effect of early feeding is unclear, but meeting full metabolic requirements in the early postoperative period may not be necessary. If the

effect is systemically mediated then benefit will be seen despite feeding distal to an anastomosis. The importance of the amount and type of feed given remains speculative.

We agree with Bozzetti and Mariani that the type of enteral nutrition and amount of intravenous fluids given to control groups may have been important. Careful consideration will have to be given to control interventions in future trials. The results from their recent trial, which showed that early postoperative enteral nutrition compared to isocaloric and isonitrogenous parenteral nutrition is beneficial, are compatible with the findings from our review.¹

Goodfellow and Everitt claim that in four of the studies included in our analysis (Hartsell et al, Ortiz et al, Stewart et al, Reissman et al) oral intake during the first 24 hours was restricted to clear enteral fluids. This is incorrect. In the trial by Hartsell et al liquid enteral nutrition was started on the day of operation, and in the other three studies oral nutrition was offered, as tolerated, to patients on their first postoperative day.

Goodfellow and Everitt also misinterpret the role of systematic reviews and meta-analysis. In systematic reviews all relevant studies meeting eligibility criteria are identified through extensive literature searches, the methodological quality is carefully assessed, and results are examined on a common scale.² If, after careful consideration, a meta-analysis is deemed appropriate, the last step consists of estimating an average effect by combining the data. Sometimes definitive answers emerge. More often systematic reviews conclude, as we did, that the available evidence is insufficient and that new trials are required. Reviews such as ours are nevertheless an important resource in evidence based health care.

Stephen J Lewis *consultant*
Department of Medicine, Addenbrooke's Hospital,
Cambridge CB2 2QQ

Matthias Egger *senior lecturer in epidemiology and public health medicine*
MRC Health Services Research Collaboration,
Department of Social Medicine, University of
Bristol, Bristol BS8 2PR

Paul A Sylvester *specialist registrar*
Department of Surgery, Bristol Royal Infirmary,
Bristol BS2 8HW

Steven Thomas *senior lecturer*
Department of Maxillofacial Surgery, University of
Bristol, Bristol BS1 2LY

1 Bozzetti F, Braga M, Gianotti L, Gavazzi C, Mariani L. Post-operative enteral versus parenteral nutrition in malnourished patients with gastrointestinal cancer: a randomised multicentre trial. *Lancet* 2001;358:1487-92.

2 Egger M, Davey Smith G. Principles and procedures. In: Egger M, Smith DG, Altman DG, eds. *Systematic reviews in health care: meta-analysis in context*. London: BMJ Books, 2001.

Anaesthetic technique during gastrointestinal surgery has postoperative effects

EDITOR—We were surprised that neither the paper by Lewis et al nor the accompanying editorial mentioned the anaesthetic technique used during gastrointestinal surgery.^{1,2} There are several ways in which anaesthesia affects the return of gastrointestinal function and thus resumption of feeding.

Epidural anaesthesia is frequently used for this type of surgery. When continued into the postoperative period epidurals have several well recognised benefits, including pain relief, a reduction in pulmonary thromboembolism and other pulmonary complications, and suppression of the neuroendocrine stress response. In addition, data confirm a reduction in postoperative ileus and improvement in gastrointestinal function after surgery, with the mean time from surgery to first flatus and first bowel movement occurring earlier in epidural groups than control groups receiving parenteral morphine. The most effective method seems to be thoracic epidural analgesia with plain bupivacaine.³

The anaesthetic technique may affect anastomotic leakage rates. Despite unopposed parasympathetic activity, epidurals do not increase leak rates.⁴ The use of neostigmine to antagonise neuromuscular blockade may increase bowel motility and result in a higher rate of anastomotic leakage. It is now recognised, however, that in many cases modern neuromuscular blocking drugs need not be antagonised or a reduced dose of antagonist can be given.⁵

Holte and Kehlet's review on preventing postoperative ileus recommends the use of continuous thoracic epidural anaesthesia for at least 48 hours and the avoidance of opioid analgesia where possible.⁶ Their multimodal approach, in conjunction with early feeding, avoidance of nasogastric intubation, and possibly the use of prokinetic drugs, surely provides the direction over the next 10 years for surgeons and anaesthetists in the management of patients after gastrointestinal surgery.

W J Fawcett *consultant anaesthetist*
wjfawcett@netscapeonline.co.uk

W E J Jewsbury *specialist registrar in anaesthetics*
Anaesthetic Department, Royal Surrey County
Hospital, Guildford GU2 7XX

1 Lewis SJ, Egger M, Sylvester PA, Thomas S. Early enteral feeding versus "nil by mouth" after gastrointestinal surgery: systematic review and meta-analysis of controlled trials. *BMJ* 2001;323:773-6. (6 October.)

2 Silk DBA, Menzies Gow N. Postoperative starvation after gastrointestinal surgery. *BMJ* 2001;323:761-2. (6 October.)

3 Holte K, Kehlet H. Postoperative ileus: a preventable event. *Br J Surg* 2000;87:1480-93.

4 Holte K, Kehlet H. Epidural analgesia and risk of anastomotic leakage. *Regional Anesthesia and Pain Medicine* 2001;26:111-7.

5 Hunter JM. Is it always necessary to antagonise neuromuscular block? Do children differ from adults? *Br J Anaesth* 1996;77:707-9.

Rest in peace, drip and suck

EDITOR—Excellent! Another nail in the coffin that lays to rest the century-long general medical fashion that motion and activity are detrimental to healing.

I should confess that "drip and suck" is not my business, it being over 30 years since, as a general surgical registrar, I last cut and cared for a patient's belly; I am now a retired orthoped. The excuse for my tongue in cheek comment is an interest in the history of how rest came to be so widely adopted as beneficial—nay, essential—to recovery from virtually all medical or surgical conditions. Whether by resting the whole patient, such as by bed rest or prolonged convalescence,

or by resting a part, as with a splint or by nasogastric suction, all specialties became so indoctrinated in this dogma that most of us had to be pushed kicking and screaming to abandon it once we were shown that it had no scientific foundation.

Until recently it would have been unthinkable to repair a hernia as a day case procedure and for the patient to return to work in a fortnight; to give treadmill exercise to a patient with an infarcted heart only days after he or she came out of intensive care; or for a mother to walk about and go home the day her baby was born. Only in the past few years have we orthopods learnt to abandon bed rest for back pain. Now we learn that we can also safely feed a patient with an anastomosis: goodbye, drip and suck!

For most conditions the fashion to rest our patients seems to have started in the latter part of the 19th century, risen to a zenith of zealous application by the second quarter of the 20th, and then taken since the 1960s to be slowly debunked. I have yet to discover when and who invented and pioneered nasogastric suction for this purpose, and what arguments they used to explain their rationale (old surgical texts are remarkably bashful in explaining why); I hope that some reader can tell me something about the inception of this technique.

Paul Moynagh *retired consultant orthopaedic surgeon*
Chase Farm Hospital, Enfield EN2 8JL
pmoynagh@argonet.co.uk

Cardiotocography v Doppler auscultation

Guidelines highlight gaps in research evidence

EDITOR—The clinical effectiveness support unit of the Royal College of Obstetricians and Gynaecologists has recently developed the guideline of the National Institute for Clinical Excellence (NICE) on the use of electronic fetal monitoring.¹ One of the potential benefits of guidelines is to highlight gaps in research evidence. As part of the guideline development process we conducted a survey of maternity units in England and Wales to evaluate the current use of electronic fetal monitoring in intrapartum care.²

The guideline recommends that in women who are healthy and have an uncomplicated pregnancy, intermittent auscultation is a suitable method of monitoring during labour. This recommendation was based on the evidence from nine randomised controlled trials that have enrolled 18 000 and are included in three systematic reviews.¹ This conclusion was reached because screening with continuous electronic fetal monitoring should not be conducted in the absence of evidence of benefit. This recommendation will perhaps have a minimal impact in practice as the survey found that only 2.5% of units report using continuous electronic fetal monitoring for all women in labour.

Most maternity units (79%), however, reported using admission cardiotocography in the initial assessment of women who are healthy and have had an otherwise uncomplicated pregnancy. When evaluating the research that exists to support the use of admission tests in populations at low risk we were surprised to find only a single cohort study that addressed this issue; this cohort study included only 1041 women and related admission tests to fetal distress.³ The trial by Mires et al is the first published randomised controlled trial to address this clinically important question. The study is also nearly four times the sample size of previous studies. As such it would have been important to publish this trial irrespective of whether the results were positive, negative, or equivocal.⁴

The results of the trial are in line with those of the earlier cohort study and do not change the recommendation in the guideline. We are aware of unease in some quarters that the guideline did not recommended admission cardiotocography and that this recommendation does not concur with current practice in many maternity centres. We hope that this knowledge will act as an incentive to researchers and clinicians to conduct and enrol patients in randomised controlled trials to answer this important question.

Jane Thomas *director*
Shantini Paranjothy *clinical research fellow*
Tony Kelly *clinical research fellow*
Josephine Kavanagh *research fellow*
 Royal College of Obstetricians and Gynaecologists,
 Clinical Effectiveness Support Unit, London
 NW1 4RG

- 1 Clinical Effectiveness Support Unit. *The use of electronic fetal monitoring. The use and interpretation of cardiotocography in intrapartum fetal surveillance*. London: Royal College of Obstetricians and Gynaecologists, 2001.
- 2 Maternal and Child Health Research Consortium. *Confidential enquiry into stillbirths and deaths in infancy: eighth annual report*. London: Department of Health, 2001.
- 3 Ingemarsson I, Arulkumaran S, Ingemarsson E, Tambyrajah RL, Ratnam SS. Admission test: a screening test for fetal distress in labor. *Obstet Gynecol* 1986;68:800-6.
- 4 Mires G, Williams F, Howie P. Randomised controlled trial of cardiotocography versus Doppler auscultation of fetal heart at admission in labour in low risk obstetric population [with commentaries by S Goldbeck-Wood, GD Murray, B-I Nesheim]. *BMJ* 2001;322:1457-62. (16 June.)

Conclusions do not recognise difference between statistical and clinical significance

EDITOR—Mires et al conclude that admission cardiotocography does not benefit neonatal outcome in women at low risk, but that it does increase the rate of operative delivery.¹ No information is given regarding the classification of the admission traces in their study. Other electronic responses, by Murphy and Chalmers (published as the next letter in this cluster),^{2,3} have addressed the likely cause of this increased operative delivery rate and the suggestion that lack of effect can be proved.

An earlier study, although not randomised, was a prospective blinded study of women designated as low risk on admission in labour and included a similar number of women to the cardiotocography arm of the current study once defined as low risk at labour onset (1041 compared with 1186).⁴ This study showed a 1% incidence of an "ominous" fetal heart rate trace on admission testing and, of these 10 infants, one died three

hours after admission, without stethoscopic auscultation detecting any fetal compromise.

This again highlights the point made by Chalmers (next letter).³ The conclusions drawn from the current study do not recognise the difference between statistical and clinical significance in an individual case, particularly when the adverse outcome is rare. Intermittent auscultation will not detect the development of all abnormalities in fetal heart rate and depends on the ability to provide adequate levels of midwifery care. The inability to take the cardiotocography off after an admission test is as much a training issue as the inability to recognise as abnormal the cardiotocography identified by the Confidential Enquiry into Stillbirths and Deaths in Infancy.⁵

Simon Grant *consultant in obstetrics and gynaecology*
 Royal Cornwall Hospital (Treliske), Truro TR1 3LJ
 simon.grant@rcht.swest.nhs.uk

- 1 Mires G, Williams F, Howie P. Randomised controlled trial of cardiotocography versus Doppler auscultation of fetal heart at admission in labour in low risk obstetric population [with commentaries by S Goldbeck-Wood, GD Murray, B-I Nesheim]. *BMJ* 2001;322:1457-62. (16 June.)
- 2 Murphy DJ. Electronic responses. Randomised controlled trial of cardiotocography versus Doppler auscultation of fetal heart at admission in labour in low risk population. *bmj.com* 2001;322 (www.bmj.com/cgi/eletters/322/7300/1457#15117).
- 3 Chalmers I. Electronic responses. Randomised controlled trial of cardiotocography versus Doppler auscultation of fetal heart at admission in labour in low risk population. *bmj.com* 2001;322 (www.bmj.com/cgi/eletters/322/7300/1457#15118).
- 4 Ingemarsson I, Arulkumaran S, Ingemarsson E, Tambyrajah RL, Ratnam SS. Admission test: a screening test for fetal distress in labor. *Obstet Gynecol* 1986;68:800-6.
- 5 Confidential Enquiry into Stillbirths and Deaths in Infancy. *1st annual report*. London: Department of Health, 1995.

All unbiased comparative studies should be published

EDITOR—Goldbeck-Wood's invitation to respond to questions raised by the first randomised comparison of cardiotocography versus auscultation of the fetal heart at admission in labour prompts me to emphasise the following points.¹ The term "negative trial" should be outlawed, as should editorial acquiescence in statements implying that it is possible to prove negatives.² For example, the *BMJ* report of the admission cardiotocography trial states that this intervention does not benefit neonatal outcome in women at low risk, yet the estimate of its effect on the incidence of hypoxic ischaemic encephalopathy is compatible with the reduction in neonatal seizures found in other randomised controlled trials of fetal monitoring during labour.³

Although it is worth making pretrial estimates of the sample sizes needed to rule in or rule out, with specified statistical confidence, effects of a magnitude likely to be relevant to patients, how many participants you needed in a trial depends on what you found, not on what you thought you would find.⁴

Statements that it is unethical to embark on controlled trials unless an arbitrarily defined level of statistical power can be assured make no sense if the alternative is acquiescence in ignorance of the effects of healthcare interventions. Unbiased estimates should be generated through randomised trials because they leave us less ignorant than we were before.⁵ The importance

of the admission cardiotocography study is that it provides the first unbiased estimates of the ranges within which the effects of this healthcare intervention are likely to lie. All unbiased comparative studies should be published, so that the totality of the relevant evidence can be evaluated in the systematic reviews that are needed by those making choices and decisions in health care and about further research. It is now possible to register and publish successive versions of trial protocols and all subsequent reports of registered studies. Reluctance to exploit this potential does not reflect scientific or technical problems but rather academic and commercial interests.

Iain Chalmers *director*
 UK Cochrane Centre, Oxford OX2 7LG
 ichalmers@cochrane.co.uk

- 1 Mires G, Williams F, Howie P. Randomised controlled trial of cardiotocography versus Doppler auscultation of fetal heart at admission in labour in low risk obstetric population [with commentaries by S Goldbeck-Wood, GD Murray, B-I Nesheim]. *BMJ* 2001;322:1457-62. (16 June.)
- 2 Chalmers I. Proposal to outlaw the term "negative trial." *BMJ* 1985;290:1002.
- 3 Thacker SB, Stroup D, Chang M. Continuous electronic heart rate monitoring for fetal assessment during labor. *Cochrane Library*. Issue 1. Oxford: Update Software, 2002.
- 4 Detsky AS, Sackett DL. When was a "negative" clinical trial big enough? How many patients you needed depends on what you found. *Arch Intern Med* 1985;145:709-12.
- 5 Lilford RJ, Thornton JG, Braunholtz D. Clinical trials and rare diseases: a way out of a conundrum. *BMJ* 1995;311: 1621-5.

Trials are underpowered

EDITOR—Goddard in her editorial acknowledged the fact that trials of routine electronic fetal monitoring are underpowered, but she did not draw the logical conclusion.¹ With her figures for intrapartum stillbirth and cerebral palsy of intrapartum origin, a trial powered (80% power, alpha 5%, two-sided) to show a 50% decrease in intrapartum stillbirth would need to recruit 83 500 women to each arm, and a trial powered to detect a 50% decrease in cerebral palsy due to intrapartum causes would require 670 000 women in each arm. The current meta-analysis has less than 19 000 cases in total.² The evidence is not, as she states, strongly against the routine use of electronic fetal monitoring. The evidence simply does not exist and, given the numbers required, it probably never will.

Routine electronic fetal monitoring halves the risk of neonatal seizures even when applied to a low risk population.³ Statements to the effect that electronic fetal monitoring has not been shown to reduce the risk of intrapartum stillbirth and cerebral palsy have the same fundamental weaknesses as statements associated with some previous public health disasters, such as there is no evidence to suggest that blood products can transmit HIV or that there is no evidence that bovine spongiform encephalopathy can be transmitted to humans. Falsely equating absence of evidence with evidence of absence does not have a glorious history and should be avoided.

Gordon C S Smith *subspecialist trainee, maternal-fetal medicine*
 Department of Obstetrics and Gynaecology,
 The Queen Mother's Hospital, Glasgow G3 8SH
 gcs4@cornell.edu

- 1 Goddard R. Electronic fetal monitoring. *BMJ* 2001;322:1436-7. (16 June.)
- 2 Thaker SB, Stroup DF, Chang M. Continuous electronic heart rate monitoring for fetal assessment during labor (Cochrane Review). *Cochrane Database Syst Rev* 2001;2:CD000063.

Evidence is not as good as NICE suggests

EDITOR—The recently published clinical guideline from the National Institute for Clinical Excellence (NICE) on the use of electronic fetal monitoring and the clinical guideline from the Royal College of Obstetricians and Gynaecologists from which the NICE guideline is derived recommend that intermittent auscultation is the most appropriate method of fetal monitoring for low risk women in labour.^{1 2}

The guideline development groups erroneously conclude that there is grade A evidence (at least one randomised controlled trial addressing the specific recommendation) to support intermittent auscultation every 15 minutes in the first stage of labour and every 5 minutes in the second stage. The College guideline, however, states that there are no studies evaluating different protocols for frequency of intermittent auscultation. Any recommendation on frequency of auscultation has to be grade C (evidence from expert committee reports or opinions or clinical experience of respected authorities, or both). This is an important point because the NICE guideline is likely to be held by the courts as the required standard of care in labour. This degree of fetal monitoring may be impossible to achieve in hard pressed labour wards and may encourage the use of continuous electronic fetal monitoring instead.³ For women in our unit who are at low risk we are currently auscultating the fetal heart every 30 minutes in the first stage and after every contraction in the second stage of labour.

J P Stuart consultant in obstetrics and gynaecology
Diana, Princess of Wales Hospital, Grimsby
DN33 2BA

- 1 National Institute for Clinical Excellence. *The use of electronic fetal monitoring: the use and interpretation of cardiotocography in intrapartum fetal surveillance*. London: NICE, 2001.
- 2 Clinical Effectiveness Support Unit. *The use of electronic fetal monitoring. The use and interpretation of cardiotocography in intrapartum fetal surveillance*. London: Royal College of Obstetricians and Gynaecologists, 2001.
- 3 Goddard R. Electronic fetal monitoring. *BMJ* 2001;322:1436-7. (16 June.)

Author of editorial's reply

EDITOR—The publication of the paper by Mires et al and my accompanying editorial have resulted in a wide range of responses from both sides of the argument. This issue has long been controversial and, as indicated in the letters by Smith and Grant, is likely to remain so as it may prove impossible to carry out a large enough trial with sufficient power to answer the key question: Can electronic fetal monitoring either on admission or during labour reduce the risk of poor obstetric outcome by identifying those infants who are stressed by labour where early intervention will improve the outcome?

Unfortunately, cardiotocography is a poor screening test, especially in low risk women, which has been widely introduced

into practice without the evidence for its value. There are wide variations in interpreting traces. It is also a test that is highly sensitive for a rare disease but has a low specificity.¹ As a consequence, the false positive rate is high, leading to a high intervention rate. Although many obstetricians such as Smith and Grant feel that any increase in caesarean rate is justified if there is perceived to be a neonatal benefit, many others disagree particularly where epidemiological data would suggest that there has been no effect on mortality. In particular, midwives, pregnant women themselves, and other professionals who may have to deal with the long term effects of our management are dismayed at the escalating intervention rate that has occurred as a consequence of using electronic monitoring in low risk women.^{2 3}

There is no grade A evidence for the form of intermittent auscultation. There are randomised controlled trials of electronic fetal monitoring versus intermittent auscultation but no up to date studies comparing different methods of intermittent auscultation.¹ The aim of the guideline was to look at electronic fetal monitoring and not intermittent auscultation. The recommendations on the form of intermittent auscultation used are therefore based on randomised controlled trials comparing electronic fetal monitoring and intermittent auscultation not different methods of intermittent auscultation. Many units do not have the staffing levels to achieve the level of auscultation suggested by the guideline. Thomas et al rightly say that a properly formed trial of which is the best method of intermittent auscultation needs to be done. Although the paper by Mires et al is flawed, it has stimulated this important debate, which will hopefully provide the impetus for a larger trial to answer this important question.

Ros Goddard specialist registrar
Royal United Hospital, Bath BA1 3NG
ros.goddard@hotmail.com

A longer version of this letter is published on bmj.com

- 1 Clinical Effectiveness Support Unit. *The use of electronic fetal monitoring. The use and interpretation of cardiotocography in intrapartum fetal surveillance*. London: Royal College of Obstetricians and Gynaecology, 2001.
- 2 Electronic responses. Randomised controlled trial of cardiotocography versus Doppler auscultation of fetal heart at admission in labour in low risk population. bmj.com 2001;322 (www.bmj.com/cgi/eletters/322/7300/1457#responses).
- 3 Electronic responses. Electronic fetal monitoring. bmj.com 2001;322 (www.bmj.com/cgi/content/full/322/7300/1436#responses).

Authors of paper's reply

EDITOR—Thomas et al recognise the importance of our study as the first randomised trial of the value of the admission cardiotocograph in low risk labour. Despite the paucity of evidence to support its use, some 80% of obstetric units in the United Kingdom have introduced the admission cardiotocograph into routine clinical practice. The evidence based guideline from the Royal College of Obstetricians and Gynaecologists on the use of electronic fetal monitoring did not recommend the use of the admission cardio-

tocograph, and the results of our trial are in line with this recommendation.¹

Grant and Chalmers raise concerns about our conclusion that the admission cardiotocograph does not benefit neonatal outcome in women at low risk. Our conclusion relates to the presence of metabolic acidosis at delivery, which was our primary outcome. Umbilical arterial acid base status is one of the recognised intermediate fetal or neonatal measures of fetal hypoxia and as such is an appropriate surrogate for fetal condition at birth.¹

We acknowledge that our sample size was not adequate to provide definitive information on secondary outcomes such as hypoxic ischaemic encephalopathy and that our findings are compatible with a reduction in this hypoxic complication. Trials to evaluate this and other uncommon outcomes would need to be much larger. Chalmers indicates that our trial provides the first unbiased estimates of the ranges within which the effects of this healthcare intervention are likely to lie. Our data will therefore inform future sample size calculations.

Grant refers to a previous study in which the only intrapartum death was in a woman with an ominous admission test in which intermittent auscultation did not identify a problem with the fetal heart pattern.² He raises concerns about the difference between statistical and clinical significance in an individual case, particularly when the adverse outcome is rare. The trial by Ingemarsson et al was not randomised, and the admission test was not available to the clinician managing the labour, with the interpretation and classification of traces being retrospectively assigned. It cannot be assumed that if the admission test had been available to the clinician that the outcome would have been different.

It is feasible that when the admission test was performed that the fetus could have already suffered irreversible hypoxic damage. In addition few details are given about this baby, which was in a low risk group defined in the study as patients recruited in the first stage of labour with a pregnancy of 34 weeks or more, practically all of whom had attended antenatal clinics. Our study showed a significantly higher incidence of operative delivery in the admission test group. As well as the fetal outcome, maternal outcome needs to be considered. Caesarean section is associated with an increased risk of maternal death compared with vaginal delivery, and both caesarean section and deliveries by forceps or ventouse are associated with increased risk of severe maternal morbidity.^{3 4} This is in addition to the increased potentially serious neonatal morbidity associated with forceps and ventouse—for example, cephalhaematoma.⁴

We recognise that further trials need to be undertaken to confirm our findings, and provide more evidence for the use of the admission cardiotocograph. In association with Thomas et al, we hope that the results of our trial will act as an incentive to researchers and clinicians to conduct and

enrol patients into randomised trials to further evaluate the role of the admission cardiocardiograph.

G J Mires senior lecturer, department of obstetrics and gynaecology
F L R Williams lecturer, department of epidemiology and public health
P W Howie professor, department of obstetrics and gynaecology
 Ninewells Hospital and Medical School, Dundee DD1 9SY

- 1 Clinical Effectiveness Support Unit. *The use of electronic fetal monitoring. The use and interpretation of cardiotocography in intrapartum fetal surveillance*. London: Royal College of Obstetricians and Gynaecologists, 2001.
- 2 Ingemarsson I, Arulkumaran S, Ingemarsson E, Tambyraja RL, Ratnam SS. Admission test: a screening test for fetal distress in labour. *Obstet Gynecol* 1986;68:800-6.
- 3 Waterstone M, Bewley S, Wolfe C. Incidence and predictors of severe obstetric morbidity: case controlled study. *BMJ* 2001;322:1089-94.
- 4 Johansen RB, Menon BK. Vacuum extraction versus forceps for assisted vaginal delivery. *Cochrane Library*. Issue 4. Oxford: Update Software, 2001.

Gastro-oesophageal reflux disease in general practice

Last year's guidelines are probably biased

EDITOR—We read with interest last year's paper by Dent et al on the management of gastro-oesophageal reflux disease in general practice.¹ Dent et al are closely connected to AstraZeneca, as is shown by the competing interests declared at the end of the article. The same company participated in the preparation of the manuscript. The article is based on the view of participants at a workshop arranged by AstraZeneca, which accordingly determined who should be invited. We suppose that the questions posted at the workshop had been made by AstraZeneca before the selected "experts" were asked to answer them by putting their finger on the button. The votes from such a highly selected panel are now, more or less, presented as a guideline for general practitioners in the treatment of gastro-oesophageal reflux disease.

It is not surprising that a pharmaceutical company is trying to sell its products, and it is not surprising that doctors closely connected to pharmaceutical companies write such a manuscript. It is, however, remarkable that a reputable journal like the *BMJ* accepts such a manuscript for publication.

Helge L Waldum professor of medicine
Eiliv Brenna professor of medicine
 Department of Intra-Abdominal Diseases, Faculty of Medicine, Norwegian University of Science and Technology, Regionsykehuset i Trondheim, N-7006 Trondheim, Norway

- 1 Dent J, Jones R, Kahrlas P, Talley NJ. Management of gastro-oesophageal reflux disease in general practice. *BMJ* 2001;322:344-7.

Utility and acceptability of Infai C13-urea breath test has been shown

EDITOR—Last year's clinical review by Dent et al on the management of gastro-oesophageal reflux disease in general practice provides another interesting perspective on this problem.¹ Little reference has, however, been made to testing for and eradication of *Helicobacter pylori* infection in

this latest review. Moayyedi et al published a meta-analysis in the *BMJ* showing a small but definite benefit to treating patients who have non-ulcer dyspepsia for *H pylori* infection.² They said that eradication of *H pylori* might be a cost effective treatment for non-ulcer dyspepsia in infected patients. The accompanying editorial by Veldhuyzen van Zanten recommended that patients younger than 50 years without symptoms should be tested for *H pylori* with a non-invasive test and if the test is positive the patient should be treated.³

Regarding testing for *H pylori*, most guidelines are recommending the urea breath test over serological testing, and according to a recent summary of the current evidence serology testing is now not considered accurate enough for a diagnosis of infection.⁴

The breath test kits are available on a FP10 prescription but are rarely used, and no studies have been published assessing their use within primary care and their acceptability to patients. We performed a small feasibility study in two general practices on Teesside to test the utility and acceptability of the Infai C13-urea breath test over one year.⁵ Eighty eight patients were included in the study, and a practice nurse performed 90% of the tests in 10 minutes or less. Patient satisfaction with the test was found to be high, according to self completed questionnaires. All but one of the patients were satisfied with how long the test took to perform and were happier coming to their general practice rather than a hospital to perform the test. No patients found the urea breath test unpleasant, and all except one would recommend the test to someone else. We concluded that the test could be easily incorporated into clinical practice. *H pylori* testing does seem to still have a place in the primary care management of gastro-oesophageal disease despite its scarce reference in this latest review.

Hugh Alberti general practitioner
 Cleveland Health Centre, 20 Cleveland Square, Middlesbrough TS1 2NX
 hughalberti@ntlworld.com

- 1 Dent J, Jones R, Kahrlas P, Talley NJ. Management of gastro-oesophageal reflux disease in general practice. *BMJ* 2001;322:344-7.
- 2 Moayyedi P, Soo S, Deeks J, Forman D, Mason J, Innes M, et al. Systematic review and economic evaluation of *Helicobacter pylori* eradication therapy for non-ulcer dyspepsia. *BMJ* 2000;321:659-64.
- 3 Veldhuyzen van Zanten SJO. Treating non-ulcer dyspepsia and *H pylori*. *BMJ* 2000;321:648-9.
- 4 Vaira D, Vakil N. Blood, urine, stool, breath, money and *Helicobacter pylori*. *Gut* 2001;48:287-9.
- 5 Heywood P, Alberti H. Utility and acceptability of the Infai C-13 breath test in general practice. *Cleveland Med J* 2001;4:116-8.

Authors' response

EDITOR—Waldum and Brenna have, in the past, expressed their concerns about profound acid inhibition, and several of their interpretations of experimental and clinical data have been discussed in a subsequent publication.^{1 2} We were surprised, therefore, to read their response to our article on the management of gastro-oesophageal reflux disease in general practice and would like to make some comments.

Firstly, although we have been connected to research undertaken by AstraZeneca, the paragraph in which we declare our competing interests also makes it clear that we have had connections with other companies who manufacture not only proton pump inhibitors but also other drugs used in the treatment of gastro-oesophageal reflux disease. It would be extraordinary if four clinicians with major research interests in the management of gastro-oesophageal reflux disease did not have some involvement in the design and conduct of clinical trials undertaken by the manufacturers of these agents.

Secondly, Waldum and Brenna are critical about basing our article on the Genval workshop. They did not participate in this workshop and do not know the process involved. The voting statements on which the recommendations were based were written independently of any input by AstraZeneca by a core group of experts that planned the scientific aspects of the workshop. Participants in this workshop were drawn from diverse backgrounds, and each played an important part in reviewing scientific data relevant to two or three workshop statements assigned to them. These reviews were presented at the workshop and discussed, and at no stage did AstraZeneca staff have any input into this discussion. The findings of the workshop were subsequently peer reviewed and published.³

Thirdly, Waldum and Brenna do not identify a single recommendation, made either in our article or by the Genval workshop, with which they take issue. Other responses to this article have been generally supportive and have also included discussion of the role of upper gastrointestinal radiology, the importance of obesity as a risk factor for gastro-oesophageal reflux disease, and the place of *Helicobacter pylori* infection and eradication, as discussed below.

Finally, Waldum and Brenna's letter criticises *BMJ* publishing policy and raises the more general question of the ethical and scientific frameworks within which research sponsored by pharmaceutical companies should be published. Given that a substantial proportion of research and development work in the United Kingdom and elsewhere is funded by the pharmaceutical industry, often in an entirely hands off mode, it is inevitable that some important and groundbreaking clinical research is supported in this way. Most journals now require a clear declaration of competing interests and of the contributions of individual authors which, together with a rigorous and transparent peer review process, should go a long way towards ensuring probity and objectivity.

Alberti has reminded us that we did not discuss *H pylori* in great detail. Although the potential role of infection with and eradication of *H pylori* in patients with dyspepsia (as distinct from gastro-oesophageal reflux disease) remains controversial, we are not aware of any strong evidence to implicate *H pylori* infection in the development or progression of reflux disease. Moayyedi et al, in their large

community based studies, have not shown a clinically relevant change in reflux symptoms after eradication of *H pylori*.⁴

Roger Jones *professor of general practice*
King's College London, London SE11 6SP
roger.jones@kcl.ac.uk

John Dent *professor of medicine*
Royal Adelaide Hospital, Adelaide, South Australia
5000, Australia
jdent@mail.rah.sa.gov.au

Peter Kahrilas *professor of medicine*
North Western University Medical School, Chicago,
IL 60611-3008, USA
p-kahrilas@northwestern.edu

Nicholas J Talley *professor of medicine, University of Sydney*
Nepean Hospital, Penrith, PO Box 63, New South
Wales 2751, Australia
talley@pnc.com.au

- 1 Waldum HL, Brenna E. Personal review: is profound acid inhibition safe? *Aliment Pharmacol Ther* 2000;14:15-22.
- 2 Yeomans ND, Dent J. Personal review: alarmism or legitimate concerns about long-term suppression of gastric acid secretion? *Aliment Pharmacol Ther* 2000;14:267-71.
- 3 Dent J, Brun J, Fendrick AM et al. An evidence-based appraisal of reflux disease management—the Genval workshop report. *Gut* 1999;44:S1-16.
- 4 Moayyedi P, Feltbower R, Brown J, Mason S, Mason J, Nathan J, et al. Effect of population screening and treatment for *Helicobacter pylori* on dyspepsia and quality of life in the community: a randomised controlled trial. Leeds HELP Study Group. *Lancet* 2000;355:1665-9.

Caesarean section rate in England and Wales

Maybe midwives were too busy

EDITOR—Dobson reports that deliveries by caesarean section in England rose from 4% in 1970 to 19% in 1999.¹ Potentially, this created substantially increased demand on postnatal care within maternity services, at just the time when many hospitals, especially in Greater London, found difficulties even in maintaining numbers of midwives at their former levels.

For 477 women in two former London health trusts, 86 aspects of maternity care were surveyed in 1999.² In general, the survey showed problems with postnatal hospital care. We re-examined the data to contrast the postnatal experiences of women after a caesarean (88, 18.4%) with those who had a “normal” vaginal delivery (340, 71.3%). We focused on 69 caesarean deliveries without any general anaesthetic (in case anaesthesia affected mothers’ recall), 34 in the first trust and 35 in the second trust. We used non-parametric statistics (χ^2 , Spearman’s rank correlation, Mann-Whitney U test), and our threshold for significance was $P < 0.05$.

Caesarean deliveries were associated with some unsurprising differences (such as greater length of stay or more problems with infant feeding) compared with a normal delivery. One key question about postnatal care in both trusts was, “Did you ever feel the midwives were too busy to spend enough time with you?” Patients’ answers to this question were highly correlated with overall ratings of their postnatal care while in hospital. Our concerns about the rising rates of caesareans were borne out in the first trust. The 34 women were significantly more

likely to feel that the midwives were often too busy and less likely to report that they had known any of these midwives before, compared with the normal group. But this was not the picture in the second trust. Those 35 women were no more likely to report their midwives did not have enough time for them than the normal group, and the service actually arranged that they were more likely to know one or more of these midwives well.

Only a few miles apart, both trusts had a similar range of clients. Organisational responses to more caesarean sections differed. For example, an independent consultancy reported that the first trust exemplified the special difficulties in recruiting and retaining staff in London. The maternity and neonatal workforce group is investigating the organisation of gold standard care.³ A forthcoming national service framework for children’s and maternity services offers midwives a fresh chance to develop their work in responsive and far sighted ways.

Woody Caan *public health specialist*
University College London, London WC1
woody@public-health.ucl.ac.uk

Peter Messent *public health specialist*
Queen Mary, University of London, London E1
Competing interests: None declared.

- 1 Dobson R. Caesarean section rate in England and Wales hits 21%. *BMJ* 2001;323:951. (27 October.)
- 2 Messent P. Evaluating women’s experiences for a maternity services liaison committee: going beyond the tick box. *Br J Midwifery* 2002 (in press).
- 3 Pearson S. Special delivery. *NHS Magazine* 2001;September: 8-11.

Caesarean section rates in Hong Kong

EDITOR—The report on the national sentinel survey of caesarean section highlights important public health issues that are not confined to England and Wales.¹ We have examined population based obstetric data in Hong Kong from 1993 to 1999, covering 99.5% of all live births in the period.² The information was obtained from the hospital authority that manages all 44 public hospitals, and from the department of health that regulates all 12 private hospitals in the territory. From 1993 to 1999, the overall annual rate of caesarean section rose steadily from 22% to 27.4%. Hong Kong has an even higher rate than all the developed countries reviewed in the national sentinel survey report, including the United Kingdom and the United States.

The figure compares the rates of caesarean deliveries in public and private hospitals in Hong Kong between 1993 and 1999. The absolute mean difference in rates of surgical delivery between public (mean public = 16.5%) and private (mean private = 45.1%) institutions was 28.6% (95% confidence interval 26.6 to 30.6, $P < 0.001$). According to the most recent data in 1999, one out of every two live births in private hospitals—which have accounted for one quarter to one third of all births in the territory during the seven year period—are caesarean deliveries. Not only has an overall upward trend in surgical births and high rates among private hospitals occurred throughout much of the 1990s, the absolute

Caesarean section rates in public and private hospitals in Hong Kong, 1993-9

gap between public and private hospitals has widened by 64%. It is difficult to accept that case mix and clinical indications alone contributed to this striking difference across the public-private divide.

In mixed medical economies where both public and private healthcare providers operate, we should be mindful of the potential for organisational, financing, and other non-medical factors to influence clinical practice unduly. A pattern is beginning to emerge from other parts of the world, documenting similar findings where private hospitals have much higher rates of caesarean section than their publicly funded counterparts.^{3,4} It would be interesting to compare the results of the national sentinel survey between NHS facilities and private hospitals.

The epidemic of caesarean section is a matter deserving international attention. Health services must be prepared to address issues such as defensive medicine, work patterns of private practitioners and institutions, and caesarean section on maternal demand underlying these rising trends.

Gabriel M Leung *clinical assistant professor*
gmleung@hku.hk

Tai-Hing Lam *head*
Irene Wong *senior research assistant*
Thuan Q Thach *statistician*

Department of Community Medicine, University of Hong Kong Medical Centre, 7 Sassoon Road, Pokfulam, Hong Kong Special Administrative Region, China

Competing interests: None declared.

- 1 Dobson R. Caesarean section rate in England and Wales hits 21%. *BMJ* 2001;323:951. (27 October.)
- 2 Leung GM, Lam TH, Thach TQ, Wan S, Ho LM. Rates of caesarean births in Hong Kong: 1987-1999. *Birth* 2001;28:166-72.
- 3 Roberts CL, Tray S, Peat B. Rates for obstetric intervention among private and public patients in Australia: population based descriptive study. *BMJ* 2000;321:137-41.
- 4 Belizan JM, Althabe F, Barros FC, Alexander S. Rates and implications of caesarean sections in Latin America: ecological study. *BMJ* 1999;319:1397-402.

Self management is the future

EDITOR—I assume that I am one of the few medical practitioners in this country who self manages his own oral anticoagulation and would like to comment on the paper by Fitzmaurice and Machin on patients undertaking self management of oral anticoagulation.¹

Firstly, the suggestion that patient demand for self management is partly fuelled by a national media advertising campaign for CoaguChek is disingenuous. I wonder how much of an NHS hospital trust

haematology department budget is derived from patients attending their warfarin clinic?

Secondly, routine performance in anticoagulation clinics in the United Kingdom may compare favourably with that in other countries, but at what hidden costs to the patients? As I have noted before, attending the warfarin clinic at my local hospital every two to three weeks would occupy 150 working days over 25 years, over half a year.²

Thirdly, testing is recommended every three to seven days and it is not clear why it is required. For the first week of my using CoaguChek in August 1997 I measured my international normalised ratio twice daily at about 8 am and 8 pm. Morning and evening readings ranged from 2.4 to 3.0 from a regular 8 pm dose of 5 mg of warfarin. In addition, on at least 10 occasions since then the ratio has been 3.1 or higher and on at least two occasions 2.0 or lower from the same drug regimen for more than seven days.

If I had attended a hospital warfarin clinic on the low days, my dose of warfarin would probably have been increased and not remeasured for at least a week, and with what consequences? I have already had one major epistaxis requiring a 6 unit blood transfusion, urgent ear, nose, and throat surgery, a week in hospital, and a month off work—all secondary to an international normalised ratio of 4.8 created by the warfarin clinic. I wonder what proportion of patients at warfarin clinics have had similar experiences?

Had I attended on a high day, my dose of warfarin may well have been reduced. I have not yet had any emboli from my aortic valve replacement, and I do not want any, thank you.

In the past three months I have altered my warfarin regimen of 5 mg a day on 11 occasions to keep within the range of 2.5–3.0. In other words, some people taking warfarin, and presumably not just consultant psychiatrists, may have significant individual variation in international normalised ratio.

Finally, the current debate on self management of anticoagulation is reminiscent of the dilemmas about insulin dependent diabetes 25 or so years ago, when I was a medical student and finger prick testing for blood glucose estimation became available. Look what has happened there.

D Hambidge *independent consultant psychiatrist*
9 Weavervale Park, Warrington Road, Bartington,
Nirthwich, Cheshire CW8 4QU

1 Price C. P. Point of care testing. *BMJ* 2001;322:1325-8. (26 May.)

2 Hambidge D. It suits the patient as well. Electronic response. *bmj.com*, 24 August 2001.

Management of chronic disease

System needs managing better

EDITOR—With reference to Wagner's editorial on meeting the needs of chronically ill people, many doctors should remember that patients with chronic diseases are not always socially deprived, are often articulate when their condition is not debilitating their cognitive function, and cannot help being ill.¹

Depression is more often brought on by the frustration of dealing with the medical system and attitudes than by the condition.

Could anyone imagine a well person being expected to recover from the effects of accidents or acute illnesses without necessary physiotherapy, treatments, or further follow up appointments? Yet it is expected of people who have a chronic condition, despite the knowledge that such incidents can exacerbate conditions. Is it because doctors are psychologically compartmentalising events into allowable chunks? Condition management is also pointless if patients cannot access the medical services at times when they need them most.

Patients also need to see the same doctor whenever possible, even if it means changing other patients' appointments. It is not an effective use of time and not fair on patients if they have to endure a succession of new faces and temperaments each time they meet with an emergency. Imagine how dreadful it must be for a patient of different cultural values, or from another country, to be facing the prospect of yet another face when English people find the system so daunting themselves.

Affordable treatment choices for chronic conditions are being withheld from people because doctors have habitual prescribing habits, hands-on treatments are still scoffed at, and general practitioners cannot associate and disassociate conditions appropriately for the benefit of their patients. There is also a denial of patients' individuality resulting from a strict adherence to scientifically unsubstantiated protocols. The system is the chronic condition that needs managing.

Carol Teasdale *patient*
Hereford HR1 1QN
ateasd5941@btinternet.com

1 Wagner EH. Meeting the needs of chronically ill people. *BMJ* 2001;323:945-6. (27 October.)

Proactive medicine is better option

EDITOR—The management of chronic disease throws up multiple problems for doctors and their patients as discussed by Wagner in his editorial on meeting the needs of chronically ill people.¹ Until recently doctors have been trained to respond to acute symptoms and done little to correct underlying causes. Patients have also been advised to attend when they have symptoms rather than when they are apparently well.

For example, a male patient with uncontrolled diabetes presents with balanitis or cellulitis but does not attend his routine clinic appointment with the nurse. He says that he did not need to see her as he was well. His diabetes control has to be squeezed into the time left after treating his presenting symptom.

If we as doctors are to manage chronic disease effectively we have to stop spending our time reacting to events and start anticipating and preventing them. We need to alter our frames of reference to stop overvaluing technical brilliance in treating acute events and instead start to reward

people for saving the need for such brilliance to be displayed.

This is an important change to medical value systems, and many doctors (and patients) will prefer to cling to the wreckage of the outdated model of acute reactive medicine. We need to develop systems of working that will allow us to make this change, and in general practice the first step must be to increase the length of consultations.

Peter Davies *general practitioner*
Mixenden Stones Surgery, Halifax HX2 8RQ
alisonlea@aol.com

1 Wagner EH. Meeting the needs of chronically ill people. *BMJ* 2001;323:945-6. (27 October.)

Life is a chronic disease

EDITOR—With reference to Wagner's editorial, my patients in a deprived coastal town in Cumbria are getting unhealthier.¹ We pour vast energy into chronic disease management. Groups with single issues forget that many patients in deprived towns acquire lists of chronic diseases. Add in the inevitable chronic disease of old age and for many of our patients, attending and managing their chronic disease(s) becomes synonymous with life itself.

Endless mildly abnormal blood results, 20 visits to the general practitioner a year, telephone follow up, and contradictory messages relayed from the different clinics and hospital letters lead to groaning records (albeit computerised) and, for even the best general practitioners, complete bewilderment about how to coordinate all this. The chances of the patients understanding even a small proportion of the issues are zero. The surgery becomes like a second home to many of them and the general practitioner and nurse parental figures. I watch as patients weave their chronic disease(s) into their lives and sometimes probably enjoy the interest of the whole process: it probably beats pottering in the garden and watching television.

We are increasingly taking over vast tracts of some patients' lives. Inactivity, poor diet, endless occupational stress, emotional incompetence caused by television and other media, alcohol intake, and smoking—these are the causes of most chronic disease and should be addressed at a higher level than the general practitioner's surgery. General practitioners are increasingly treating life itself as a disease. There must be more to chronic disease than endless clinics, blood tests, and contact with the health service: it is just not sustainable.

Graeme Mackenzie *general practitioner*
Maryport Health Services, Alneburgh House,
Ewanrigg Road, Maryport, Cumbria CA15 8EL
g.mackenzie@eidosnet.co.uk

1 Wagner EH. Meeting the needs of chronically ill people. *BMJ* 2001;323:945-6. (27 October.)

Rapid responses

Correspondence submitted electronically
is available on our website