Website: bmj.com
Email: letters@bmj.com

Unhappy doctors

Medical education must reassert its claim to academic high ground

EDITOR—Ham and Alberti place the current crisis in medicine in context.¹ Let us hope that the public, the profession, and the government can rise to the challenges that they laid down. They showed remarkable restraint in describing the politicisation of the NHS, but I was surprised that they did not emphasise the crucial importance of medical education and the role of our medical schools in understanding what is going wrong with medicine.

The General Medical Council has repeatedly called for radical reforms of the curriculums in undergraduate medicine, some of which have now been implemented. But whether our medical schools can instil the personal attributes and professional attitudes that are necessary for graduating students to engage with the complex sociopolitical agenda described by Ham and Alberti is less clear. The teaching and assessment of attitudes is a real challenge, but if we continue to rely on role modelling then medical education will remain hopelessly out of step with the changes taking place in society and the profession. Medical schools have a crucial role in preparing their

students for participation in the new compact described by Ham and Alberti.

In an attempt to bolster a reduced and demoralised medical workforce in the United Kingdom new medical schools are being opened and student numbers increased in the old ones. At the same time higher education funding is being squeezed as never before. Many established medical schools are reeling from the financial consequences of the 2001 research assessment exercise. Staffing cuts are inevitable, and many will probably fall on clinical academics, who give most to undergraduate education; those who work for research deans obsessed by their ratings in the research assessment exercise, however, will probably be spared.

It must be a matter of public concern that higher education funding streams to support undergraduate medical education are being diverted to prop up faltering research enterprises. Medical education now needs to reassert its claim to the academic high ground in our medical schools, on which so much depends.

Roger Jones *Wolfson professor of general practice*
Guy's, King's and St Thomas's School of Medicine,
London SE11 6SP
roger.jones@kcl.ac.uk

¹ Ham C, Alberti KGM. The medical profession, the public and the government. *BMJ* 2002;324:838-42. (6 April.)

Advice to authors

We prefer to receive all responses electronically, sent directly to our website. Processing your letter will be delayed unless it arrives in an electronic form.

We are now posting all direct submissions to our website within 24 hours of receipt and our intention is to post all other electronic submissions there as well. All responses will be eligible for publication in the paper journal.

Responses should be under 400 words and relate to articles published in the preceding month. They should include ≤5 references, in the Vancouver style, including one to the BMJ article to which they relate. We welcome illustrations.

Please supply each author's current appointment and full address, and a phone or fax number or email address for the corresponding author. We ask authors to declare any competing interest. Please send a stamped addressed envelope if you would like to know whether your letter has been accepted or rejected.

Letters will be edited and may be shortened.

bmj.com
letters@bmj.com

Alongside this traditional reactive work there has been a developing drive towards proactive care of specific diseases. Politically this is represented by national service frameworks. Historically it is based on the idea of upstream intervention in the river of disease model, which sees acute problems as needless complications of undertreated risk factors. If someone is overweight, hypertensive, and smoking why wait for the myocardial infarction before you do anything about it?

The difficulty with this approach is getting patients interested in it. What worries patients is not really what they should be worried about. Few are particularly bothered about their obesity, their excess alcohol intake, or other harmful behaviours. Even fewer of those who are bothered are bothered enough to do anything about it.

In the new compact we need to decide whether general practice is to continue on a reactive model responding to excess worry about minor illnesses or whether it is to throw its efforts into energetic treatment of the real threats to health. Until this conflict is resolved it is difficult for general practitioners to know what they should be aiming to achieve in their consultations. If we cannot appreciate our achievements we will become demoralised and unhappy.

National service frameworks or 48 hour access: whose choice is it?

Peter Davies *general practitioner*
Mixenden Stones Surgery, Halifax HX2 8RQ
npgdavies@doctors.org.uk

¹ Ham C, Alberti KGM. The medical profession, the public and the government. *BMJ* 2002;324:838-42. (6 April.)

Unhappy doctors may become sick doctors

EDITOR—There is a difference between unhappy and unwell. Brandon, a mental health professional with first hand experience of mental illness, wrote: "we are happiest when unconcerned and not knowing whether we are happy or not."¹

Edwards et al and Ham and Alberti have written of the widespread unhappiness associated with the collective loss of confidence among clinicians, the rising demands on them, and the growing conflict between clinicians.^{2,3} When organisations like the NHS prove unsupportive of dedicated staff experiencing work related stress, some will cross the invisible frontier into illness and potentially into life limiting disability.

The Mental Health Foundation's "out at work" study has shown how often people struggling with mental problems conceal

General practice must decide whether to be more proactive

EDITOR—Ham and Alberti talk of a compact between doctors, patients, and the state.¹ To set up a compact it is necessary to have some role specifications; at present these are vague in primary care.

About 30 years ago general practice was exclusively concerned with reacting to symptoms and making a diagnosis from them. Nowadays patients still tend to get worried about illnesses. Usually their symptoms are due to minor self limiting diseases, and doctors do more work treating distress about the symptoms than they do treating the illnesses themselves.

Doctors often complain that their surgeries are full of patients with "trivial" illness. The danger in filling our surgeries with them is that we miss the seriously ill patients as we are distracted by too many patients worrying needlessly. One current political demand is for 48 hour access, but I suspect that this will simply fill our surgeries with more minor illness.

their illness from their employers. Medicine is not the only caring profession that might benefit from better occupational health services: in a study of nearly 500 social workers with depression four fifths attributed their illness to work causes such as overload and exhaustion.⁴

In the distinct population of doctors, generic employment initiatives such as *Improving Working Lives* might succeed as health protection for clinicians who are now unhappy but well.⁵ Two common conditions (clinical depression and alcohol dependence) increase the risk of suicide, accidents, sickness absence, and a premature exit from the workforce. When just these conditions are considered, probably thousands of clinicians have unrecognised but treatable conditions. There is a duty of care towards our colleagues.

This year the BMA has supported our research efforts to understand those doctors who seek help for their psychological distress. Except for anecdotal subsamples from the General Medical Council and various specialist counsellors, practically nothing is known about that much larger population of doctors who are struggling to cope with a mental disorder invisible to colleagues, employers, patients, and sometimes themselves.

Two things are urgently needed to plan an acceptable, responsive, and effective service for sick doctors. These are a mental health needs assessment conducted across the profession and a health impact assessment for early detection and intervention techniques. The Department of Health's consultation in 1999 (*Supporting Doctors, Protecting Patients*) assumed the equation sick = irresponsible or dangerous. Let's just start with the assumption that sick = sick.

Woody Caan *professor of public health*
School of Health Care Practice, Anglia Polytechnic University, Chelmsford CM1 1LL
a.j.mcvicar@anglia.ac.uk

Judith Stanton *senior registrar in public health medicine*
International Centre for Health and Society, University College London, London WC1E 6BT

- 1 Brandon D. *Tao of survival, Spirituality in social care and counselling*. Birmingham: Venture Press and British Association of Social Workers, 2000.
- 2 Edwards N, Kornacki MJ, Silversin J. Unhappy doctors: what are the causes and what can be done? *BMJ* 2002;324:835-8. (6 April.)
- 3 Ham C, Alberti KGM. The medical profession, the public, and the government. *BMJ* 2002;324:838-42. (6 April.)
- 4 Manthorpe J, Stanley N, Caan W. Is depression a workplace issue? *Professional Social Work* 2002;Feb:12-3.
- 5 Abbott S. Valuing doctors. *NHS Magazine* 2002;Apr:22-3.

Regarding sick people as consumers changes their role

EDITOR—By subscribing to the concept of sick people being consumers, Edwards et al further the global political-economic project to define all aspects of human life in commercial terms—and thereby contribute to the very malaise they describe.¹ Diseases are more correctly described as the consumers in this context. The traditional compact pits doctor and patient (as more or less equal allies) against the disease and requires a degree of stoicism from both partners.

The commercial model, however, replaces the traditional model ((doctor + patient) *v* disease) with another (consumer *v* (disease + doctor)); this means that the doctor becomes accountable for the effects of disease, and the now solitary consumer (the patient) confronts a potential conspiracy. The medical discontent occurs irrespective of national boundaries because the new commercial paradigm is itself an aspect of globalisation.

Who will benefit? A working hypothesis identifies profit for transnational pharmaceutical and health insurance industries, together with states hoping to offload medical costs, if a patient's trust in a doctor is replaced by a consumer's faith in personal expenditure.

David H Bevan *senior lecturer*
St George's Hospital, London SW17 0QT
dbevan@sghms.ac.uk

- 1 Edwards N, Kornacki MJ, Silversin J. Unhappy doctors: what are the causes and what can be done? *BMJ* 2002;324:835-8. (6 April.)

At least some of the unhappiness is due to excessive accountability

EDITOR—Edwards et al ask why doctors are unhappy and what can be done.¹ Perhaps excessive accountability is a reason.

As a professor of surgery and honorary consultant surgeon I am attempting to compile a comprehensive list of the various bodies that are assessing/appraising/validating my performance and activity. I would be grateful if readers would peruse the list below and inform me of any glaring omissions; after all, I would hate to think I have forgotten some.

- *University*
Internal quality assurance committee
Staff review and development committee
Annual university appraisal
Quality Assurance Agency
Research assessment exercise
Peer review of teaching
Research governance committee
- *Hospital*
Preregistration house officer review for postgraduate dean
Senior house officer review for postgraduate dean
Specialist registrar review for postgraduate dean
Royal College of Surgeons hospital visit for basic surgical training
Royal College of Surgeons hospital visit for specialist registrar
Clinical governance committee
Continuing professional development committee
Professional advisory panel
Clinical audit committee
Annual appraisal
- *National*
General Medical Council revalidation procedures
UK Council for Regulation of Healthcare Professionals
National Clinical Assessment Authority
National Care Standards Commission

Commission for Health Improvement
National Patient Safety Agency

Irving Taylor *professor of surgery*
Department of Surgery, University College London, London W1W 7EJ
irving.taylor@ucl.ac.uk

- 1 Edwards N, Kornacki MJ, Silversin J. Unhappy doctors: what are the causes and what can be done? *BMJ* 2002;324:835-8. (6 April.)

Nurses need more resources to do their job

EDITOR—I was a nurse for some years. My response to Buchan's editorial on global nursing shortages is to say that factors such as flexible working hours and continuous professional development may help, but there are more fundamental problems.¹

I gave up nursing in the mid-1980s. I would have gone back like a shot in the following few years if I could have worked somewhere in the NHS that had adequate resources to do the job properly. I have checked with another colleague who left the wards more recently, and she shares this view.

During my time as a student in the NHS and then later as a staff nurse in oncology I had to leave alone patients who were dying, alert, and terrified. And be late giving essential analgesia. I never had the time to offer any proper emotional support, or the care that some very sick or scared patients needed. A large chunk of my time was wasted on endlessly searching for scarce basic resources such as pillows.

If anyone thinks the resourcing problems are new, they're wrong. I got fed up with statements from the management, at least once a year, along the lines of, "There's a funding crisis; we know we've told you that before, but this time it's much worse." After 10 years of that you lose hope.

I note that the references on nurse retention in the editorial are from American and Canadian studies. I hope that there are, or soon will be, some British NHS studies too. And that their findings are acted on—if they aren't it's just another waste of money.

I hope that things have improved since my friend and I left but suspect that they haven't.

Fiona McLean *health care information officer*
British Library, London NW1 2DB
fiona.mclean@bl.uk

- 1 Buchan J. Global nursing shortages. *BMJ* 2002;324:751-2. (30 March.)

BMA negotiator calls for more male medical students

Working women seem to be scapegoats in medicine

EDITOR—It seems that sexism is operating in the BMA. Holden says that there is a sound case for biasing medical selection in favour of male applicants because "you do not get 35-40 years of service from the females."¹

It is a biological fact that women have babies and therefore take some time off work. Holden would solve the problem of understaffing in medicine by preventing women entering the medical profession in such large numbers. Let us follow this line of reasoning to its logical conclusion.

It is a biological fact that diabetes and heart disease are more prevalent in people with a south Asian ethnic background.² Many Asian doctors die of cardiovascular disease while in service, before reaching their retirement age. Is Holden going to suggest that medical schools should bias their entry away from south Asian applicants because they may not fulfil 40 years' service?

But why stop here? Others who may take career breaks are those who are interested in medical politics. Negotiating on behalf of one's colleagues may be a noble calling, but it's hardly delivering 40 years of service.

Increasing patient expectations, new technologies, and the spectre of legal action have increased stress at work. It is not surprising that many doctors, male and female, are taking time out or retiring early. The solution is to change the working conditions of the profession, not to place irrational restrictions on those who wish to enter it.

Society has changed. Once men were willing to be present at their children's conception and graduation but rarely in between. The increasing number of women in medicine has brought about changes that have benefited patients and colleagues alike. It has helped to create a more humane and representative profession and has created pressure for a working ethos that is more likely to preserve the mental and physical health of its practitioners.

Our negotiators should be pressing for improved child care, improved access to occupational health, some protection from malicious legal claims, a slowing of the pace of administrative change, and, above all, sensible resourcing for a 21st century NHS. Instead, working women have been chosen to be the scapegoat.

Susan Walker *locum general practitioner*
Brooklands Avenue Medical Centre, 7 Brooklands
Avenue, Cambridge CB2 2BB
walksh30@genie.co.uk

1 Moore W. BMA negotiator calls for more male medical students. *BMJ* 2002;324:754. (30 March.)

2 Bhopal R. Epidemic of cardiovascular disease in South Asians. *BMJ* 2002;324:625-6. (16 March.)

Plus ça change...

EDITOR—Here we go again. The difficulties in keeping qualified women at work continue to be a problem, but the solutions do not seem to be getting any more imaginative.¹ A few important questions need to be asked before women are excluded from medicine. Do they make good doctors? Do they make better doctors as a group than men do? Do patients, or particular subgroups of patients, prefer female to male doctors? Is there a human rights issue involved in denying someone access to education when the place is given to another candidate with weaker credentials?

If the answers to these questions suggest that there is an advantage in keeping women in medicine the important question is how to achieve that. Living and working in Malaysia, I do not know a single woman doctor (of whom there are many) who has stopped working in order to care for her family. Many prefer to stay away from demanding hospital specialties and opt for general practice, without on call work. They can do this because of a combination of family support and access to cheap child care. Women will stay in work if the burden of child care does not fall entirely on their shoulders, as it seems to do in the United Kingdom. Energy should focus less on keeping women out of medicine and more on how to keep them in it.

Surely if women candidates for medical school are stronger than male candidates it would be more logical to call for a national initiative to keep women at work and men at home. Failing that option, maybe there could be different forms of part time work for fathers and mothers, and increased access to child care in preschool state run nurseries could be provided.

Madeleine Kathleen Piper *general practitioner in solo practice*
Klinik Dr Madeleine, Damai Point, Kota Kinabalu,
88 764 Kota Kinabalu, Malaysia
maddykp@tm.net.my

1 Moore W. BMA negotiator calls for more male medical students. *BMJ* 2002;324:754. (30 March.)

Access to invasive procedures can be audited by ethnic group

EDITOR—Feder et al reported findings from the appropriateness of coronary revascularisation (ACRE) study showing that among patients deemed clinically appropriate for coronary artery bypass grafting, south Asians are less likely than white patients to undergo procedures.¹ They pointed out the lack of a mechanism to monitor ethnic differences in invasive management of coronary disease in the NHS. The quality of hospital episode statistics has improved significantly in many trusts since the ACRE study. In 1999-2000 the quality index for our local trust data on ethnic group was 92.8% compared with a national average of 65.4%.² This contrasts with a rate of invalid coding in 1995-6 of 62.9%.³

We conducted an audit of all patients admitted with unstable angina or acute myocardial infarction in a local trust between July 1999 and June 2000. Information was obtained on age, sex, ethnic group, and Carstairs socioeconomic deprivation category derived from postcode of residence. Record linkage to subsequent hospital episode statistics provided follow up information on procedures up to 18 months after admission. Data on 53 south Asians and 1556 white patients were analysed using logistic regression models.

We found higher access to angiography and coronary artery bypass grafting among south Asian compared with white patients in

the univariate analysis. We found no difference in access to angiography (odds ratio = 0.94, 95% confidence interval 0.49 to 1.83) or coronary artery bypass grafting (1.27, 0.62 to 2.62) between south Asian and white patients after controlling for age, sex, and Carstairs deprivation category.

Despite the small sample size, lack of control for severity, or appropriateness we have shown that access can be audited by ethnic group using routine data. Many trusts may currently have the ability to carry out such audit, especially in areas with high ethnic minority populations. It may not be long before national audit of ethnic data on access to procedures is possible.

Ibrahim I Abubakar *specialist registrar in public health*

David Kanka *consultant in public health medicine*
Public Health Laboratory Service Communicable
Disease Surveillance Centre Eastern, Institute of
Public Health, Cambridge CB2 2SR
iad21@medschl.cam.ac.uk

1 Feder G, Crook AM, Magee P, Banerjee S, Timmis A, Hemingway H. Ethnic differences in invasive management of coronary disease: prospective cohort study of patients undergoing angiography. *BMJ* 2002;324:511-6. (2 March.)

2 Department of Health. *HES data quality report—1999/00*. www.doh.gov.uk/hes/data_quality/dq99/rgn99.html (accessed 5 Mar 2002).

3 Aspinall PJ. The mandatory collection of data on ethnic group of inpatients: experience of NHS trusts in England in the first reporting years. *Public Health* 2000;114:254-9.

Hip protectors

Carpets can be used to reduce injury from falls

EDITOR—It takes about 40 years of wearing hip protectors to prevent one fracture, according to the editorial by Cameron,¹ so other strategies may be helpful. Hip protectors look like institutional underwear—white, plain, and unfashionable. Making them more attractive might improve their acceptability. The alternative approach is to pad the floor. Carpet improves gait in older people² and in a small observational study dramatically reduced serious injury from falls.³ Perhaps it's time for the Axminster trial of hip protection.

Jed Rowe *consultant geriatrician*
Well Balanced Clinic, Moseley Hall Hospital,
Birmingham B13 8JL
jedrowe@geriatrictry.freemove.co.uk

1 Cameron ID. Hip protectors. *BMJ* 2002;324:375-6. (16 February.)

2 Wilmott M. The effect of a vinyl floor surface and a carpeted floor surface upon walking in elderly hospital in-patients. *Age Ageing* 1987;16:119-20.

3 Healey F. Does flooring type affect risk of injury in older in-patients? *Nursing Times* 1994;90:40-1.

Integrated approach to care of older people is necessary

EDITOR—We wish to raise the contentious issue of funding in the context of the management of falls in care homes. Critical success factors for the use of hip protectors are accurate fitting and compliance, as mentioned in the editorial by Cameron¹; these can be difficult in care homes now that the prevalence of dementia is increasing. Furthermore, wearing hip protectors may be a

confounding factor in continence and an impediment to the promotion of continence. A more environmental approach in care homes might be as effective and less intrusive.

The mechanical assault on a hip from a fall will depend on the properties of the flooring. Floor coverings in institutional settings are inevitably a compromise between homeliness, clinical appropriateness (largely the maintenance of cleanliness), and cost. Floor coverings are available with cushioning, as has been used for reducing head injury in children's playgrounds. Environmental design and technology have much unexploited potential for unobtrusive risk management.

People will certainly need three sets of protectors, with an estimated initial outlay of £120 per person, and several replacements will be required over time; that, together with the added care to ensure appropriate placement and compliance, greatly adds to the cost. Although this may be justified, the question of who pays remains unclear. Presently, we are occasionally obliged to provide hip protectors from social care money; investment by health services is sporadic.

In the United Kingdom the justification for and funding of hip protectors and environmental protection cuts across health technology assessment and the work of the National Institute for Clinical Excellence. Perhaps eventually funding will be withdrawn from orthopaedic trauma and rehabilitation programmes to facilitate this preventive care.

Hip protectors show that an integrated approach to the health and care of older people is necessary to yield outcome improvement.

Clive Bowman *medical director*

Ian Ireland *head of quality*

BUPA Care Services, Horsforth, Leeds LS18 4UP
bowmanc@bupa.com

1 Cameron ID. Hip protectors *BMJ* 2002;324:375-6. (16 February.)

Acceptability of hip protectors was 35% at six months in the community

EDITOR—Cameron identifies the low use of hip protectors as one of the main drawbacks to their more general use.¹ Initial compliance rates of 50-60% have been reported, but, as Cameron points out, little research has been done into the acceptability of the protectors among patients living in the community.^{2,3} We carried out a small study to assess acceptability in such a group.

Forty nine patients with osteoporosis living in inner city Belfast were provided with hip protectors (Safehip). After six months the project nurse completed a questionnaire to assess the rate of use and the factors influencing this. The acceptance rate was initially 53% (26/49) and fell to 35% (17/49) at six months. The patients rated the protectors poorly on bulkiness, heat, and the tightness of the waistband; these were also the factors that most strongly influenced their decisions on wearing them. They also disliked the plastic cover and the noise made

on movement when the protectors were in place. Overall appearance, appearance under tight clothes, and looking after the protectors were rated more favourably.

Many patients said that it was difficult to get the hip protectors on and off (19 rated it generally difficult or very difficult, and 20 commented that it was difficult or very difficult to get them on and off to go to the toilet). Similar factors have been noted in other studies.²⁻⁴ Few of the patients were prepared to wear the hip protectors at night.

Those who wore the hip protectors were almost three times more likely than those who did not to believe that they offered protection from hip fractures (86% (25/29) *v* 30% (6/20)). Two thirds of patients wearing the protectors felt more confident about going out. Most patients (46) stated that they would be likely or very likely to recommend the protectors to a friend.

Some problems were identified in patients who forgot the hip protectors in a nursing home, so forgetting them is likely to be more of a problem with patients living at home. Compliance may be improved in this group if primary care workers in contact with the patients remind them that hip protectors should be used every day.

Jeni M McAughey *general practitioner*

Skegoneill Health Centre, 195 Skegoneill Avenue, Belfast BT15 3LL
jmmcaughey@doctors.org

Margaret McAdoo *project nurse, North and West*

Belfast Locality Consortium
Springfield Road Surgery, Springfield Road, Belfast BT12 7AH

1 Cameron ID. Hip protectors. *BMJ* 2002;324:375-6. (16 February.)

2 Hindso K. Prevention of hip fractures using external hip protectors. Risk factors for falls, hip fractures, and mortality; and evaluation of the consequences of fear of falling among older orthopaedic patients. Copenhagen: University of Copenhagen, 1998. (PhD thesis.)

3 Villar TMA, Hill P, Inskip H, Thompson P, Cooper C. Will elderly rest home residents wear hip protectors? *Age Ageing* 1998;27:195-8.

4 Cameron I, Quine S. External hip protectors: likely non-compliance among high risk elderly people living in the community. *Archives of Gerontology and Geriatrics* 1994;19:273-81.

Rate of adherence is 42% at three months in residential homes

EDITOR—With reference to Cameron's editorial, we have previously shown that patients' adherence with hip protectors is achievable at a level that is likely to result in reduction of hip fractures but were unable to assess usage in subjects with cognitive dysfunction, confusion, or incontinence because of ethical issues in a controlled trial.^{1,2} The establishment of a hip protector service in Dorset has allowed us to study these factors in an observational study.

Residential homes in Poole were approached, offering their residents hip protectors. Assessments were made of falls risk (modified STRATIFY score), cognitive function, confusion, and urinary continence. Residents were offered hip protectors free of charge. Assessment of adherence was made after 1 week and 3, 6, and 12 months. Forty seven residential homes with 783 beds are taking part. Over 880 residents have been

identified, mean age 88 (SD 7) years; the female:male ratio is 4.6:1. Of the 873 residents who were offered protectors, 645 expressed an interest and 612 were wearing them at one week. Of the 753 residents so far assessed at three months, 340 were wearing protectors. At three months 180 out of 292 incontinent residents compared with 158 out of 369 (43%) residents who were continent were wearing protectors, as well as 170 out of 291 (58%) residents with dementia compared with 169 out of 474 (35%) without dementia, and 89 out of 141 (63%) residents who were always confused compared with 150 out of 292 (51%) of residents who were sometimes confused and 95 out of 227 (42%) who were never confused ($P < 0.001$, Mann-Whitney test). There was correlation between the risk of falling and adherence (STRATIFY score 0 = 33%, 1 = 45%, 2 = 52%, 3 = 49%, 4 = 73%, and 5 = 82%).

Extrapolating the results to all residents suggests 42% adherence at three months, a rate greater than in our previous study and likely to result in a reduced incidence of hip fracture. The higher adherence among cognitively impaired subjects, confused subjects, and subjects at high risk of falling supports our concept that hip protectors are worn by subjects at greatest risk of fracture.³

Paul W Thompson *consultant rheumatologist*

Poole Hospital, Poole, Dorset BH15 2JB
pwt@dial.pipex.com

Carol Jones *osteoporosis prevention officer*

Osteoporosis Dorset, Bournemouth, Dorset BH1 4JQ

Tracy Villar *consultant physician, care of the elderly*

Poole Hospital NHS Trust, Poole, Dorset BH15 2JB

This work is administered by Osteoporosis Dorset (Registered Charity No 1023507) and funded by Dorset Health Authority and Poole Social Services).

1 Cameron ID. *BMJ* 2002;324:375-6. (16 February.)

2 Villars TMA, Hill P, Inskip H, Thompson PW, Cooper C. Will elderly rest home residents wear hip protectors? *Age Ageing* 1998;27:195-8.

3 Thompson PW, Jones C, Dawson JA, Davies F. Adherence with hip protectors in elderly persons requiring domiciliary care is greater in fallers than non-fallers. *Age Ageing* 2000;29:459.

Cardiothoracic surgeons do a good job

EDITOR—Partridge provides no evidence to support his claims that cardiothoracic surgeons are less likely than thoracic surgeons to stage the extent of lung cancer objectively and assess the patients' fitness; he points out that they have "competing demands from a waiting list for coronary artery surgery."¹ He also says that relying on cardiothoracic surgeons to do lung resections may mean that some do too few to maintain competence.

Partridge states that five year survival figures are better in those who operate more often, referring to a study showing an association between improved survival and more resections done in the hospital, with numbers performed by individual surgeons not analysed.² In three quarters of the hospi-

tals included in the study less than 22 lung cancer resections were performed a year, with the best results being obtained in units where the annual total was over 66 (only 3% of the hospitals). This is unlikely to reflect practice on this side of the Atlantic, where almost all lung cancer resections are performed in specialised thoracic or cardiothoracic units, where the mean number of resections a year is 72 (www.scts.org/file/thoracicregister1999-00.xls).

All cardiothoracic surgeons appointed in the United Kingdom undergo training in thoracic surgery and sit the same specialist examination as thoracic surgeons. All are audited in the same national database of the Society of Cardiothoracic Surgeons of Great Britain and Ireland, the marker operation for surgeon specific mortality being lobectomy for primary lung cancer. This has shown that the results of cardiothoracic surgeons are as good as those of thoracic surgeons, with a mortality of 2.5% and no difference related to the volume of operations done (T Treasure, personal communication). Thoracic surgery is well ahead of other surgical specialties in national audit and assessment of surgeon specific mortality, solely because of its association with cardiac surgery and their joint society.

We do need more surgeons to manage benign and malignant thoracic diseases, but it seems absurd to discourage the many surgeons who are already doing a good job merely because they also operate on the heart.

Alastair NJ Graham *consultant cardiothoracic surgeon*
Royal Victoria Hospital, Belfast BT12 6BA
alastair.graham@royalhospitals.n-i.nhs.uk

1 Partridge MR. Thoracic surgery in a crisis. *BMJ* 2002;324:376-7. (16 February.)

2 Bach PB, Cramer LD, Schrag D, Downey RJ, Gelfand SE, Begg CB. The influence of hospital volume on survival after resection for lung cancer. *N Engl J Med* 2001; 345:181-8.

Informaticist services for policymakers will differ from those studied

EDITOR—We were pleased to see Greenhalgh et al's formal evaluation of two quite different informaticist services.¹ Our own experience in facilitating evidence supported policymaking suggests that some of the points raised apply to policy as well as to practice, though additional points must be considered. Our developing approach has some commonality with both projects but is aimed at policymaking bodies comprising health professionals and lay representatives rather than groups of practitioners.

We decided that a "laboratory test service" approach would not be appropriate in policymaking, although as an academic unit we found it intuitively attractive, at least initially. We were obliged to recognise that health policy issues rarely resolve into a question that can be answered in three stages. Often, several questions need to be

tackled simultaneously as policymakers need to consider local processes, structures, times scales, and concerns.^{2,3}

The outcome is a set of options rather than an answer. Over time we have increasingly favoured an approach closer to the Basildon model. We aim to arrive at best accepted wisdom by accessing authoritative sources of pre-appraised guidance,⁴ particularly those whose methodological approach to evidence appraisal is both explicit and systematic.

We have worked closely with local policy-making bodies and have designed dissemination tailored to each issue. The close relation between our academic unit and service departments of the NHS locally has made this possible. Recently we have addressed diverse questions, such as the health risks from household waste incinerators, the cost effectiveness of photodynamic treatment for macular degeneration, and the organisation of antenatal screening programmes.

Our experience suggests that facilitation of evidence based policymaking should be underpinned by an acknowledgement of the complexity of policy. We have to accept that we may sometimes need to compromise the technical quality of the answers: if a decision is needed quickly it may be appropriate to base it on easily accessible authoritative guidance rather than on a comprehensive review of all available evidence including appraisal of primary research. We believe that the adoption of a pragmatic approach can achieve a more systematic use of evidence than is usual in the NHS.⁵

The question of whether our approach, which focuses on the service dimension, produces good policy decisions needs evaluating. We believe that the two dimensions of a health informatics service, as described by Greenhalgh et al, will prove to be an important component of the methodology. We also agree that such an evaluation requires a developmental rather than a conventional approach to research.

Helen Thornton-Jones *senior lecturer in health services research*
H.Thornton-Jones@hull.ac.uk

Susan Hampshaw *research associate*
Department of Public Health and Primary Care, University of Hull, Willerby HU10 6NS

Andrew Taylor *health economist*
Effectiveness Facilitation Unit, East Riding and Hull Health Authority, Willerby HU10 6DT

1 Greenhalgh T, Hughes J, Humphrey C, Roger S, Swinglehurst D, Martin P. A comparative case study of two models of a clinical informaticist service. *BMJ* 2002;324:524-9. (2 March.)

2 Dopson S, Locock L, Chambers D, Gabbay J. Implementation of evidence-based medicine: evaluation of the promoting action on clinical effectiveness programme. *J Health Serv Res Policy* 2001;6:23-31.

3 Lomas J. Connecting research and policy. *Canadian Journal of Policy Research* 2000;1:140-4.

4 Thornton-Jones H, Hampshaw S, Soltani H, Madhok R. Reviewing local screening programmes—a worthwhile exercise. *Journal of Clinical Governance* (in press).

5 Milbank Memorial Fund. *Better information, better outcomes. The use of health technology assessment and cost effectiveness data in health care purchasing decisions in the United Kingdom and the United States, July 2000*. New York: MMF, 2000. (www.milbank.org/000726purchasing.html)

Attitudes towards general practice are changing

EDITOR—Bowler and Jackson say in their paper on the career intentions of general practice registrars that the government's national plan may underestimate the numbers of new doctors needed.¹ They found that many general practice registrars intend to work part time or as non-principals.

Our own studies of all medical graduates in particular graduation years show important trends in recent years, away from full time principalships. We reported a large fall in the percentage of young medical graduates who want a career in general practice; a large increase in the percentage of general practitioners who work part time, mainly because many more women than men now enter general practice; and increasing reluctance on the part of general practitioners to enter into traditional principalships soon after completion of training.²

As well as quantifying change we need to understand why it is happening. Recently we reported some of the reasons for these changes as given in comments from respondents.³ There seems little doubt that general practice is now regarded by some as a less prestigious, and less interesting, career than specialist hospital medicine. Continual administrative changes, increases in workload, and an increasing "complaints culture" have reduced morale and job satisfaction for some established general practitioners. These negative elements of general practice are picked up as disincentives by some younger doctors who might consider general practice as a career. But it is also considered easier to work in general practice on a part time or flexible basis and some doctors, particularly women, currently choose it for that reason. Given the increasing percentage of women in the medical workforce, part time working will increase. The result is a departure of full time general practitioners, some at retirement age and some earlier, to be replaced by a high percentage of part timers.

We see the way forward as twofold. Firstly, to increase recruitment, a cultural change is needed among medical educators to promote general practice as a career choice that is as attractive as hospital practice. Secondly, a greater variety of contractual arrangements is needed for those already in general practice, to reflect doctors' desires for more flexible working patterns. It is clear that the government is aware of the impact of increasing female entry to medical school and its likely consequences for part time working and of the need for flexible working practices.^{4,5} Whether recognition translates into sufficient change remains to be seen.

Trevor Lambert *study coordinator*
trevor.lambert@dphpc.ox.ac.uk

Julie Evans *social scientist*
julie.evans@dphpc.ox.ac.uk

Michael Goldacre *director*
michael.goldacre@dphpc.ox.ac.uk

UK Medical Careers Research Group, Institute of Health Sciences, Oxford OX3 7LF

- 1 Bowler I, Jackson N. Experiences and career intentions of general practice registrars in Thames deaneries: postal survey *BMJ* 2002;324:464-5. (23 February.)
- 2 Lambert TW, Evans J, Goldacre MJ. Recruitment of UK-trained doctors into general practice: findings from national cohort studies. *Br J Gen Pract* 2002 (in press).
- 3 Evans J, Lambert TW, Goldacre MJ. GP recruitment and retention: a qualitative analysis of doctors' comments about training for and working in general practice. *Royal Coll Gen Pract* 2002 (occasional paper 83).
- 4 Department of Health. *Planning the medical workforce: third report, medical workforce standing advisory committee*. London: Stationery Office, 1997.
- 5 Department of Health. *Improving working lives for doctors*. London: Stationery Office, 2001.

Animal models can assist medicine in humans

EDITOR—The debate about whether we can learn from animals is far more subtle than the rapid responses to Roberts et al's systematic review of fluid resuscitation after haemorrhage suggest.^{1,2} It depends on the problem and on the species of the animal; the lessons of comparative medicine are most useful when the model has sufficient similarities to suggest its relevance and enough differences to make it informative. Blanket assertions that we cannot learn from animals are naive, but many generalisations about humans fail when we look at ethnic differences and genetic predisposition.

Several odd points arise in Roberts et al's systematic review.

Firstly, the authors confuse treatment aimed at increasing circulating volume (for example, with solutions based on isotonic sodium) with treatment with hypertonic sodium solutions, whose benefit in shock probably rests on pharmacological properties other than their minuscule and transient impact on circulating volume.

Secondly, the review contains no trials from species routinely treated for haemorrhage by veterinary surgeons. There is a substantial literature on the use of hypertonic saline to treat haemorrhage in animals, notably cats, dogs, and horses. Treatment of haemorrhage is rare in sheep and exceedingly so in pigs.

The review makes the usual assumption that research in animals is synonymous with studies in laboratory animals, mainly rats. The transected tail model in rats is particularly misleading as the transection of the major artery is likely to predispose to rebleeding once arterial pressure starts to rise. This may be relevant to some forms of clinical haemorrhage, but I doubt if it is a good model for most. I also doubt that studies with controls that have been denied any fluid resuscitation after haemorrhage are scientifically or ethically justifiable nowadays.

Comparative medicine reminds us that humans are not unique; they are animals of a particularly interesting type. Only when we realise this can we start to think sensibly about questions such as how much salt mammals need and how much is detrimental, and why some animals are susceptible to extremely high blood pressure yet resistant to its damaging effects. We can also capitalise on the fact that spontaneous tumours in

animals, equivalent to those in humans, may be a useful intermediate step—between tumours induced in experimental rodents and tumours in human patients—in evaluating new treatments. Awareness of such opportunities is too low in the medical community.

Alastair R Michell *professor of comparative medicine*
University of London, St Bartholomew's Hospital
Medical School, Department of Pharmacology,
London EC1M 6BQ
bobmichell@hotmail.com

1 Electronic responses. Does animal experimentation inform human healthcare? *bmj.com* 2002. (bmj.com/cgi/eleletters/324/7335/474; accessed 26 April 2002.)

2 Roberts I, Kwan I, Evans P, Haig S. Does animal experimentation inform human health care? Observations from a systematic review of international animal experiments on fluid resuscitation. *BMJ* 2002;324:474-6. (23 February.)

Staphylococci are unlikely to cause acute dental infections

EDITOR—O'Rourke et al report on a patient who, after root canal therapy, developed an epidural abscess from which *Staphylococcus aureus* was grown.¹ They contend that there is a causal association, but we believe that this is unlikely. Bacteraemia often occurs after dental procedures, and metastatic infection can follow. Infective endocarditis is a good example of the phenomenon.

S aureus is not commonly isolated from dental root canals, periapical infections, or periodontal pockets.²⁻⁴ In the reported case there is no microbiological evidence to confirm oral *S aureus* infection.

This is important, and not just for semantic reasons. Reported as it is, the case serves to reinforce the misheld belief that dental infections are caused by staphylococci and can therefore be treated with flucloxacillin. Acute dental infections are best treated with other agents such as phenoxymethylpenicillin (penicillin V), amoxicillin, erythromycin, clindamycin, or metronidazole.⁵ Life threatening orofacial infection may follow inappropriate antibiotic treatment of odontogenic infection, especially if the infected teeth are not extracted and pus drained.

Noel F F Ribeiro *locum registrar in maxillofacial surgery*
Maxillofacial Unit, North Manchester General
Hospital, Crumpsall, Manchester M8 5RB
noel_ribeiro@doctors.org.uk

Gary C S Cousin *consultant maxillofacial surgeon*
Blackburn Royal Infirmary, Blackburn BB2 3LR

1 O'Rourke F, Barker R, Khan S. Minerva. *BMJ* 2002;324:686. (16 March.)

2 Brauner AW, Conrads G. Studies into the microbial spectrum of apical periodontitis. *Int Endod J* 1995;28:244-8.

3 Williams BL, McCann GF, Schoenkecht FD. Bacteriology of dental abscesses of endodontic origin. *J Clin Microbiol* 1983;18:770-4.

4 Haake SK. Periodontal microbiology. In: Carranza FA, Newman MG, eds. *Clinical periodontology*. 8th ed. Philadelphia: WB Saunders, 1996:84-103.

5 Lewis MA, MacFarlane TW, McGowan DA. Antibiotic susceptibilities of bacteria isolated from acute dentoalveolar abscesses. *J Antimicrob Chemother* 1989;23:69-77.

Expert shows how pens should be held

EDITOR—Having married a doctor many years ago, I am aware that doctors are generally without fault. But Minerva's request for instruction on how to pick up a pen reminds me that many medical practitioners could profit from advice on how to hold one once they have managed to pick it up.¹

Correct

Writer's cramp

Diagrams showing correct way to hold a pen and way that may cause writer's cramp

The figure shows two possible writing positions for a pen, and experimentation will show which affords a relaxed hold and which is likely to give writer's cramp. Any change of hold is likely to take a month's practice before it feels natural.

Unfortunately, this advice will not affect the legibility of the handwriting, on which subject I would not presume to advise doctors.

Richard Middleton *chairman, Society of Scribes and Illuminators*
37 Grove Lane, Barrow upon Soar, Leicestershire
LE12 8NP
r.middleton@freuk.com

1 Minerva. *BMJ* 2002;324:798. (30 March.)

bmj.com

Rapid responses

Correspondence submitted electronically is available on our website