Website: bmj.com
Email: letters@bmj.com

Suturing *v* conservative management of hand lacerations

To suture or not to suture?

All lacerations need to be examined thoroughly

EDITOR—Quinn et al show that conservative treatment is faster and less painful for small uncomplicated lacerations of the hand.¹ However, we think that lacerations to the hand, no matter how small, must be examined thoroughly to exclude injuries to tendons, nerves, or joints. The authors make no comment on the mechanism of injury, which is extremely important. A knife stab laceration or glass injury to the hand would make exploration of the wound mandatory. An unimpressive skin wound may hide a remarkable amount of damage to deep structures.² Similarly, injuries caused by thin slivers of glass produce unimpressive skin wounds but commonly divide flexor tendons and nerves in the forearm.³

In emergency settings we think that it is crucial to take a good history from the patient about the mechanism of injury and to examine the patient thoroughly before deciding on further management of hand lacerations, albeit suturing or conservative management. In our plastic surgery unit the nurse practitioners who refer cases of hand trauma to us have all been on a hand trauma study day organised by our department. If the mechanism of injury raises any suspicion of a tendon or nerve injury, patients are referred to us and their wounds formally explored in an operating theatre.

Beryl A De Souza plastic surgery registrar
bds@dr.com

Mohammed Shibu consultant plastic surgeon
Graham Moir consultant plastic surgeon
Nigel Carver consultant plastic surgeon
Department of Plastic Surgery, Barts and the Royal London Trust, Royal London Hospital, London E1 1BB

- 1 Quinn J, Cummings S, Callahan M, Sellers K. Suturing versus conservative management of lacerations of the hand: randomised controlled trial. *BMJ* 2002;325:299-300. (10 August.)
- 2 Schwager RG, Smith JW, Goulian D. Small deep forearm lacerations. *Plast Reconstr Surg* 1975;55:190-4.
- 3 Joseph KN, Kalus AM, Sutherland AB. Glass injuries of the hand in children. *Hand* 1981;13:113-9.

Hand lacerations should be explored before conservative treatment

EDITOR—We are surprised by the publication of the article by Quinn et al on conservative treatment of small hand lacerations.¹ Their conclusions will come as no surprise to surgeons who treat hand injuries often, who recognise that wounds on the palmar aspect of the hand heal well if left open.² Skin defects in the palm are often left to heal by secondary intention after surgery for Dupuytren's disease, with excellent results.³

Our concern is that this paper trivialises small hand lacerations, ignoring the fact that lacerations such as that shown in the front cover photograph may injure any of the underlying soft tissue and bony structures in the finger. In their method, they do not state who made the judgment that there was no associated neurovascular, tendon, or bone injury. This cannot be excluded in this type of wound unless a careful history of the mechanism of injury is taken, a radiograph is obtained, and the wound is explored under local anaesthesia by someone experienced enough to make this judgment.

If neglected, injuries to these structures will usually result in permanent functional disability. Wounds that have penetrated and contaminated the flexor tendon sheath can lead to devastating infection with massive soft tissue loss and all of its sequelae. Similarly, those which have penetrated the joint capsule may lead to septic arthritis.

Small hand lacerations, along with many other conditions in emergency departments, are seen and treated by (through no fault of their own) junior and inexperienced doctors and, increasingly, nurse practitioners. This paper sends out a message that is likely to result in more patients having treatable hand injuries neglected, with regrettable and entirely avoidable consequences. Encouraging such wounds to be treated conservatively is unlikely to benefit the patients or the medical staff treating them, but should keep the lawyers busy.

Roderick Dunn specialist registrar
roderick.dunn@virgin.net

Stuart Watson consultant
West of Scotland Regional Plastic Surgery and Burns Unit, Canniesburn Hospital, Glasgow G61 1QL

- 1 Quinn J, Cummings S, Callahan M, Sellers K. Suturing versus conservative management of lacerations of the hand: randomised controlled trial. *BMJ* 2002;325:299-300. (10 August.)

- 2 Brown PW. Open injuries of the hand. In: Green DP, Hotchkiss RN, Pederson WC, eds. *Green's operative hand surgery*. Philadelphia: Churchill Livingstone, 1999:1612-4.
- 3 McCash CR. The open palm technique in Dupuytren's contracture. *Br J Plast Surg* 1964;17:271-80.

Incisions are not lacerations

EDITOR—Quinn et al in their paper on the management of lacerations are guilty of a failure to use the correct nomenclature in describing wounds accurately.¹ For, although they refer to lacerations of the hand, it is clear from the photographs used to illustrate the paper, both on the front and inside the journal, that these are incised wounds. Indeed, one of them has all the appearances of being a penetrating incised wound or stab wound from a single edged weapon.

Without going into detailed definitions in respect of the differences between lacerations and incisions, in my experience, full thickness wounds to the hands are more commonly incised wounds, which are the result of contact with a sharp bladed object or implement. They are much less likely to be lacerations, which are the result of splitting or tearing of the skin as a result of some form of blunt force trauma. Unfortunately this paper seems to be a further example of doctors, and not always junior doctors, using the incorrect nomenclature in describing wounds accurately.

This is an issue that has been raised in the past by Milroy and Rutty and Norfolk and Stark.^{2,3} Both sets of authors made it clear that the accurate description of wounds and the use of correct nomenclature are of considerable importance, particularly in assessing the causation of wounds, which is clearly of considerable relevance in medicolegal issues.

In many respects this is a sad indictment of present day undergraduate medical training, which is devoid of any input in forensic medicine. This use of incorrect nomenclature is on the increase and causes considerable problems and arguments in court.

Advice to authors

We would like to receive all responses electronically at our website. Please send your letter directly to bmj.com as a rapid response to a published article.

All rapid responses will be considered for publication in the paper journal; authors will be notified by email if their rapid response has been accepted, but not otherwise.

For more detailed advice please see bmj.com/advice/sections.shtml#letters

bmj.com

Without the reintroduction of formal training in forensic medicine for medical students these are problems that are going to continue increasing.

Alistair J Irvine *forensic medical examiner*
Medico-Legal Services, Neasless Farm, Sedgefield,
Stockton on Tees, Cleveland TS21 3HE
alistair.irvine@which.net

- 1 Quinn J, Cummings S, Callahan M, Sellers K. Suturing versus conservative management of lacerations of the hand: randomised controlled trial. *BMJ* 2002;325:299-300. (10 August.)
- 2 Milroy CM, Rutty GN. If a wound is "neatly incised," it is not a laceration. *BMJ* 1997;315:1312.
- 3 Norfolk GA, Stark MM. The future of clinical forensic medicine. *BMJ* 1999;319:1316-7.

Authors' reply

EDITOR—Most hand lacerations are incised wounds from sharp objects, and we apologise to those who were offended that we did not differentiate between incised wounds and blunt lacerations. Most hand wounds in emergency departments are incised, small, superficial, and uncomplicated wounds. Trained practitioners can provide excellent wound care and routinely assess these wounds to determine whether an underlying injury or foreign body exists.

Our trial excluded patients with wounds suspicious for underlying injury, and no underlying injuries were missed on the patients enrolled in the study. De Souza's recommendations to have trained people assess all hand wounds and refer suspicious cases were precisely the methods followed in our study. These are the standards of care in any emergency department in North America and hopefully in the United Kingdom as well.

Contrary to Dunn and Watson's comments, our study does not trivialise small hand lacerations, and we are concerned that anyone would conclude this. All wounds in our study received the same initial care regardless of wound closure (sutures or a simple dressing). This care included history taking and examination by a trained practitioner followed by appropriate general wound care. The only difference in the two groups was the method of closing the wound. We think that underlying injuries may be missed regardless of the closure method used. In addition, small lacerations closed without thought or proper examination are at risk of becoming occult injuries and are also likely subject to a higher risk of infection.¹

We admit that we could have better described the wound care techniques used in this study. We assumed that most readers were knowledgeable about this or had access to the numerous articles and texts on the topic,² but our assumption in no way condones ignoring small hand lacerations.

Finally, the *BMJ* unfortunately selected photographs to accompany our article that were neither from our study nor reviewed by us before publication. It is hard to comment on the eligibility of such lacerations from photographs, but the location and apparent depth of the wound make it likely that such a wound would have been ineligible. The pictured wound is misleading and may have caused some of the concerns generated by hand surgeons.

James Quinn *associate clinical professor of medicine*
Karen Sellers *research coordinator*
University of California, San Francisco,
505 Parnassus Avenue, Box 0208, San Francisco,
CA 94143-0208, USA

- 1 Edlich RF, Panek PH, Rodeheaver GT. Physical and chemical configuration of sutures in the development of surgical infection. *Ann Surg* 1973;177:679-87.
- 2 Singer AJ, Hollander JE, Quinn JV. Evaluation and management of traumatic lacerations. *N Engl J Med* 1997;337:1142-8.

Risk factor thresholds

Threshold is £37 000 per QALY

EDITOR—Although I agree with most of Law's and Wald's conclusions with regard to risk factors, I cannot agree that, as a result, treatment thresholds do not exist.¹ With regard to the risk of coronary heart disease, the recent joint British recommendations recommend starting treatment of high blood pressure at an absolute 10 year risk of coronary heart disease of 15%, and of a high lipid ratio at 30%.² Neither these recommendations nor those of the Standing Medical Advisory Committee explain why these particular thresholds have been set. (Neither do they mention when treatment should be stopped.) But could or should it have something to do with cost? I think that, despite the article's title, Law and Wald acknowledge this implicitly by saying that people at high risk should be targeted.

There have been several published cost effectiveness analyses of lipid lowering drugs. The report from Pickin et al puts the cost per (presumably good quality) year of life gained of treating coronary heart disease risk above 3% per year at £8200, which they describe as of comparable cost effectiveness to many treatments in wide use.³ They say, however, that treatment below this level is unlikely to be affordable. The de facto threshold currently being used by the National Institute of Clinical Excellence is considerably higher—about £37 000 per QALY.

Ethical questions such as the value the NHS and other health systems should place on preventive rather than immediately lifesaving care remain largely undiscussed. Perhaps that is why so many authors overlook that resources are scarce and so thresholds must always exist. Isn't it time that this collective blind spot was removed?

Michael A Soljak *consultant in public health medicine*
Strategic Intelligence Unit, North West London
Health Authority, London W1T 7HA
michael.soljak@nwlha.nhs.uk

- 1 Law MR, Wald NJ. Risk factor thresholds: their existence under scrutiny. *BMJ* 2002;324:1570-6. (29 June.)
- 2 British Cardiac Society, British Hyperlipidaemia Association, British Hypertension Society, British Diabetic Association. Joint British recommendations on prevention of CHD in clinical practice. *Heart* 1998;80(suppl 2):S1-29.
- 3 Pickin DM, McCabe CJ, Ramsay LE, Payne N, Haq IU, Yeo WW, et al. Cost effectiveness of HMG-CoA reductase inhibitor (statin) treatment related to the risk of CHD and cost of drug treatment. *Heart* 1999;82:325-32.

Hypothesis is dangerous nonsense

EDITOR—Law and Wald discuss risk factor thresholds,¹ having established in a previous paper on the time lag hypothesis that the risk factors in French men aged 45-64² are:

- Total cholesterol concentration 6.1 mmol/l
- Systolic blood pressure (their previous "best blood pressure risk factor") 150 mm Hg
- High density lipoprotein concentration 1.3 mmol/l
- Smoking 32%

Add to this data from the British Heart Foundation:

- Average body mass index 26.6
- Those never exercising 32%
- Rate of coronary heart disease 128/100 000/year

The risk factors in British men aged 45-64 are:

- Total cholesterol concentration 6.2 mmol/l
- Systolic blood pressure 148 mm Hg
- High density lipoprotein concentration 1.3 mmol/l
- Smoking 29%
- Average body mass index 26.6
- Those never exercising 24%
- Rate of coronary heart disease 487/100 000/year

Can Law and Wald fit these figures onto their semilogarithmic scale? The suggestion that no levels of any risk factor in the Western world are currently normal, and that what we call a normal blood pressure is actually high and should be lowered, is dangerous nonsense.

Are Law and Wald aware of data from Framingham, which show that falling cholesterol concentrations are directly associated with an increased risk of coronary heart disease?³ Are they aware of research from Japan that shows a completely inverse relation between rising cholesterol concentrations and deaths from coronary heart disease?⁴ Hundreds of papers contradict the association between raised cholesterol concentrations and death from coronary heart disease.

Shestov in his lipid clinics study in Russia even showed an inverse relation, with higher rates of coronary heart disease in patients with hypocholesterolaemia. The Honolulu study shows that, in people older than 50, a low cholesterol concentration is by far the most important risk factor for premature death.⁵ Law and Wald did not show one curve relating to cholesterol lowering—the J shaped curve of total mortality with 5.2 mmol/l at the bottom of that curve (figure).

J shaped curve of total mortality v total cholesterol concentration based on data from Jacobs et al, *Circulation* 1992;86:1046-60.

Law and Wald are effectively suggesting that there is no non-dangerous blood pressure or cholesterol concentration and that, therefore, almost everyone in the Western world should be given some kind of drug treatment. This is dangerous nonsense, and we should not be afraid to say so.

Malcolm E Kendrick *medical director*
Adelphi Lifelong Learning, Adelphi Mill,
Bollington, Macclesfield SK10 5JB
malcolm@llp.org.uk

- 1 Law MR, Wald NJ. Risk factor thresholds: their existence under scrutiny. *BMJ* 2002;324:1570-6. (29 June.)
- 2 Law MR, Wald NJ. Why heart disease mortality is low in France: the time lag explanation [with commentaries by M Stampfer, E Rimm, D J P Barker, J P Mackenbach, and A E Kunst]. *BMJ* 1999;318:1471-80.
- 3 Anderson KM, Castelli WP, Levy D. Cholesterol and mortality: 30 years of follow-up from the Framingham study. *JAMA* 1987;257:176-80.
- 4 Okayama A, Ueshima H, Marmot MG, Nakamura M, Kita Y, Yamakawa M. Changes in total serum cholesterol and other risk factors for cardiovascular disease in Japan 1980-1989. *Int J Epidemiol* 1993;22:1038-47.
- 5 Schatz IJ, Masaki K, Yano K, Chen R, Rodriguez BL, Curb JD. Cholesterol and all-cause mortality in elderly people from the Honolulu Heart Program: a cohort study. *Lancet* 2001;358:351-5.

Pharmacological treatment should not be determined by age alone

EDITOR—Law and Wald provide evidence for the absence of thresholds in the relation between risk factor and disease.¹ They conclude that people with high absolute risk will benefit from a reduction in risk factors whatever their initial risk factor level. They say that in people without cardiovascular disease, intervention to change risk factors could be introduced when a person's risk of a disease event over the following few years exceeds a specified value. Risk could be estimated from age alone or age and sex. Individuals at high risk should receive drug treatment to modify all important reversible risk factors simultaneously.

Although combination pharmacological cardiovascular risk factor reduction for the whole elderly population represents an ideal strategy for the pharmaceutical industry it has numerous limitations as a public health strategy.

(1) By setting an absolute risk threshold for treatment (determined by age only) the proposed strategy still targets only those at high risk of disease (the tail of the population distribution curve of risk factors) and therefore can only have a minimal effect on the overall burden of disease in the population.

(2) There will be major opportunity costs in treating the whole elderly population with combination drugs and consequently other preventive, treatment, care and rehabilitation services for elderly people would be constrained.

(3) The authors' statement that lower limits of thresholds (such as blood pressure), beyond which harm will arise, are not reached by current drug treatment is false—the risks of polypharmacy in elderly people are significant^{2,3} and would be increased.

(4) Even in elderly people, absolute risk of coronary heart disease may not reach 3% per year without additional risk factors such as smoking and diabetes. According to the

Framingham coronary risk prediction score (www.nhlbi.nih.gov/about/framingham/riskmen.pdf), a man aged 70-74, total cholesterol 5.18-6.21 mmol/l, high density lipoprotein 1.17-1.29 mmol/l, and blood pressure (140-159)/(90-99) has a risk of coronary heart disease of 2.5% per year; for a woman the risk is only 1.3%.

Targeting people who have had a vascular event or who have diabetes with treatment to reduce risk factors is an appropriate "high risk" strategy. The only appropriate strategy, as challenging as it may be, to reduce the risk of vascular disease in the rest of the population (who cause most of the burden of disease) is to reduce the average levels of risk factors in the population, through the promotion of a healthy diet, exercise, and smoking cessation. Proposals to use drug treatment for primary prevention in the whole population over a certain age should be resisted.

Steven M Laitner *specialist registrar, public health*
National Screening Committee, Institute of Health Sciences, Oxford OX3 7LF
slaitner@ntlworld.com

- 1 Law MR, Wald NJ. Risk factor thresholds: their existence under scrutiny. *BMJ* 2002;324:1570-6. (29 June.)
- 2 Beyth RJ, Shorr RI. Epidemiology of adverse drug reactions in the elderly by drug class. *Drugs Aging* 1999;14:231-9.
- 3 Malhotra S, Karan RS, Pandhi P, Jain S. Drug related medical emergencies in the elderly: role of adverse drug reactions and non-compliance. *Postgrad Med J* 2001; 77:703-7.

Authors' reply

EDITOR—We pointed out that within the range of values of common risk factors for disease in our population there is no evidence of a threshold below which further modification of risk factors yields no further reduction in the risk of disease. None of the correspondents has provided any evidence to the contrary.

Soljak accepts the absence of biological thresholds but mentions other thresholds such as the costs per year of life gained. With low cost drugs (many useful drugs have come off patent or are shortly to come off patent) such financial thresholds could largely disappear.

We believe that Kendrick's views are incorrect and have been refuted in detail elsewhere.^{1,2} The J shaped curve relating to total mortality shows a combination of two different processes. The risk of occlusive cardiovascular disease increases with increasing serum cholesterol concentration, but many fatal chronic diseases lower serum cholesterol concentration in their early stages.²

Laitner believes that there are limitations to our view that reduction of risk factors needs to be offered on a much wider basis using drugs if necessary. Of course, the preventive treatment would have to be effective, reasonably safe, inexpensive, and sufficiently simple for people to take widely without frequent medical examination or monitoring. We agree that the assessment of such a preparation would have to be undertaken in elderly people and a quantitative assessment of benefits and risks determined.

As Laitner points out, selecting a cut-off point for absolute risk such as 3% per year (a very high risk) means that many who will die of cardiovascular disease would be below the risk threshold. This is the main reason for simply using age as the selection criterion.

The important point in our paper is that there are no biological thresholds that should limit the extent to which risk factors are reduced in the community, whether through drug treatment or diet.

Malcolm Law *professor of preventive medicine*
m.law@qmul.ac.uk

Nicholas J Wald *professor and head*
Wolfson Institute of Preventive Medicine,
Department of Environmental and Preventive
Medicine, Barts and The London, Queen Mary's
School of Medicine and Dentistry, London
EC1M 6BQ

- 1 Heart Protection Study Collaborative Group. MRC/BHF Heart protection study of cholesterol lowering with simvastatin in 20 536 high-risk individuals: a randomised placebo-controlled trial. *Lancet* 2002;360:7-22.
- 2 Law MR, Thompson SG, Wald NJ. Serum cholesterol reduction and health: assessing possible hazards. *BMJ* 1994;308:373-9.

Most patients depressed by cancer do not need drugs

EDITOR—In the ABC of psychological medicine Peveler et al discuss depression in medical patients.¹ As a haematologist I often have to respond to depressive reactions in patients with malignant disease who are confronted with a poor prognosis and difficult decisions about treatment. Some degree of this reactive demoralisation is normal and has to be acknowledged. It is psychological in nature and different from major depression as a disease of the brain.² Sometimes the problem is distinguishing depressed patients with tumours from patients who are depressed because of tumours.

Most of the studies included in the meta-analysis of drug treatment taken from the Cochrane review by Gill and Hatcher had to struggle with this distinction.³ This analysis did not compare tricyclic antidepressants with selective serotonin reuptake inhibitors, as stated in the caption of the table; rather it compared drug treatment (tricyclic antidepressants, tetracyclic antidepressants, and selective serotonin reuptake inhibitors) with placebo. The reflected benefit is therefore neither the benefit of antidepressive treatment in major depression (with medical comorbidity) nor the benefit in patients with reactive symptoms. The first two lines of the table should not be taken as evidence that every symptom of depression in a patient with cancer warrants drug treatment.

Meta-analyses of selective serotonin reuptake inhibitors versus other antidepressants are also available in the *Cochrane Library* and show no major difference in efficacy and only a modest advantage for selective serotonin reuptake inhibitors in terms of fewer dropouts.^{4,5}

Fortunately, most patients with cancer and depressive adjustment disorders respond to a supportive environment and

counselling and do not need drug treatment. The observed depressive reaction—for example, after the disclosure of malignancy—should not be taken as an excuse to withhold the truth from patients or to overmedicalise the condition. I sometimes wonder whether patients who seem to show no emotional reaction at all to bad news are in fact the more abnormal.

Ulrich S Schuler *senior haematologist*
Department of Medicine I, University Hospital Carl Gustav Carus, Fetscherstrasse 74, D-01307 Dresden, Germany
schuler@mk1.med.tu-dresden.de

- 1 Peveler R, Carson A, Rodin G. Depression in medical patients. *BMJ* 2002;325:149-52. (20 July.)
- 2 Angelino AF, Treisman GJ. Major depression and demoralization in cancer patients: diagnostic and treatment considerations. *Support Care Cancer* 2001;9:344-9.
- 3 Gill D, Hatcher S. Antidepressants for depression in medical illness. *Cochrane Database Syst Rev* 2000;CD001312.
- 4 Geddes JR, Freemantle N, Mason J, Eccles MP, Boynton J. SSRIs versus other antidepressants for depressive disorder. *Cochrane Database Syst Rev* 2000;CD001851.
- 5 Barbui C, Hotopf M, Freemantle N, Boynton J, Churchill R, Eccles MP, et al. Selective serotonin reuptake inhibitors versus tricyclic and heterocyclic antidepressants: comparison of drug adherence. *Cochrane Database Syst Rev* 2000;CD002791.

Management of infertility: one stop clinic may offer solution

EDITOR—In their review of the management of infertility Cahill and Wardle report that most investigations to establish a cause of subfertility are simple to undertake.¹ In our experience, these investigations are protracted for many couples, requiring several outpatient visits and often inpatient laparoscopy for the female partner. At a time of understandable stress for the couple, the assessment process can be frustrating for all concerned.

Since February 2000 we have attempted to rationalise the investigation of infertility using a "one stop" approach as used for conditions such as dyspepsia and menorrhagia (A Taylor et al, meeting of the British Society for Gynaecological Endoscopy, Portsmouth, May 2002). Couples who fulfilled our selection criteria were invited to the one stop fertility clinic.

History, examination, and review of blood tests previously organised were followed by pelvic ultrasonography (to look for polycystic ovaries, adnexal masses, and uterine fibroids), hysteroscopy (to look for endometrial polyps, submucous fibroids, and uterine anomalies), and culdoscopy with hydrotubation (to look for adhesions, endometriosis, and tubal disease).

Culdoscopy (telescopic examination of the pouch of Douglas via the posterior fornix) was first introduced to gynaecological practice in 1941 but was superseded by laparoscopy.² Recently a refined version using narrow instruments and saline irrigation to investigate infertility has been reintroduced and carried out under local anaesthesia.³ This technique allows for clear visualisation of the pouch of Douglas and adnexa, as well as assessment of tubal patency by hydrotubation. Although culdoscopy is more invasive than ultrasonography

in the female partner,⁴ it can exclude pelvic endometriosis and adhesions, which are important causes of infertility.

To date, 130 of the 211 (61%) couples referred to the clinic met our selection criteria, and 87 of this group have been seen. Of these, 70 patients (80%) completed the three diagnostic procedures successfully and 26 patients (30%) were found to have pelvic pathology. The average time for the three procedures was 41 (SD 17) minutes. The investigations were well tolerated, and there were no serious complications. Patient feedback was positive, the availability of immediate results being particularly appreciated.

While a one stop approach is not suitable for all couples, and not all couples opt for outpatient investigation, we believe that a one stop fertility clinic offers a rational, efficient, and potentially cost effective alternative to the traditional investigation of infertility.

Malini Sharma *clinical research fellow*
Alex Taylor *clinical research fellow*
Amina Al Khouri *visiting clinical fellow*
Natasha Goumenou *visiting clinical fellow*
Panos Tsirkas *visiting clinical fellow*
Peter Scott *clinical research fellow*
Adam Magos *consultant gynaecologist*
Minimally Invasive Therapy Unit and Endoscopy Training Centre, University Department of Obstetrics and Gynaecology, Royal Free Hospital, London NW3 2QG
gynendo@rfc.ucl.ac.uk

- 1 Cahill DJ, Wardle PG. Management of infertility. *BMJ* 2002;325:28-32. (6 July.)
- 2 Decker A, Decker W, Milowsky J. Anesthetic technique for culdoscopy examination: preliminary report. *Am J Obstet Gynecol* 1950;59:455-7.
- 3 Gordis S, Campo R, Rombauts L, Brosens I. Transvaginal hydro-laparoscopy as an outpatient procedure for infertility investigation. *Hum Reprod* 1998;13:99-103.
- 4 Kelly SM, Sladkevicius P, Campbell S, Nargund G. Debate continued: Investigation of the infertile couple: a one-stop ultrasound-based approach. *Hum Reprod* 2001;16:2481-4.

Collaboration is key to preventing syphilis

EDITOR—The letter by Clark et al about the epidemiological data represented in our paper on the resurgence of infectious syphilis implies that we did not consult with local colleagues in preparing our paper and that the situation in Manchester has thereby been "misrepresented" to the detriment of the genitourinary medicine clinics dealing with the increased workload.^{1,2} In fact the collaboration between the Communicable Disease Surveillance Centre and colleagues in genitourinary medicine on the frontline was close and productive, as evidenced by the fact that representatives from clinics at each outbreak site participated in all aspects of preparing the report.

Data from enhanced surveillance systems, such as that instituted in Manchester, provide more sensitive information. By collaborating with local clinicians, we made every effort to ensure that figures obtained from each site were as current as possible at the time of submission. However, the explosive nature of the outbreak, coupled with the attendant reporting and publishing delays, may have contributed to the disparity in the data noted by Clark et al. Rather than suggesting that any of the outbreaks had

ended, we aimed to raise awareness among health professionals about the resurgence of infectious syphilis in England and to reiterate key aspects of its clinical management and public health control and the central importance of clinics.

Through its national and regional offices the Communicable Disease Surveillance Centre has continued to work with local partners in identifying and managing these and other outbreaks of sexually transmitted diseases in England. This robust collaboration between genitourinary medicine, public health, microbiology, and the voluntary sector has raised awareness of outbreaks and facilitated the development of national guidelines on controlling outbreaks of sexually transmitted diseases³; a new national enhanced surveillance programme for infectious syphilis; revision of clinical guidelines on syphilis management; additional investment into local prevention initiatives; and research aimed at developing new diagnostic assays for infectious syphilis. This process has been overseen by a national advisory committee, which has formal representation from a range of stakeholders in sexual health.

We sincerely hope that our hard pressed colleagues in genitourinary medicine will continue to recognise the added value of working with their public health colleagues in investigating outbreaks of sexually transmitted disease. We also hope that they will continue to collaborate with the Communicable Disease Surveillance Centre, a process that was extremely helpful in the syphilis outbreaks.

Kevin A Fenton *consultant epidemiologist*
HIV/STI Division, PHLS Communicable Disease Surveillance Centre, London NW9 5EQ
kfenton@phls.org.uk

Lorraine Doherty *senior medical officer*
Department of Health, Social Services and Public Safety, Belfast BT4 3SJ

- 1 Clark P, Cook PA, Lighton L, Syed Q, Bellis MA. Don't forget syphilis. *BMJ* 2002;325:775. (5 October.)
- 2 Doherty L, Fenton KA, Jones J, Paine TC, Higgins SP, Williams D, et al. Syphilis: old problem, new strategy. *BMJ* 2002;325:153-6. (20 July.)
- 3 Guidelines for managing outbreaks of sexually transmitted infections at a local, district or regional level. An outbreak plan. Available at www.phls.co.uk/topics_az/hiv_and_sti/guidelines/managing_outbreaks_of_sti.htm

Are inactivation procedures for blood products good or bad?

EDITOR—In their report on recent developments in transfusion medicine, Regan and Taylor mention that, in future, infectious diseases transmitted by cellular blood products might be prevented by inactivation procedures.¹ All pathogen inactivation procedures for cellular blood products involve one step, and failure cannot be compensated by a second step.

Owing to technical constraints, validation of procedures is limited to five to seven logs. Without screening (or a virus carrier escaping the screening process), however, a platelet concentrate from a hepatitis B virus carrier might contain up to 5×10^{10} virus particles per ml. Parvovirus B19 load might exceed 10^{13}

viruses per platelet concentrate. Moreover, validation of inactivation procedures relies mainly on the comparability of the model viruses to the relevant human viruses (for example, hepatitis B, hepatitis C), which is a matter of constant debate. Thus, implementation of inactivation does not allow nucleic acid amplification tests to be given up but might even pave the way for additional such tests to limit the pathogen load.

All inactivation procedures affect not only the pathogen but also to some degree the active ingredient. It could be argued that subtle differences not detected by clinical trials do not matter, but recent experience with solvent/detergent treated plasma has shown the contrary.^{2,3}

If pathogen inactivation works by modifying pathogens' nucleic acids, residual amounts of the inactivating substance in the product might also affect the nucleic acids of the transfusion recipient, resulting in possible mutagenicity and carcinogenicity—there is said to be no threshold for mutagenicity and carcinogenicity. Manufacturers try to rule out these side effects by extensive pharmacological and toxicological studies precisely defined by the regulatory authorities. But can these studies also exclude very low incremental risks of carcinogenicity in humans?

The risk of dying from infectious diseases through cellular blood products is vanishingly small: about two in 1 million transfusions (one in 5 million for HIV and one in 600 000 for bacteria,⁴ while no cases of hepatitis C virus have occurred in Germany since the polymerase chain reaction was introduced). I think that pathogen inactivation is only justified if it has a net benefit—if death by malignancies and other side effects through the inactivated product does not exceed the number of patients saved from death by infection. Pathogen inactivation might seem attractive for the public and politics. Establishing its benefit will be a challenge for manufacturers and regulatory authorities.

Gregor Caspari *head, transfusion unit*
Institut für Transfusionsmedizin, 14770
Brandenburg an der Havel, Germany
gregor.caspari@dgn.de

- 1 Regan F, Taylor C. Blood transfusion medicine. *BMJ* 2002;325:143-7. (20 July.)
- 2 De Jonge J, Groenland TH, Metselaar HJ, IJzermans JN, van Vliet HH, Visser L, et al. Fibrinolysis during liver transplantation is enhanced by using solvent/detergent virus-inactivated plasma (ESDEP®). *Anaesth Analg* 2002;94:1127-31.
- 3 Kerr K. Plasma's fatal flaw. New variety on LI connected to liver deaths. *Newsday (Nassau and Suffolk edition)* 2002 May 6:A.06. www.newsday.com (accessed 1 October 2002).
- 4 Perez P, Salmi LR, Folléa G, Schmit J-L, de Barbeyrac B, Sudre P, et al. Determinants of transfusion-associated bacterial contamination: results of the French BACTHEM case-control study. *Transfusion* 2001;41:862-72.

Weber's test demystified

Physics renders Weber's test not so mysterious ...

EDITOR—Weber's test is a positive Weber's test in the normal ear in unilateral sensorineural hearing loss but in the affected ear in unilateral conductive hearing

loss—has baffled many neurologists and ear, nose, and throat surgeons for some time; I wonder whether my explanation would convince him.¹

The word conduction is used confusingly by ear, nose, and throat surgeons and neurologists to describe normal transmission of sound from the outside world to the ear. Everything apart from the sensorineural aspect of the hearing is thought to be conductive. The sound is normally conducted (transmitted) through the air through the external ear into the middle ear. This makes air a better sound conductor (when defined this way) than a solid object.

It is, however, clear to any engineering student that bone or any denser object is a better conductor of sound than is air. When the tuning fork is placed directly on the bone, there is no significant sound transmission from the tuning fork directly through the air. The conduction being tested is that through bone to the inner ear.

The air medium in the ear, being a less efficient transmitter of sound, results in sound energy loss at the interface of bone and air. The resultant sound energy to the inner ear is therefore less. If you have a more solid (denser) object in the ear (which would have resulted in conduction deafness (as defined by doctors) the sound conduction is actually better. Less energy is lost, and the sound is localised to that side in Weber's test.

If both ears are blocked but with different materials with different conductive properties, positive results in Weber's test would localise to the side with the denser and therefore better sound conducting material. You could test this by blocking your ear with one finger and the other with another material, comparing the sounds and comparing each with air. Nice little study for a neurologist, I say.

Chima E Mbubaegbu *consultant orthopaedic surgeon*
Homerton University Hospital NHS Trust, London
E9 6SR
chima.mbubaegbu@ntlworld.com

1 Weatherall MW. The mysterious Weber's test. *BMJ* 2002;325:26. (6 July.)

... and a collaborative group of otorhinolaryngologists reports its findings

EDITOR—Weber's test is indeed mysterious, and the filler by Weatherall had our ear, nose, and throat department, as well as an eminent visiting professor from Glasgow, occupied for the best part of a week.¹ Literature searches and audiological experiments were conducted. Arcane, dusty journals were dusted off.

The simulation of conductive hearing impairment by occluding the ear with a finger suggested by Weatherall is the basis of the Bing test. A tuning fork is placed on the skull, and the patient indicates when the sound disappears. A normal conductive mechanism is demonstrated if the sound is heard again when the ear canal is occluded.²

The so called occlusion effect described by Tonndorf et al in 1966 is responsible for this phenomenon.³ Sound conducted

through bone causes the cochlea, the ossicular chain, and the air in the external auditory canal to vibrate. Some lower frequency sound, as produced by the 512 Hz tuning fork, escapes from the canal. When the ear is occluded, these frequencies cannot escape and the sound seems to become louder.

We conducted an experiment to determine whether the exclusion of environmental noise or the occlusion effect was responsible for Weatherall's observation. Baseline binaural air and bone thresholds were obtained at 500 Hz on a subject in a soundproof audiometric booth. Both ears were occluded with modified tympanometry earpieces (blocked with plasticine), without causing a measurable conductive loss. The ears were then occluded with foam "ear defenders," which caused a 25 dB conductive impairment. In both instances the bone conduction thresholds improved by 15 dB.

Having proved that an occlusion effect occurs independently of a conductive loss, we then introduced background noise to assess its influence. Binaural masking noise (similar to environmental noise) was presented over headphones, first occluding with modified tympanometry earpieces and then with ear defenders. Bone conduction thresholds were reassessed when the masking noise was audible despite the conductive impairment. In both occluded ear conditions the masking noise did not affect the improvement in the bone conduction thresholds.

We therefore concluded that it is the occlusion effect, rather than elimination of environmental sound, that is responsible for the improved bone conduction threshold when occluding a normal ear. Other explanations are required for aetiologies of conductive loss other than occlusion. Middle ear effusion and ossicular chain disruptions cause a "mass loaded" middle ear, with lowering of the inherent resonant frequency. Ossicular chain fixation causes a phase shift in the sound wave. Both cause preferential transmission of lower frequencies to the cochlea.²

Gary Kroukamp *registrar*
Department of Otorhinolaryngology, Faculty of Health Sciences, University of Stellenbosch, 7505
South Africa

On behalf of Carel Van Wyk, Adriaan Pentz, Ola Basson, Zak Mansab, Amal Alabdulla, Rory Attwood, James Look, and George Browning (visiting professor), all of the ear, nose, and throat department.

- 1 Weatherall MW. The mysterious Weber's test. *BMJ* 2002;325:26. (6 July.)
- 2 Gelfand SA. *Essentials of audiology*. New York: Theimme, 1997.
- 3 Tonndorf J, Campbell RA, Bernstein L, Reneau JP. Quantitative evaluation of bone conduction in cats. *Acta Otolaryngol* 1966;213(suppl):10-38.

Rapid responses

Correspondence submitted electronically
is available on our website