Media influence on suicide

Television programme makers have an ethical responsibility

EDITOR—We agree with Hawton and Williams that training courses for careers in the media offer the potential for improved portrayal of suicide in the media.¹ The media, however, clearly can affect many facets of health related behaviour.

We recently reported the effect of the death from cervical cancer of a character (Alma) in the television soap opera *Coronation Street* on the NHS cervical screening programme in the north west of England.^{2,3} Our studies showed an excess of 14 000 cervical smear tests performed as a result of the storyline (a 21% increase on the previous year), although only 2000 of them were in women whose test was overdue or who had had no previous smear test. The remaining 12 000 smear tests were performed on women attending for an early, unscheduled test or who were due a smear test anyway and brought their appointment forward.

The large increase in the number of smear tests led to a strain on local laboratories, with the time taken to report results increasing to beyond acceptable quality assurance limits—a factor likely to provoke excess anxiety in women. We also found that many women were prompted to attend for a cervical smear test because the storyline made them worry.

This anxiety generating approach contrasts with current initiatives to encourage women to make an informed choice about screening.

Television programme makers should realise the power of such stories not only to achieve maximal viewing figures but also to cause fear and anxiety, as well as the consumption of scarce healthcare resources. Those responsible for promoting health need to engage programme makers in a full ethical debate.

Andy Howe *specialist registrar in public health*
andyhowe@clara.co.uk

Vicci Owen-Smith *specialist registrar in public health*
Department of Clinical Strategy, Greater Manchester Strategic Health Authority, Manchester M60 7LP

Judith Richardson *director of public health*
South Manchester Primary Care Trust, Withington Hospital, West Didsbury M20 2LR

³ Richardson J, Owen-Smith V, Howe A. The effect of "Alma's" death on women attending for a cervical smear: a questionnaire survey. *J Public Health Med* 2002;24:305-6.

Media influence behaviour

EDITOR—Hawton and Williams's study provides evidence that supports an argument for (increased) awareness among media producers and editors about their potential influence on the public over health matters.¹ Howe et al in response note the effect on the NHS cervical screening programme of a television character's death (Alma in *Coronation Street*) from cervical cancer (letter above). We examined the impact of the same television story and related tabloid newspaper reporting on inquiries made to CancerBACUP's helpline (the United Kingdom's leading cancer information charity, www.cancerbacup.org.uk).

The change in the volume and content of calls during the three months between the story "breaking" in the *Sun* newspaper and Alma's eventual death, compared with calls received in the three months before the story, supports Hawton and Williams's work.¹ Peaks in inquiries about cervical cancer occurred on three occasions, directly coinciding with developments in the storyline (figure). The story triggered up to 300 additional weekly enquiries to CancerBACUP. Evidence corroborated Hawton and Williams's suggestion of a similarity between media stimulus and the viewer in terms of age, sex, and nationality.¹ Data

showed a slight increase in the proportion of calls from women in their 50s and 60s; the actor who played Alma was 63 years old.

In the three months before the storyline just 6.7% of people telephoning CancerBACUP about cervical cancer said that they had first heard about the charity via a newspaper or the television. This percentage rose to 41.8% while the storyline was in progress. Furthermore, the proportion of inquiries from homemakers and retired people more than doubled compared with calls during the previous three months.

Our research has questions left unanswered. What, for example, became of the people whom the *Coronation Street* story worried but who did not contact CancerBACUP? Did they contact other information services or their general practitioner, or did they remain concerned, probably unnecessarily, about the risk of cervical cancer? What is clear is the potential for media to impinge on the viewing public in a marked, and in this case measurable, way.

The trend for giving information at the end of potentially delicate television programmes seems not only responsible but necessary. In our study making such information available led to the use of a cancer helpline by a broad and, in part, previously untapped group.

Rachel Hardyman *research officer*
CancerBACUP, London EC2A 3JR
rhardyman@cancerbacup.org

Geraldine Leydon *lecturer and MRC fellow*
Cancer and Public Health Unit, London School of Hygiene and Tropical Medicine, London WC1E 7HT

Inquiries about cervical cancer to CancerBACUP telephone helpline, April-June 2001

¹ Hawton K, Williams K. The influence of media on suicide. *BMJ* 2002;325:1374-5. (14 December.)

² Howe A, Owen-Smith V, Richardson J. The impact of a television soap opera on the NHS cervical screening programme in the north west of England. *J Public Health Med* 2002;24:299-304.

1 Hawton K, Williams K. The influence of media on suicide. *BMJ* 2002; 325:1374-5. (14 December.)

Media's role is double edged

EDITOR—The alarming escalation of charcoal burning suicide in Hong Kong supports Hawton and Williams's call for guidelines on the reporting of suicide.¹ We previously reported on the emergence of this new method of suicide in Hong Kong.²

In November 1998 a 35 year old woman committed suicide by burning charcoal in a barbeque grill in her sealed and cramped apartment. Compared with jumping, which accounts for most local suicide deaths, suicide by burning charcoal was romanticised as an easy and comfortable way of dying. The incident was pictorially reported in the media. Two months after its appearance charcoal burning became the third commonest method of suicide in Hong Kong, where carbon monoxide poisoning was previously uncommon.³ In 2001 it replaced hanging as the second commonest method of suicide, accounting for 25% of all deaths from suicide.

People committing suicide by charcoal burning were often middle aged and were portrayed as debt ridden because Hong Kong was experiencing its first recession after two decades of economic boom. Charcoal burning was represented as a legitimate way out of the person's financial predicaments. Our coroner's court findings were highly consistent with these media representations.²

The influence of the media on suicide is contentious and has not been taken seriously in public health.^{3,4} In the case of suicide by charcoal burning in Hong Kong, concerns have been raised by researchers and healthcare workers about the potentially contagious impact of media reporting. However, news editors have expressed reservations and remain unconvinced about the guidelines published by the World Health Organization on suicide reporting.

The first suicide intervention and prevention centre in China, funded by the Chinese government, was opened in December in Beijing. Media publicity was rightly adopted as one of the key strategies for raising public awareness.⁵ None the less, given that the handy use of pesticides claimed most of the suicide deaths in rural China, the possible effect of such publicity deserves monitoring too.

Kathy P M Chan *adjunct associate professor*
chanpm@ha.org.hk

Dominic T S Lee *associate professor*

Sing Lee *professor*

Department of Psychiatry, Chinese University of Hong Kong, Shatin, New Territories, Hong Kong Special Administrative Region, China

Paul S F Yip *director*

Centre for Suicide Research and Prevention, University of Hong Kong, Hong Kong Special Administrative Region

1 Hawton K, Williams K. The influence of media on suicide. *BMJ* 2002; 325:1374-5. (14 December.)

2 Lee D, Chan K, Lee S, Yip P. Burning charcoal: A novel and contagious method of suicide in Asia. *Arch Gen Psychiatry* 2002; 59:293-4.

3 Stack S. The effect of the media on suicide: Evidence from Japan 1955-1985. *Suicide Life Threat Behav* 1996; 26:132-42.

4 Phillips DP. The influence of suggestion on suicide: substantive and theoretical implications of the Werther effect. *Am Sociol Rev* 1974; 39:340-54.

5 Lee S, Kleinman A. Suicide as resistance in Chinese society. In: EJ Perry EJ, Selden M, eds. *Chinese society: change, conflict and resistance*. London: Routledge, 2000:221-40.

Death risk other than from suicide is raised in self harm

EDITOR—Jenkins et al report on continuing suicide risk after deliberate self harm.¹ They use their findings to argue that clinicians should pay close attention to continuing risk of suicide in people with a history of deliberate self harm. Their findings, in a cohort from the late 1970s, are similar to findings from a 1981 Scottish discharge cohort.²

Using the Scottish linked dataset we followed up a cohort of 8304 people discharged over a 13 year period from Scottish general hospitals after deliberate self harm. We found that the greatest number of deaths from suicide or undetermined cause were in the five years after discharge. In the third five year period, however, the ratio of observed self harm to expected self harm was 5.33 (95% confidence interval 3.26 to 8.23) for men and 9.46 (5.61 to 14.95) for women. Homicides and accidental deaths were also raised.

We endorse the advice by Jenkins et al that clinicians should pay attention to suicide risk but think that their method may have concealed another important clinical implication. They note that people who had consumed alcohol at the time of the initial episode were less likely to be traced. They also censored the 13 deaths in their cohort that were not attributed to definite or probable suicide.

In the Scottish cohort, we examined deaths by suicide and undetermined cause, and deaths by other causes. Altogether 214 people died by suicide or undetermined cause during the follow up period, 196 more deaths than expected. Nine other categories of illness, however, accounted for 780 deaths, 344 more than would have been expected at general population rates. Natural causes, therefore, were responsible for more excess deaths than were suicides.

We identified a higher risk of digestive system disease, respiratory and circulatory disease, and cancers. The pattern indicates to us that alcohol, as well as unhealthy lifestyles and possibly impaired access to medical care, may be important in this group of people. Clinicians should pay attention to alcohol use and physical health as well as suicide risk in people with a history of deliberate self harm.

Cameron Stark *honorary senior lecturer*

Highlands and Islands Health Research Institute, Inverness IV2 3ED
crs@hihri.abdn.ac.uk

David Hall *consultant psychiatrist*

Dumfries and Galloway Primary Care NHS Trust, Crichton Royal Hospital, Dumfries DG1 4TG

Anthony Pelosi *consultant psychiatrist*

Lanarkshire PC NHS Trust, Strathclyde Hospital, Motherwell ML1 3BW

1 Jenkins GR, Hale R, Papanastassiou M, Crawford MJ, Tyrer P. Suicide rate 22 years after parasuicide: cohort study. *BMJ* 2002; 325:1155. (16 November.)

2 Hall DJ, O'Brien F, Stark C, Pelosi A, Smith H. Thirteen year follow-up of deliberate self-harm, using linked data. *Br J Psychiatry* 1998; 172:239-42.

Low dosage tricyclic antidepressants in depression

Giving low dose tricyclics is not justified by evidence

EDITOR—The meta-analysis by Furukawa et al must be considered downright naughty.¹ While masquerading as a contribution to an academic debate about appropriate antidepressant treatment it actually does little more than endorse the widely held prejudice in favour of using low dose tricyclics in depression. Published in a widely read UK general medical journal, it will inevitably encourage a practice that is not encouraged by either the Royal College of Psychiatrists or the American Psychiatric Association.

The subject of optimal dosage of tricyclics remains controversial, but the value of treatment with tricyclic antidepressants at standard dosage compared with placebo is abundantly clear. As Furukawa et al concede but perhaps do not emphasise sufficiently, the evidence for low dose tricyclic antidepressants is of generally poor quality. Many of the trials used in their analysis took place before standardised diagnostic or outcome criteria were commonplace.

What is less clear is the motivation for their undertaking. Fluoxetine is now available in generic preparations for about £7 per month, significantly reducing the financial advantage of older antidepressants. Some evidence supports the use of tricyclics in severe depression, but presumably the authors would not recommend low dose tricyclics in such patients. The chronic severe nature of depression would make it unethical to recommend a treatment without a secure evidence base. Being charitable, one can only hope that the therapeutic advice was added to add colour to an otherwise unexceptional meta-analysis.

Hugh M Jones *specialist registrar*

Ward DB2, Maudsley Hospital, London SE5 8AZ
hugh.jones@iop.kcl.ac.uk

Competing interests: HMJ has received a small grant for a research study on olanzapine by Eli Lilly.

1 Furukawa AT, McGuire H, Barbui C. Meta-analysis of effects and side-effects of low dosage tricyclic antidepressants in depression: systematic review. *BMJ* 2002; 325:991-5. (2 November.)

Evidence to change current guidelines is insufficient

EDITOR—The study by Furukawa et al consists of two separate analyses looking at quite different things so it is difficult to draw any firm conclusions.¹

The first meta-analysis of 35 studies indicates that tricyclic antidepressants given at low dosage may improve certain symptoms, some of which may be secondary to

depression. This has already been established, and, as respondents on *bmj.com* have pointed out, low dose tricyclics are commonly used to treat chronic pain, insomnia, and anxiety, whether these symptoms occur secondary to a depressive illness, as part of self limiting adjustment and anxiety disorders, or independently.²

The question that remains unanswered is whether low dose tricyclics can adequately treat an episode of clinical depression. Furukawa et al are careful to say only that their meta-analysis shows that low dose tricyclics are more effective than placebo.

The second meta-analysis, which examines whether low dose tricyclics are as effective as recommended doses in the treatment of depressive illness, is only six studies with 551 patients. The authors acknowledge some of the weaknesses in their analysis (heterogeneity of the studies, the quality of some studies, the adequacy of blinding, and the lack of operational criteria and interview schedules to diagnose depression). Respondents on *bmj.com* have also pointed out that the cut-off point for a therapeutic dose is incorrect as most authorities recognise that 125 mg, not 100 mg, is the minimum recommended dose.²

The only useful result to come out of this small analysis is therefore that patients do not tolerate higher doses as well as low doses of tricyclics. Again, this is an already established finding and adds nothing new to the debate.

My concern is that this meta-analysis will be misinterpreted and will deter clinicians from prescribing recommended doses of tricyclics and thereby fail to treat depressive illness adequately. A report in *Hospital Doctor*, that the use of low dose tricyclics in depression is "vindicated," confirms my fear.³

Imad M Ali *locum consultant psychiatrist*
East Glamorgan Mental Health Unit, Church
Village, Rhondda Cynon Taff, South Wales
CF38 1AB
imad.ali@virgin.net

Competing interests: IMA has received funding from Eli Lilly for a study and a few small honorariums for talking to professionals on the management of depression, as well as having attended conferences and symposia, some of which were sponsored by various pharmaceutical companies.

1 Furukawa AT, McGuire H, Barbui C. Meta-analysis of effects and side-effects of low dosage tricyclics antidepressants in depression: systematic review. *BMJ* 2002;325:991-5. (2 November.)

2 Electronic responses. Meta-analysis of effects and side-effects of low dosage tricyclics antidepressants in depression. *bmj.com* 2002. www.bmj.com/cgi/eletters/325/7371/991 (accessed 29 Jan 2003).

3 Hitche L, Roberts R. Use of low-dose tricyclics in depression "vindicated." *Hospital Doctor* 2002 November 14:10.

Non-superiority does not equal equivalence

EDITOR—In their comprehensive meta-analysis of tricyclic antidepressants, Furukawa et al addressed the important question of dose-response.¹ This meta-analysis provides a further opportunity to learn about the concept of equivalence in meta-analysis.

A key question that practitioners will ask of this meta-analysis is whether low doses of tricyclics are equivalent to standard doses.

However, in the absence of a clearly devoted research question, it remains unclear whether this review was intended as a superiority or equivalence trial. This should be clarified for practitioners as these concepts are not immediately intuitive.

The message that low dose tricyclics may or may not be as effective as standard dosage tricyclics was correctly asserted in the original paper. However, Furukawa misleadingly says on *bmj.com* that an earlier Canadian meta-analysis had shown that selective serotonin reuptake inhibitors in standard dosage are as effective as tertiary amine antidepressants in low and standard dosages.^{2,3} In contrast to the earlier statement by Furukawa et al, indicating non-superiority between standard and low dose tricyclics in their meta-analysis, this statement claims that the Canadian meta-analysis showed that selective serotonin reuptake inhibitors are as effective as tricyclics.

These concepts, non-superiority and equivalence, are keenly different.⁴ Equivalence cannot be safely defined simply as overlapping confidence intervals, as is inferred above from the Canadian study, which did not explicitly set out to prove equivalence. Rather, if proving equivalence is the intention, then objective criteria should be defined a priori, as for example, 95% confidence intervals within the bounds of 0.90 to 1.10 for an aggregate relative effect size of 1.00.

Theory aside, how shall practitioners interpret this meta-analysis? As Furukawa et al have said, standard dosage is not significantly better than low dosage. But this does not infer that low or standard dosage is better or worse than the other. Equally important, neither does it mean they are the same. The bottom line is that we remain uncertain whether standard and low doses are the same or different. The outer limits of the 95% confidence intervals define just how different they might be, and the possibility of clinically important differences cannot be ruled out. Much larger studies are needed to bring these bounds within the threshold of an acceptable definition of equivalence.⁵

Janet E Martin *project leader, evidence-based prescribing initiative*
London Health Sciences Centre, London, ON,
Canada N6A 4G5
janet.martin@lhsc.on.ca

Competing interests: None declared.

1 Furukawa AT, McGuire H, Barbui C. Meta-analysis of effects and side effects of low dosage tricyclics antidepressants in depression: systematic review. *BMJ* 2002;325:991-5. (2 November.)

2 Furukawa TA. Re: low dose tricyclics are not justified by the evidence. *bmj.com* 2002. bmj.com/cgi/eletters/325/7371/991#28044 (accessed 5 Feb 2002).

3 Trindade E, Menon D. *Selective serotonin reuptake inhibitors (SSRIs) for major depression. Part 1: Evaluation of the clinical literature (report 3E)*. Ottawa: Canadian Coordinating Office for Health Technology Assessment, 1997.

4 Greene WL, Concato J, Feinstein AR. Claims of equivalence in medical research: are they supported by the evidence? *Ann Intern Med* 2000;132:715-22.

5 Edmonds ML, Camargo CA Jr, Brenner BE, Rowe BH. Replacement of oral corticosteroids with inhaled corticosteroids in the treatment of acute asthma following emergency department discharge: a meta-analysis. *Chest* 2002;121:1798-805.

Authors' reply

EDITOR—Jones questions our motivation for undertaking this meta-analysis. It was sim-

ple. We surveyed psychiatric practices in Japan and found that benzodiazepines were often given with antidepressants and that tricyclic antidepressants were often given in "inadequate dosage." We condemned these practices.¹

Then, however, we wondered if there was enough evidence to say this. We therefore undertook two meta-analyses, one of combined treatment with antidepressants and benzodiazepines and another of tricyclics at low dosage. The first one found that our reproof may be wrong: the combined treatment brought about more response than antidepressant alone up to four weeks.² The second meta-analysis again showed that our reprimand (along with similar claims by many others) was ill founded.

However, some of Ali's concerns are understandable. For lack of space, we could not report all sensitivity analyses examining possible confounders.

Firstly, we limited our analyses to those using operationalised criteria; the results were essentially identical.

Secondly, we conducted several subgroup analyses (planned a priori) on older people, patients in primary care, and psychiatric patients. The results were all compatible with the overall results. The details of these sensitivity and subgroup analyses will soon be available in the *Cochrane Library*.

We agree with Martin's argument on non-superiority and equivalence. The plausible ranges of absolute benefit increase of response for standard dosage over low dosage, on the basis of the 95% confidence intervals of relative risks and the average control event rates, are 6% (95% confidence interval -4% to 14%) at four weeks and -4% (-24% to 9%) at six to eight weeks; that of absolute risk increase of dropout owing to side effects is 9% (4% to 12%).

Thus clinically important increases in benefit on standard dosage over low dosage cannot be ruled out. At the same time, the increases point to a roughly equal number of patients who drop out because of side effects. We remain uncertain whether standard and low doses are the same or different, but it seems clear to us that we should not try to increase the dosage above 100 mg or 125 mg up front for patients with depression.

So, we must not turn the argument upside down, as does Jones. If a meta-analysis does not support a certain practice recommended by a guideline, is it this part of the guideline that may be wrong or is it the meta-analysis?

Toshi A Furukawa *professor of psychiatry*
Department of Psychiatry, Nagoya City University
Medical School, Mizuho-cho, Mizuho-ku, Nagoya
467-8601, Japan
furukawa@med.nagoya-cu.ac.jp

Corrado Barbui *lecturer in psychiatry*
Department of Medicine and Public Health,
Section of Psychiatry, University of Verona,
Ospedale Policlinico, I-37134 Verona, Italy

Hugh McGuire *trials search coordinator*
Cochrane Collaboration Depression, Anxiety and
Neurosis Group, Health Services Research, King's
College Institute of Psychiatry, London SE5 8AF

Competing interests: TAF has received fees for speaking from several pharmaceutical companies, some of which manufacture various types of antidepressants such as paroxetine, fluvoxamine, and milnacipran.

- 1 Furukawa TA, Kitamura T, Takahashi K. Treatment received by depressed patients in Japan and its determinants: naturalistic observation from a multi-center collaborative follow-up study. *J Affect Disord* 2000;60:173-9.
- 2 Furukawa T, Streiner DL, Young LT. Antidepressant plus benzodiazepine for major depression. *Cochrane Library*. Issue 3. Oxford: Update Software, 2002.

Regulatory bodies can root out corruption

EDITOR—Wilmshurst described institutional corruption in medicine.¹ In Queensland the Nursing Act enables the Queensland Nursing Council to appoint inspectors. Acts regulating other health professions, including medicine, also have similar powers. Inspectors can require the production of documents, etc, and can enter and search for evidence of breaches of the act. The inspectors can also apply to a magistrate for a warrant, and it is an offence to obstruct an inspector.

These are fairly draconian powers, but they were born out of an era of widespread corruption of the state government and the enormous difficulty a subsequent inquiry had in rooting out offenders.

Obviously, the council does not use these powers as a first recourse. It is also my perception that universities here have very good relations with regulatory bodies and would probably cooperate fully with a legitimate investigation. Usually the appropriate councils will investigate complaints confidentially and bring material into the public arena only if a matter goes to a hearing. Thus there is generally no damage to an institution's reputation until that stage, by which time they may have deserved the damage.

I suggest that, if Wilmshurst's complaints are indicative of a common practice, amendments be made to the legislation regulating medicine along the lines of the Queensland act. The powers do not have to be used, but they may be useful in reminding recalcitrants and those who would obstruct investigations that failure to cooperate is not acceptable.

With the General Medical Council attempting to improve the reputation of the profession after the scandals of recent years, it does not need the behaviours described by Wilmshurst.

Francis J Morrissey senior lecturer
School of Nursing Sciences, James Cook University,
Townsville, Queensland 4811, Australia
Joe.Morrissey@jcu.edu.au

- 1 Wilmshurst P. Institutional corruption in medicine. *BMJ* 2002;325:1232-5. (23 November)

Mortality from methicillin resistant *Staphylococcus aureus*

Welsh surveillance data show plateau

EDITOR—Crowcroft and Catchpole conclude from a study of death certificates that

Bacteraemias due to *Staphylococcus aureus* reported in Wales, 1991-2001

infections due to methicillin resistant *Staphylococcus aureus* are an increasing cause of mortality in England and Wales.¹ That this was likely to have been the case in Wales over the period of their study, 1993-8, is supported by surveillance data, which show a large rise in bacteraemias caused by *S aureus* between 1991 and 1997 (figure).

This can largely be attributed to increasing occurrence of methicillin resistant *S aureus* as a cause of these infections from 1993, although a smaller rise in methicillin sensitive *S aureus* bacteraemias was also recorded. The degree to which this increase in staphylococcal bacteraemias has affected outcomes in relation to the underlying clinical conditions is, however, unknown, and for the reasons pointed out by Liggett and Swift in their electronic response cannot be accurately inferred from data derived from death certificates.²

The Welsh surveillance data do not support the statement that the number of infections caused by methicillin resistant *S aureus* is continuing to increase every year in England and Wales and the supposition that mortality is also continuing to rise. The number of bacteraemias in Wales has changed little since 1997, and laboratory reports relating to overall numbers of isolates from infected and colonised patients reached a plateau in 2000 after a comparatively small rise between 1998 and 2000 (5650 to 6031 reports).

Presumably the steady state that has been achieved reflects the interaction of the innate transmissibility and virulence of the *S aureus* clones circulating; the ability to prevent transmission and infection; and the selective pressure exerted by usage of antibiotics. To shift this steady state and reduce the overall burden of disease caused by *S aureus* new initiatives that prevent transmission and infection and reduce the selective pressure from using antibiotics will need to be introduced on a wide scale.

Anthony J Howard director, Public Health
Laboratory Service in Wales
University Hospital of Wales, Cardiff CF14 4XW

Mari Morgan senior scientist (epidemiology)
Communicable Disease Surveillance Centre Wales,
Cardiff CF14 3QX

David N Looker director
Rhyl Public Health Laboratory, Glan Clwyd District
General Hospital, Rhyl, Denbighshire LL18 5UJ

- 1 Crowcroft NE, Catchpole M. Mortality from methicillin resistant *Staphylococcus aureus* in England and Wales: analysis of death certificates. *BMJ* 2002;325:1390-1. (14 December)

- 2 Liggett AK, Swift B. Death certificates are an unreliable source of data. Electronic response to: Mortality from methicillin resistant *Staphylococcus aureus* in England and Wales. *bmj.com* 2002. bmj.com/cgi/eletters/325/7377/1390#27826 (accessed 12 Feb 2003).

Is death by disease or treatment?

EDITOR—The past practice of isolating patients with severe neutropenia arising from cancer chemotherapy was abandoned when studies showed that isolation did not prevent the spread of hospital infections to these immunologically suppressed patients. Isolation protocols increased mortality even in this predominantly younger population. It was hypothesised that the bother of donning gloves and gowns deterred hospital staff from entering the patient's isolation room, thereby depriving the patient of effective communication with staff, as well as limiting professional observation and physical examination.

This begs the question to what extent death is caused by methicillin resistant *Staphylococcus aureus* itself as opposed to the isolation treatment.

Isolation protocols not only include encumbrances on professional staff from physically accessing a patient but, more importantly, severely inhibit patients from leaving their hospital bedroom, limiting both their physical activity and appetite. Out of sheer boredom the patient spends most of the day horizontal in bed. The ravages of excessive bedrest were noted by Asher—not least, hypostatic pneumonia, hypercalcaemia, osteopenia, faecal impaction, bed sores, and depression.¹

Few studies have been done on the effectiveness of hospital isolation techniques in preventing the spread of antibiotic resistant organisms, and none are convincing. Elderly people who survive hospital acquired infection with methicillin resistant *S aureus* are often discharged in a weakened, deteriorated state to a nursing home, where their condition improves. Fortunately, most nursing homes do not isolate these people, and yet spread to other residents, who by definition are chronically ill or disabled, seems to be low.

As pointed out by Crowcroft and Catchpole,² 86% of those who died were elderly people. Isolation increases their invisibility. This is a double whammy for elderly people, who tolerate physical inactivity the least well and, at the same time, are the most likely to have their signs and symptoms not recognised by doctors. Perhaps in hospitals the wellbeing of older people takes a back seat to the threat of illness and death to younger people, hindering the investigation of the benefits and adverse effects from the intervention of isolation.

Janice E Lessard chief
Department of Geriatrics, St. Joseph's Health Centre,
30 The Queensway, Toronto, ON, Canada M6R 1B5
Candy@istar.ca

- 1 Asher R. The dangers of going to bed. *BMJ* 1947;iii:967-8.
 2 Crowcroft NS, Catchpole M. Mortality from methicillin resistant *Staphylococcus aureus* in England and Wales: analysis of death certificates. *BMJ* 2002;325:1390-1. (14 December.)

Managing acute renal colic

Intramuscular diclofenac should be avoided

EDITOR—The article by Wright et al on managing the pain of ureteric colic in the community has raised some concerns.¹ The paper implies that a strategy of administering intramuscular diclofenac in the community would reduce the need for admitting patients to emergency departments. Although we welcome any attempt to reduce the number of unnecessary admissions to emergency departments, we raise the following points about using intramuscular diclofenac.

Firstly, diclofenac causes pain on injections with a concomitant rise in serum phosphokinase activity. Cases of necrotising fasciitis have been recorded.²

Secondly, other non-steroidal anti-inflammatory drugs may be used, such as sublingual piroxicam or intramuscular ketorolac, that may prove as effective as diclofenac without damaging tissues.^{3,4}

Thirdly, cyclo-oxygenase 2 inhibitors may also prove effective in the treatment of renal colic and confer benefits in terms of their gastric protective effects. While clinical information is lacking in this area we believe that it is worthy of further analysis.

The defence unions have stated that intramuscular injections of diclofenac into the thigh are the least safe. The Committee on Safety for Medicines has received reports concerning this issue. None of the cases that were opened were defensible, and one of the recommendations suggested that alternative routes of administration be used where possible.⁵ Such practice must therefore be able to withstand scrutiny in the event of a complaint or litigation.

Our trust has taken the step of withdrawing intramuscular diclofenac from the trust formulary. We prefer to use rectal diclofenac or one of the cyclo-oxygenase 2 inhibitors.

A Hindle consultant in anaesthesia and acute pain management
 gasdochindle@aol.com

R Spedding consultant in accident and emergency medicine
 Warrington Hospital, Warrington Cheshire
 WA5 1QG

- 1 Wright PJ, English PJ, Hungin APS, Marsden SNE. Managing acute renal colic across the primary secondary interface: a pathway of care based on evidence and consensus. *BMJ* 2002;325:1408-12. (14 December.)
 2 Pillans PI, O'Connor N. Tissue necrosis and necrotising fasciitis after intramuscular administration of diclofenac. *Ann Pharmacother* 1995;29:264-6.
 3 Supervia A, Pedro-Botet J, Nogues X, Echarte JL, Minguez S, Iglesia ML, et al. Piroxicam fast dissolving form versus diclofenac in the treatment of acute renal colic: a double blind controlled trial. *Br J Urol* 1998;81:27-30.
 4 Stein A, Ben Dov D, Finkel B, Mecz Y, Kitzes R, Lurie A. Single dose intramuscular Ketorolac versus Diclofenac for pain management in renal colic. *Am J Emerg Med* 1996;14:385-7.
 5 Hegan T. Non steroidal, needles and negligence. *Medical Protection Society casebook*. Leeds: Medical Protection Society, 2002:12-13.

Perpetuating a urological myth

EDITOR—Wright et al include maintaining a higher than normal fluid intake among their recommendations for the management of acute renal colic.¹ Such advice is often given to patients with renal stone disease by medical practitioners.

However, this recommendation is based on expert consensus rather than any sound evidence. Foster et al previously pointed out that increasing fluid intake can produce nausea and vomiting. Moreover, it may cause a diuresis, which in the presence of an acute urinary obstruction can lead to increased pain, extravasation of urine, and formation of a urinoma, while having no effect on the passage of the stone.²

Encouraging patients with renal calculi to drink more fluid is of no proved benefit and may be detrimental. We question the value of this piece of advice in the otherwise useful protocol.

James P B O'Connor senior house officer
 jpboc@breathe.com

Philip M T Weston consultant
 Department of Urology, Pinderfields General,
 Wakefield WF1 4DG

- 1 Wright PJ, English PJ, Hungin APS, Marsden SNE. Managing acute renal colic across the primary-secondary care interface: a pathway of care based on evidence and consensus. *BMJ* 2002;325:1408-12. (14 December.)
 2 Foster MC, Upsdell SM, O'Reilly PH. Urological myths. *BMJ* 1990;301:1421-3.

Establish reason why purulent rhinitis is persistent first

EDITOR—The editorial by Arroll and Kenealy was a helpful and interesting summary of the role of systemic antibiotic treatment in managing acute purulent rhinitis.¹ The authors define the condition as persisting for up to 10-14 days before treatment may be indicated. Starting treatment at this time is, however, to be discouraged without a diagnosis. This is particularly true when the rhinitis is unilateral or bloodstained.

Persisting unilateral purulent rhinitis in a child may be caused by a nasal foreign body requiring removal.² If the discharge is bilateral and recurrent over a long time the resulting adenoiditis and nasal obstruction may warrant consideration of adenoidectomy.

In adults neoplasia is a diagnosis of exclusion if purulent rhinitis persists, particularly if it is unilateral and bloodstained.³

The article does not offer important axiomatic information to non-specialists, who may be asked to offer an opinion on such a symptom fairly often. If purulent rhinitis persists for more than two weeks, the question must be, what is the diagnosis—not, are antibiotics appropriate?

Peter J Robb honorary secretary, British Association for Paediatric Otorhinolaryngology
 Ear, Nose, and Throat Department, Ashted Hospital, Ashted, Surrey KT21 2SB

- 1 Arroll B, Kenealy T. Antibiotics for acute purulent rhinitis. *BMJ* 2002; 325:1311-12. (7 December.)
 2 Ransome J. Foreign bodies in the nose. In: Kerr AG, ed. *Scott-Brown's Otolaryngology*. Butterworths: London, 1987:276-9.

- 3 Harrison DFN. Differential diagnosis of skull base neoplasms involving the anterior and middle cranial fossa. In: Cummings CW, Fredrickson JM, Harker LA, Krause CJ, Schuller DE, eds. *Otolaryngology head and neck surgery*. St Louis: Mosby, 1986:3389-401.

G7 countries are not to blame for Africa's brain drain

EDITOR—Ferriman's news item describes the three initiatives taken at the recent meeting of the Global Forum for Health Research in Tanzania, designed to change the face of health research and health care in Africa and other developing countries.¹ But I cannot see the point in criticising the G7 countries on the grounds that they are saving money on training costs because health professionals decide to leave the failed countries in the developing world to settle in G7 countries.

In 1996 the Nigerian Medical Forum, in its presentation to the Vision 2010 Health Subcommittee of the federal government of Nigeria, pointed out the reasons why professionals of all types leave the country. Similar reasons may apply to other African countries—namely, the failure of governments in developing countries to create the enabling environment and provide the tools needed for their professionals to practise their expertise. It is not simply a matter of salaries because when countries such as Nigeria showed an interest in the progress of the professions, all its professionals who trained abroad returned home as soon as they collected their certificates in any G7 country. If the developing countries have come to notice their loss through the brain drain of their professionals, I suggest they begin to put their resources and investment in the professions in their countries.

Unfortunately, it may be too late to entice back those who have left and settled in G7 countries, but at least they may dissuade those at home from leaving. The good news, as a survey conducted by the Nigerian Medical Forum in 1997-8 showed, is that many of these experts from developing countries are prepared to give their time and expertise to their home country, and at no charge, if only their safety can be guaranteed and they can deliver care to their people.

Joseph Ana volunteer trustee-director, Nigerian Medical Forum
 65 Warden Hill Road, Luton LU2 7AE
 nmf.nmf@virgin.net

Competing interests: JA is a pioneer volunteer trustee-director of the Nigerian Medical Forum, a registered UK charity since 1991.

- 1 Ferriman A. Where there is despair, hope. *BMJ* 2002;325:1194. (23 November.)

bmj.com

Letters appearing here are an edited selection of rapid responses originally posted on bmj.com

We ask for all letters to the editor to be submitted as rapid responses via bmj.com

For advice see: bmj.com/rapidresponses