from debt relief programmes directly to schools to increase enrolment and minimise transaction costs. Moreover, it would be cost effective to direct funds from the newly established Global Fund to Fight AIDS, Tuberculosis, and Malaria to maternal and infant healthcare programmes and to education. Young children without both parents should be given the highest priority. If children are cared for by their adolescent siblings, then such adolescents should be the first to qualify for vocational training. In the absence of a cure for AIDS, international agencies have an important role in providing technical advice as well as funds for maintaining the current levels of economic activity in sub-Saharan African countries.

Funding: The authors' field trip was supported by the World Bank. The opinions expressed here are those of the authors and should not be attributed to any organisation.

Competing interests: None declared.

- 1 Buve A, Caraël M, Hayes RJ, Auvert B, Ferry B, Robinson NJ, et al. Multi-centre study on factors determining differences in rate of spread of HIV in sub-Saharan Africa: methods, and prevalence of HIV infection. *AIDS* 2001;15:55-14.
- 2 Falkner F, Holzgreve W, Schloo TH. Parental influences on postnatal growth: overview and pointers for research. *Eur J Clin Nutr* 1994;48:15-24.
- 3 Bhargava A. Modelling the effects of maternal nutritional status and socioeconomic variables on the anthropometric and psychological indicators of Kenyan infants from age 0-6 months. *Am J Phys Anthropol* 2000;110:89-104.
- 4 Bhargava, A. Family planning, gender differences and infant mortality: evidence from Uttar Pradesh, India. *J Econometrics* 2003;112:225-40.
- 5 Spira R, Lepage P, Msellati P, Van de Perre P, Leroy V, Simonon A, et al. Natural history of human immunodeficiency virus type 1 infection in children: a five-year prospective study in Rwanda. *Pediatrics* 1999;104:1-9.
- 6 Wiktor S, Ekpin E, Karon J, Nkengasong J, Maurice C, Severin S, et al. Short course oral zidovudine for prevention of mother-to-child transmission of HIV-1 in Abidjan, Cote d'Ivoire: a randomised trial. *Lancet* 1999;353:781-5.
- 7 Demographic and Health Survey. *Demographic and health survey, Ethiopia 2000*. Washington, DC: Macrointernational, 2000.
- 8 World Bank. *World development indicators*. Washington, DC: World Bank, 2001.
- 9 Meekers D. Going underground and going after women: Trends in sexual risk behavior among gold miners in South Africa. *Int J STD AIDS* 2000;11:21-6.
- 10 Demographic and Health Survey. *Demographic and health survey, Malawi 2000*. Washington, DC: Macrointernational, 2000.
- 11 Demographic and Health Survey. *Demographic and health survey, Tanzania 1999*. Washington, DC: Macrointernational, 1999.
- 12 Deininger K. Does cost of schooling affect enrolment by the poor? *Universal primary education in Uganda*. Washington, DC: World Bank, 2001.
- 13 Ter-Minassian T, Gupta S, Clements B, Guin-Siu MT, de Mello L. *Debt relief in HIPC: fiscal policy issues*. Washington, DC: International Monetary Fund: 2001.
- 14 Kinsman J, Nakiyingi J, Kamali A, Carpenter L, Quigley M, Pool R, et al. Evaluation of a comprehensive school-based AIDS education programme in rural Mawaka, Uganda. *Health Educ Res* 2001;16:85-100.
- 15 Ndubani P, Hojer B. Sexual behavior and sexually transmitted diseases among young men in Zambia. *Heal Pol Plan* 2001;16:107-12.
- 16 Nzoika C. Perspectives of adolescent boys on the risks of unwanted pregnancy and sexually transmitted infections: Kenya. *Reprod Health Matters* 2001;9:108-17.
- 17 Machel JZ. Unsafe sexual behavior among school girls in Mozambique: a matter of gender and class. *Reprod Health Matters* 2001;9:82-90.
- 18 Bhargava A, Jamison D, Lau L, Murray C. Modeling the effects of health on economic growth. *J Health Econ* 2001;20:423-40.
- 19 Marchat JM, Nasir J, Ramchandran V, Shah MK, Tyler G, Zhao L. *Results of the Nigeria firm survey*. Washington, DC: World Bank, 2001.
- 20 World Bank Madagascar. *Education and training in Madagascar*. Washington, DC: World Bank, 2001.

(Accepted 17 March 2003)

Tackling India's HIV epidemic: lessons from Africa

Malcolm Potts, Julia Walsh

India stands on the brink of a major HIV epidemic. However, by examining where public health initiatives went wrong in Africa, the international community may be able to help India avoid the devastating effects seen in Africa

The rapid spread of HIV in sub-Saharan Africa is one of the greatest failures in the history of public health. Given our detailed understanding of HIV and the natural course of AIDS, the virus should have been controllable. Yet in some African countries 20% of people aged over 15 are HIV positive and 70% of them will eventually die from AIDS.¹ India shares some of the same risk factors as Africa, including a similar pattern of health expenditure, an uneven health infrastructure, and prevalent high risk sexual behaviours (table A, bmj.com).² By 2010 the number of HIV infections in India is predicted to rise from 4 million to 20-25 million¹⁻³ We discuss 10 important lessons from Africa that could limit the spread of HIV in India.

Methods

The views expressed in this paper are based on current literature reviews, economic analyses of the Berkeley International Group (<http://big.berkeley.edu>), and extensive personal experience working on HIV, reproductive health, programme management, and international finance.

Involve high risk groups in all phases of programmes

HIV infection begins in the core groups of commercial sex workers, intravenous drug users, and men who have sex with men (table B, bmj.com). Mathematical models show both the benefits of early intervention and the

Number of cases of HIV infection that could be averted with \$2m AIDS budget spent on preventive measures or antiretroviral drugs (ARV) from different sources⁵

See also editorial by Ammann

Bay Area International Group, School of Public Health, University of California, Berkeley, CA 94720-7360, USA

Malcolm Potts
Bixby professor,
population and family
planning

continued over

BMJ 2003;326:1389-92

Further tables and figures giving HIV statistics are available on bmj.com

Institute for Human Development,
University of California, Berkeley

Julia Walsh
*adjunct professor,
international health
and maternal and
child health*

Correspondence to:
M Potts
pottsmalcolm@yahoo.com

importance of focusing on these core groups.⁴⁻⁷ India remains a traditional society: its laws forbidding homosexuality derive from British legislation in the 19th century,⁸ and the sex industry relies on bribing the police to operate. The high risk groups are outside the political, social, and economic mainstream. Leaders must overcome prejudice and be prepared to include these groups in designing well funded programmes.

Focus on cost effective preventive programmes

Poverty is a major contributor to the spread of HIV and thus considered an important focus for preventive programmes. However, the prevalence of HIV can double within high risk groups in six months and in the general population in three to five years, while socio-economic change takes decades to achieve. Moreover, poverty does not always drive the spread of HIV. Botswana is relatively wealthy and over 90% of women are educated, but it has a high prevalence of HIV. As long as the AIDS community highlights poverty, it allows senior political leaders to claim that only socio-economic change will cure the disease and to avoid controversial interventions, such as distributing condoms to unmarried people.

The most appropriate immediate strategy for India is a targeted approach in core groups. Its cost effectiveness is well established.⁵⁻⁹⁻¹⁰ Substantial numbers of cases can be averted by management of sexually transmitted diseases in commercial sex workers, blood screening, voluntary counselling and testing, treatment of sexually transmitted diseases in the general population, and providing antiretroviral drugs at childbirth (figure, table).⁵ The cost of treating one person with antiretroviral drugs for a year (at full price) is equivalent to that of preventing almost 50 cases (figure).⁵

Secure adequate supplies of condoms and antibiotics

In sub-Saharan Africa, too little attention was given to the supply of condoms. In 2000, fewer than 650 million condoms were distributed (about four for each adult man).¹¹⁻¹² The United Nations Population Fund estimates that at least \$1bn (£626m, €881m) is needed for condoms to control AIDS, and the gap between demand and supplies is growing.¹³ If we assume that people will spend up to 1% of their income on buying condoms, 95% of India's population cannot afford the \$10.65 a year needed for manufacture, promotion, and distribution of condoms.¹⁴

India has the advantage of a large, competitive, technically competent pharmaceutical industry, and the gov-

ernment is subsidising distribution of condoms to low income families. This subsidy should be extended to antibiotics for controlling sexually transmitted diseases. The international donor community should ensure that priority is given to funding condoms and antibiotics before funding other programmes.

Invest only in projects that can be fully implemented

Africa is awash with pilot projects, yet no donor agency or national government has set up a systematic programme of basic interventions for all sex workers in every large city. Since most possible interventions have already been tested more than once, the need for additional small scale studies is questionable. Larger scale, relatively simple programmes that provide basic services to all should be given priority over sophisticated, labour intensive programmes covering a few people. The available resources should be allocated equally across all the sites that need intervention. The sophistication of programmes should be determined by the amount of money available for each site. Although the amount might be relatively small, the programmes could still have a big impact.

Pilot studies of interventions should not be funded if the resources are not available to implement the intervention nationwide, however promising it may seem in theory. The worst case scenario would be one where India's many non-governmental organisations divide the limited resources to create carefully crafted, custom built programmes and end up having no overall impact on the disease.

Include traditional health practitioners in control programmes

In areas where the epidemic has spread beyond the initial high risk groups, programmes providing interventions for the general population are essential.¹⁵ Rural medical practitioners, nurses, Western trained doctors, and other health professionals will have to be included. Many African countries have resisted the non-medical prescription of antibiotics. In India, rural medical practitioners often have a formal training in Ayurvedic or other traditional medicine but are not formally permitted to prescribe antibiotics. It is essential to recognise the role of India's private sector, which provides care for most rural poor people (see bmj.com), and realign prescription regulations to reflect reality.

Reconsider the structure and work of international donors

The Global Fund for AIDS, Tuberculosis, and Malaria now controls most of the funding for tackling HIV and AIDS. However, the \$2-3bn available amounts to less than one quarter of the annual projected needs for controlling AIDS.¹⁶

Underspending of donated funds is common and helps curtail the resources available and deter future allocations from the donor. Implementing agencies spend a great deal of time preparing proposals; donors require time to analyse unsolicited proposals, to prepare, and to review requests for proposals. A more cost effec-

Cost per life year saved by HIV prevention programmes

Programme	Cost/life year saved (\$)
Blood screening	3.35
Management of STDs for sex workers	3.95
Voluntary counselling and testing	22.03
Treatment of STDs for general population	22.32
Prevention of mother to child transmission:	
Zidovudine	213.66
Nevirapine	11.24

STD=sexually transmitted disease.

tive strategy would be for governmental and foundation donors to set achievable output goals, specify how much money they wish to allot to this particular area, and then invite evidenced based proposals from possible implementing agencies. This would draw on the rich experience of agencies in both developing and developed countries, simplify the work of donors, and pre-empt the second guessing of the donor's goals. Support for output based services should ensure that money follows results and increase the efficiency of both non-governmental and governmental programmes.

Large international meetings waste resources

Large meetings cost a great deal in airfares and living and opportunity costs. The communication of ideas and initiation of new collaborations at such meetings has fallen over the years, and the meetings have become platforms for non-evidence based lobbying.¹⁰ Meetings that deal with specific topics and focus on science are likely to remain useful.

Confront lobbying for increased use of antiretroviral drugs

Infected people in rich countries who benefit from antiretroviral drugs form a compelling lobby for extending treatment elsewhere. However, individuals at risk of infection do not lobby for investment in prevention of HIV and AIDS. When prevention programmes succeed, it is impossible to say that any one person is alive because of them.¹⁰ Any objective effort to allocate the limited resources available to confront the HIV epidemic in India will have to take these asymmetries in lobbying into account.

One emerging lesson is that money spent on antiretroviral drugs is money removed from prevention, and vice versa. The Global Fund for AIDS, Tuberculosis, and Malaria has allocated 60% of its first \$378m of grants to support HIV projects, and 21 of 28 countries receiving grants will use this money to purchase antiretroviral drugs.¹⁷ These drugs are difficult to use (except to prevent mother to child transmission) and, even at the greatly reduced prices, are very expensive, especially when the necessary testing, monitoring, and counselling costs are included (figure). In extreme cases they may even encourage increased rates of unsafe sexual behaviour.¹⁸

Even if there are further massive reductions in drug prices, the costs will remain beyond the reach of individual and community subsidy in much of India. Nevertheless, there will be intense emotional pressure to mount token subsidised antiretroviral programmes. The people who benefit will probably be those who are most educated or have access to specialist care. The example of Thailand, where an increase in antiretroviral drug costs took money away from prevention budgets and was associated with an increase in HIV infections, must shore up our commitment to prevention.¹⁹

Ensure policies are based on latest evidence

In hindsight, the investment Africa made in safeguarding the blood supply rather than focusing on high risk

Commercial sex workers in India

groups was probably over enthusiastic. In the 1980s, the scientific evidence that other sexually transmitted diseases facilitate the transmission of HIV was not acted on quickly enough. Evidence is coming out of Uganda that sexual abstinence and reduction in the number of sexual partners can help reduce prevalence of HIV.²⁰ Religions such as Islam, Christianity, and Hinduism emphasise certain aspects of sexual abstinence and reproductive health. Religious organisations could therefore be used to help prevention alongside programmes to distribute condoms and treat sexually transmitted disease.

Attention needs to be given to the increasingly strong evidence that male circumcision slows transmission of HIV. In a recent study in Uganda, 30% of uncircumcised men became infected from their HIV positive female partners compared with none of the circumcised men.²¹ Finding ways to offer circumcision to Hindu men (who are generally not circumcised) could slow transmission of HIV and other sexually transmitted diseases.^{21 22}

Despite its devastating effect, HIV is a fragile, difficult to transmit, and easily destroyed virus. A microbicide that women can use secretly and effectively to protect themselves against HIV infection would be valuable.²³ Western managers estimate that development of a 60% effective microbicide would cost \$775m. Although this is achievable, the microbicide would not reach the market before 2007 at the earliest, which will be too late to contain the spread of HIV in India. A concerted effort should be made to use these resources in ethical research frameworks that fit the enormous risk of death in developing countries. The suggestion that lemon juice may be an effective microbicide should also be investigated.²⁴ Decisions about which programmes to implement should be based on cost effectiveness. Comparable and rigorously collected data are needed on implementation costs and on consumers' willingness to pay for prevention and therapy.

Increase national and global budgets now

Government investment in AIDS prevention in all countries has been a story of too little too late. Large investments at the outset of the epidemic will slow progress more than those at a later stage. India

Summary points

The number of cases of AIDS in India will probably exceed 20 million by 2010.

The limited resources should be used for large cost effective programmes to decrease spread of the disease

Money spent now will be much more effective than money spent later in the epidemic

Adequate supplies of condoms and antibiotics must be secured

Prevention should be given priority over antiretroviral treatment

currently invests R300 crore (£43.5m) annually, much less than needed.¹⁶ This simple lesson in epidemiology needs to be understood by external donors, the Indian government, and foundations. The Macroeconomic Commission on Health urges a large increase in donor funds to confront AIDS and other important diseases. Such increases are both justified and achievable, given the size of the world economy.²⁵

We thank Mike Musante, Russell Green, Stephanie Heise, Elliot Marseille, and Ndola Prata for their contributions to analysis, editing, and graphical presentation.

Contributors: MP conceived the analysis and led the writing of the paper. JAW led the overall data collection and helped write the paper. MP is the guarantor.

Funding: Fred H Bixby Endowment, and grants from the Gates and Hewlett Foundations.

Competing interests: None declared.

- 1 UNAIDS. Report on the global HIV/AIDS epidemic. Geneva: UNAIDS, 2000/2002.
- 2 Hawkes S, Santhya KG. Diverse realities: sexually transmitted infections and HIV in India. *Sex Transm Infect* 2002;78(suppl1):i31-9.
- 3 National Intelligence Council (CIA). *The next wave of HIV/AIDS: Nigeria, Ethiopia, Russia, India, and China*. Washington, DC: CIA, 2002. (ICA 2002-04 D). www.cia.gov/nic/pubs/index.htm (accessed 10 March 2003).
- 4 Boily MC, Lowndes C, Alary M. The impact of HIV epidemic phases on the effectiveness of core group interventions: insights from mathematical models. *Sex Transm Infect* 2002;78(suppl 1):i78-90.

- 5 Masaki E, Green R, Greig F, Walsh J, Potts M. Cost effectiveness in HIV prevention versus treatment for resource scarce countries: setting priorities for HIV/AIDS management. <http://big.berkeley.edu/research.workingpapers.htm> (accessed 10 March 2003).
- 6 Lowndes CM, Alary M, Meda H, Gnintoungbe CA, Mukenge-Tshibaka L, Adjovi C, et al. Role of core and bridging groups in the transmission dynamics of HIV and STIs in Cotonou, Benin, West Africa. *Sex Transm Infect* 2002;78(suppl 1):i69-77.
- 7 Stover J, Walker N, Garnett GP, Salomon JA, Stanecki KA, Ghys PD, et al. Can we reverse the HIV/AIDS pandemic with an expanded response? *Lancet* 2002;360:73-7.
- 8 International Lesbian and Gay Association. Situation of homosexuals in India: report for the Swedish Embassy by a Delhi law firm. September 9, 1997. www.ilga.org/Information/asia_pacific/situation_of_homosexuals_in_india.htm (accessed 6 May 2003).
- 9 Creese A, Floyd K, Alban A, Guinness L. Cost-effectiveness of HIV/AIDS interventions in Africa: a systematic review of the evidence. *Lancet* 2002;359:1635-47.
- 10 Marseille E, Hofmann PB, Kahn JG. HIV prevention before HAART in sub-Saharan Africa. *Lancet* 2002;359:1851-6.
- 11 Shelton JD, Johnston B. Condom gap in Africa: evidence from donor agencies and key informants. *BMJ* 2001;323:139.
- 12 United Nations Population Fund. Securing essential supplies. www.unfpa.org/supplies/index.htm (accessed 12 May 2003).
- 13 McNeil D. Global war against AIDS runs short of vital weapon: donated condoms. *New York Times* 2002 October 9.
- 14 Green R. Empty pockets: estimating ability to pay for family planning. International Health Economics Association third international conference, July 22 - 25, 2001, University of York. <http://big.berkeley.edu/research.workingpapers.htm> (accessed 10 March 2003).
- 15 Nagelkerke N, Jha P, de Vlas S, Korenromp EL, Moses S, Blanchard JF, et al. Modelling HIV/AIDS epidemics in Botswana and India: impact of interventions to prevent transmission. *Bull World Health Organ* 2002;80:89-96.
- 16 Jha P, Mills A, Hanson K, Kumaranayake L, Conteh L, Kurowski C, et al. Improving the health of the global poor. *Science* 2002;295:2036-9.
- 17 The Global Fund to Fight AIDS, Tuberculosis and Malaria. Global Fund announces first grants [press release, 26 April 2002]. www.globalfundatm.org/journalists/press%20releases/NY_250402.html (accessed 6 May 2003).
- 18 Chen SY, Gibson S, Katz MH, Klausner JD, Dilley JW, Schwarcz SK, et al. Continuing increases in sexual risk behavior and sexually transmitted disease among men who have sex with men: San Francisco, California 1999-2001. *Am J Pub Health* 2002;92:1387.
- 19 De Sam Lazaro F. AIDS challenge in Thailand. Segment on NewsHour with Jim Lehrer. www.pbs.org/newshour/bb/asia/july-dec02/aids_10-22.html (accessed 10 March 2003).
- 20 Parkhurst JO. The Ugandan success story? Evidence and claims of HIV-1 prevention. *Lancet* 2002;360:78-80.
- 21 Szabo R, Short RV. How does male circumcision protect against HIV infection? *BMJ* 2000;320:1592-4.
- 22 Gray RH, Kiwanuka N, Quinn TC, Sewankambo NK, Serwadda D, Mangan FW, et al. Male circumcision and HIV acquisition and transmission: cohort studies in Rakai, Uganda. Rakai Project Team. *AIDS* 2000;14:2371-81.
- 23 Microbicide Initiative. *Mobilization for microbicides: the decisive decade*. New York: Rockefeller Foundation, 2002.
- 24 LemonAIDS. www.aids.org.au/ (accessed 12 May 2003).
- 25 Attaran A, Sachs J. Defining and redefining international donor support for combating the AIDS pandemic. *Lancet* 2000;357:57-61.

(Accepted 10 April 2003)

Submitting articles to the *BMJ*

We are now inviting all authors who want to submit a paper to the *BMJ* to do so via the web (<http://submit.bmj.com>).

We have introduced Benchpress, our new web based manuscript tracking system, with the aim of streamlining our processes and providing better, quicker information for authors, reviewers, and editors.

Benchpress is a website where authors deposit their manuscripts and editors go to read them and record their decisions. Reviewers' details are also held on the system, and when asked to review a paper reviewers will be invited to access the site to see the relevant paper. The system is secure, protected by passwords, so that authors see only their own papers and reviewers see only those they are meant to. The system is run by Highwire Press, who host bmj.com, and is already being used by 30 journals, including most of the BMJ Publishing Group's specialist journals.

For authors in particular the system offers several benefits. The system provides all our guidance and forms and allows authors to

suggest reviewers for their paper—something we'd like to encourage. Authors get an immediate acknowledgement that their submission has been received, and they can watch the progress of their manuscript. The record of their submission, including editors' and reviewers' reports, remains on the system for future reference.

Anyone with an internet connection and a web browser can use the system.

As with all new systems we expect a few teething problems, but the system itself offers extensive help, and the *BMJ*'s editorial office is geared up to help authors and reviewers if they get stuck. We see Benchpress as part of our endeavour to improve our service to authors and reviewers and, as always, we'd welcome feedback.

Benchpress is accessed via <http://submit.bmj.com> or via a link from bmj.com