Clinical review

Extracts from “BestTreatments”

Bulimia nervosa

Regina Z Lilly

BMJ Unified,
London WC1H 9JR
Regina Z Lilly
medical writer

rzlilly@netscape.net

BMJ 2003;327:380-3

Description

Bulimia nervosa is an eating disorder that affects women 9-10 times more often than men. It has three key features:

- Intense preoccupation with body weight and shape
- Repetitive episodes of binge eating (uncontrollably eating a large quantity of food in a defined period—usually less than two hours)
- Routinely taking extreme measures to prevent weight gain, including self induced vomiting, fasting, excessive exercise, and misuse of laxatives and diuretics.

The condition is characterised by the patient's inability to control the urge to indulge in binge eating. Medical complications are caused by the weight reducing behaviours, particularly vomiting and laxative misuse. The frequency of these behaviours is, on average, at least twice a week for three months. Episodes may be spontaneous or planned and can be triggered by stress, intense hunger, or dysphoria. Patients are usually in the normal weight range.

Treatment

Despite the availability of effective treatments, few people with bulimia nervosa seek help. One study found that only 1 in 10 people with the condition are in treatment.¹

Treatments that are likely to work

Antidepressants

Antidepressants can reduce binge eating and purging, and improve depressive symptoms and attitude towards food. There is not enough evidence on how long the benefits last or for comparing the effects of different classes of antidepressants. Antidepressants and cognitive behaviour therapy are equally effective for bulimia nervosa, though dropout rates are higher with antidepressants, probably due to their side effects. For information on drug doses, see www.besttreatments.org/bulimianervosa.

Cognitive behaviour therapy

As with antidepressants, cognitive behaviour therapy can reduce binge eating and purging, and improve depressive symptoms and attitude towards food. It may be preferable to medication in patients who are sensitive to, or wish to avoid, side effects of medication. Cognitive behaviour therapy and other psychological treatments seem to be equally good at treating binge eating, although cognitive behaviour therapy is better at improving depression scores. There is some evidence that cognitive behaviour therapy has long term benefits in treating bulimia, but there is insufficient evidence about patients' characteristics that predict response.

Combination treatment with antidepressants and psychotherapy

Remission rates of some (but not all) symptoms are higher with combination treatment than with either treatment alone. Patients are more likely to withdraw from combined treatment or antidepressants alone than from psychotherapy alone. Some trials of combination treatment with antidepressants and psychotherapy have excluded patients with severe comorbid conditions such as depression and substance abuse, yet these conditions are common in patients seen in a clinical setting.

Other psychotherapies

Other psychotherapies, such as interpersonal psychotherapy and dialectical behaviour therapy (see definitions on www.besttreatments.org/bulimianervosa), are better than control (being on a waiting list for treatment) at improving binge eating and other symptoms of bulimia. The few trials of these psychotherapies have been hampered by small sample sizes and lack of blinding. It remains unclear which of these many techniques works best.

Treatments that need further study

Antidepressants as maintenance

There is not enough evidence on whether antidepressants have a role as maintenance therapy. Two very small randomised controlled trials (total of 18 patients) found that maintenance treatment with a tricyclic antidepressant was no better than placebo.

This extract is from www.besttreatments.org, a shared information resource for US patients and doctors that is based on Clinical Evidence (www.clinicalevidence.com). More information, including additional references for this BMJ article, is at www.besttreatments.org/bulimianervosa

Newer antidepressants

There have been no randomised controlled trials of the newer antidepressants mirtazapine, reboxetine and venlafaxine in treating bulimia nervosa.

Selective serotonin reuptake inhibitors other than fluoxetine

The only selective serotonin reuptake inhibitor shown to work in bulimia nervosa is fluoxetine.

This extract will remain available on bmj.com. The rest of the BestTreatments website is currently available only to US doctors and patients who are members of United Healthcare Group health plans in the United States.

Competing interests: None declared.

- 1 Welch SL, Fairburn CG. Sexual abuse and bulimia nervosa: three integrated case control comparisons. *Am J Psychiatry* 1994;151:402-7.

Commentary: Putting the research into practice

Phillipa Hay

Bulimia nervosa occurs in about 1% of young women in Western countries. It is often chronic and adversely affects quality of life.¹ The rate of diagnosis is very low, and community and primary care studies have shown that treatment is often delayed.^{2,3} An important contributing factor to this delay is the secrecy and shame around binge eating, self induced vomiting, and laxative misuse.⁴

Bulimia nervosa should be suspected in young and middle aged women who present with weight and shape concerns (whether they are normal or overweight)³ or with common comorbid conditions such as depression, anxiety, or substance use disorders.⁵ People who are underweight may have bulimic behaviours, but in current classificatory schemes the diagnosis of anorexia nervosa takes precedence in such patients.⁴ People with bulimia may present with physical effects of the illness, such as dental caries, menstrual irregularities, and unexplained hypokalaemia.⁵ When asked directly about bulimic symptoms in a treatment setting, patients are usually relieved to have the opportunity to talk openly about them.

Selecting treatments

Cognitive behaviour therapy (following the comprehensive treatment manual of Fairburn and colleagues) is the treatment of choice.⁶ Because of difficulty in accessing specialist therapists, self help modes of delivering cognitive behaviour therapy have been developed. These involve patients following a treatment manual themselves, usually out of a book (such as Cooper's *Bulimia Nervosa and Binge Eating: A Guide to Recovery*⁷). Such pure self help approaches are helpful, but the effects tend to be more modest than cognitive behaviour therapy guided by a therapist.⁸

Antidepressants are also helpful in the short term, but there is insufficient evidence for their role as maintenance therapy.⁸ The best evidence is for high dose fluoxetine.⁹ A useful strategy is to start with a low dose and then increase it weekly up to 60 mg daily. Fluoxetine may be associated with nausea and weight loss, and these side effects need to be monitored particularly carefully in people with bulimia.

A combination of antidepressants and psychotherapy is more effective than either treatment alone.⁸ This approach may be helpful for people with very

severe symptoms, or those who have an incomplete response to treatment.

Pitfalls in using the evidence

The evidence on treatment has several gaps. There have been too few follow up studies of antidepressants and only a few trials have compared cognitive behaviour therapy with other psychotherapies. Rates of remission from binge eating in outcome assessments have been about 33-43%, pointing to scope for improvement in current treatments. The best approach to managing comorbid obesity is unclear, as restrictive dieting can exacerbate binge eating.

When to refer

Many patients with bulimia nervosa can be treated effectively in primary care, but specialist referral should be considered for any of the following:

- Failure to respond to adequate trials of treatment
- Significant comorbid medical or psychiatric illness or other complications (such as pregnancy)
- Risk of self harm
- Need for clarification of diagnosis or advice on treatment.⁵

Competing interests: PH has received reimbursement for attending symposiums from Solvay, Bristol-Myers Squibb, Janssen-Cilag, and Pfizer and fees for organising education from Bristol-Myers Squibb, Pfizer, and Lundbeck.

- 1 Hay PJ. Quality of life and bulimic eating disorder behaviours: findings from a community-based sample. *Int J Eat Disord* 2003;33:434-42.
- 2 Fairburn CG, Welch SL, Norman PA, O'Connor ME, Doll HA. Bias and bulimia nervosa: how typical are clinic cases? *Am J Psychiatry* 1996;153:386-91.
- 3 Hay PJ, Marley J, Lemar S. Covert eating disorders: the prevalence, characteristics and help-seeking of those with bulimic eating disorders in general practice. *Prim Care Psychiatry* 1998;4:95-9.
- 4 American Psychiatric Association. *Diagnostic and statistical manual of mental disorders*. 4th ed. Washington, DC: American Psychiatric Association, 1994.
- 5 American Psychiatric Association. Practice guidelines for the treatment of patients with eating disorders (revision). *Am J Psychiatry* 2000;157(suppl 1):1-39.
- 6 Fairburn CG, Harrison PJ. Eating disorders. *Lancet* 2003;361:407-16.
- 7 Cooper PJ. *Bulimia nervosa and binge eating: a guide to recovery*. London: Robinson, 1995.
- 8 Hay PJ, Bacaltchuk J. Bulimia nervosa. In: *Clinical Evidence*. Issue 9. London: BMJ Publishing, 2003:1388-98.
- 9 Fluoxetine Bulimia Nervosa Collaborative Study Group (FBNCSSG). Fluoxetine in the treatment of bulimia nervosa. A multi-center, placebo controlled trial. *Arch Gen Psychiatry* 1992;49:139-47.

Department of Psychiatry,
University of Adelaide, Royal Adelaide Hospital,
Adelaide SA 5001, Australia
Phillipa Hay
senior lecturer
phillipa.hay@adelaide.edu.au

Commentary: Information for patients: What happens during cognitive behaviour therapy for bulimia

Alison Tonks

BMJ Unified,
London WC1H 9JR
Alison Tonks
freelance medical
editor
atonks@bmj.com

Cognitive behaviour therapy is a kind of psychotherapy. Sometimes it's just called therapy. During cognitive behaviour therapy you talk to a trained therapist about your problems. Most people with bulimia see a therapist regularly for about 20 weeks, but you can see a therapist for longer if you need to. Each meeting with the therapist usually lasts about an hour. Your therapist could be a psychologist, a psychiatrist, a psychiatric nurse, a psychotherapist, or even a family doctor.

Three phases of therapy

Cognitive behaviour therapy for bulimia is often split into three phases.

Phase one

You start by learning to reduce bingeing and dieting. This takes about seven to nine weeks. Usually, you keep a diary of how often you eat and how often you binge, purge, or exercise. You might also write down your thoughts about food in your diary.

The first phase is all about allowing yourself to eat. For example, if you never eat breakfast or lunch, you might agree to eat a small breakfast every day (yogurt and an apple, say) as part of your treatment.

You also learn basic facts about bulimia and treatment, and how these affect your body. Here's one important fact: most people do not put on weight during cognitive behaviour therapy.

Phase two

You start talking more about what you eat, not when. You slowly branch out so that you're eating a lot of different foods, even ones you've craved or thought were banned, like doughnuts. When foods are not forbidden in your mind, you'll probably find that you don't crave them.

In this phase, you begin talking about your feelings, including feelings about your weight and eating. You

may feel desperate to be thin, or you may think that losing weight will make you happy. You and your therapist might explore why you feel this way. Your therapist will help you find healthier ways to feel good about yourself. If you binge on food because you feel stressed, your therapist will help you deal with that too.

Phase three

By phase three, you are likely to be feeling better and bingeing less. During the last few weeks you learn how to keep it that way. You also learn what to do if the symptoms of bulimia come back.

What's involved in the treatment

Therapists use many different techniques to help people change their thinking and behaviour. Your therapist will teach you some of them. And you'll have homework to do between sessions. Writing in your food diary might be one kind of homework. Eating a small meal that you would normally skip could be another. Homework is an important part of cognitive behaviour therapy.

Treatment doesn't end when you stop going to a therapist. You'll be able to use the techniques you've learned by yourself. You can be your own therapist whenever you feel bulimia taking over again. So the treatment may be especially good for people who want to help themselves get better.

There's even a kind of "do it yourself" or self help cognitive behaviour therapy. Your therapist gives you a book with information and homework exercises in it. You work through the book on your own for a few weeks, but go to the therapist every now and then for advice and encouragement.

Competing interests: None declared.

Commentary: A patient's story: "My life revolves around food and exercise"

A 32 year old woman describes how bulimia nervosa has affected her.

What is life like with bulimia?

Living with an eating disorder is a daily struggle. I feel it's worse than a drug addiction because you have to deal with food on a daily basis.

My life revolves around food and exercise. Because of my abuse of diet pills and purging, I had a stroke when I was 23. I now have headaches. I am at risk of having another stroke, and this time I have a high chance of not coming out of it. Emotionally, it's a daily

battle. I'm depressed because I want to eat, and I'm depressed because I know if I do eat, I'll get fat and gain all the weight back that I have lost.

Everyone around me is terrified that I may die from this, and it has put a lot of stress on my marriage. I have no bedroom life anymore because I refuse to let my husband touch me or even look at my body. My kids are affected greatly by it because I usually have no energy to do anything with them, and when I do have energy, I am staying busy to burn the calories I have put in my body.

What would you tell other people?

If you start to feel like you need to lose weight and you're taking it to extremes, go and find someone to talk to. This eating disorder can and will kill you. If you have an eating disorder, seek help now, before it's too late. The longer you live with an eating disorder, the harder it is to give it up. Furthermore, I think people need to know that:

- People with eating disorders aren't just trying to get attention

- People with eating disorders use their eating disorders to cope with stress
- Getting better and becoming healthy is an everyday battle, as you know that you can relapse and have to start all over.

If you know anyone who you think has an eating disorder, try to talk to them in a way that they won't think they are being ganged up on, and talk to a person you feel like you can trust. I wish I had sought treatment early on.

Lesson of the week

Poor glycaemic control caused by insulin induced lipohypertrophy

Tahseen A Chowdhury, Valerie Escudier

Good glycaemic control reduces the risk of complications of diabetes.¹ Secondary failure of oral hypoglycaemic treatment is common in patients with type 2 diabetes, and thus insulin treatment is often needed to improve glycaemic control. Indeed, in the UK prospective diabetes study 38% of patients with type 2 diabetes needed insulin treatment after 10 years.¹ The reasons for poor glycaemic control are many; and despite insulin treatment many patients with diabetes have poor glycaemic control.² Lipohypertrophy is characterised by a benign "tumour-like" swelling of fatty tissue secondary to subcutaneous insulin injections. We describe two cases in which poor glycaemic control was directly related to insulin induced lipohypertrophy, recognition of which led to major improvements in glycaemic control.

Case reports

Case 1

A 37 year old woman had been given a diagnosis of type 1 diabetes when she was 7 years old. She was treated with soluble insulin twice daily. She was transferred to our diabetes unit in 2000. She had experienced problems with fluctuating blood glucose concentrations, recurrent hyperglycaemia, and frequent unpredictable hypoglycaemia, despite compliance with diet and regular self monitoring of blood glucose. At her most recent annual review she was noted to have mild background retinopathy but no other microvascular or macrovascular complications of diabetes. Results of lipid, urea and electrolytes, and thyroid function tests were normal, but her glycated haemoglobin was 9.1% (normal range 3.6% to 5.1%). She was treated with Human Actrapid (Novo Nordisk) at a total daily dose of 30 units a day and Human Insulatard (Novo Nordisk) 26 units at night, both administered through a pen device. After her latest annual review she was referred to the diabetes specialist nurse for further follow up in an attempt to improve her glycaemic control.

The diabetes specialist nurse noted that she had fluctuating blood sugar concentrations, ranging from

2.0 mmol/l to 18.9 mmol/l, and unpredictable hypoglycaemic episodes occurring three or four times a week, with good awareness. The nurse reviewed her injection technique and noted significant lipohypertrophy at the sites of her abdominal injections. She was advised to avoid these sites for future injections and to reduce her insulin dose by 10%. Over the next six months her glycaemic control improved, with home tests showing blood sugar concentrations ranging from 3.4 mmol/l to 9.8 mmol/l before meals and 6.7 mmol/l to 12.3 mmol/l after meals. The frequency of occurrences of hypoglycaemia reduced to less than once a month. Her glycated haemoglobin fell to 6.8% within three months, despite the reduction in insulin dose.

Case 2

A 56 year old woman was seen in our diabetes clinic with poor control of her diabetes. She had been given a diagnosis of type 2 diabetes 10 years earlier and was changed to insulin treatment four years earlier because of secondary failure of oral hypoglycaemic treatment, when her glycated haemoglobin was 9.5%. She was treated with Human Mixtard 30 insulin (Novo Nordisk) twice daily at a total daily dose of 64 units and achieved reasonable control of her diabetes within six months of insulin treatment (glycated haemoglobin 6.4%). At her most recent annual review, however, it was noted that her glycaemic control was deteriorating—with home blood sugar concentrations ranging between 3.8 mmol/l and 24.5 mmol/l and occasional unpredictable hypoglycaemia. Her treatment was changed to Human Actrapid and Human Insulatard in a basal bolus dosing regimen at a total daily dose of 52 units a day. Despite good dietary compliance, regular self monitoring of blood sugar concentrations, and adjustment of her insulin dose she still had erratic glucose concentrations and poor control, with a haemoglobin A_{1c} of 8.9%. There was no evidence of microvascular or macrovascular complications of diabetes, and lipid, urea and electrolytes, and thyroid function tests all gave normal results.

She was seen by a diabetes specialist nurse who reviewed her injection technique and injection sites and noted that she had severe lipohypertrophy. She

Look for lipohypertrophy in all patients being treated with insulin, particularly in patients with erratic glycaemic control

Mile End Diabetes Centre, The Royal London Hospital, London E1 4DG
Tahseen A Chowdhury
consultant in diabetes
Valerie Escudier
senior diabetes specialist nurse

Correspondence to:
T A Chowdhury
Tahseen.Chowdhury@bartsandthelondon.nhs.uk

BMJ 2003;327:383-4