reviews

BOOKS • CD ROMS • WEBSITES • MEDIA • PERSONAL VIEWS • SOUNDINGS • MINERVA

The Clinician's Practical Guide to Attention-Deficit/Hyperactivity Disorder

Marianne Mercugliano, Thomas Power, Nathan Blum

Brookes Publishing, \$39.95, pp 368
ISBN 1 55766 358 0

Rating: ★★★★★

Attention-deficit hyperactivity disorder must surely have a remarkable status within medical practice. It is the most common neurodevelopmental disability, affecting 3-5% of children of school age, yet most of those receiving medical care have to be looked after by clinicians without any formal training in the condition. Aimed at primary care physicians (in the United

States), this book has, as its starting point, the recognition that many paediatricians who are struggling with these extremely complex children, have limited access to specialist advice.

This scenario will be familiar to British community and developmental paediatricians, who are also experiencing the consequences of the disorder increasingly being understood as a neurodevelopmental disorder rather than just a behavioural one. It can hardly be surprising that, particularly in the United States, the problem of "overdiagnosis" and subsequent "overprescription" of stimulants has appeared. On both sides of the Atlantic there is an urgent need for a comprehensive text that will enhance clinicians' confidence to practise safely and independently when necessary.

As an evidence based text, this may well fill the gap, being well supplied with resources to photocopy—such as parental and school questionnaires and advice for teachers on interventions—and backed up with extensive references and practical information for ordering resource material.

Its developmental approach is a rare feature in books on the subject, and this marks it as a paediatric text rather than a psychiatric one. In the initial section on diagnosis and

assessment, clinicians are helped to distinguish between behaviours that are a problem but within the normal range and those that cause an impairment of functioning. This is achieved by describing the core symptoms of the disorder at different stages of childhood and adolescence. At each developmental stage the symptoms are described in three levels of severity, allowing the reader to judge whether a particular manifestation is a developmental variation, problem, or disorder. Similarly, an unusual and welcome inclusion is a description of the normal evolution through childhood of so called neurological soft signs. These are a very common finding in attention-deficit hyperactivity disorder and sometimes difficult to interpret.

The chapter on medication, describing the use in some detail of about 25 drugs, gave me some concern as it seemed to encourage exploration of a far wider range of psychotropic agents than I, a fairly experienced clinician, would consider appropriate. This criticism apart, however, this is a quality text with much to help British paediatricians in dealing with this difficult problem.

Daphne V Keen consultant developmental paediatrician, Community Child Health, St George's Hospital, London

Antimony in Medical History

R Ian McCallum

Pentland Press, £15, pp 125
ISBN 1 85821 642 7

Rating: ★★★

Antimony for medical use has a history stretching back to ancient times. But the use of such a potentially toxic purgative has provoked bitter disagreement among practitioners, inspired vitriolic satire, and despatched countless men and women to an early grave. Recently, the presumed connection between antimony and cot deaths has added yet another chapter to the saga. Advanced by some early modern healers as a near miraculous panacea, it was also revered by alchemists because of its uncanny

power to purify gold. By 1678, over 106 standard remedies contained antimonials of various sorts; even more could be found in any 19th century medicine chest.

The history of antimony is important not only in its own right but also for the insight it offers into the changing intellectual milieu of medicine and pharmacy. Like theriac, the sovereign remedy with which it might profitably have been compared, antimony exercised a fascination well beyond the pages of the *Materia Medica*. Professor McCallum has mined a rich seam of evidence from Scotland, the United States, and France, as well as England, and illuminates his text with an apt choice of quotations.

He is generous with hard facts but less forthcoming in the matter of context or explanation. Readers unfamiliar with the tenets of Galenic and Paracelsian medicine, or with the close relation between physics, astrology, alchemy, and religion, may find some passages difficult to understand.

Since the use of antimony became a battlefield over which members of the English medical profession campaigned against empirics and apothecaries, it would have been useful to explore the wider social and economic issues at stake. One of the most

interesting of the many excellent illustrations reproduced here depicts a 17th century healer offering an antimony cup containing tartar emetic to her patient. She is repulsed by his guardian angel, who favours a learned physician skilled in the art of medicine.

The use of such overt propaganda against meddlesome women merits further comment, especially as "irregulars" such as Grace, Lady Mildmay, were at this time practising a promiscuous mix of Galenic and Paracelsian medicine heavily reliant on mineral substances. Any preparation that gave credence to the unlettered or unlicensed posed a serious threat to the status quo.

McCallum rightly emphasises the problems facing men and women who had to deal with human suffering in an age before the clinical advances that we now take for granted. If the cure was infinitely worse than the disease, the prospect of a treatment capable of eliminating all corruption from the body was surely irresistible. In the last resort, the triumph of hope over experience may explain why antimony enjoyed such enduring popularity.

Carole Rawcliffe reader in the history of medicine, University of East Anglia, Norwich

Champion of Women and the Unborn: Horatio Robinson Storer, MD

Frederick N Dyer

Science History Publications,
\$39.95, pp 626
ISBN 0 88135 266 7

Rating: ★★

Horatio Robinson Storer (1830-1922) is a historical figure to cause indignation among supporters of a woman's right to choose contraception and abortion. He motivated the American Medical Association into a "physicians' crusade" against abortion in 19th century United States. The resulting antiabortion legislation did untold harm or untold good, depending on your viewpoint, and the Supreme Court did not revoke much of it until 1973.

He was as much against contraception, holding that all available methods—including "use of a sponge, cold injections, the employment of condoms or the so-called womb guard, or that worst measure of all, incomplete intercourse"—contributed to uterine disease. In 1870 he discounted evidence from

the Oneida community that their common practice of delayed or suppressed ejaculation was harmless.

It would be easy to dismiss Storer as an example of the repressive attitudes of his time, but he would have seen himself as a progressive, advocating the best medical care for women in a world that often mistreated them. He used his considerable surgical skill to pioneer improvements in gynaecology and was the first physician in the United States to give a complete collegiate course on diseases of women as distinct from midwifery. He urged the use of chloroform to relieve the pain of childbirth and deplored the policy of refusing to admit unmarried women to lying-in hospitals. He was also a champion of the oppressed, whether black citizens or immigrant Europeans.

Storer was born into a Protestant family in Boston, though he converted to Catholicism at the age of 49. The boundaries between natural science and medicine were less distinct than now, and his father, David Humphreys Storer, was a keen ichthyologist who became professor of obstetrics and medical jurisprudence at Harvard and president of the American Medical Association (Horatio was to become vice-president). Revealing anecdotes about his famous contemporaries are delights in a biography that contains a lot of less interesting detail.

Storer seems to have been one of those people who live long and exert great influence despite apparent chronic ill health. The list of his achievements is considerable. But despite the author's best efforts, he does not emerge from these pages as an attractive figure. Every quarrel is documented, but

women's voices, apart from his devoted daughter's, are almost absent. Storer disapproved of female physicians, especially after an unhappy collaboration at an all female hospital, although he mellowed towards them in later years. He married three times, but we can only guess at his real feelings about women and what they thought of him. His first wife seems to have had a severe mental illness related to menstruation.

His descriptions of female patients are at odds with present day sympathies, which is exactly why they should be read by those with a special interest in the history of women's bodies and the eternal debate over the morality of abortion.

Jeanette Cayley senior clinical medical officer in family planning and reproductive health care, Featherstone Road Clinic, Southall

The BMJ Bookshop will endeavour to obtain any books reviewed here. To order contact the BMJ Bookshop, BMA House, Tavistock Square, London WC1H 9JR.
Tel: 020 7383 6244, Fax: 020 7383 6455
email: orders@bmjbookshop.com
Online: www.bmjbookshop.com
(Prices and availability subject to change by publishers.)

Reviews are rated on a 4 star scale
(4=excellent)

WEBSITE OF THE WEEK

NHS on the web Although a search engine is probably the first internet tool that any web newcomer will make use of, there are other ways to find what you're looking for. Using your internal search engine—your brain—to make an informed guess about the words that might be in the web address, or uniform resource locator (URL), is one way to find what you want.

Lots of people have been doing this to try to track down NHS services, according to the NHS director of planning Alasdair Liddell. According to Mr Liddell, the NHS's newest website, www.nhs.uk, was getting thousands of hits each day even before it had received any content—presumably by people trying to guess their way to NHS Direct Online (www.nhsdirect.nhs.uk). This top level domain will form a crucial part of the senior NHS management's strategy of imposing greater coherence and quality on the electronic output from NHS organisations. The site itself illustrates some of the difficulties of this approach: although the home page has the NHS white and blue look and commits no serious web faux pas, the pages that it links to—the NHS organisations in Scotland, Wales, and Northern Ireland, and NHS Direct itself—all have substantially different looks, content, and feel.

Mr Liddell also mentioned the acquisition of Digerati, a suite of new web tools that will be made available to NHS users at "low or no cost," and new central web hosting services that, he claimed, "would cut duplication and effort" together with a new coherent scheme of URL addressing based on the convention of www.nhs.uk/service_name. For example, general practitioners who wish to remind themselves of some aspect of the minutiae within the *Statement of Fees and Allowances* (the "Red Book") will be able to consult the online version at www.nhs.uk/redbook.

This is all well and good, but some doubt must remain as to whether such intense centralisation will work within the context of the web.

Douglas Carnall
BMJ
dcarnall@bmj.com

Hit parade

These articles scored the most hits on the BMJ's website in the week of publication

FEBRUARY

- 1 ABC of heart failure: Acute and chronic management strategies**
2000;320:559-62
52 939 hits
- 2 Recent advances: Tropical medicine**
2000;320:490-4
48 903 hits
- 3 ABC of heart failure: Management: digoxin and other inotropes, β blockers, and antiarrhythmic and antithrombotic treatment**
2000;320:495-8
48 200 hits
- 4 ABC of heart failure: Non-drug management**
2000;320:366-9
36 539 hits
- 5 ABC of heart failure: Management: diuretics, ACE inhibitors, and nitrates**
2000;320:428-31
34 564 hits
- 6 Recent advances: Intensive care**
2000;320:358-61
28 148 hits
- 7 Advances in managing chronic disease**
Editorial 2000;320:525-6
24 008 hits
- 8 Doctor as murderer**
Editorial 2000;320:329-30
19 859 hits
- 9 Regular review: Tumour markers in malignancy**
2000;320:424-7
15 371 hits
- 10 Cholesterol and strokes**
Editorial 2000;320:459-60
12 843 hits

PERSONAL VIEW

Deaths on the operating table

The consequences of a patient dying during surgery have made headline news in the medical press recently. The message appeared unequivocal in the *BMJ*, *Hospital Doctor*, and *BMA News Review*. An operating list should stop if a patient dies on the operating table, and the surgeon should not operate for the rest of that day.

The *BMJ* reported Professor Sir Alfred Cuschieri as saying, "My own view is that the death on the operating table of the patient is a harrowing experience for a surgeon. In my view, the surgeon is emotionally and mentally not in the frame of mind to continue to operate that day." It was also stated that the sheriff conducting the inquiry agreed with this view. The president of the Royal College of Surgeons of Edinburgh was also quoted as saying, "We can understand the pressure that single surgeons are under, but I think there would be a very strong feeling that, when a surgeon loses a patient, he should not continue operating that day."

I was recently involved in treating two severely injured patients from a road traffic accident. The first died during emergency surgery in the small hours of the morning. The second needed emergency surgery for four open long bone fractures. I can recall several deaths during surgery in the 17 years since I qualified. I cannot honestly say that I was unable or unfit to continue working after any of these unfortunate events.

In view of the media comments, however, I sought advice from a medical defence organisation at about 5.30 am. Both my registrar, also a member of the same defence organisation, and I were advised not to operate further and to inform our trust accordingly. Fortunately, a colleague was available to carry on operating. This advice has been considered "totally untenable" and setting "a very dangerous precedent" by Sir Alfred (personal communication) as his comments applied only to elective surgery. This was not made clear in the selective reporting of the case. Subsequent advice from another medical defence organisation, based on the same set of facts, was to continue operating; a clear dichotomy of opinion.

In this age of evidence based medicine, is it logical to stop operating after any death on the operating table? As far as I am aware, there is no evidence either way. Perhaps there should be before issuing any guide-

lines or getting the counsellors in. If surgeons are found to be traumatised after a death on the operating table are the anaesthetists unaffected? The cause of death may be multifactorial and unrelated to the actual surgical procedure.

What would happen if the patient arrested as the drapes were being put on the patient immediately before the incision was made? Surely the anaesthetists would feel

more traumatised than the surgeons: the subject has been described as one of the greatest taboos of modern anaesthesia literature. Nursing and other staff may be similarly affected. How susceptible individuals would be under such circumstances would become apparent only after the event. Some individuals

may require more time off work than others. The day off suggested so far is hardly based on any science.

As the effect of a death in theatre on the team involved is currently unknown, surely it is illogical to differentiate between elective and emergency surgery. Research in the future may discover that any such death may cause significant dysfunction in individuals. If so, then the consequences of preventing the surgical and anaesthetic team from working are profound. How would military field hospitals function during armed conflicts? The same logic would apply to NHS staff after major civilian disasters.

Every major hospital dealing with emergency surgery would require at least two on-call teams in susceptible specialties just in case a death occurred during surgery. As this is an uncommon event, it will have no significant impact in terms of the European Union working time directive. However, the increase in the on-call workload is likely to have a detrimental effect on waiting times for our patients.

The issue clearly requires urgent and reasoned discussion so that sensible conclusions can be reached. The consequences of rashness are immense.

Mel Jones consultant orthopaedic and trauma surgeon, Bangor

If you would like to submit a personal view please send no more than 850 words to the Editor, BMJ, BMA House, Tavistock Square, London WC1H 9JR or email editor@bmj.com

It is illogical to differentiate between elective and emergency surgery

The cause of death may be multifactorial and unrelated to the actual surgical procedure

SOUNDINGS

How to party

At an academic dinner recently two professors remarked that their grown up children had advised them to get a life. This advice had been given in all seriousness and, in one case, had been reinforced by a postcard saying, "WORK is for people who don't know how to party." "It's true," the professor added sadly, "I really don't."

Such ignorance is new. Older professors always used the word "fun" in their retirement speeches, usually when referring to the second world war or academic life in the 1960s. Sooner or later medical schools will realise that those bon viveurs were actually more productive than their workaholic successors.

When this happens, partying will come back into fashion. Indeed, it will become compulsory. We shall have a national RAE (Revelry Assessment Exercise) involving detailed questionnaires on party quality and quantity. We shall agonise about how many parties to declare.

We shall need published guidelines. At present there is no basis for evidence based partying. What follows is a draft for consultation.

The invitation: Email is boring. Have your invitation sung by a stripping lab technician or encoded on a genetic fingerprint.

The venue: Be original. Choose somewhere that will be stimulatingly unfamiliar to academics—the outpatient clinic, for example. Inexpensive decorations can be made from bunches of used airline tickets.

Games: Academics love to see themselves as Machiavellian plotters. Get things going with a box of rubber daggers. Everyone has to stab everyone else in the back and when stabbed has to writhe in agony on the floor. (Keep an eye on the stilettos to make sure that nobody slips in a real one.)

The food: Bring back memories of your days as a junior. Install a hospital vending machine with sinister looking pasties. And why not retune the microwave to sound like a mobile phone?

The music: Hip disc jockeys with strange names and huge loudspeakers can be easily hired. Most are medical students but a few are ear, nose, and throat surgeons drumming up trade.

The drugs: Having trouble recruiting volunteers for your latest phase III trial? Make sure your doormen have a good supply of consent forms.

Remember, these are only guidelines. Enjoy.

James Owen Drife professor of obstetrics and gynaecology, Leeds