Table 1. Efficiency of quantification of DRG mass by the *n*-octyl- β -D-glucopyranoside and Triton X-100 micellar methods in the presence of varying amounts of endogenous lipid extract

Efficiency of conversion of 330 pmol of *sn*-1-stearoyl-2-arachidonoylglycerol was assayed in the presence of four concentrations of endogenous cell lipid prepared from confluent monolayers of Swiss 3T3 fibroblasts. The assays were performed as described in Fig. 1. The determinations were performed in triplicate and values are expressed as means $\pm$ S.D.

Number of cells used to prepare endogenous lipid added to assay	Efficiency of phosphorylation of <i>sn</i> -1-stearoyl-2-arachidonoylglycerol (%)	
	1.5 % <i>n</i> -Octyl- β -D-glucopyranoside	0.3 % Triton X-100
2.25×10^5	80.4 ± 52.1	101.2 ± 2.1
4.50×10^5	60.4 ± 9.6	97.3 ± 16.8
7.50×10^5	4.45 ± 10.1	80.8 ± 7.6
2.25×10^6	0.4 ± 14.0	38.8 ± 8.5

phosphorylation of exogenous DRG added to increasing amounts of crude lipid extract derived from confluent Swiss 3T3 fibroblasts has been assessed for the two detergent systems. The original *n*-octyl β -D-glucoside assay did not support complete conversion at any lipid concentration, giving maximally 80.4 ± 52.1 % conversion at 2.25×10^5 fibroblasts but only 0.4 ± 14.0 % at 2.25×10^6 cells (this range is equivalent to the cell numbers found in 6.25 – 62.5 cm² of confluent cell monolayer). In comparison we found the Triton X-100 micelle capable of supporting higher levels of conversion at all cell lipid concentrations. Full conversion of the DRG added was observed at lipid concentrations equivalent to $(2.25$ – $4.5) \times 10^5$ cells, although at higher cell number the efficiency of conversion did fall, e.g. to 38.8 ± 8.5 % at 2.25×10^6 cells (Table 1). However, it was possible to recover near quantitatively 100 pmol–1 nmol of phosphatidic acid in the presence of a lipid extract equivalent to 2.25×10^6 Swiss 3T3 fibroblasts that had been included in a sham assay. This would seem to demonstrate that the inability to quantify DRG fully at high lipid concentrations was not due to decreased recovery of [³²P]phosphatidic acid in the presence of high levels of cellular lipid, but rather to an inhibition of diacylglycerol kinase activity towards its DRG substrate. Thus, we feel it is apparent that the Triton X-100 system is capable of complete kinase-catalysed phosphorylation of the DRG present in cell lipid extracts, provided the amount of lipid that is used for the determination of DRG mass is carefully titrated to avoid the use of concentrations that would prevent complete quantification of the DRG present.

By employing the adapted assay and taking care to titrate the amount of tissue used in the determination, we have assayed the DRG mass in cell lipid extracts where previously we were unable to gain satisfactory results. It would seem that similar improvements in DRG mass quantification in other tissues should be possible where previous application of the 'Preiss' assay has been problematic. However, we propose that the following points are considered when initially attempting to quantify DRG mass.

1. If DRG mass is to be measured in unstimulated tissue, it is necessary to assess the mass from a standard curve constructed in the absence of lipid. For the measurement to be accurate, the efficiency of phosphorylation of both the standard and experimental lipid samples must be identical. This problem can be addressed by ensuring the phosphorylation is performed with

100 % efficiency in both the standard curve and lipid extract. Table 1 shows that this is possible for the curve; however, the lipid extracts must be carefully titrated such that the component DRG is also quantitatively phosphorylated. This can be assessed by the addition of exogenous DRG and quantification of the [³²P]phosphatidic acid produced above basal levels.

2. It remains necessary to ensure that the changes in DRG mass can be quantified fully in the presence of lipid extract, i.e. the presence of cellular lipid should not affect the linearity of the standard curve since this would prevent accurate quantification of the changes in DRG mass. This can be determined by assessing the efficiency of phosphorylation of standard DRG in the presence of lipid and requires continued linearity over the range of DRG mass likely to be found experimentally.

3. If the specific nature of the tissue prevents use of the assay with 100 % efficiency in the lipid extracts, the assay may still be used to measure basal DRG levels providing the efficiency of phosphorylation in a standard curve constructed in the presence of cellular lipid is linear. Quantification of DRG in the cell lipid extracts can then be made by correcting for the difference in efficiency of phosphorylation in standard curves constructed in the absence and presence of cellular lipid.

These three basic criteria have been applied with the 0.3 % (w/v) Triton X-100 method to quantify fully DRG in crude lipid extracts from both Swiss 3T3 fibroblasts and A10 aortic vascular smooth muscle cells. There are, however, cell types, such as adipocytes, where the amount of cellular lipid is too high for accurate measurement of the DRG component. For such samples it has proven necessary to partially purify the DRG by t.l.c. before quantification in the assay. However, we feel application of the modified method will allow quantification of DRG mass levels in many cell and tissue types where previously the results have lacked accuracy, sensitivity and precision.

The assay was adapted during work funded by the Cancer Research Campaign and Wellcome Trust. A. P. is a recipient of a University of Glasgow Postgraduate Scholarship from the Medical Faculty, University of Glasgow.

Andrew PATERSON, Robin PLEVIN and Michael J. O. WAKELAM

Molecular Pharmacology Group, Department of Biochemistry, University of Glasgow, Glasgow G12 8QQ, U.K.

Neugebauer, J. M. (1990) *Methods Enzymol.* **182**, 239–253

Preiss, J., Loomis, C. R., Bishop, W. R., Stein, R., Neidel, J. E. & Bell, R. M. (1986) *J. Biol. Chem.* **261**, 8597–8600

Thompson, N. T., Tateson, J. E., Randall, R. W., Spacey, G. D., Bonser, R. W. & Garland, L. G. (1990) *Biochem. J.* **271**, 209–213

Walsh, J. P. & Bell, R. M. (1986) *J. Biol. Chem.* **261**, 6239–6247

Received 12 August 1991

Quantification of diradylglycerols: a reply

Paterson *et al.* (1991) have experienced difficulties in employing the *sn*-1,2-diacylglycerol assay developed by our laboratory (Preiss *et al.*, 1986). Their report focuses attention on factors known to be important to successfully quantify cellular diradylglycerols (DRG). The assay employs *E. coli* diacylglycerol kinase (DGK) in β -octylglucoside mixed micelles containing a phospholipid cofactor, such as cardiolipin, to quantify the amount of *sn*-1,2-diacylglycerol present in the lipid extracts from cells. The assay is based on the ability of the *E. coli* DGK to quantitatively convert *sn*-1,2-diacylglycerol to [³²P]phosphatidic acid. That the

E. coli DGK quantitatively recognizes alkylacylglycerols as well as diacylglycerols has been previously reported (Dougherty *et al.*, 1989). Further, the fact that the mixed micellar assay can be overloaded at high levels of cellular lipids is thoroughly predictable (Preiss *et al.*, 1986, 1987). The points raised by Paterson *et al.* agree with out previously published description of what constitutes appropriate controls for the assay (Preiss *et al.*, 1987). We believe that their solutions to the technical problems encountered may be acceptable, but we have some reservations as described below. Other aspects of the assay also are discussed.

The initial communication by Preiss *et al.* (1986) described an efficient and sensitive assay for the stereospecific quantification of diacylglycerols from platelets, hepatocytes, normal fibroblasts, and transformed fibroblasts. The data were in excellent agreement with those values obtained by other investigators who had used different quantitative methods (Bell *et al.*, 1979; Hughes *et al.*, 1984; Rittenhouse-Simmons, 1979; Bocckino *et al.*, 1985). A chapter in *Methods in Enzymology* (1987) further detailed the method. In this article some frequently encountered technical problems and ways to overcome these problems were described (Preiss *et al.*, 1987). As stated in that chapter, impurities in either the [γ - 32 P]ATP or the β -octylglucoside used in the assay can result in problems in diacylglycerol measurements. Non-quantitative conversion of DRGs to the corresponding [32 P]-phosphatidic acids can be the result of poor quality of the β -octylglucoside. Impurities present in commercially available β -octylglucoside can inhibit the *E. coli* DGK, and can result in failure to solubilize completely the DRG present. It is essential to recrystallize the β -octylglucoside prior to use. The quality of the [γ - 32 P]ATP employed should be assessed. A protocol described by our laboratory (Preiss *et al.*, 1987) or another reliable method should suffice.

Another potential problem in determining the cellular DRG concentrations concerns the improper quantification or inadequate quality of the diacylglycerol standard. Inaccurate quantification of the diacylglycerol standard will lead to systematic errors in the estimated values obtained for cellular DRGs. As described by our laboratory (Preiss *et al.*, 1986), the concentration of the diacylglycerol standard can be accurately determined by a simple ester assay (Stern & Shapiro, 1953). Delivery of the diacylglycerol standard with a Hamilton syringe is advised. Additionally, the *sn*-1,2-diacylglycerol standard should be analysed periodically to assess isomerization to 1,3-diacylglycerol.

The concentration and type of detergent employed in the assay also can affect accuracy. As Paterson *et al.* have discovered, 0.3% (w/v) Triton X-100 more readily solubilized the lipids from a very large number of cells (2.25×10^6 Swiss 3T3 fibroblasts) than did 1.5% (w/v) β -octylglucoside. They suggest that the Triton X-100 micelles may be less susceptible to lipid overloading, although at the very high concentration of cell lipids examined even the Triton X-100 assay conditions did not support complete conversion of the DRGs. The β -octylglucoside micelles present at 1.5% (w/v) can be overloaded with the total lipids from a large number of cells. The amounts of lipid and detergent must be adjusted to assure solubilization and conversion to phosphorylated products. Triton X-100 is eliminated as the detergent of choice because the *E. coli* DGK phosphorylates Triton X-100 (Schneider & Kennedy, 1976). The presence of phosphorylated Triton X-100 products, comigrating with authentic phosphatidic acid, may account for the observations of Paterson *et al.* that the efficiencies of diacylglycerol phosphorylation in the presence of Triton X-100 exceed 100%.

The appropriate controls as outlined by Preiss *et al.* (1987) ensure accurate DRG determination. The three points that Paterson *et al.* offer for consideration when making initial DRG measurements have been described previously by our laboratory

(Preiss *et al.*, 1987) and constitute the appropriate controls. The ability of the enzyme to phosphorylate several substrates other than diacylglycerol has not impeded the efforts of our laboratory and others to make accurate lipid determinations. On the contrary, the broad substrate specificity of the enzyme, coupled with simple chemical treatments, has allowed quantification of diacylglycerols, alkylacylglycerols and ceramides from many cell types and tissues.

Some modifications to the original assay can increase the sensitivity. Increasing the specific activity of the radiolabelled ATP substrate will increase the sensitivity of the assay for DRGs (Preiss *et al.*, 1987). Preincubating the DGK membranes with Mg^{2+} -ATP will help to prevent DRG-dependent enzyme inactivation (Walsh & Bell, 1986; Wright *et al.*, 1988), and decrease the background counts by removing endogenous DRG present in the *E. coli* membranes.

We do not recommend some of the modifications suggested by the Paterson group. Cardiolipin is the most effective lipid cofactor for DGK, and the use of phosphatidylserine as the lipid cofactor leads to poorer enzyme activation (Walsh & Bell, 1986). Substitution of cardiolipin with phosphatidylserine is not recommended. Sonication of the lipids for 30 min at 4 °C may promote the isomerization of 1,2-diacylglycerols to 1,3-diacylglycerols (Serderavich, 1967), and the 1,3-isomer is not a substrate for the enzyme; consequently, this modification also is not recommended. Likewise, t.l.c. of the lipid samples (from lipid-rich cells) prior to assay might also promote acyl migration, and should be avoided if possible.

Two further points should be considered when using the DRG assay. Although we have not performed a complete analysis of the efficiency of phosphorylation of all diacylglycerol species, we have found that this efficiency is lower for long chain and C_{18} disaturated diacylglycerol species. Substitution of Triton X-100 mixed micelles for the β -octylglucoside mixed micelles does not improve this efficiency. However, when reactions are run at an elevated temperature (45 °C), quantitative conversion of these species is achieved. Finally, we have found that some brands of glass test tubes are inhibitory to the DGK reaction. We recommend that only Pyrex glass tubes be employed.

Roy A. BORCHARDT,* W. Robert BISHOP,†
Steve B. BOCKKINO,‡ Carson R. LOOMIS,‡
Daniel M. RABEN,§ J. Kevin RAMER,*
Paul P. VAN VELDHoven|| and Robert M. BELL*

*Department of Biochemistry, Duke University Medical Center, Durham, NC, U.S.A., †Molecular Pharmacology Section, Schering-Plough Research, Bloomfield, NJ, U.S.A., ‡Sphinx Pharmaceutical Corporation, Durham, NC, U.S.A., §Department of Physiology, The Johns Hopkins University School of Medicine, Baltimore, MD, U.S.A., and ||Katholieke Universiteit Leuven, Leuven, Belgium

Bell, R. L., Kennerly, D. A., Stanford, N. & Majerus, P. W. (1979) *Proc. Natl. Acad. Sci. U.S.A.* **76**, 3238–3241

Bocckino, S. B., Blackmore, P. F. & Exton, J. H. (1985) *J. Biol. Chem.* **260**, 14201–14207

Dougherty, R. W., Dubay, G. R. & Niedel, J. E. (1989) *J. Biol. Chem.* **264**, 11263–11269

Hughes, B. J., Rye, K. A., Pickford, L. B., Barritt, G. J. & Chalmers, A. H. (1984) *Biochem. J.* **222**, 535–540

Paterson, A., Plevin, R. & Wakelam, M. J. O. (1991) *Biochem. J.* **280**, 829–830

Preiss, J. E., Loomis, C. R., Bishop, W. R., Stein, R., Niedel, J. E. & Bell, R. M. (1986) *J. Biol. Chem.* **261**, 8597–8600

Preiss, J. E., Loomis, C. R., Bell, R. M. & Niedel, J. E. (1987) *Methods Enzymol.* **141**, 294–300

Rittenhouse-Simmons, S. (1979) *J. Clin. Invest.* **63**, 580–587

Serderavich, B. (1967) *J. Am. Oil Chem. Soc.* **44**, 381–393

Schneider, E. G. & Kennedy, E. P. (1976) *Biochim. Biophys. Acta* **441**, 201–212

- Stern, I. & Shapiro, B. (1953) *J. Clin. Pathol.* **6**, 158–160
 Walsh, J. P. & Bell, R. M. (1986) *J. Biol. Chem.* **261**, 6239–6247
 Walsh, J. P. & Bell, R. M. (1986) *J. Biol. Chem.* **261**, 15062–15069
 Wright, T. M., Rangan, L. A., Shin, H. S. & Raben, D. M. (1988) *J. Biol. Chem.* **263**, 937–9380

Received 2 October 1991

Part of the phospho group of pyridoxal phosphate may titrate over the pH range 5–8 in aspartate aminotransferase

The ^{31}P n.m.r. spectrum of aspartate aminotransferase (EC 2.6.1.1) shows a pK of about 6.5 (the value varies with isoenzyme and with salt concentration) in the pH-dependence of the chemical shift of the phosphorus atom of the pyridoxal phosphate. When this was first observed, it was attributed to titration between $-\text{PO}_3\text{H}^-$ and $-\text{PO}_3^{2-}$; it was soon realized, however, that the change in shift was too small for this conversion (Mattingly *et al.*, 1982; Korpela *et al.*, 1987a,b; Schnackerz *et al.*, 1989; Sanchez-Ruiz *et al.*, 1991). These three groups of authors therefore concluded (1) that the group was in the $-\text{PO}_3^{2-}$ form throughout the pH range 5–8 [Sanchez-Ruiz & Martinez-Carrion (1986) and Sanchez-Ruiz *et al.* (1991) qualified this by saying merely that it was mainly in the dianionic form], and (2) that an ionization of pK 6.5 elsewhere in the protein was affecting the phosphorus, e.g. by changing O–P–O bond angles.

Although the arguments were convincing that complete titration of $-\text{PO}_3\text{H}^-$ to $-\text{PO}_3^{2-}$ would give a much larger change in chemical shift than that observed, no one argued whether a fraction of the group might not possess the observed pK. Dixon & Tipton (1973) pointed out that, to a close approximation,

Scheme 1. The ionization of the postulated acid

The pK values marked on the scheme are those of the groups, and those to the right are the titration pK values, which are so far apart that they differ negligibly from the molecular values. It may be noted that although the higher of the two group values for the phospho group is 5.8, it is with a pK of 6.5 that a fifth of it titrates; this titration pK is dominated by the value for $-\text{AH}$, since most of this value is contributed by the dissociation of $-\text{AH}$.

ionizing groups, if capable of interacting with others, can titrate in fractions. Dixon *et al.* (1991) recently showed that this statement is exact, and not merely an approximation. Thus a fraction of one group will have one pK, and the rest will have another (or other fractions will have others if the group interacts with more than one other ionizing group). The one H^+ released by the protein with each pK is made up of the fractions from different groups that possess this pK. Hence a small fraction of the phospho group might indeed be in the $-\text{PO}_3\text{H}^-$ form, and change the shift of the phosphorus atom as it titrates to form $-\text{PO}_3^{2-}$.

Fig. 1. Curves of the pH-dependences of the relative concentrations of a dibasic acid

The relative concentrations of the ionic forms of the acid $\text{HA}-\text{PO}_3\text{H}^-$ are shown: (1) $[\text{HA}-\text{PO}_3\text{H}^-]$; (2) $[\text{HA}-\text{PO}_3^{2-}]$; (3) $[\text{A}^--\text{PO}_3\text{H}^-]$; (4) $[\text{A}^--\text{PO}_3^{2-}]$; (5) = (2) + (3), i.e. the total concentration of the singly dehydrated form; (6) = (2) + (4), i.e. the total concentration of the form $-\text{PO}_3^{2-}$; (7) = (3) + (4), i.e. the total concentration of the form $-\text{A}^-$. These curves are obtained by assigning the $-\text{PO}_3\text{H}^-$ group a pK of 2.1 (intersection a) when the other group is in the form $-\text{AH}$, and 5.8 (intersection c) when the other group is in the form $-\text{A}^-$ (i.e. $-\text{AH}$ greatly stabilizes the dianionic form of the phospho group, e.g. by hydrogen bonding, lowering its pK greatly in comparison with a normal phosphate), and assigning the $-\text{AH}$ group a pK of 2.7 (intersection b) when the phospho group is in the $-\text{PO}_3\text{H}^-$ form, and 6.4 (intersection d) when the phospho group is in the $-\text{PO}_3^{2-}$ form, as shown in Scheme 1. The assignment of a difference of 0.6, i.e. $\log_{10}(4)$, between the pK values of the two groups gives a 4:1 ratio between the two monohydrated forms, so that 20% of the group $-\text{PO}_3\text{H}^-$ titrates with one pK, which is 6.5 for the actual values of group constants chosen (curve 6). The remaining 80% of the H^+ released with this titration pK comes from the group $\text{HA}-$; the remaining 80% of the phospho group titrates with the pK of 2.