REVIEW ARTICLE

Molecular genetics of actin function

Emma S. HENNESSEY*, Douglas R. DRUMMOND† and John C. SPARROW‡

Department of Biology, University of York, York YO1 5DD, U. K.

INTRODUCTION

Actin is found in all eukaryotes and plays a fundamental role in many diverse and dynamic cellular processes. Despite being so ubiquitous, actin isoforms exhibit an unusually high degree of amino acid sequence similarity. This poses the question of how actin can fulfil such diverse functions in different organisms, yet retain such a high degree of sequence conservation? In this Review we examine the impact molecular genetics has made on our understanding of the relationship between actin amino acid sequence and function. We refer the reader to other reviews for more details of actin structure, biochemistry and cell biology than space allows us to include [1–10].

Actin is a globular protein (G-actin) which polymerizes into filaments (F-actin) [6] for most of its biological functions [7]. F-actin is a major component of the cytoskeleton [3]. The control of F-actin assembly and turnover is important for many cellular processes such as motility, morphological changes, cell division and intracellular movements. Many proteins control actin polymerization and depolymerization within cells. These include proteins which sever or cap actin filaments, nucleate polymerization, cross-link and stabilise F-actin (reviewed in [3–5]). Others bind G-actin, maintaining up to 50% of cytoplasmic actin as monomers [8] and can respond to extracellular signals by releasing the G-actin which then polymerizes [11]. Cytoskeletal actin may localize some metabolic pathways within the cell (reviewed in [12]). In addition to roles in non-muscle cell motility and contractile processes, such as cell division (reviewed in [9,10]), actin is a major component of myofibrils. In myofibrils, F-actin is assembled into the thin filaments which interact with the myosin heads of the thick filaments to produce contraction (for reviews see [13,14]).

Molecular genetic techniques can be used to study the relationships between a protein's functions and its amino acid sequence by examining the effects of sequence variation. Gene cloning and sequencing and, more recently, the polymerase chain reaction (PCR) have accelerated the accumulation of new actin sequences in the databases. These naturally occurring isoforms can be used to assess the significance of amino acid sequence variation for functional differences between actins. Information on the importance of specific amino acids in actin can also be obtained from induced 'man-made' mutations recovered from genetic studies. Actin mutations have been described in the yeast *Saccharomyces cerevisiae* [15,16], the fruitfly *Drosophila melanogaster* [17–22], the nematode *Caenorhabditis elegans* [23,24] and cultured human cells [25,26]. Mutations made in cloned actin genes have been studied after expression either by transcription and translation *in vitro* [27–36] or *in vivo* following transformation

of suitable organisms such as *S. cerevisiae* [37–43], *Dictyostelium discoideum* [44], *D. melanogaster* [20,33,45–49], or cultured cell lines [25,26,28,50].

A most significant development has been the determination of the atomic structure of actin from crystals of an actin–DNase I complex [51; reviewed in 1,2]. Atomic structures of complexes with the G-actin binding protein, profilin [52], the muscle protein, myosin [53] and gelsolin fragment-1 [54] will soon provide information on these physiologically important actin–ligand interactions. Most existing mutants were made without the benefit of the atomic structure of actin. It is therefore timely to consider the effects of actin mutants with reference to this atomic structure.

COMPARISON OF NATURALLY OCCURRING ACTINS

Conservation of actin sequences

Comparison of protein sequences can identify conserved regions and reveal evolutionary relationships. Sequence conservation implies evolutionary constraint, often identifying regions of functional importance. Actin sequences have been reviewed [55,56] and are much more conserved than other proteins, such as globins, found in the same species [56].

In an alignment of the 81 unique actin protein sequences taken from the recent OWL database version 15.1 [57], 17.4% of positions in the consensus sequence are invariant in all actins, 35.6% of positions are invariant or have a difference in only one of the 81 actins, and 40.4% of positions are invariant or have what are considered conservative changes (D. R. Drummond, unpublished work). For most positions the amino acids are identical in the majority of actin sequences and at least 95% of actins have the same amino acid at 62.9% of positions. The most divergent sequences (44.9%) are those from the protozoan *Euplotes crassus* and carrot actin 2. Skeletal muscle α -actins in human, mouse, rat, rabbit and chicken are identical, as are the cytoplasmic β -actins in human, mouse, rat, cow and chicken.

The high conservation of actin sequences means that, unlike other proteins, one cannot identify important sites as isolated regions with the most conserved sequences. Rather it suggests constraints throughout the actin structure. Thus regions of sequence variation may identify sequences where the constraints are reduced or which are important for isoform-specific functions.

Actin phylogenetic trees ([55,56,58,59]; Figure 1) produce relatively few groupings which can be considered significant [56,59], a consequence of the high sequence conservation of actins. The major divisions correspond to conventional taxonomic classifications. For example, plants and animals form distinct groups.

* Present address: Microbial Genetics Group, School of Biological Sciences, University of Sussex, Falmer, Brighton BN1 9QG, U. K.

† Present address: CRC Medical Oncology Unit, Southampton General Hospital, Southampton SO9 4XY, U. K.

‡ To whom correspondence and reprint requests should be addressed. The actin amino acid sequence alignment mentioned in the text as (D. R. Drummond, unpublished work), details of the sequences used, references etc. will be provided on request by J. C. S. Please make requests either by fax (to +44 904 432860) or by e-mail to JCS1@uk.ac.york.vaxa.

Figure 1 Actin protein tree

A phylogenetic tree of currently available actin protein sequences from the OWL database version 15.1 [57] was produced with the program Clustal V [191] which uses the neighbour-joining method of Saitou and Nei [192]. Actin names are described below. The percentage divergence is marked in italics on the branches. The reproducibility of the tree was determined using 1000 bootstrap replicas. The percentage of replicas containing a given branch is circled next to the branch. Only values greater than 80% are reported. The tree is unrooted but, as is common practice, the root can be placed on the midpoint of the longest branch, which is to *Euplotes*. Key: skel, skeletal; card, cardiac; smth, smooth; cyt, cytoskeletal. *Xenia*, *Xenopus laevis* (African clawed frog); *Xentr*, *X. tropicalis* (Western clawed frog); *Xenbo*, *X. borealis* (Kenyan clawed frog); *S* clava, *Styela clava*; *S* plicata, *Styela plicata* (sea squirts); *Artemia* sp. (brine shrimp); *C* elegans, *Caenorhabditis elegans* (nematode); *Hydra*, *Hydra attenuata*; *Dros*, *Drosophila melanogaster* (fruit fly); *Bombyx*, *Bombyx mori* (silkworm); *Aplysia*, *Aplysia californica* (California sea hare); *Arabidopsis*, *Arabidopsis thaliana* (mouse-ear cress); *Pisaster*, *Pisaster ochraceus* (sea star); *Amoeba*, *Acanthamoeba castellanii* (amoeba); *E* histolyt, *Entamoeba histolytica*; *Strpu*, *Strongylocentrotus purpuratus* (purple sea urchin); *Strfn*, *S. franciscanus* (sea urchin); *Taenia*, *Taenia solium*; *Dicty*, *Dictyostelium discoideum* (slime mould); *Physarum*, *Physarum polycephalum* (slime mould); *S* pombe, *Schizosaccharomyces pombe* (fission yeast); *S* cerevisiae, *Saccharomyces cerevisiae* (baker's yeast) and *S. carlsbergensis* (lager beer yeast); *Absidia*, *Absidia glauca* (zygomycete); *Aspergillus*, *Aspergillus nidulans*; *Thermomyces*, *Thermomyces lanuginosa* (*Humicola lanuginosa*); *K* lactis, *Kluyveromyces lactis* (yeast); *Phyin*, *Phytophthora infestans* (potato late blight fungus); *Phyme*, *P. megasperma* (potato pink rot fungus); *Candida*, *Candida albicans* (yeast); soybean, *Glycine max*; *P* falcip, *Plasmodium falciparum*; *Tetth*, *Tetrahymena thermophila*; *Tetpy*, *T. pyriformis*; *T* brucei, *Trypanosoma brucei*; *Oxytricha*, *Oxytricha nova*; *Euplotes*, *Euplotes crassus*. Details of sequences and references obtained from the OWL database [57] will be provided on request (see footnote). Actins listed as human are common to the following species; skel: mouse, rat, rabbit and chicken; card: mouse, chicken; smth A: mouse, rat, cow; smth G: mouse, rat; cyt G: mouse, rat, cow; cyt B: mouse, rat, cow, chicken.

Most species have several actin genes [60–71] with the largest numbers occurring in plants [68,69]. These multigene families encode different isoforms but in *Physarum polycephalum* [60], the protozoan *Taenia solium* [80] and *Caenorhabditis elegans* [71] multiple actin genes encode the same amino acid sequence. A number of lower eukaryotes, e.g. *Aspergillus nidulans* [72], *Saccharomyces cerevisiae* [73–75], and the protozoan *Tetrahymena thermophila* [76] have only a single actin gene, though some also contain genes for 'actin-like' proteins (see below). In metazoans actin multigene families have been maintained throughout evolution with a relatively low rate of change compared to many other proteins. Frequently members of actin gene families are more conserved between than within species. For example, rice actin 1 is more similar to *Arabidopsis* actin than any other rice actin [77]. The two carrot actins (dicot) are

closer to maize (monocot) than to soybean (dicot) [78], suggesting that the formation of different actin gene families predated the divergence of monocots and dicots [55,56]. Especially when actins from different species are more similar than within a multigene family the possibility of isoform-specific functional differences must be considered.

Muscle actins from several chordate species form a group distinct from their cytoplasmic actins, suggesting that the emergence of separate muscle and cytoplasmic actins predated chordate speciation [58]. Indeed, vertebrate cytoplasmic actins are more similar to arthropod actins than to vertebrate muscle actins [55], indicating a very ancient common origin for actin genes [59]. Within arthropods, *Drosophila melanogaster* (Diptera) and *Bombyx mori* (Lepidoptera) muscle actins form a group distinct from their cytoplasmic actins [59]. *Artemia* (crustacean) actins

Figure 2 Ribbon and cylinder diagram of the actin structure, redrawn from [51]

The numbers indicate residues at either end of α -helices (cylinders) or β -sheet strands (arrows). A molecule of ATP is indicated by the wire diagram and the metal atom as a sphere. The orientation is the same as that shown in Figures 3(a) and 3(b). Note: the loop formed by amino acids 241–247 should be in front of the cylinder (α -helix) formed by amino acids 203–216.

divide between these two groups and it will be interesting to determine if these actins are muscle or cytoplasmic specific. Conserved amino acids that define different groups have been proposed for vertebrates [58], insect muscle [59] and plants [79]. The functional significance of these amino acids remains to be tested.

Intriguingly, the amino acids at which mammalian skeletal muscle actins differ from β - and γ -actins occur in three-dimensional clusters in the atomic structure [2]. The largest cluster, involving amino acids 1–6, 103, 129, 357 and 354, occurs in the lower right hand corner of subdomain 1 (Figures 2, 3a and 3b, coloured red). This region contains the proposed binding site of many actin binding proteins, including myosin [1,2]. Variations in this region may be adaptations to optimize the binding of different actin-binding proteins. Other clusters occur in the central β -sheet of subdomain 3 (residues 153, 162, 176 and 298) and in the region between subdomains 3 and 4 (residues 225, 259, 266 and 271). Subdomains 2 and 4 are conserved [2]. Of the nine amino acid substitutions which define the insect muscle group of actins, only four occur at positions included in those which define the mammalian skeletal muscle actins [59] and only Ile-76

is conserved in both groups. The insect muscle-specific substitutions do not cluster in the actin structure.

Although most actins are highly conserved in sequence and length (374–376 amino acids), proteins with more divergent sequences, known collectively as 'actin-like' proteins, have been identified recently in several species [80–83]. In addition to its single actin gene, *ACT1*, *Saccharomyces cerevisiae* contains the *ACT2* gene which encodes a protein of 391 amino acids that is 53.6% divergent from *ACT1* actin [80]; *Schizosaccharomyces pombe* has an *ACT2* protein (427 amino acids) which is 52.6% diverged from its *ACT1* actin [81]. The two *ACT2* proteins are 64.8% diverged from each other. Genetic experiments show that *ACT1* cannot substitute for *ACT2*, or vice versa, arguing that both genes are essential for cell viability and have different functions [73,80,81]. Two identical actin-like sequences, with an identity of about 55% compared to standard actins, but only 40% identity to the *ACT2* genes of *S. cerevisiae* and *S. pombe*, have been isolated and studied in man [82] and dog [83]. These sequences, named respectively actin-RPV (related protein vertebrate) and contractin, are associated with microtubule based cellular motor systems. Actin-RPV is a component of the

dynactin complex [82] and cetractin is associated with the centrosomes [83]. Actin-like proteins have also been reported in *Aspergillus*, *Physarum*, *Caenorhabditis* and the prokaryotic cyanobacter *Spirulina platensis* [80]. Actin-like proteins are probably ubiquitous and the evidence so far indicates that they have different functions from standard actins.

In proteins with a conserved core of residues, the structure of one protein is usually a reasonable model of the main structural features of other members of the group [84,85]. Superimposition of the actin-RPV/cetractin sequence onto the rabbit actin structure shows that regions corresponding to the actin core are more conserved [83]; similarly, the increased length of the two *ACT2* proteins derives from insertions in positions corresponding to surface loops in the rabbit actin structure [51,80,81]. This contrasts with substitutions in standard actins which occur with equal frequency in surface loops and other parts of the molecule. Like all actins, the 'actin-like' proteins conserve a region corresponding to the ATP binding site [51,80,81].

Comparison of actin isoform function

Functional differences between isoforms have been sought both in intact cells and by biochemical analysis [86]. Actin isoforms segregate into different structures within cells [87–94], suggesting they have specialized functions. Purified isoforms, and actins from different species, can differ in thermodynamic stability [31,95] and polymerization characteristics [32,96–99]. Actins from different species also differ in their affinity for profilin [100,101], DNase I [102–104], α -actinin and tropomyosin [104].

In binding assays of non-muscle actins [96,106–108] or rabbit muscle actin isoforms [91,109–111] with various rabbit muscle myosins, significant differences in affinities were found. The higher affinity of skeletal muscle actin compared to gizzard actin for both gizzard and enteric myosins suggests that the higher affinity is an intrinsic property of the actin, rather than the myosin [112]. In all cases the $V_{\max}$ for the actin-activated myosin ATPase was the same. Only in comparisons of actin and myosin from scallop and rabbit skeletal muscle was a difference in $V_{\max}$ found: homologous systems gave higher $V_{\max}$ values, but no difference in the affinity of each actin for either myosin [113]. Since such comparisons are often conducted in non-physiological conditions and any differences can often be eliminated by varying pH or ionic strength [111], it is unclear if any of these *in vitro* differences are relevant to the situation *in vivo* [111]. Furthermore, proteins which interact with actin will have coevolved [86,113]. Thus functional differences found with heterologous mixtures of actins and other proteins are probing not just the function of actin but also that of the other protein isoforms as well.

The physiological significance of isoform differences can be tested by substituting one actin for another in a whole cell, plant or animal. A mutation in the promoter sequence, which regulates expression, of the mouse cardiac actin gene reduces its expression to 16.5% of its normal level [114]. There is a corresponding increase in expression of the α -skeletal muscle actin gene. Despite containing predominantly the skeletal muscle actin isoform the heart functions normally, suggesting that the two isoforms are functionally equivalent. However, isoform substitution was incomplete and the low levels of cardiac actin may have been sufficient for any isoform-specific functions. Isoform substitution can be achieved by genetic transformation with cloned genes. Although no difference was found in the incorporation of cardiac and cytoplasmic actin isoforms into the cytoskeleton of cultured non-muscle cells [115], transformation of adult rat cardiomyocytes with different actin isoforms showed isoform-specific

effects on both their incorporation into the myofibrils and cell morphology [116]. In *Drosophila melanogaster*, of nine mutations encoding single amino acid substitutions in the *Act88F* actin gene corresponding to isoform-specific amino acid differences, only two of the mutations altered muscle structure [117]. However, two of three mutant *Act88F* gene constructs with multiple isoform-specific amino acid changes produced altered muscle structure, suggesting that as more changes are made, actins become functionally nonequivalent. Undoubtedly substitution of a complete actin isoform is desirable since it will contain multiple actin amino acid changes, which have co-evolved. Despite only 90% similarity the chicken β -actin is able to substitute for the yeast *ACT1* actin without loss of viability, although the cells exhibited an altered morphology, slower growth and temperature-sensitive lethality [118].

Functional isoform differences between some members of actin gene families offers a partial explanation for the evolution of actin multigene families. Although comparisons of isoform sequences and functions are a valuable source of information about actin, we need to look further than the study of naturally occurring actins to unravel the role of individual amino acids in actin function.

Actin processing

Actins are post-translationally modified. Most actins have a methylation of His-73 and modification of the N-terminus by removal of one or two residues followed by acetylation of the new N-terminus [3]. Actins are unusual in the type of amino acids which occur at the N-terminus [119] and the processing enzyme is specific for actin [34].

Methylation of His-73 occurs in all actins examined, except for that from *Naegleria gruberi*, whose actin binds DNase I less well than other actins and has a lower affinity compared to rabbit actin for scallop myosin S1 fragment but the same $V_{\max}$ for the actin-activated myosin ATPase activity [120]. His-73 is not close to the DNase I-binding loop [51], though both are in subdomain 2 (see Figures 2 and 3d). It is therefore unclear if these differences are due to a lack of methylation or to other amino acid differences in *Naegleria* actin. Substitution of His-73 with Arg or Tyr in human skeletal muscle actin expressed *in vitro* had no effect on the ability of the actins to co-polymerize or bind DNase I [28]. However, it is uncertain whether wild-type actin expressed *in vitro* and used as a control was methylated [32,121]. The same mutant actins, transformed into simian COS-1 cells, had no effect on cell growth [28]. No effects of His-73 methylation on actin functions are known.

Inhibition of N-terminal processing *in vitro* can alter actin polymerization [32]. However, yeast cells normally lack part of the N-terminal processing activity [38] and completely blocking all N-terminal processing does not affect the stability of actins expressed in yeast [38]. Non-yeast actins expressed in yeast are not fully processed yet polymerize *in vitro* [38,39]. Actin from *A. castellani* is also unprocessed at its N-terminus *in vivo* [122]. Indeed, although it is frequently inferred from sequence data that all actins can be N-terminally processed, this has been confirmed experimentally to occur *in vivo* in only a few cases.

The *Drosophila* flight muscle-specific *Act88F* actin undergoes two further modifications. First, the *mod⁺* gene, which is not an actin gene, encodes a modifier which alters the charge of the actin [45] yet *mod⁻* flies, which lack this modification, are fully flighted and have a wild-type wingbeat frequency [32]. Second, a proportion of the *Act88F* actin forms a stable 1:1 conjugate with ubiquitin, called arthrin [123]. The roles of these actin modifications are unknown. In general, despite some small changes

detected *in vitro*, no clear functions have emerged for post-translational processing.

ARTIFICIALLY INDUCED ACTIN MUTANTS

Mutants obtained by selection for flightlessness or uncoordinated movement have led to the recovery of actin mutants in *Drosophila melanogaster* [17–22] and *Caenorhabditis elegans* [23,24]. Serendipitous actin mutants have also been recovered in a melanoma [124] and transformed tissue culture cells [125]. However, these are inefficient ways to recover new actin mutants. First, mutants of other proteins often produce the same or similar phenotype. Second, severe actin mutations are often lethal, limiting selection of actin mutants to systems where the requirement for a specific isoform is not vital, or where conditional mutations, such as temperature-sensitive alleles, can be obtained. Finally, many cells and organisms express multiple actins so recovery of mutants may be difficult.

Protein engineering approaches remove many of these problems and, by *in vitro* mutagenesis of cloned actin genes, permit any amino acid substitution to be made at any point in actin. The only limits are our ability to formulate specific questions about actin at the level of individual amino acids. After mutagenesis the actin must be expressed. This has not been easy, in part because all eukaryotic cells contain actin leading to problems of separating mutant from endogenous actins. Only recently has the expression and purification of mutant actins in sufficient quantities for *in vitro* studies been achieved (see below).

An advantage of expression for *in vivo* analysis is that the mutant actin is in a normal environment where its ability to assemble into cellular structures and interact with a whole range of actin-binding proteins may be tested. However, it may be difficult to pinpoint which actin interactions are affected. For such studies it is useful if the mutant actin gene is the only one expressed since the expression of other actin isoforms may mask the effects of the mutation.

In the following section we describe the systems used for expression and study of actin mutants *in vivo*. Each has advantages and disadvantages but together they provide a wide choice for investigating different aspects of actin structure and function.

Drosophila melanogaster

The fruitfly genome contains six actin genes; two encode cytoplasmic actins and the remaining four are muscle-specific [62]. Mutations have been recovered only in the *Act88F* actin gene which is expressed specifically in the indirect flight muscles [62,126] and encodes all the muscle actin found in these tissues [see 123,127 for review]. *Act88F* mutations affect flight ability but not the viability of the flies, so the *Act88F* gene provides a unique system for studying the effects of actin mutations on muscle assembly and function.

Apart from *Act88F* mutants recovered as flightless flies following whole fly mutagenesis, others have been made by *in vitro* mutagenesis and inserted into the fly genome by P-element transformation [20,33,45–49]. In flies, the effects of the mutations can be analysed by flight testing, by microscopy of the flight muscles and by mechanical testing of muscle fibres [47,128]. Partially purified *Act88F* mutant actins have been used in biochemical studies [129].

Most *Act88F* mutations show a dominant flightless phenotype. This can be explained by a requirement for two wild-type gene copies for flight. However, there are many examples where this is clearly not the case. For instance, flies containing a mutant

gene and two wild-type copies may still be flight defective and show aberrations of the myofibrillar filament lattice [20–22, 33,48]. This is a common feature of actin mutants in other organisms as well as *Drosophila*, and is known as ‘antimorphism’ or ‘dominant negative complementation’. The usual explanation is that the mutant monomers interfere or ‘poison’ assembly of wild-type monomers into F-actin by retaining some capacity to bind wild-type monomers. Antimorphism thus indicates that the actin mutation has not destroyed all actin binding sites. The importance of actin polymerization for biological function undoubtedly explains the frequent antimorphism of actin mutants (Table 1) in flies and other systems. The incorporation of perhaps only one mutant monomer may be sufficient to ‘cap’ and prevent growth of an actin filament even in the presence of a large excess of wild-type monomers. The ease of genetically manipulating the gene copy number in organisms such as yeast and *Drosophila* means that antimorphic effects can be identified and studied, providing valuable information on mutant actin function *in vivo*.

Heat-shock proteins, a ubiquitous cellular response to stress and damage to proteins [130,131] are induced constitutively in *Drosophila* by certain *Act88F* mutants [19,20,33,126]. Actin is one of very few proteins for which heat-shock-protein-inducing mutants have been found [132]. Intriguingly, actin has some structural similarity with hsc70 (heat-shock cognate 70kDa protein) [133,134].

Caenorhabditis elegans

This nematode has been extensively used for molecular genetic studies of muscle development and function. Many muscle protein genes were identified by mutations with phenotypic effects of paralysis, twitching or uncoordinated movements [24,135]. The genome contains four actin genes but only regulatory mutants of these genes have been characterized to date [71]. The actin genes have been cloned and with a gene transformation system [136] *C. elegans* may provide another model system for the *in vivo* study of actin mutations.

Dictyostellium discoideum

The genetics of myosin and other cytoskeletal proteins is well developed in this slime mould [137–141]. No actin mutants have been recovered by selecting for abnormal phenotypes, probably because the genome contains at least 17 actin genes [67]. This system has been used to express and purify large quantities of actin following *in vitro* mutagenesis of the *Dictyostellium Act15* gene [44]. Charge changes were introduced into the *Act15* gene by site-directed mutagenesis to facilitate separation of the mutant from the endogenous actins.

Saccharomyces cerevisiae

The yeast *S. cerevisiae* provides a powerful system for the study of actin mutants and cell biology. Many mutations of the single actin gene, *ACT1*, are now known (see Table 1), although none were recovered by their phenotypic effects. *ACT1* mutants made by *in vitro* mutagenesis are easily transformed into the yeast genome and the phenomenon of homologous recombination can be used to disrupt the endogenous wild-type *ACT1* copy while inserting the mutant gene copy into the yeast genome [15,73,142]. This requires that the mutant alleles permit cell survival.

Mutant actins can be obtained from *S. cerevisiae* in sufficient quantities for biochemical studies either by replacing the wild-type *ACT1* copy [38,40] or using yeast expression vectors to express heterologous actins e.g. chicken β -actin [37,39] in cells

Table 1 Actin mutants

Table 1 summarizes current actin mutants residue-by-residue from the N- to C- termini. The actin gene in which each mutation occurs, the system used to express it and any mutant effects are given. The mutants are named using the one letter code with the original amino acid first, its number, then the new amino acid; 'Term' denotes a translation stop codon; *in vitro*, expression in rabbit reticulocyte lysate; *ACT1*, yeast actin gene; *Act88F*, *Drosophila* flight-muscle specific gene; *Act15*, an actin gene from *Dictyostelium*, Hs α and Hs β , human α -skeletal and β -actin genes; and Ch β , chicken β -actin gene; pure, actin was purified after expression; NDG, mobility on non-denaturing gels; thermodynamic stability, as determined on urea gradient gels; =, as well as wild-type actin; >, better than wild-type actin; <, less than wild-type actin; (+), normal; (-), abnormal; structure, muscle appearance by light or electron microscopy; V_{max} , for myosin ATPase; K_m , for the binding of actin to myosin; ATPase, myosin ATPase; motility, velocity in the *in vitro* motility assay; mechanics, mechanical response of dissected myofibrils; DNase (DNase I), ATP, S1 (S1 fragment of myosin), profilin or tropomyosin, binding to each ligand relative to wild-type actin; F-actin, production of actin filaments; processing, N-terminal processing; heat, cold sensitive, yeast's inability to grow at high or low temperatures; antimorphic, mutant actin preventing wild-type actin from functioning normally *in vivo*; hypomorphic, produces normal actin but at a reduced level; hsp, induces heat-shock protein synthesis; ADP-ribosylation, actin is a substrate for *Clostridium* toxin ADP-ribosylating activity.

Position	Expression	Gene	Results	Reference
COD,N,H	<i>in vitro</i>	Hs α	(-) Processing	34
COS,G,F,Y	<i>in vitro</i>	Hs α	Not processed	34
del0	<i>in vitro</i>	Hs α	(+) Processed	30
D1H, D1H/D4H	<i>Dictyostelium</i> , pure	<i>Act15</i>	< Motility; < V_{max} ; < ATPase	44
del1-12	<i>in vitro</i>	<i>Act88F</i>	> Profilin; = ATP; = DNase; zero ADP-ribosylation	35,147
D2N/E4Q	Yeast, pure	<i>ACT1</i>	< ATPase; = polymerization	40
	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth; = processing; (-) F-actin	40
del2-4	Yeast, pure	<i>ACT1</i>	< ATPase; = polymerization	40
	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth; (-) processing; (-) F-actin	40
D2V	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	142
D2A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive cold, heat sensitive; < spore viability	42
D3A,H,N	<i>in vitro</i>	Hs α	= Polymerization; = S1; = DNase; = NDG	26
D3N	<i>in vitro</i>	Hs α	(-) Processed	34
del3-4, D3K/D4K, D3A/D4A	Yeast, pure	Ch β	= DNase; = polymerization; (-) S1	39
del3-4, D3A/D4A	Yeast, pure	Ch β	< ATPase; < velocity; = tropomyosin	156
D3K/D4K	Yeast, pure	Ch β	< ATPase; no velocity; = tropomyosin	156
E4V,A	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	42,142
D4E	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(+) Structure	117
G6A	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(-) Structure	117
G6A/A7T	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Flightless; (-) structure	46
I10V	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(+) Structure	117
D11E,N, D11N/N12D	<i>in vitro</i>	Hs α	= Polymerization; = S1; < DNase; < NDG; < processing	29
D11H	<i>in vitro</i>	Hs α	< Polymerization; = S1; < DNase; < NDG; < processing	29
D11Q,K	Yeast, <i>in vivo</i>	<i>ACT1</i>	Dominant lethal	142
D11A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Partial dominant lethal	42
D11E	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth; viable spores	178
D11N	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth; no viable spores	178
D24H/D25H	<i>Dictyostelium</i> , pure	<i>Act15</i>	Zero motility	193
D24A/D25A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive cold, heat sensitive	42
R28C	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Strong antimorph; weak hsp induction	17,19
R28L	Cells	Mouse A ^x	Causes melanoma cell-line	50
P32L	Yeast, <i>in vivo</i>	<i>ACT1</i>	Temperature sensitive	15
G36E/E83D	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(-) Structure	49
G36E/E83D/G245D	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(+) Structure; flighted; hsp induction	49
	Cells	Hs β	(+) Morphology; < incorporation into structures; < DNase	25
R37A/R39A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive cold, heat sensitive	42
K50A/D51A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive cold, heat sensitive	42
D56A/E57A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive heat sensitive; = cold sensitive	42
A58T	Yeast, <i>in vivo</i>	<i>ACT1</i>	Temperature sensitive	15
K61A/R62A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Partial dominant lethal	42
R68A/E72A	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	42
H73Y	<i>in vitro</i>	Hs α	< Polymerization; = DNase; = processing	28
H73Y,R	Cells	Hs α	(+) Growth	28
H73R	<i>in vitro</i>	Hs α	= Polymerization; = DNase; = processing	28
I76F	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Strongly antimorphic; weak hsp induction; (-) Z discs	17,19,117
T79Term	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	No actin accumulation	21
D80A/D81A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive cold, heat sensitive	42
E83A/K84A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive cold, heat sensitive	42
E93K	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(-) Structure; no Z-discs	22
E93A/R95A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive lethal	42
E99H/E100H	<i>Dictyostelium</i> , pure	<i>Act15</i>	< Motility	193
E99A/E100A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive heat sensitive; = cold	42
R116A/E117A/K118A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive heat sensitive; = cold	42
S129V	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(+) Structure	117
V139M	Cells, <i>in vivo</i>	β -actin	Partially responsible for cytochalasin resistance	185
D154A/D157A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Partial dominant lethal	42
F169Y/A260S	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(-) Structure	117
F169Y/C257T	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(+) Structure	117
L176M	<i>in vitro</i>	<i>Act88F</i>	> Profilin; = ATP; = DNase; = NDG; (+) ADP-ribosylation	35,147
R177Q	<i>in vitro</i>	<i>Act88F</i>	> Profilin; = ATP; = DNase; = NDG; zero ADP-ribosylation	35,147
R177A/D179A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive heat sensitive; = cold	42

R183A/D184A	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	42
D187A/K191A	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	42
K191M, K191M/C374A	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	142
E195A/R196A	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	42
E205A/R206A/E207A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Possibly dominant lethal	42
R210A/D211A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive weak heat sensitive; = cold	42
K213A/E214A/K215A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive cold; heat sensitive; low spore viability	42
D222A/E224A/E226A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive heat sensitive; = cold	42
T234S	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(+) Structure	117
E237A/K238A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Partial dominant lethal	42
E241A/D244A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Partial dominant lethal	42
G245D	Cells	Hs β	< Polymerization	178
	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Antimorphic	48
	Cells	Hs β	Stable; (-) morphology at high expression levels	25
	Yeast, pure	Ch β	< Polymerization; < velocity; = S1; > K_m	151
G245K	Yeast, pure	Ch β	< Polymerization; = S1; = velocity	151
E253A/R254A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Partial dominant lethal	42
R256A/E259A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive cold; heat sensitive	42
del269-272	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	179
E270A/D275A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive lethal	42
V278T	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(+) Structure	117
D286A/D288A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive lethal	42
I289F	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Hypomorphic; myofibrillar degeneration	49
R290A/K291A/E292A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive lethal	42
A295D	Cells, <i>in vivo</i>	β -actin	Partially responsible for cytochalasin resistance	185
T304G/M305T/Y307Term	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Unstable; hsp inducing	21
E311A/R312A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive cold, heat sensitive; < spore viability	42
E316K	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Flightless; almost = muscles; (-) mechanics; weak hsp induction	33,42
E316K	<i>in vitro</i>	<i>Act88F</i>	= Polymerization; = NDG; < thermodynamic stability	31,33
			> Profilin; = ATP; = DNase	35
K326A/K328A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Possibly dominant lethal	42
E334K	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(-) Structure; weak hsp induction	33
E334K	<i>in vitro</i>	<i>Act88F</i>	= Polymerization; = NDG; = thermodynamic stability	31,33
			> Profilin; = ATP; = DNase	35
E334A/R335A/K336A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive lethal	42
K336M,Q	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	142
V339I	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	No actin accumulation	33
V339I	<i>in vitro</i>	<i>Act88F</i>	= Polymerization; = NDG; < thermodynamic stability	31,33
			> Profilin; = ATP; = DNase	35
Y356Term	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Antimorph, some actin (?); strong hsp; nuclear swelling	18,21
W356Y,H	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	142
K359A/E361A	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	42
Q360E	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(+) Structure	117
E360H/E361H	<i>Dictyostelium</i> , pure	<i>Act15</i>	= Motility	193
E361A/D363N	<i>in vitro</i>	<i>Act88F</i>	> Profilin; = ATP; = DNase; = NDG	35
Y362C,I	<i>in vitro</i>	<i>Act88F</i>	> Profilin; < ATP; < NDG; = DNase	35
Y362V,L	<i>in vitro</i>	<i>Act88F</i>	< Profilin; < ATP; < NDG; = DNase	35
Y362F	<i>in vitro</i>	<i>Act88F</i>	= Profilin; = ATP; = NDG; = DNase	35
D363H/E364H	<i>Dictyostelium</i> , pure	<i>Act15</i>	= Motility	193
D363Y	<i>in vitro</i>	<i>Act88F</i>	= Profilin; = ATP; = NDG; = DNase	35
D363H	<i>in vitro</i>	<i>Act88F</i>	> Profilin; = ATP; = NDG; < DNase	35
D363A/E364A	Yeast, <i>in vivo</i>	<i>ACT1</i>	Recessive heat sensitive; = cold	42
E364K	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(-) Structure; strong hsp induction	33
E364K	<i>in vitro</i>	<i>Act88F</i>	= Polymerization; = thermodynamic stability; < NDG	31,33
			> Profilin; < ATP; < DNase	35
G366S	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Antimorphic; strong hsp induction	20,21
G366D	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(-) Structure; strong hsp induction	33
G366D	<i>in vitro</i>	<i>Act88F</i>	= Polymerization; = thermodynamic stability; < NDG	31,33
G366D,P,A	<i>in vitro</i>	<i>Act88F</i>	> Profilin; < ATP; = DNase	35
G368S	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	(+) Structure	117
G368E	<i>Drosophila</i> , <i>in vivo</i>	<i>Act88F</i>	Flightless; almost (+) muscles; (-) mechanics	47
G368E	<i>in vitro</i>	<i>Act88F</i>	= Polymerization; = NDG; = thermodynamic stability	31,33
			= Profilin; < ATP; = DNase	35
G368T,S,Q	<i>in vitro</i>	<i>Act88F</i>	= Profilin; = ATP; = DNase; = NDG	35
G368K	<i>in vitro</i>	<i>Act88F</i>	> Profilin; = ATP; = DNase; = NDG	35
G368Term	<i>in vitro</i>	<i>Act88F</i>	> Profilin; < ATP; = DNase; = NDG	35
R372H	<i>in vitro</i>	<i>Act88F</i>	= Polymerization; = NDG; = thermodynamic stability	31,33
			> Profilin; > ATP; = DNase	35
K373Term	Yeast, <i>in vivo</i>	<i>ACT1</i>	Lethal	142
C374A	Yeast, <i>in vivo</i>	<i>ACT1</i>	(+) Growth	142
C374Term	Yeast, <i>in vivo</i>	<i>ACT1</i>	Temperature sensitive	142
P375Term	Yeast, <i>in vivo</i>	<i>ACT1</i>	Temperature sensitive	142

Figure 3 Three-dimensional structure of actin

Coloured ribbon diagram and spacefilling models of actin drawn using the coordinates of the actin–DNase I crystal structure [51] and displayed using the Quanta molecular graphics program (Molecular Simulations Inc.). Only the actin is shown. (a) and (b) show a ribbon diagram and spacefilling model respectively in which the four structural subdomains are shown in different colours. Subdomain 1 (residues 1–32, 70–144 and 338–375) is in red; subdomain 2 (residues 33–69) is yellow; subdomain 3 (residues 145–180 and 270–337) is blue; subdomain 4 (residues 181–269) is green. In (b) the isolated Ca^{2+} at the top right, shown in orange, is bound to DNase I and has been included to orientate the view. The ATP is shown in pink. Panels (c), (d) and (e) show the distribution of polarity on the 'front', 'back' and 'right-hand side' faces of actin respectively. The amino acid residues are coloured by polarity. Red, acidic; blue, basic; white, nonpolar; yellow, polar. Also shown are: pale green, ATP; orange, Ca^{2+} ; purple, methylhistidine-73. The isolated Ca^{2+} from the DNase I is shown in pale green. Panel (f); the interior of actin around amino acids Asp-11 and Val-339. The atoms are coloured by type: red, oxygen; blue, nitrogen; green, carbon; pink, metal ion; yellow, phosphate. Arrows point to both Asp-11 and Val-339 and the approximate orientation is looking from the left-hand side of (e). Panel (g), a surface view of Tyr-362 and Phe-124. Atoms are coloured by type: red, oxygen; blue, nitrogen; green, carbon. Arrows point to both Tyr-362 and Phe-124 and the approximate orientation is looking towards (e). Panel (h), a surface view around amino acid Gly-245. The atoms are coloured by type: red, oxygen; blue, nitrogen; green, carbon. An arrow points to Gly-245. Approximate orientation is looking directly down on the top of the molecule. Panel (i), a surface view of amino acids 269–272. The atoms are coloured by type: red, oxygen; blue, nitrogen; green, carbon; yellow, sulphur. Arrows point to Met-269, Glu-270 and Ala-272. Orientation is as in (d). Panel (j), a surface view of Gly-366. The atoms are coloured by type: red, oxygen; blue, nitrogen; green, carbon. The approximate orientation is looking towards (e). Arrows point to Gly-366 and Ser-368.

with a wild-type genomic *ACT1* copy. Yields are limited in the latter system because actin overexpression is lethal [143] and must be regulated to maintain levels of the transformed actin equivalent to the endogenous actin [37,39]. The lack of N-terminal processing of actin in yeast [38] may be a disadvantage in the study of non-yeast actins.

Cultured cells

Cultured human fibroblasts [25,26,50] and simian COS-1 cells [28] have been transformed with mutant actin genes to look for effects on their growth and morphology. Partially purified preparations of a mutant actin from human fibroblasts have been used for *in vitro* polymerization studies [26]. Insect cells are commonly used to express foreign proteins following baculovirus transfection [144] but do not yet appear to have been used for expression of mutant actins.

Escherichia coli

The *Dictyostelium Act8* gene has been expressed in *E. coli* [145,146] but the actin forms inclusion bodies and only a small amount can be extracted in its native state. This is no doubt related to the ease with which native actin is irreversibly denatured. A cytoplasmic chaperone protein recently isolated from rabbit reticulocyte lysate catalyses the folding of denatured β -actin expressed in *E. coli* [147]. This may solve the problems of expressing native actin in *E. coli* for biochemical studies and allow suitable mutants to be used to study actin folding.

Expression *in vitro*

The expression of actin mutants from cloned and mutagenized DNA by *in vitro* transcription and translation in rabbit reticulo-

cyte lysate has been used to produce small quantities of radioactive actin. Although it may not be fully processed [32,121], such actin behaves the same as the product of the same gene expressed *in vivo* [32]. Mutant actins have been expressed from the genes for human α -skeletal muscle actin [28–30], rat brain cytoplasmic actin [27] and *Drosophila Act88F* [31–33,35,36,148] and used in co-polymerization and protein binding studies.

Although the costs are prohibitive for making large amounts of actin this way, expression is quickly and easily achieved and binding assays can be used to screen rapidly large numbers of mutations and pinpoint those of interest for more detailed study.

ACTIN MUTANTS

Table 1 summarizes the mutants residue-by-residue from the N- to C- termini, listing their effects *in vivo* and *in vitro*. We will consider actin functions in turn, describing the effects of mutants with reference to the actin structure where possible. Some mutants inevitably affect more than one function and will be found under more than one heading.

The atomic structure of actin [1,51] is an essential framework for interpreting mutant effects and is shown in Figures 2 and 3(a)–3(j). By comparing this structure with the extensive biochemical data from studies of actin by chemical modification, chemical crosslinking, spectroscopy etc. and electron microscope reconstructions of F-actin, F-actin–tropomyosin and filaments decorated with the myosin S1 fragment [149], potential binding sites can now be identified with some certainty [1,2,51].

Actin has four structural subdomains [51] (Figures 2, 3a and 3b). Subdomains 1 and 2 form what was previously known as the 'small domain'; subdomains 3 and 4 the 'large domain'. The chemical nature of amino acid side chains on the 'front', 'back' and 'right-hand side' surfaces of actin are shown in Figures 3(c), 3(d) and 3(e). These illustrate several interesting features. The

DNase I binding loop (top right in Figure 3c and top left in Figure 3d) contains several non-polar residues which form a hydrophobic contact with DNase I [51]. Long 'stripes' of non-polar residues in the middle of the front of the molecule suggest possible contact sites with other proteins (Figure 3c). There are also regions of stacked negative and positive charges visible on both the front and back surfaces. The N- and C-termini in subdomain 1 contain a cluster of negative charge (bottom right in Figure 3c and bottom left in Figure 3d). Methylhistidine-73 is located between two acidic groups and close to the conserved residue at Arg-177 (Figure 3d). The ATP is almost completely buried in the cleft between the so-called large and small domains, with only the purine ring visible (Figure 3c). Since there is ATP/ADP exchange some conformational changes must occur. Any opening of the cleft would involve breaking the salt bridge between Arg-62 and Glu-207 which links subdomains 2 and 4 (Figure 3c). Actin is a disc-like molecule and the side views present a very small aspect (Figure 3e).

STUDIES IN VITRO

Polymerization

In F-actin there are actin-actin contacts between the four structural subdomains [1,2,150] and it is not surprising to find extensive parts of the monomer surface involved in these contacts. From the model of F-actin [1,150] the contact sites have been identified as amino acids 41–50 (the DNase I loop), 110–112, 166–169, 195–197, 243–245, 264–273, 286–289, 322–325 and 375. The actin sequence alignment does not indicate that these sites are more highly conserved than other regions of the sequence. Only Gln-111, Gly-168, Asp-244, Gly-245, Asp-288, Arg-290 and Pro-322 are invariant. The conserved residues, with only occasional conservative substitutions, are Met-44, Pro-112, Asp-286, Ile-287, Ile-289, Lys-291, Thr-324, Met-325, Lys-326 and Phe-375. In the light of this it may seem surprising that actins from different species co-polymerize. However, this alignment is based on a large number of actins from a much broader range of species than have been used for co-polymerization studies. From their phenotypic effects *in vivo* it has been proposed that the alanine substitutions at residues 37, 39, 61, 62, 205, 244, 286 and 288 in the yeast *ACT1* gene affect the actin-actin contact surfaces [41,42] although these are not all within the contact sites proposed by modelling F-actin.

Two different actins containing the same G245D mutation both had reduced ability to polymerize [26,151] as did the G245K mutation [151]. Gly-245 is involved in actin-actin contacts [1,151] so the deleterious effects of these substitutions on polymerization are not unexpected.

Mutant actins, expressed *in vitro*, can be tested for co-polymerization with non-mutant carrier actin, added to achieve the critical concentration for polymerization. Although the interpretation of these experiments has been questioned [36], a number of mutants are able to co-polymerize as well as wild-type actin expressed *in vitro* [28,29,33,36] (Table 1). None of these mutants were in actin-actin contact sites.

Although there is no evidence of the N-terminus forming actin contacts in F-actin [53,150,152–154], complete N-terminal processing appears essential for normal polymerization [32]. Replacement of Asp-11 by non-acidic amino acids in a human α -skeletal muscle actin expressed *in vitro* prevented N-terminal processing, but in only one case did the actin have reduced ability to polymerize [29]. The lack of any polymerization effects of mutants at Asp-11 is surprising given their effect on protein conformation [155] and the involvement of this amino acid with metal binding (see below). The contradictory evidence for a role

of N-terminal processing in polymerization might arise from the use of heterologous or homologous actins for the co-polymerizations in the different studies [32]. Since there is no biochemical evidence for a direct role of the N-terminus in polymerization, these effects probably arise from a conformational change which affects a distant actin-actin contact site.

Motility *in vitro* and myosin binding

The ability of mutant actins to function as part of the actomyosin motor have been studied by mechanics experiments on demembrated muscle fibres of *Drosophila* mutants [47], the use of *in vitro* motility assays [39–44,151,156] and assays of the actin-activated Mg-ATPase of myosin or myosin fragments [129,151,156,44]. The *in vitro* motility assays use purified actin and myosin which can be obtained from both muscle and non-muscle systems. By fixing one protein, usually myosin or myosin fragments, to a surface and adding F-actin stabilized with phalloidin and made fluorescent with a rhodamine label the velocity of the filaments can be measured, which is believed to correspond to the V_{max} of myosin ATPase [157].

Electron microscopic reconstruction of S1-decorated actin [149], the G- and F-actin structures [51,150] and biochemical data from many laboratories have been used to locate the myosin binding site, or at least that of the rigor state, on the outer face of subdomain 1 (Figures 2 and 3) [1,2]. The imminent publication of the atomic structure of the myosin S1 fragment means that modelling of actin and myosin S1 structures to the constraints of the electron microscopic reconstructions of S1-actin is now possible and the details of their interaction should soon be known.

Subdomain 1 is formed by residues 1–32, 70–144 and 338–372. While actin with the first 47 amino acids cleaved from the N-terminus is still able to bind myosin and activate its ATPase activity [154] albeit with reduced *in vitro* motility [153], the N-terminal peptide alone (amino acids 1–47) can also activate myosin ATPase [158,159]. The N-terminus forms contacts with myosin heavy chain [160,161] in the presence of nucleotide [161]. The proposed binding site includes the positions of known myosin-binding residues (in rabbit skeletal muscle) of Asp-1, Glu-2, Asp-3, Glu-4, Asp-24, Asp-25, Arg-28, Glu-93, Arg-95 and Lys-336. The hydrophobic surface of the α -helix 338–348 is probably a major contact site but due to the chemical nature of the residues these interactions would be refractory to most crosslinking chemicals. Crosslinking [162] and n.m.r. [163] studies implicate the 360–364 region in the binding of the A1 light chain from the vertebrate skeletal myosin. Actin mutants have been used to examine the importance of acidic groups at the N-terminus in myosin binding, a particularly variable region in different actin isoforms. Removal of acidic residues at positions within the first four amino acids reduced myosin S1 binding, myosin ATPase activation and *in vitro* motility [39,40,44,193]. Interestingly, substitutions with histidine at E360/E361 and D363/E364 resulted in wild-type motility suggesting the presence of acidic groups at the C-terminus is not necessary for motility [193]. This is surprising given the close proximity of the N-terminus.

Substitution of Gly-245 with aspartic acid in chicken β -actin gave an actin which bound S1 (decoration of F-actin filaments) but had a slower velocity in the motility assay and reduced affinity for myosin [151]. Replacement with lysine resulted in an actin with normal S1 binding and *in vitro* motility [151]. There is no evidence from biochemical studies of myosin-binding sites in this region, suggesting that the mutations cause long-range conformational effects on the whole actin molecule.

G-actin binding proteins

DNase I and profilin are members of a large group of proteins which bind to G-actin and prevent it from polymerizing [3].

DNase I

All actins tested bind DNase I with the exceptions of those from *Entamoeba histolytica* [102] and *Tetrahymena thermophila* [104]. From the atomic structure of the actin–DNase I complex the contacts between the proteins are well established [1,51]. Amino acids Thr-203 and Glu-207 (subdomain 4) form the minor contact site; amino acids 39–46 and 61–64 (subdomain 2) the major contact site. The *E. histolytica* and *T. thermophila* actins contain Thr-203 and Glu-207, but substitutions in amino acids 39–46 in these actins may explain the absence of DNase I binding. Although the DNase I binding of many actin mutants has been tested [28,29,35,39], only replacements at D11 [29], G36E/E83D/G245D [25], D363 and E364 [35] reduced DNase I binding. Since the positions of the mutant amino acids do not coincide with the contact sites the effects are probably due to conformational changes [29]. Actin with residues 1–15 deleted had normal DNase I binding [35] and the different results obtained with this and the D11 mutants suggest that distortion caused by non-aspartate residues at position 11 is more disruptive than removal of the first 15 amino acids. The triple mutant G36E/E83D/G245D may have a direct effect since amino acid 36 is particularly close to the DNase I binding site. However, it is unknown if the G36E mutation alone is sufficient to reduce binding and given the disruptive effects of Gly-245 mutations, reduced DNase I binding may not be surprising.

Profilin

Biochemical crosslinking studies have shown that profilin binds to the C-terminal region of actin in subdomain 1 [164]. Deletion of amino acids 1–12 of actin increased profilin binding [35] and this could be due to a direct effect on the profilin binding site at the C-terminus given the close proximity of the N- and C-termini [51]. Changes at positions 176, 177 and 316–372 increased profilin binding whilst others at position 362 had decreased affinity for profilin [35]. The mutations between 362 and 372 are in, or close to, the profilin binding site [164] so this probably explains their altered profilin binding. Other positions are distant from the profilin binding site so their effects must produce long-range effects. Since all these mutants had wild-type binding to DNase I (Table 1), any conformational change must be localized and this is supported by the failure to detect conformational changes in these mutants on non-denaturing gels [31].

ATP binding and actin conformation

ATP or ADP binds in the cleft between the large and small actin domains [51] and makes a number of different contacts. Subdomain 1 contacts the metal ion and the phosphates of ATP [51]. The ATP or ADP is closely associated with a divalent cation (Mg^{2+} in native actin, Ca^{2+} in the actin–DNase I crystals) and the binding of both ATP and the metal ion are essential in maintaining actin's conformation [165–168]. The divalent cation is enclosed by the β - and γ -phosphates of ATP, and the charged residues Asp-11, Asp-154, Lys-18 and Gln-137. The phosphates themselves bind the β -bends formed by residues 14–16 and 157–159. Lys-18 forms a complex with the oxygens of the α - and β -phosphates. The adenine is held in a hydrophobic pocket formed between subdomains 3 and 4. Residues 302–308 form

one side of this pocket, with Gly-302 hydrogen bonded to the α -phosphate. Tyr-306 closes the back of the pocket and the upper part is formed by Glu-214 and Lys-213. Our alignment of actin sequences (D. R. Drummond, unpublished work) shows that of the amino acids forming or close to the ATP-binding site residues 18, 156–159, 213–214, 300–301 and 304–305 are invariant; amino acids 14, 212, 215, 302, 306 are conserved or have conservative substitutions.

The relative contribution of each contact to the binding of ATP can only be determined experimentally. Only one study has mutated amino acids directly involved in ATP and metal ion binding [29]. Although ATP binding was not measured directly, replacement of Asp-11 by non-acidic groups caused a conformational change in the actin. Since Asp-11 binds to the metal ion [51] (Figure 3f) which complexes with the ATP and is required for actin stability, the effects of Asp-11 mutants may be explained by reduced metal-ion binding.

Another study has looked directly at the ability of mutants to bind ATP [35]. Although none of the substitutions were in ATP contact sites [51], several mutants had reduced ATP binding and intriguingly one had increased binding. This suggests that these mutants affect ATP binding via a change in actin structure and indeed, all mutants with reduced ATP binding also had altered conformations [31,33,35]. Of five substitutions at position Tyr-362, only replacement by Phe had normal conformation and ATP binding suggesting that an aromatic group is required at this position. The flat aromatic ring of Tyr-362 is stacked against the ring of Phe-124 (Figure 3g) and this might explain the inability of other amino acids to substitute at these positions.

ADP-ribosylation

Actin is a substrate for various ADP-ribosylating toxins from *Clostridium* strains [169–171]. Once modified, the actin is prevented from polymerizing [169,172] and G-actin ATPase activity is blocked [173]. Protein chemistry identified Arg-177 as the site of ADP-ribosylation [174,175]. Toxins from *Clostridium botulinum* and *Clostridium perfringens* differ in their substrate specificity (see [176]) and it has been proposed that Leu-176 [175] or the N-terminus [177] might be responsible for this specificity.

Wild-type *Drosophila* Act88F actin can be ribosylated *in vitro*, whilst mutant R177Q actin is not, confirming Arg-177 as the site of modification [148]. Mutant L176M actin was modified by all toxins, indicating that this residue does not define substrate specificity. An actin in which the first 12 amino acids were deleted was not ribosylated by any toxin, implying that this region may be important for recognition by the toxins.

STUDIES IN VIVO

Cell biology

Actin mutants will be essential in elucidating the roles of actin and actin-binding proteins in all aspects of cell biology. The involvement of actin in a great variety of structures and functions explains the variety of phenotypic effects within a single cell type and between different cell types.

Mutants of the yeast *ACT1* gene produce a number of phenotypes. Many are lethal, temperature-sensitive for growth or spore production [15,38,41,42,142,178,179]. Dominant lethal mutations have been detected [42] and must be antimorphic. Some temperature-sensitive mutants show, at non-permissive temperatures, an aberrant distribution of actin filaments which causes a uniform deposition of chitin at the cell surface rather than at the bud sites, accumulation of secretory vesicles and cell enlargement [16]. Such effects have shown that actin is involved

in vectorial protein transport and bud formation. In addition *ACT1* mutants may also show osmosensitivity [16].

Deletion of residues 269–272 in *ACT1* had no effect on yeast growth [179] despite a proposal that they form a surface loop which stabilizes an adjacent α -helix [51]. The atomic structure of this region is shown in Figures 2 and 3(i). Deletion of the C-terminus (Lys-Cys-Phe-COOH) is lethal to yeast while removal of Phe-375 or Cys-374/Phe-375 produces a temperature-sensitive phenotype [142], indicating the importance of the C-terminus. These residues are not represented in the actin structure [51].

The interpretation of the cellular effects of mutant actins depends on biochemical data. In yeast, since actins can be readily purified in the quantities necessary for biophysical and biochemical assays such information should be easily obtained. For instance, proposals that many of the mutations made by alanine scanning [41,42] are in residues close to putative actin–actin contacts (see above) [1] should be easily testable.

Actin mutants are responsible for a melanoma phenotype in mice [124] and the tumorigenic potential of an *in vitro* transformed human fibroblast line [25,26,50]. The mutant β -actin, G245D, responsible for the transformation of human fibroblasts, was stable and caused alterations in cell shape, motility and distribution of cytoskeletal actin [180,181]. These effects are probably due to the decreased polymerization ability of the mutant actin [25]. Transformation of cells with the triple mutant G36E/E83D/G245D did not affect cell morphology, perhaps because the actin was less readily incorporated into cell structures than wild-type actin [25]. While actin mutations are uncommon in neoplasia, effects on actin expression and the cytoskeleton are usual following tumorigenic transformation [182–184].

One actin double mutant could be useful in the study of the action of cytochalasin, a fungal toxin which is known to inhibit actin polymerization in normal cells. Cells containing V139M/A295D actin are resistant to cytochalasin [185]. The substituted amino acids are neither adjacent nor are they in known actin–actin contact sites. Mutants such as V139M/A295D may also be useful in studying the processes of actin polymerization and depolymerization.

Muscle assembly

Although many actin mutants appear to have normal functions when expressed *in vitro* or *in vivo* in yeast, this is rarely the case when they are expressed in muscle. In *Drosophila* all mutants of the *Act88F* gene recovered by their flightless phenotype result in complete absence of muscle or disrupted structure [22,46,117] (Table 1). This might be accounted for if selection for flightless mutants enriches for actin mutations with a strong phenotypic effect. However, most mutants derived by *in vitro* mutagenesis have similar effects, though a minority affect myogenesis only slightly, especially when the single amino acid substitutions correspond to *Drosophila* actin isoform differences [33,46,47,117].

Since the *Act88F* actin gene is expressed specifically in the indirect flight muscles [62,126], encodes all the muscle actin found in these muscles [123] and a lack of function does not affect fly viability, its mutations have proved especially useful for studies of myofibril assembly. *Act88F* mutants can be divided into at least five phenotypic groups [186]. There are null mutants which produce no actin, mutants with low actin accumulation, mutants which accumulate actin but form no myofibrils, mutants producing myofibrils with no Z-discs and those giving normal or near normal myofibrillar structure. It is difficult to identify the precise effects of these mutants on actin structure and function due to the complex situation *in vivo* and, until recently, the

difficulty in obtaining sufficient actin for biochemical studies. One study has investigated the effects observed *in vivo* using *in vitro* methods. A range of actin mutants, with different *in vivo* effects, all showed the same ability to polymerize [33] or bind to *Drosophila* myofibrils *in vitro* [36] as wild-type: the differences seen *in vivo* were not reproduced *in vitro*, though doubts have been expressed about what was actually being measured in these experiments [36]. However, electron microscopy (M. Stark and J. C. Sparrow, unpublished work) has shown that all the mutants used [33] make thin filaments *in vivo* although to very different degrees. Experiments *in vitro* may reflect the situation *in vivo*, but may not be sensitive enough to detect subtle, biologically significant differences between mutant actins. F-actin forms the core of thin filaments and actin mutants which do not form myofibrils will frequently affect actin polymerization. However, the interaction of actin with other proteins, such as the tropomyosins and troponins, must be important for thin filament assembly. Tropomyosin has been implicated in stabilizing F-actin *in vivo* [187,188]. The myofibrils of the *Act88F* mutant E93K have thin filaments but no Z-discs and it has been proposed that this mutant may reduce binding of a Z-disc protein, perhaps α -actinin [22]. Biochemical data [189] suggests there is an α -actinin binding site close to amino acid 93 in rabbit actin which may include residues 112–125 [190].

Some *Act88F* mutants show both a strong heat shock response and a strong antimorphic effect on muscle structure or function [17,18,20,21,33,49,126] but this is not always the case [19,49]. The common features of heat shock-inducing actin mutants are reduced ATP binding [35] and the presence of multiple conformers on non-denaturing gels [29], which suggests that the binding of heat shock proteins to the mutant actins [33] *in vivo* is a response to localized changes in conformation. This effect may be quite specific since most strong heat shock-inducing mutations found so far cluster around amino acids 364–366. Indeed, thermodynamic stability of mutant actins, measured by denaturation with urea *in vitro* [31], does not correlate with muscle structure, antimorphism or heat shock induction.

Muscle function

The development of techniques with which to perform mechanical experiments on isolated, demembrated fibres from the indirect flight muscles of *Drosophila* [47,128] made it possible to study the effects of actin mutants on muscle contraction, providing alternative information to that increasingly available from the *in vitro* motility assays. However, a prerequisite for such mechanical experiments is *Act88F* mutants which produce normal, or near normal, muscle structure. To date this criterion has been satisfied by only a small number of mutants. These include the chimaeric actin genes made by replacing *Act88F* sequences with corresponding regions from other actin genes (E. A. Fyrberg, personal communication) and single amino acid substitutions. The latter include isoform-specific changes [46,117] and the novel substitutions G368E and E316K [33,47].

The G368E and E316K *Act88F* gene mutations produce normal or near normal muscle structure but alter the kinetics of the crossbridge cycle [47]. Homozygous G368E flies fly poorly with a reduced wingbeat frequency; mechanical experiments on skinned flight muscles [47] show a reduced rate constant for the delayed tension increase, a distinctive feature of this type of muscle. Actin purified from G368E mutant flies had reduced S1 binding [129] which may be explained by the proximity of amino acid 368 to the myosin light chain binding site [162,163]. The E316K mutant does not fly but mechanical experiments on skinned fibres show an increased rate constant for delayed

tension rise [47]. This mutation lies within subdomain 3, precluding a direct effect on myosin binding, but the proximity of the tropomyosin binding site [1,2] suggests that E316K may have an effect on thin filament activation. The E93K mutation was originally isolated as a dominant flightless mutation [22] and produces myofibrils containing parallel arrays of thick and thin filaments, which lack Z-discs and any other sarcomeric pattern. Fibre mechanics, electron microscopy and proteolytic protection of the myosin head by actin binding all performed under rigor conditions were used to show that the E93K mutant actin can bind myosin heads in rigor. This was perhaps surprising given the proximity of Glu-93 to the myosin binding site. Unfortunately, the aberrant mutant muscle structure of E93K flight muscles prevents meaningful mechanical studies.

CONCLUSIONS AND FUTURE PROSPECTS

There is great interest in actin both because of its dynamic involvement in many aspects of cell biology and because of its role as part of a major protein motor system. Our understanding of many subcellular processes and muscle contraction will depend upon knowing much more about the relationships between the amino acid sequences of actins, their structure and function. Actin is remarkable for the diversity of its biochemistry. Few other proteins have binding sites for such a large range of proteins in addition to having enzymic activity. Undoubtedly many actin-binding proteins share binding sites and this will have acted to constrain allowable variation, reducing the rate and range of actin evolution.

The current sequence database contains 103 different actin gene sequences, representing 81 different protein sequences [57]. The high sequence similarity should permit the amino acid substitutions to be modelled within the atomic structure. These, together with our increasing certainty about the location of specific binding sites on actin from structural and biochemical studies, will provide useful pointers to the functional significance of variation. For many of these actins only the DNA sequences are known and the biochemical properties of the proteins have not been studied. Functional comparisons between such isoforms related to their three-dimensional homology could provide very useful information and lead to a more detailed analysis in which amino acid changes are made in some of the isoform-specific residues.

A large number of mutations have already been made in actin (Table 1). What has been learnt overall? Perhaps the most surprising feature for such a highly conserved protein is that it appears remarkably robust. Most mutations, even non-conservative amino acid substitutions, retain actin structure and function. This is important for genetic analysis, since it suggests that despite the high sequence conservation and the well known lability of actin *in vitro*, it is possible with mutants to explore one part of the molecule without destroying others. Of course, as found in protein engineering studies of other proteins, there are already examples of mutants in one part of the molecule having long-range effects. Such effects are important since they raise questions about the dynamic structure of the molecule which are not apparent from the static picture provided by X-ray crystallography. The fact that actin contains multiple binding sites will permit different parts of a mutant actin to be readily monitored for such long-range conformational changes.

The phenotypic effects of actin mutants are diverse, deriving from the different systems used and the variety of actin functions. However, the highly conserved nature of actin means that mutants with effects in one system are likely to have effects in another. This has been exploited so far for only a small number

of mutants. For example, the G245D mutation oncogenically transforms human fibroblasts [26], disrupts *Drosophila* indirect flight muscle development and function [48] and affects actin movement in the *in vitro* motility assay [151]. There is clearly much to be gained from studying both the existing and new mutants in a multiplicity of systems.

Genetic and molecular biological approaches have much to offer in understanding not only actin, but also in analysing the biological processes in which it is involved. It is premature to assess what contributions the genetic studies of actin can make to our understanding of actin function, since only recently have the major advances been made that will permit a directed protein engineering assault on questions relating to actin sequence and function. These advances include the resolution of the atomic structure of G-actin [51], the modelling of F-actin [150] and the development of systems in which cloned genes mutated by site-directed mutagenesis can be expressed to produce mutant actins in sufficient quantities for biochemical studies. For the first time deductions from the atomic structure and biochemical data can be used to ask testable questions about the role of specific amino acids. Initially this is likely to focus on those interactions for which atomic structures are available. In addition novel assay systems, such as *in vitro* motility and mechanical experiments on muscle fibres from flies transformed with actin mutants, are now available with which to study actin's role in movement and contraction. Many of the elements are now in place to take the genetical analysis of actin function into exciting areas of actin biochemistry and cell biology.

We thank Professor Peter A. Rubenstein, Professor Kazuo Sutoh, Dr Roger Karlsson and Dr Ken Wertman for generously providing prepublication results and reprints; Dr Leslie Leinwand, Dr Eric A. Fyrberg and Mary C. Reedy for sending reprints; Dr Rod Hubbard and Stuart Priest for help with the molecular graphics; Susan Sparrow for Figure 2; Pete Crosby for photography; Miss Anne Lawn and Frank Katzer for helping to prepare this paper and Dr Simon Hardy, Dr Phil Hunt, Dr Simon Watkins, Dr David White, Jon Clayton and Azam Razzaq for critical reading of the manuscript and many helpful suggestions. This work benefitted from the use of the SEQNET computing facility at SERC Daresbury. The authors were supported during this time by SERC research grants.

REFERENCES

- Holmes, K. C. and Kabsch, W. (1991) *Curr. Opin. Struct. Biol.* **1**, 270–280
- Kabsch, W. and Vandekerckhove, J. (1992) *Annu. Rev. Biophys. Biomol. Struct.* **21**, 49–76
- Pollard, T. D. and Cooper, J. A. (1986) *Annu. Rev. Biochem.* **55**, 987–1035
- Hartwig, J. H. and Kwiatkowski, D. J. (1991) *Curr. Opin. Cell Biol.* **3**, 87–97
- Matsudaira, P. (1991) *Trends Biochem. Sci.* **16**, 87–92
- Carlier, M.-F. (1991) *J. Biol. Chem.* **266**, 1–4
- Korn, E. D., Carlier, M.-F. and Pantaloni, D. (1987) *Science* **238**, 638–644
- Korn, E. D. (1982) *Physiol. Rev.* **62**, 672–737
- Bray, D. (1992) *Cell Movements*, Garland Publishing Inc., New York and London
- Pollard, T. D. (1990) *Curr. Opin. Cell Biol.* **2**, 33–40
- Lassing, I. and Lindberg, U. (1985) *Nature (London)* **314**, 472–474
- Masters, C. (1984) *J. Cell Biol.* **99**, 222s–225s
- Cooke, R. (1990) *Curr. Opin. Cell Biol.* **2**, 62–66
- Geeves, M. A. (1991) *Biochem. J.* **274**, 1–14
- Shortle, D., Novick, P. and Botstein, D. (1984) *Proc. Natl. Acad. Sci. U.S.A.* **81**, 4889–4893
- Novick, P. and Botstein, D. (1985) *Cell* **40**, 405–416
- Mogami, K. and Hotta, Y. (1981) *Mol. Gen. Genet.* **183**, 409–417
- Karlik, C. C., Couto, M. D. and Fyrberg, E. A. (1984) *Cell* **38**, 711–719
- Karlik, C. C., Saville, D. L. and Fyrberg, E. A. (1987) *Mol. Cell. Biol.* **7**, 3084–3091
- Hiromi, Y., Okamoto, H., Gehring, W. J. and Hotta, Y. (1986) *Cell* **44**, 293–301
- Okamoto, H., Hiromi, Y., Ishikawa, E., Yamada, T., Isoda, K., Maekawa, H. and Hotta, Y. (1986) *EMBO J.* **5**, 589–596
- Sparrow, J., Reedy, M., Ball, E., Kyrtatas, V., Molloy, J., Durston, J., Hennessey, E. and White, D. (1991) *J. Mol. Biol.* **222**, 963–982
- Kenyon, C. (1988) *Science* **240**, 1448–1453
- Waterston, R. H. (1989) *Cell Motil. Cytoskel.* **14**, 136–145

- 25 Leavitt, J., Ng, S.-Y., Aebi, U., Varma, M., Latter, G., Burbeck, S., Kedes, L. and Gunning, P. (1987) *Mol. Cell Biol.* **7**, 2457–2466
- 26 Taniguchi, S., Sagara, J. and Kakunaga, T. (1988) *J. Biochem. (Tokyo)* **103**, 707–713
- 27 Saborio, J. L. and Palmer, E. (1981) *J. Neurochem.* **36**, 1659–1669
- 28 Solomon, L. R. and Rubenstein, P. A. (1987) *J. Biol. Chem.* **262**, 11382–11388
- 29 Solomon, T. L., Solomon, L. R., Gay, L. S. and Rubenstein, P. A. (1988) *J. Biol. Chem.* **263**, 19662–19669
- 30 Sheff, D. R. and Rubenstein, P. A. (1989) *J. Biol. Chem.* **264**, 11491–11496
- 31 Drummond, D. R., Hennessey, E. S. and Sparrow, J. C. (1991) *Biochem. J.* **274**, 301–303
- 32 Hennessey, E. S., Drummond, D. R. and Sparrow, J. C. (1991) *Eur. J. Biochem.* **197**, 345–352
- 33 Drummond, D. R., Hennessey, E. S. and Sparrow, J. C. (1991) *Mol. Gen. Genet.* **226**, 70–80
- 34 Sheff, D. R. and Rubenstein, P. A. (1992) *J. Biol. Chem.* **267**, 2671–2678
- 35 Drummond, D. R., Hennessey, E. S. and Sparrow, J. C. (1992) *Eur. J. Biochem.* **209**, 171–179
- 36 Hennessey, E. S., Harrison, A., Drummond, D. R. and Sparrow, J. C. (1992) *J. Muscle Res. Cell Motil.* **13**, 127–131
- 37 Karlsson, R. (1988) *Gene* **68**, 249–257
- 38 Cook, R. K., Sheff, D. R. and Rubenstein, P. A. (1991) *J. Biol. Chem.* **266**, 16825–16833
- 39 Aspenström, P. and Karlsson, R. (1991) *Eur. J. Biochem.* **200**, 35–41
- 40 Cook, R. K., Blake, W. T. and Rubenstein, P. A. (1992) *J. Biol. Chem.* **267**, 9430–9436
- 41 Wertman, K. and Drubin, D. G. (1992) *Science* **258**, 759–760
- 42 Wertman, K., Drubin, D. G. and Botstein D. (1992) *Genetics* **132**, 337–350
- 43 Kron S., Drubin, D. G., Botstein, D. and Spudich J. A. (1992) *Proc. Natl. Acad. Sci. U.S.A.* **89**, 4466–4470
- 44 Sutoh, K., Ando, M., Sutoh, K. and Toyoshima, Y. Y. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **88**, 7711–7714
- 45 Mahaffey, J. W., Coutu, M. D., Fyrberg, E. A. and Inwood, W. (1985) *Cell* **40**, 101–110
- 46 Reedy, M. C., Beall, C. and Fyrberg, E. (1989) *Nature (London)* **339**, 481–483
- 47 Drummond, D. R., Peckham, M., Sparrow, J. C. and White, D. C. S. (1990) *Nature (London)* **348**, 440–442
- 48 Sakai, Y., Okamoto, H., Mogami, K., Yamada, T. and Hotta, Y. (1990) *J. Biochem. (Tokyo)* **107**, 499–505
- 49 Sakai, Y., Okamoto, H., Mogami, K., Matsuo, H. and Hotta, Y. (1991) *J. Biochem. (Tokyo)* **109**, 670–673
- 50 Sadano, H., Taniguchi, S., Kakunaga, T. and Baba, T. (1988) *J. Biol. Chem.* **263**, 15868–15871
- 51 Kabsch, W., Mannherz, H. G., Suck, D., Pai, E. F. and Holmes, K. C. (1990) *Nature (London)* **347**, 37–44
- 52 Schutt, C. E., Lindberg, U., Myslik, J. and Strauss, N. (1989) *J. Mol. Biol.* **209**, 735–746
- 53 Forné, T. and Chaussepied, P. (1992) *J. Muscle Res. Cell Motil.* **13**, 233
- 54 Mannherz, H. G., Gooch, J., Way, M., Weeds, A. G. and McLaughlin, P. J. (1992) *J. Mol. Biol.* **226**, 899–901
- 55 Hightower, R. C. and Meagher, R. B. (1986) *Genetics* **114**, 315–332
- 56 Meagher, R. B. (1991) *Int. Rev. Cytol.* **125**, 139–163
- 57 Bleasby, A. J. and Wootton, J. C. (1990) *Protein Eng.* **3**, 153–160
- 58 Miwa, T., Manabe, Y., Kurokawa, K., Kamada, S., Kanda, N., Bruns, G., Ueyama, H. and Kakunaga, T. (1991) *Mol. Cell. Biol.* **11**, 3296–3306
- 59 Mounier, N., Gouy, M., Mouchiroud, D. and Prudhomme, J. C. (1992) *J. Mol. Evol.* **34**, 406–415
- 60 Adam, L., Laroche, A., Barden, A., Lemieux, G. and Pallotta, D. (1991) *Gene* **106**, 79–86
- 61 Tobin, S. L., Zulauf, E., Sanchez, F., Craig, E. A. and McCarthy, B. (1980) *Cell* **19**, 121–131
- 62 Fyrberg, E. A., Mahaffey, J. W., Bond, B. J. and Davidson, N. (1983) *Cell* **33**, 115–123
- 63 Haymer, D. S., Anleitner, J. E., He, M., Thanaphum, S., Saul, S. H., Ivy, J., Houtchens, K. and Arcangeli, L. (1990) *Genetics* **125**, 155–160
- 64 Mounier, N. and Prudhomme, J. C. (1991) *Insect Biochem.* **21**, 523–533
- 65 Vandekerckhove, J. and Weber, K. (1978) *J. Mol. Biol.* **126**, 783–802
- 66 Sawtell, N. M., Hartman, A. L. and Lessard, J. L. (1988) *Cell Motil. Cytoskel.* **11**, 318–325
- 67 Vandekerckhove, J. and Weber, K. (1980) *Nature (London)* **284**, 475–477
- 68 Meagher, R. B. and McLean, B. G. (1990) *Cell Motil. Cytoskel.* **16**, 164–166
- 69 Baird, W. V. and Meagher, R. B. (1987) *EMBO J.* **6**, 3223–3235
- 70 Campos, A., Bernard, P., Fauconnier, A., Landa, A., Gómez, E., Hernández, R., Willms, K. and Laclette, J. P. (1990) *Mol. Biochem. Parasitol.* **40**, 87–94
- 71 Krause, M., Wild, M., Rosenzweig, B. and Hirsh, D. (1989) *J. Mol. Biol.* **208**, 381–392
- 72 Fidel, S., Doonan, J. H. and Morris, N. R. (1988) *Gene* **70**, 283–293
- 73 Shortle, D., Haber, J. E. and Botstein, D. (1982) *Science* **217**, 371–373
- 74 Gallwitz, D. and Sures, I. (1980) *Proc. Natl. Acad. Sci. U.S.A.* **77**, 2546–2550
- 75 Ng, R. and Abelson, J. (1980) *Proc. Natl. Acad. Sci. U.S.A.* **77**, 3912–3916
- 76 Cupples, C. G. and Pearlman, R. E. (1986) *Proc. Natl. Acad. Sci. U. S. A.* **83**, 5160–5164
- 77 McElroy, D., Rothenberg, M., Reece, K. S. and Wu, R. (1990) *Plant Mol. Biol.* **15**, 257–268
- 78 Stranathan, M., Hastings, C., Trinh, H. and Zimmerman, J. L. (1989) *Plant Mol. Biol.* **13**, 375–384
- 79 Nairn, C. J., Winsett, L. and Ferl, R. J. (1988) *Gene* **65**, 247–257
- 80 Schwob, E. and Martin, R. P. (1992) *Nature (London)* **355**, 179–182
- 81 Lees-Miller, J. P., Henry, G. and Helfman, D. M. (1992) *Proc. Natl. Acad. Sci. U.S.A.* **89**, 80–83
- 82 Lees-Miller, J. P., Helfman, D. M. and Schroer, T. A. (1992) *Nature (London)* **359**, 244–246
- 83 Clark, S. W. and Meyer, D. J. (1992) *Nature (London)* **359**, 246–250
- 84 Lesk, A. M. and Chothia, C. (1986) *Philos. Trans. R. Soc. London B* **317**, 345–356
- 85 Chothia, C. and Lesk, A. M. (1986) *EMBO J.* **5**, 823–826
- 86 Rubenstein, P. A. (1990) *Bioessays* **12**, 309–315
- 87 Rubenstein, P., Ruppert, T. and Sandra, A. (1982) *J. Cell Biol.* **92**, 164–169
- 88 McKenna, N., Meigs, J. B. and Wang, Y.-L. (1985) *J. Cell. Biol.* **100**, 292–296
- 89 Denning, G. M., Kim, I. S. and Fulton, A. B. (1988) *J. Cell Sci.* **89**, 273–282
- 90 Otley, C. A., Kalnoski, M. H. and Bulinski, J. C. (1988) *Cell Motil. Cytoskel.* **9**, 337–348
- 91 Shires, A. K. and Rubenstein, P. A. (1989) *Cell. Motil. Cytoskel.* **14**, 263–270
- 92 DeNofrio, D., Hoock, T. C. and Herman, I. M. (1989) *J. Cell Biol.* **109**, 191–202
- 93 Skalli, O., Gabbiani, F. and Gabbiani, G. (1990) *Exp. Cell Res.* **187**, 119–125
- 94 Peng, I. and Fischman, D. A. (1991) *Cell Motil. Cytoskel.* **20**, 158–168
- 95 Strzelecka-Golaszewska, H., Venyaminov, S. Y., Zmorzynski, S. and Mossakowska, M. (1985) *Eur. J. Biochem.* **147**, 331–342
- 96 Gordon, D. J., Boyer, J. L. and Korn, E. D. (1977) *J. Biol. Chem.* **252**, 8300–8309
- 97 Strzelecka-Golaszewska, H., Zmorzynski, S. and Mossakowska, M. (1985) *Biochim. Biophys. Acta* **828**, 13–21
- 98 Próchniewicz, E. and Yanagida, T. (1981) *J. Biochem. (Tokyo)* **89**, 1215–1221
- 99 Khaitlina, S. Y. (1986) *FEBS Lett.* **198**, 221–224
- 100 Larsson, H. and Lindberg, U. (1988) *Biochim. Biophys. Acta* **953**, 95–105
- 101 Ohshima, S., Abe, H. and Obinata, T. (1989) *J. Biochem. (Tokyo)* **105**, 855–857
- 102 Meza, I., Sabanero, M., Cazares, F. and Bryan, J. (1983) *J. Biol. Chem.* **258**, 3936–3941
- 103 Fulton, C., Lai, E. Y., Lamoyi, E. and Sussman, D. J. (1986) *J. Protozool.* **33**, 322–327
- 104 Hirono, M., Kumagai, Y., Numata, O. and Watanabe, Y. (1989) *Proc. Natl. Acad. Sci. U.S.A.* **86**, 75–79
- 105 Woolley, D. E. (1972) *Arch. Biochem. Biophys.* **150**, 519–530
- 106 Gordon, D. J., Eizenberg, E. and Korn, E. D. (1976) *J. Biol. Chem.* **251**, 4778–4786
- 107 Uyemura, D. G., Brown, S. S. and Spudich, J. A. (1978) *J. Biol. Chem.* **253**, 9088–9096
- 108 Green, C. and Schekman, R. (1982) *Mol. Cell. Biol.* **2**, 1270–1278
- 109 Sheetz, M. P., Painter, R. G. and Singer, S. J. (1976) *Biochemistry* **15**, 4486–4492
- 110 Elce, J. S., Elbrecht, A. S. U., Middlestadt, M. U., McIntyre, E. J. and Anderson, P. J. (1981) *Biochem. J.* **193**, 891–898
- 111 Mossakowska, M. and Strzelecka-Golaszewska, H. (1985) *Eur. J. Biochem.* **153**, 373–381
- 112 Strzelecka-Golaszewska, H. and Sobieszek, A. (1981) *FEBS Lett.* **134**, 197–202
- 113 Hue, H. K., Labbé, J.-P., Harricane, M.-C., Cavadore, J.-C., Benyamin, Y. and Roustan, C. (1988) *Biochem. J.* **256**, 853–859
- 114 Garner, I., Minty, A. J., Alonso, S., Barton, P. J. and Buckingham, M. E. (1986) *EMBO J.* **5**, 2559–2567
- 115 Gunning, P., Ponte, P., Kedes, L., Hickey, R. J. and Skoultschi, A. I. (1984) *Cell* **36**, 709–715
- 116 Perriard, J.-C., von Arx, P., Bantle, S., Eppenberger, H. M., Eppenberger-Eberhardt, M., Messerli, M. and Soldati, T. (1993) *Soc. Exp. Biol. Symp.* **46**, 219–235
- 117 Reedy, M. C., Beall, C. and Fyrberg, E. (1991) *Biophys. J.* **59**, 187a
- 118 Karlsson, R., Aspenstrom, P. and Bystrom, A. S. (1991) *Mol. Cell Biol.* **11**, 213–217
- 119 Rubenstein, P., Smith, P., Deuchler, J. and Redman, K. (1981) *J. Biol. Chem.* **256**, 8149–8155

- 120 Sussman, D. J., Sellers, J. R., Flicker, P., Lai, E. Y., Cannon, L. E., Szent-Gyorgyi, A. G. and Fulton, C. (1984) *J. Biol. Chem.* **259**, 7349–7354
- 121 Redman, K. and Rubenstein, P. A. (1981) *J. Biol. Chem.* **256**, 13226–13229
- 122 Redman, K. L., Martin, D. J., Korn, E. D. and Rubenstein, P. A. (1985) *J. Biol. Chem.* **260**, 14857–14861
- 123 Ball, E., Karlik, C. C., Beall, C. J., Saville, D. L., Sparrow, J. C., Bullard, B. and Fyrberg, E. A. (1987) *Cell* **51**, 221–228
- 124 Taniguchi, S., Kawano, T., Kakunaga, T. and Baba, T. (1986) *J. Biol. Chem.* **261**, 6100–6106
- 125 Leavitt, J. and Kakunaga, T. (1980) *J. Biol. Chem.* **255**, 1650–1661
- 126 Hiromi, Y. and Hotta, Y. (1985) *EMBO J.* **4**, 1681–1687
- 127 Sparrow, J., Drummond, D., Peckham, M., Hennessey, E. and White, D. (1991) *J. Cell Sci. Suppl.* **14**, 73–78
- 128 Peckham, M., Molloy, J. E., Sparrow, J. C. and White, D. C. S. (1990) *J. Muscle Res. Cell Motil.* **11**, 203–215
- 129 Geeves, M. A., Drummond, D. R., Hennessey, E. S. and Sparrow, J. C. (1991) *Biophys. J.* **59**, 419a
- 130 Burdon, R. H. (1986) *Biochem J.* **240**, 313–324
- 131 Lindquist, S. and Craig, E. A. (1988) *Annu. Rev. Genet.* **22**, 631–677
- 132 Parker-Thornburg, J. and Bonner, J. J. (1987) *Cell* **51**, 763–772
- 133 Flaherty, K. M., McKay, D. B., Kabsch, W. and Holmes, K. C. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **88**, 5041–5045
- 134 Hightower, L. E. (1991) *Cell* **66**, 191–197
- 135 Anderson, P. (1989) *Annu. Rev. Genet.* **23**, 507–525
- 136 Fire, A. (1986) *EMBO J.* **5**, 2673–2680
- 137 DeLozanne, A. and Spudich, J. A. (1987) *Science* **236**, 1086–1091
- 138 Gerisch, G., Segall, J. E. and Wallraff, E. (1989) *Cell Motil. Cytoskeleton* **14**, 75–79
- 139 Noegel, A. A., Leitung, B., Witke, W., Gurniak, C., Harloff, C., Hartmann, E., Wiesmuller, E. and Schleicher, M. (1989) *Cell Motil. Cytoskeleton* **14**, 69–74
- 140 Manstein, D. J., Titus, M. A., DeLozanne, A. and Spudich, J. A. (1989) *EMBO J.* **8**, 923–932
- 141 Manstein, D. J., Ruppel, K. M., Kubalek, L. and Spudich, J. A. (1991) *J. Cell Sci. Suppl.* **14**, 63–65
- 142 Johannes, F.-J. and Gallwitz, D. (1991) *EMBO J.* **10**, 3951–3958
- 143 Huffaker, T. C., Hoyt, M. A. and Botstein, D. A. (1987) *Annu. Rev. Genet.* **21**, 259–284
- 144 Miller, L. K. (1988) *Annu. Rev. Microbiol.* **42**, 177–199
- 145 Frankel, S., Condeelis, J. and Leinwand, L. (1990) *J. Biol. Chem.* **265**, 17980–17987
- 146 Frankel, S., Sohn, R. and Leinwand, L. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **88**, 1192–1196
- 147 Gao, Y., Thomas, J. O., Chow, R. L., Lee, G.-H. and Cowan, N. J. (1992) *Cell* **69**, 1043–1050
- 148 Just, I., Hennessey, E. S., Drummond, D. R., Aktories, K. and Sparrow, J. C. (1993) *Biochem. J.* **291**, 409–412
- 149 Milligan, R. A., Whittaker, M. and Safer, D. (1990) *Nature (London)* **348**, 217–221
- 150 Holmes, K. C., Popp, D., Gebhard, W. and Kabsch, W. (1990) *Nature (London)* **347**, 44–49
- 151 Aspenstrom, P., Engkvist, H., Lindberg, U. and Karlsson, R. (1993) *Eur. J. Biochem.* **207**, 315–320
- 152 DasGupta, G., White, J., Phillips, M., Bulinski, J. C. and Reisler, E. (1990) *Biochemistry* **29**, 3319–3324
- 153 Schwyter, D. H., Kron, S. J., Toyoshima, Y. Y., Spudich, J. A. and Reisler, E. (1990) *J. Cell Biol.* **111**, 465–470
- 154 Kiessling, P., Schick, B., Polzar, B. and Mannherz, H. G. (1992) *J. Muscle Res. Cell Motil.* **13**, 263
- 155 Rubenstein, P. A., Solomon, L. R., Solomon, T. and Gay, L. (1989) *Cell Motil. Cytoskeleton* **14**, 35–39
- 156 Aspenstrom, P., Lindberg, U. and Karlsson, R. (1993) *FEBS Lett.* **303**, 59–63
- 157 Kron, S. J. and Spudich, J. A. (1986) *Proc. Natl. Acad. Sci. U.S.A.* **83**, 6272–6276
- 158 Kögler, H., Moir, A. J. G., Trayer, I. P. and Rüegg, J. C. (1991) *FEBS Lett.* **294**, 31–34
- 159 Van Eyk, J. E. and Hodges, R. S. (1991) *Biochemistry* **30**, 11676–11682
- 160 DasGupta, G., White, J., Cheung, P. and Reisler, E. (1990) *Biochemistry* **29**, 8503–8508
- 161 DasGupta, G. and Reisler, E. (1992) *Biochemistry* **31**, 1836–1841
- 162 Sutoh, K. (1982) *Biochemistry* **21**, 3654–3661
- 163 Trayer, I. P., Trayer, H. R. and Levine, B. A. (1987) *Eur. J. Biochem.* **164**, 259–267
- 164 Vandekerckhove, J. S., Kaiser, D. A. and Pollard, T. D. (1989) *J. Cell Biol.* **109**, 619–626
- 165 Asakura, S. (1961) *Arch. Biochem. Biophys.* **92**, 140–149
- 166 Kasai, M., Nakano, E. and Oosawa, F. (1965) *Biochim. Biophys. Acta* **94**, 494–503
- 167 Kasai, M. and Oosawa, F. (1968) *Biochim. Biophys. Acta* **154**, 520–528
- 168 Faustich, H., Merkler, I., Blackholm, H. and Stournaras, C. (1984) *Biochemistry* **23**, 1608–1612
- 169 Aktories, K., Bärmann, M., Ohishi, I., Tsuyama, S., Jakobs, K. H. and Habermann, E. (1986) *Nature (London)* **322**, 390–392
- 170 Aktories, K. and Just, I. (1990) in *ADP-Ribosylating Toxins and G-Proteins* (Moss, J. and Vaughan, M., eds), pp. 79–95, American Society for Microbiology, Washington DC
- 171 Aktories, K., Wille, M. and Just, I. (1992) *Curr. Top. Microbiol. Immunol.* **175**, 97–113
- 172 Schering, B., Bärmann, M., Chatwal, G. S., Geipel, U. and Aktories, K. (1988) *Eur. J. Biochem.* **171**, 225–229
- 173 Geipel, U., Just, I. and Aktories, K. (1990) *Biochem. J.* **266**, 335–339
- 174 Vandekerckhove, J., Schering, B., Bärmann, M. and Aktories, K. (1987) *FEBS Lett.* **225**, 48–52
- 175 Vandekerckhove, J., Schering, B., Bärmann, M. and Aktories, K. (1988) *J. Biol. Chem.* **263**, 696–700
- 176 Aktories, K. and Wegner, A. (1989) *J. Cell Biol.* **109**, 1385–1387
- 177 Mauss, S., Chapponnier, C., Just, I., Aktories, K. and Gabbiani, G. (1990) *Eur. J. Biochem.* **194**, 237–241
- 178 Cook, R. K. and Rubenstein, P. A. (1989) *J. Cell Biol.* **109**, 271a
- 179 Rubenstein, P. A., Cook, R. K. and Babcock, G. (1991) *J. Cell Biol.* **115**, 160a
- 180 Vandekerckhove, J., Leavitt, J., Kakunaga, T. and Weber, K. (1980) *Cell* **22**, 893–899
- 181 Leavitt, J., Bushar, G., Kakunaga, T., Hamada, H., Hirakawa, T., Goldman, D. and Merrill, C. (1982) *Cell* **28**, 259–268
- 182 Friedman, E., Verderame, S., Winawer, S. and Pollack, R. (1984) *Cancer Res.* **44**, 3040–3050
- 183 Goldstein, D. and Leavitt, J. (1984) *Cancer Res.* **45**, 3256–3261
- 184 Leavitt, J., Gunning, P., Kedes, L. and Jariwalla, R. (1985) *Nature (London)* **316**, 840–842
- 185 Ohmori, H., Toyama, S. and Toyama, S. (1992) *J. Cell Biol.* **116**, 933–941
- 186 Sparrow, J. C., Drummond, D. R., Hennessey, E. S., Clayton, J. D. and Lindegaard, F. B. (1993) *Soc. Exp. Biol. Symp.* **46**, 111–129
- 187 Hitchcock-deGregori, S. E., Sampath, P. and Pollard, T. D. (1988) *Biochemistry* **27**, 9182–9185
- 188 Weigt, C., Schoepper, B. and Wegner, A. (1990) *FEBS Lett.* **260**, 266–268
- 189 Mimura, N. and Asano, A. (1987) *J. Biol. Chem.* **262**, 4717–4723
- 190 Lebart, M.-C., Mejean, C., Roustan, C. and Benyamin, Y. (1993) *J. Biol. Chem.*, in the press
- 191 Higgins, D. G. and Sharp, P. M. (1989) *Comput. Appl. Biosci.* **5**, 151–153
- 192 Saitou, N. and Nei, M. (1987) *Mol. Biol. Evol.* **4**, 406–425
- 193 Johara, M., Toyoshima, Y. Y., Ishijima, A., Kojima, H., Yanagida, T. and Sutoh, K. (1993) *Proc. Natl. Acad. Sci. U.S.A.* **90**, 2127–2131