REVIEW ARTICLE

Interleukin-6 and the acute phase response

Peter C. HEINRICH,*§ José V. CASTELL† and Tilo ANDUS‡

*Institut für Biochemie der RWTH Aachen, Pauwelsstraße, 5100 Aachen, Federal Republic of Germany,

†Unidad de Hepatología Experimental, Centro de Investigación, Hospital La Fe, Avenida de Campanar 21, 46009 Valencia, Spain, and ‡Medizinische Universitätsklinik Freiburg, Hugstetter Straße 55, 7800 Freiburg, Federal Republic of Germany

INTRODUCTION

The acute phase response

The acute phase response is the answer of the organism to disturbances of its homeostasis due to infection, tissue injury, neoplastic growth, or immunological disorders. The acute phase response is thought to be beneficial to the injured organism with the aim of restoring the disturbed physiological homeostasis. A number of reviews have dealt in detail with the pathophysiological changes during the acute phase response [1–9].

As schematically shown in Fig. 1, it consists of a **local reaction** at the site of injury characterized by a number of responses such as aggregation of platelets and clot formation, dilation and leakage of blood vessels, and an accumulation and activation of granulocytes and mononuclear cells, which in turn release acute phase cytokines. In addition, activated fibroblasts and endothelial cells are able to produce cytokines. These mediators act on specific receptors on different target cells leading to a **systemic reaction** characterized by fever, leukocytosis, increase in erythrocyte sedimentation rate, increases in secretion of ACTH and glucocorticoids, activation of complement and clotting cascades, decreases in serum levels of iron and zinc, a negative nitrogen balance, and by dramatic changes in the concentration of some plasma proteins. These proteins are named **acute phase proteins**.

Acute phase proteins

The acute phase protein patterns vary from one species to another. C-reactive protein and serum amyloid A, for example, show the highest increases during an acute phase response in man, whereas in the rat, α_2 -macroglobulin and α_1 -acid glycoprotein are the acute phase proteins with the most spectacular changes (Table 1). In many species plasma levels of fibrinogen, haptoglobin, α_1 -antichymotrypsin and α_1 -antitrypsin increase during an acute phase response [1–9]; simultaneously, albumin and transferrin concentrations decrease. These changes in plasma levels of acute phase proteins are preceded by corresponding alterations in mRNA concentrations. In addition, increases in the rates of secretion of some acute phase proteins have been observed.

The earliest description of the acute phase response came from the ancient Greeks [10]. They already observed an increased sedimentation rate of erythrocytes in blood of severely ill patients. As we know today, this increase is due to elevated plasma concentrations of fibrinogen and other acute phase proteins [11]. The term ‘acute-

Fig. 1. Acute phase response of the organism

phase’ was introduced in 1941 by Abernethy & Avery [12], who described the properties of sera from patients with febrile infectious diseases. These sera contained an acute phase protein (C-reactive protein), which was induced during febrile infections.

Search for hepatocyte-stimulating factors

In 1951 it was shown by Miller *et al.* [13] that liver is the major organ for the synthesis of acute phase proteins. Since the acute phase proteins are synthesized in the liver and injury to another part of the body results in increased synthesis of acute phase proteins, the existence of hormone-like mediators was proposed. For many years investigators have been searching for such mediators. Since leukocytes are the cells first attracted at the sites of injury, they were examined as candidates for the synthesis and release of such mediators. Indeed, leukocyte super-

Table 1. Acute phase plasma proteins in human and rat

Species	Increase			No change	Decrease
	10–100-fold	2–10-fold	≤ 2-fold		
Human	C-reactive protein Serum amyloid A	α_1 -Proteinase inhibitor α_1 -Acid glycoprotein α_1 -Antichymotrypsin Fibrinogen Haptoglobin	Caeruloplasmin C3 of complement α_2 -Anti-plasmin C ₁ -inactivator	α_2 -Macroglobulin Haemopexin Serum amyloid P Prothrombin	Inter- α -antitrypsin Transferrin α_1 -Lipoprotein Prealbumin Albumin
Rat	α_2 -Macroglobulin α_1 -Acid glycoprotein	Fibrinogen Haptoglobin Cysteine proteinase inhibitor	α_1 -Proteinase inhibitor Caeruloplasmin Prekallikrein Haemopexin C-reactive protein	α_1 -Macroglobulin Antithrombin III Serum amyloid P Prothrombin	α_1 -Inhibitor 3 Transferrin Prealbumin Albumin

natants were shown to be able to stimulate the synthesis of numerous acute phase proteins in rats, rabbits and mice [14–16]. Among the leukocytes, monocytes/macrophages turned out to be the most important producers of mediators of the acute phase response [17]. Ritchie & Fuller described and partially characterized a monocyte-derived polypeptide involved in the regulation of acute phase protein synthesis, which they called hepatocyte-stimulating factor [18]. In the following years several laboratories attempted to purify and characterize the hepatocyte-stimulating factor produced by monocytes [19–29], rat Kupffer cells [30–32], human keratinocytes [19] and leukaemia cell lines [21,27]. The molecular mass of the hepatocyte-stimulating factor from human monocytes was estimated by gel chromatography to be 23–30 kDa [19–21,25–29]. An isoelectric point of approx. 5.4 has been determined [27]. The production of hepatocyte-stimulating factor by human monocytes was found to be regulated by lipopolysaccharide in a dose-dependent manner [26].

Discrimination of hepatocyte-stimulating factor from interleukin-1 and tumour necrosis factor α

Since human monocytes also release IL-1 and TNF α after stimulation with lipopolysaccharide, it was investigated whether these two monokines could act as hepatocyte-stimulating factors. The availability of recombinant human IL-1 and TNF α allowed such studies. Whereas the synthesis of rat haemopexin, complement C3, haptoglobin, α_1 -acid glycoprotein, albumin, murine serum amyloid A, and P, and human metallothionein was regulated by IL-1 [33–38,39], the synthesis of the majority of acute phase proteins could not be stimulated by IL-1 [20–22,26,28,29,34,36,38]. Recombinant human TNF α showed a very limited effect on a subset of acute phase proteins [28,29,34–36].

Identification of interleukin-6/B-cell stimulatory factor-2 as hepatocyte-stimulating factor

After the discrimination of IL-1 and TNF α from hepatocyte-stimulating factor, interferons were examined as potential hepatocyte-stimulating factors. Highly purified interferon- β contained hepatocyte-stimulating activity, absent from recombinant human interferon- β_1 , interferon- α , and recombinant human interferon- γ , indicating that hepatocyte-stimulating factor copurified, but was distinct from, interferon- β_1 [40]. Gauldie *et al.* [41]

reported that antibodies raised against interferon- β ($\beta_1 + \beta_2$) from fibroblasts inhibited hepatocyte-stimulating activity derived from human monocytes.

In 1986/87 interferon- β_2 was shown to be identical with B-cell stimulatory factor-2, hybridoma plasmacytoma growth factor, 26 kDa protein [42–44] and the name interleukin-6 was proposed for this molecule [45] (Table 2). Using recombinant B cell stimulatory factor-2, Gauldie *et al.* [41] presented the first evidence that hepatocyte-stimulating factor is identical with IL-6. BSF-2/IL-6 induced the synthesis of β -fibrinogen, α_1 -antichymotrypsin, α_1 -acid glycoprotein and haptoglobin in HepG2 cells, and α_2 -macroglobulin and cysteine protease inhibitor in rat hepatocytes. Furthermore, in rat hepatocyte primary cultures [38], or in the rat hepatoma cell lines Fao and H-35 [40,46–48], IL-6 led to a dose- and time-dependent regulation of α_2 -macroglobulin, β -fibrinogen, cysteine protease inhibitor, α_1 -acid glycoprotein, angiotensinogen and albumin. A concentration of 30 pM of recombinant human IL-6 resulted in a half-maximal acute phase response in hepatocyte primary cultures. A maximal response was achieved at a concentration of 1 nM recombinant human IL-6 [38]. IL-6 also had a stimulatory effect on the secretion of acute phase proteins [38].

Since *in vivo* stimulated monocytes synthesize and secrete IL-6, IL-1 and TNF α , the possible interaction of these cytokines was studied. IL-1 was able to inhibit the stimulatory effect of IL-6 on the synthesis of α_2 -macroglobulin, cysteine protease inhibitor and fibrinogen in rat hepatocyte primary cultures, while TNF α did not affect the action of IL-6 [36,38,41]. This finding suggests a complex regulation of acute phase protein synthesis.

Induction of rat acute phase proteins by interleukin-6 *in vivo*

After the demonstration that IL-6 is a potent inducer of acute phase protein synthesis in rat hepatoma cells and in adult rat hepatocytes, the validity of these observations remained to be demonstrated under *in vivo* conditions. The administration of recombinant human IL-6 to rats led to increases in the hepatic mRNA levels of α_2 -macroglobulin, β -fibrinogen, cysteine protease inhibitor and α_1 -acid glycoprotein and to a simultaneous decrease in albumin mRNA concentrations [49]. When the time course of acute phase protein mRNA induction after the administration of recombinant human IL-6 was

Table 2. Interleukin-6 synonyms

Name	Abbreviation	References
Hepatocyte stimulating factor	HSF	[40,41,47,57]
B cell stimulatory factor-2	BSF-2	[64,65]
Interferon β_2	IFN β_2	[66,67]
26 kDa protein		[68,69]
Hybridoma-plasmacytoma growth factor	HPGF	[70,71]
Myeloid blood cell differentiation-inducing protein	MGI-2A	[72]

compared with that after intramuscular injection of turpentine (commonly used for the induction of the acute phase response) a more rapid induction of the β -fibrinogen and α_2 -macroglobulin mRNA levels was observed in the case of recombinant human IL-6 [49]. This delay in acute phase response could be due to an indirect action of turpentine, where a cascade of reactions including activation of inflammatory cells and the formation, secretion and transport of acute phase mediators from the local abscess to the liver had to proceed, whereas recombinant human IL-6 acted directly on the hepatocytes. These results were recently confirmed [50]. Also in mice, induction of serum amyloid A, complement factor B and fibrinogen was activated after the injection of recombinant human IL-6 [51,52].

ACTIONS OF INTERLEUKIN-6

Interleukin-6 is the major regulator of the acute phase response in human hepatocytes

First evidence for IL-6 as an inducer of the acute phase response in the human system came from work with human cell lines [41,53–58]. As shown in Table 3, upper part, increases in protein synthesis of β -fibrinogen, α_1 -antichymotrypsin, caeruloplasmin, haptoglobin, α_1 -acid glycoprotein, α_1 -antitrypsin, and complement factor B are observed upon stimulation of HepG2 cells with recombinant human IL-6. However, no induction of the major human acute phase proteins, C-reactive protein and serum amyloid A, could be observed in IL-6-stimulated HepG2 or Hep3B cells. The combination of IL-6 and IL-1 was required to stimulate C-reactive protein and serum amyloid A synthesis in Hep3B cells [54]. Only in NPLC/PRF/5 cells could an IL-6-dependent stimulation of C-reactive protein and serum amyloid A synthesis be detected [54]. First evidence for IL-6 as the major inducer of acute phase proteins in humans came from studies with human hepatocytes in primary culture. In this system IL-6 was capable of inducing C-reactive protein and serum amyloid A synthesis in a dose- and time-dependent manner (Figs. 2a and 2b) [59] (Table 3, lower part). Since several investigators have found that IL-1 [5,33–37,46,60] and TNF α [34,35] were inducers of acute phase protein synthesis, comparative studies on the induction of various acute phase proteins by the three monokines have been carried out in human hepatocyte primary cultures [61,62]. While IL-6 stimulated the full spectrum of acute phase proteins seen in inflammatory states in man, IL-1 as well as TNF α had only a moderate effect on the positive acute phase proteins [62]. Both inhibited the synthesis of β -fibrinogen, albumin and transferrin and failed to induce serum amyloid A and C-reactive protein [62]. These

data strongly suggest that IL-6 plays a key role in human acute phase protein synthesis.

Recently IL-6 has been found to stimulate synthesis of the acute phase proteins complement factor B and C3 in human skin fibroblasts [63].

Pleiotropic action of interleukin-6

Besides its effect on hepatocytes, IL-6 acts on several other target cells (Fig. 3, and briefly reviewed in [73,74]). The biological functions of IL-6 under different names have been studied in various laboratories with a variety of cell systems (Table 2). When sequence data and specific antibodies became available after the purification and cDNA cloning of these 'different factors', they turned out to be identical. To unify the nomenclature, Poupart *et al.* [45] and Yasukawa *et al.* [75] proposed the name interleukin-6 for this molecule. Although this name is somewhat misleading, because interleukin-6 is synthesized by and acts on leukocytic and non-leukocytic cells, it was accepted at a conference of the New York Academy of Sciences and the National Foundation for Cancer Research in New York in December 1988 by the majority of the investigators in the field [76].

Action of interleukin-6 on B- and T-lymphocytes, plasmacytoma and B cell hybridomas. IL-6 acts directly on B cells activated by IL-4 and IL-5 and induces immunoglobulin M, G, and A production, i.e. IL-6 plays an essential role in the terminal differentiation of B cells into immunoglobulin-secreting cells (Fig. 3) [64,65,77–79]. IL-6 does not act on normal resting B cells, because these cells do not express IL-6 receptors [80]. Although activated B cells express IL-6 receptors, and respond to IL-6 by antibody production and secretion, IL-6 does not exert any growth-promoting activity on these cells.

Also *in vivo*, IL-6 has been shown to augment primary and secondary antibody responses in mice [81]. The strongest effect of IL-6 was observed in the secondary response [81]. After transfection with Epstein–Barr virus, B cells have been described to respond to IL-6 with proliferation in a paracrine manner [82]. IL-6 functions as a potent growth factor in plasmacytoma cells [70,71, 83–86]. IL-6 and IL-3 synergistically promote the proliferation and differentiation of malignant plasma cell precursors in multiple myeloma [87]. Myeloma cells from patients produce IL-6 and express its receptors. Antibodies against IL-6 inhibited the growth of these cells, suggesting an autocrine loop involved in oncogenesis of human myelomas [85]. Autocrine loops have also been proposed for the astrocytoma cell line U 337 and the myeloid cell line HL 60 [77], since both express IL-6 and IL-6 receptor [77]. Human and murine IL-6 are very potent growth factors for B cell hybridoma cells [70,71,

Table 3. Regulation of human acute-phase protein synthesis by IL-6, IL-1 and TNF α

Key: + + +, + +, +, increase in synthesis; 0, no change; —, inhibition of synthesis; *, augments the action of IL-6; n.d., not determined.

Cell type	Acute phase protein	IL-6	IL-1	TNF α	References
HEPG2	Fibrinogen	+ + +	—	n.d.	[41,57]
	α_1 -Antichymotrypsin	+ +	+	+	[35,41,56,57]
	Caeruloplasmin	+ +	n.d.	n.d.	[55]
	Haptoglobin	+	+ +	n.d.	[41,57]
	α_1 -Acid glycoprotein	+	+ +	n.d.	[41,57]
	α_1 -Antitrypsin	+	n.d.	n.d.	[55]
	Complement factor B	+	+	+	[35,58]
	Complement factor C3	—	+	+	[35]
	Metallothionein	n.d.	+	n.d.	[39]
	Albumin	—	—	—	[35,56,58]
	α -Fetoprotein	—	n.d.	n.d.	[55]
	α_2 -HS-glycoprotein	—	—	n.d.	[56]
	Transferrin	n.d.	—	—	[35]
HEP 3B	Caeruloplasmin	+ +	n.d.	n.d.	[54,55]
	α_1 -Antitrypsin	+	0	n.d.	[35,54,55]
	α -Fetoprotein	—	n.d.	n.d.	[55]
	C-reactive protein	0	n.d.	n.d.	[54]
	Serum amyloid A	0	n.d.	n.d.	[54]
	Complement factor B	n.d.	+	n.d.	[35]
	Complement factor C3	n.d.	+ +	+	[34,35]
NPLC/PRF/5	α_1 -Antichymotrypsin	n.d.	+	n.d.	[35]
	C-reactive protein	+	n.d.	n.d.	[54]
	Serum amyloid A	+	n.d.	n.d.	[54]
Adult human hepatocytes	C-reactive protein	+ + +	0*	0	[59,61,62]
	Serum amyloid A	+ + +	0*	0	[59,61,62]
	Haptoglobin	+ +	0	0	[59,62]
	α_1 -Antichymotrypsin	+ +	0	0	[59,62]
	Fibrinogen	+ +	—	—	[59,62]
	α_1 -Antitrypsin	+	0	0	[59,62]
	α_1 -Acid glycoprotein	+	0	0	[59,62]
	α_2 -Macroglobulin	0	0	—	[59,62]
	Fibronectin	—	0	0	[59,62]
	Transferrin	—	—	—	[59,62]
	Albumin	—	—	—	[59,62]

83,84,88–92] and can replace feeder cells. By subcloning various IL-6-dependent hybridoma cells Aarden and coworkers [93] established a stable cell line (B9), which can be used in a highly sensitive IL-6 assay detecting pg amounts of IL-6. This bioassay is highly specific for IL-6. The only known weak cross-reacting cytokine is murine interleukin-4 [93]. A similar B cell hybridoma clone (MH60.BSF2) suitable for the detection of IL-6 has been obtained by Kishimoto's group [94].

IL-6 is an important helper factor in primary antigen-receptor-dependent T cell activation and subsequent proliferation of the activated T cells [95–101]. This IL-6 action is synergized by IL-1 [99,100] and has been found to be IL-2-dependent [95,96,97]; however, other investigators did not find such an IL-2-dependency [99–101]. Similar observations have been made for thymocytes and T cell lines, i.e. IL-6 acts synergistically with

IL-1 [93,102,103] and stimulates proliferation of the thymocytes via an IL-2-dependent and -independent pathway [104,105]. Also in the late phase of cytotoxic T-cell generation IL-6 seems to enhance the action of IL-2 [106,107].

From the papers referred to in this section it becomes clear that IL-6 is involved in the function of the immune system by stimulating humoral as well as cellular defence mechanisms. Furthermore, IL-6 primes the human neutrophil and monocyte oxidative burst response [108].

Interleukin-6 is involved in haematopoiesis. IL-6 enhances the proliferation of multipotential haematopoietic progenitors in the human [109–111], murine [112–118] and rat [119] systems. A synergistic action of IL-6 with IL-3 [109,112–116], IL-4 [117], GM-CSF [111] and M-CSF [110] has been described. It has been suggested that

Fig. 2. Acute phase response of adult human hepatocytes stimulated by recombinant human IL-6

Human hepatocytes obtained from small surgical biopsies of healthy individuals were stimulated with IL-6 in the presence of dexamethasone. After labelling cells with [35 S]methionine the various acute phase proteins were immunoprecipitated and quantified. (a) Dose-dependence; (b) time course.

Fig. 3. Pleiotropic actions of interleukin-6

IL-6 shortens the G_0 period of haematopoietic stem cells by stimulating the proliferation of these cells [109,112,116].

Action of interleukin-6 on the growth and differentiation of various cell types. In addition to the function of IL-6 as a growth and differentiation factor in the haematopoietic and immune system, IL-6 has been described to stimulate proliferation of human [120,121] and murine [72,122,123] leukaemia and human lymphoma cells [124]. Interestingly, IL-6 also exerts a strong inhibition on the growth of human fibroblasts [125], of leukaemia [126,127], lymphoma [126] and breast carcinoma cells [126], and of human endothelial cells in culture [128]. Since fibroblasts and endothelial cells synthesize IL-6, the growth of these cells may be controlled via a negative autocrine feedback loop. As described above, a positive autocrine feedback loop was observed in plasmacytoma cells [85]. Further investigations of these two differently regulated systems might lead to a better understanding of growth control and oncogenesis. Furthermore, IL-6 plays a role in the expression of class I histocompatibility antigens in human fibroblasts [129].

In murine myeloleukaemic M1 cells and in human histiocytic U 937 cells, IL-6 induces morphological and functional differentiation into macrophages [130]. IL-6 promotes maturation of murine megakaryocytes [131]. Recently IL-6 has also been found to be involved in neuronal differentiation. Pheochromocytoma PC 12 cells change morphologically to neurite-extending cells [132]. In pituitary cells [133] and in the rat *in vivo* [134] IL-6 triggers the release of ACTH. Since ACTH stimulates glucocorticoid synthesis and glucocorticoids suppress IL-6 synthesis in monocytes, a bidirectional communication system regulating IL-6 levels exists between the neuroendocrine and the immune system.

Besides IL-1 and $TNF\alpha$, IL-6 has been found to be an endogenous pyrogen [93]. Intravenous injection of human recombinant IL-6 into rabbits led to a rapid onset

of fever reaching a peak at 60 min after injection [93]. Although IL-6 initially was described as an interferon, its antiviral activity is weak [66,125,135–137] and several groups of investigators could not detect any antiviral activity at all [44,69,138–140].

STRUCTURE OF THE IL-6 GENE

The human IL-6 gene is located on the short arm of chromosome 7 p15–p21 [141–143], and the murine IL-6 gene maps to the proximal region of chromosome 5 [144]. Chromosomal abnormalities of human chromosomes 7 and 5 have been found to be associated with a variety of haematopoietic disorders, among them multiple myeloma.

Both human and mouse IL-6 genes consist of five exons and four introns [145,146]. This gene organization is highly homologous to that of the G-CSF gene, an indication for the possible evolution from a common ancestor gene. Three initiation sites for transcription differently used in different cell types have been found [145]. At the 5' boundary of the IL-6 gene between –225 and –113, motifs similar to the regulatory elements in the human *c-fos* gene exist and may contain the major *cis*-acting elements responsible for the activation of the IL-6 promoter [146,147]. Furthermore, several putative enhancer and mediator responsive elements for cyclic AMP, glucocorticoid receptor, and activator binding protein-1 (AP-1) are present in the highly conserved 5' flanking region of the IL-6 gene of man and mouse [146].

STRUCTURE OF IL-6 PROTEIN

The nucleotide sequences and the corresponding amino acid sequences obtained by cDNA cloning of the IL-6 mRNA from human fibroblasts [69,135], from transformed human T cells [65] and from human monocytes [148,149] turned out to be identical.

The 1.3 kb mRNA is translated into a 212-amino-acid

precursor protein with a molecular mass of 26 kDa. For this reason it was formerly called 26K-protein [68]. After removal of a 28-amino-acid signal peptide, the resulting 184-amino-acid protein, containing two possible *N*-glycosylation sites, is *N*- and *O*-glycosylated and subsequently secreted [150–154]. The IL-6 secreted by monocytes shows a similar glycosylation pattern as that from fibroblasts, consisting of five or six different forms, which are the result of different *N*- and *O*-glycosylation [150–154]. Furthermore, the fibroblast and monocyte IL-6 is phosphorylated at several serine residues [155]. No sulphation was observed [155].

Although the unglycosylated IL-6 is biologically active, these post-translational modifications may play a role in tissue-specific functions and/or in plasma half-life.

Murine IL-6 contains 211 amino acids, including a signal peptide of 24 residues. No *N*-glycosylation, but several *O*-glycosylation sites were identified in this sequence. The sequence similarity to human IL-6 is 65 % at the cDNA level [156] and 42 % at the protein level [92]. In spite of the low similarity it is remarkable that both IL-6 species are biologically active in heterologous systems.

Human granulocyte colony stimulating factor (G-CSF) is the only presently known protein with some sequence similarity to human IL-6 [65]. Moreover, the cysteine locations in human and murine IL-6 match very well with those in G-CSF [65,157].

REGULATION OF INTERLEUKIN-6 BIOSYNTHESIS

The multifunctional cytokine IL-6 is synthesized by many different cells after appropriate stimulation (Table 4). Also, several tumour cell lines produce IL-6 spontaneously (Table 5). However, endothelial cells, fibroblasts and monocytes/macrophages are probably the major sources of IL-6 during systemic inflammation. Evidence for this assumption is based on the following facts: (a) these cells are highly abundant and widely distributed in the organism, (b) they have a high capacity to synthesize and secrete IL-6, and (c) IL-6 production can be triggered by several stimulators (Table 4). From the data summarized in Table 4, it can be seen that monocytes/macrophages are preferentially stimulated by bacterial lipopolysaccharide, whereas fibroblasts and endothelial cells respond better to the endogenous cytokines IL-1 and TNF α otherwise abundantly synthesized by monocytes/macrophages, suggesting that they play a key role in the production of IL-6.

In agreement with the results obtained from the experiments *in vitro* summarized in Tables 4 and 5 are the observations of elevated IL-6 serum levels after intravenous injection of lipopolysaccharide [185,186] or TNF α [187,188] into human volunteers and during bacterial infection [189].

The cytokines IL-1 and TNF α , which stimulate the synthesis of IL-6 in human fibroblasts, act via two second messenger systems: protein kinase C [190] and cyclic AMP [163,164]. Furthermore, IL-6 synthesis is controlled by steroid hormones. Dexamethasone has been described to inhibit IL-6 production in human monocytes [23] and in the astrocytoma cell line U373 MG [181]. Oestradiol-17 β impaired IL-6 formation in human endometrial stromal cells [178].

Using IL-6 promoter sequences fused to the reporter

gene chloramphenicol acetyltransferase transfected into HeLa cells, Ray *et al.* [147] located the major *cis*-acting regulatory elements responsible for the activation by IL-1, TNF α , epidermal growth factor, viruses, phorbol ester and cyclic AMP in the region between –225 and –113 of the human IL-6 gene.

Finally, it has been observed in several laboratories that the incubation of fibroblasts with cycloheximide leads to an accumulation of IL-6 mRNA [68,69,160]. This finding may be explained by the presence of two blocks of AU-rich sequences in the 3' untranslated region of IL-6 mRNA containing the AUUUA motif thought to be involved in the degradation of mRNA [162,191]. The AUUUA motif has been found in the 3' flanking region of many short-lived cytokine mRNAs [192].

Since IL-6 is found in cerebrospinal fluid [193–195] and in synovial fluid [196–198] during bacterial and viral infection of the central nervous system and during inflammatory arthritis, other cells, such as astroglia, microglia and synoviocytes, may be important sources of IL-6 during local inflammation.

HEPATOCYTES AS TARGET CELLS FOR INTERLEUKIN-6

As described above, IL-6 acts on many different target cells *in vitro*. *In vivo*, the liver seems to be the major target organ for IL-6. Both biological activity and radioactivity of ¹²⁵I-labelled recombinant human IL-6 injected intravenously into rats disappeared very rapidly from the circulation. The kinetics of clearance is biphasic and consists of a rapid initial elimination corresponding to a half-life of about 3 min and of a second slower decrease corresponding to a half-life of about 55 min [199]. Recombinant human IL-6 associates with a plasma protein resulting in a complex with β - γ mobility upon cellulose acetate electrophoresis [199]. In Kishimoto's laboratory α_2 -macroglobulin has been identified as a carrier for IL-6 [200]. IL-6 bound to α_2 -macroglobulin retained its biological activity and became resistant to proteases [200]. However, in our laboratory (J. V. Castell & P. C. Heinrich, unpublished results) α_2 -macroglobulin could not be identified as a binding protein for IL-6.

An analysis of the organs involved in the clearance of recombinant human IL-6 showed that the liver is the major target organ for IL-6. At 20 min after injection about 80 % of the ¹²⁵I-labelled recombinant human IL-6 had disappeared from the circulation and was found in liver [199]. Within the liver, ¹²⁵I-labelled recombinant human IL-6 was exclusively localized on the surface of parenchymal cells, suggesting the existence of an IL-6 receptor on the plasma membrane of hepatocytes [199]. As expected from the pleiotropic action of IL-6, IL-6 receptors have been found on the surface of various cell lines [201,202]. IL-6-receptor cDNA has recently been cloned [203].

INTERACTION OF INTERLEUKIN-6 WITH ITS HEPATIC RECEPTOR AND SIGNAL TRANSDUCTION

Interleukin-6 receptor

The IL-6-receptor cDNA was obtained from a cDNA library constructed from poly(A) RNA of the human natural killer cell line YT [203]. The IL-6-receptor cDNA encoded a protein consisting of 468 amino acids including

Table 4. Regulation of IL-6 biosynthesis

Key: ↑↑↑, very strong stimulation; ↑↑, strong stimulation; ↑, weak stimulation; 0, no effect; ↓, inhibition.

Cells	Stimulators/ inhibitors	Effect	References
Monocytes/macrophages	Lipopolysaccharide	↑↑↑	[18,26,150,152]
	HIV virus	↑↑↑	[158]
	Human interleukin-1 α/β	↑↑	[150,152]
	Interferon γ	↑	[153]
	Granulocyte/macrophage colony stimulating factor	↑	[153]
	Substance P and K	↑	[159]
	Tumour necrosis factor	0/↑	[150,151,152]
	Glucocorticoids	↓	[133]
Human fibroblasts	Human interleukin-1 α/β	↑↑↑	[45,70,71,160–165]
	Viruses	↑↑↑	[165,166]
	poly(I)·poly(C)	↑↑↑	[68,69,71,163]
	Tumour necrosis factor α	↑↑	[71,125,129,161,162,164]
	Bacteria/yeast	↑/↑↑	[165]
	Interferon α	↑	[161,162]
	Interferon β_1	0/↑	[160,162,167]
	Epidermal growth factor	↑	[147]
	Platelet-derived growth factor	0/↑	[167]
	Lipopolysaccharide	0/↑	[168]
Human endothelial cells	Interleukin-1 α/β	↑↑↑	[128,165,169,170]
	Tumour necrosis factor α	↑↑	[128,169]
	Lipopolysaccharide	0/↑	[128,165,169]
Human keratinocytes	Interleukin-1 α/β	↑	[171]
B-lymphocytes	Bacteria	↑	[172–174]
T-lymphocytes	Bacteria	↑	[172]
Synoviocytes	Interleukin-1	↑↑↑	[175,176]
	Tumour necrosis factor α	↑↑	[175]
	Lymphotoxin	↑	[175]
Amnion cells	Poly(I)·Poly(C)	↑	[177]
Endometrial stromal cells	Interleukin-1 α	↑↑↑	[178]
	Interleukin-1 β		[178]
	Lipopolysaccharide	0	[178]
	Oestradiol-17 β	↓	[178]
Bone marrow stromal cells	Interleukin-1	↑↑	[115]
Astrocytes	Interleukin-1 α/β	↑↑↑	[179]
	LCMV virus	↑↑↑	[179]
	Tumour necrosis factor α	↑↑	[179]
	Macrophage colony stimulating factor	↑	[179]
Microglia cells	LCMV virus	↑↑↑	[179]
	Macrophage colony stimulating factor	↑↑	[179]
	Interleukin-1 α/β	0/↑	[179]
	Tumour necrosis factor α	0/↑	[179]

a signal peptide of about 19 amino acids and a domain of 90 amino acids that resembles a domain of the immunoglobulin superfamily. The cytoplasmic part of 82 amino acids lacks a tyrosine kinase domain, unlike other growth factor receptors. There are six potential *N*-linked glycosylation sites, five in the extracellular and one in the intracellular domain. Binding constants of between 10^{10}

and 10^{11} M for 125 I-labelled recombinant human IL-6 were determined in several T- and B-cell, myeloid and astrocytoma cell lines [201,202]. Except for the plasma cell line U266 (11000 sites/cell) the number of IL-6 receptors per cell in these cell lines is low (1000) or very low (< 200) [201].

Little is known about the hepatic IL-6 receptor. In the

Table 5. IL-6 producing tumour cells and cell lines

Cells	References
HTLV-1-transformed T-cells (TCL-Na1)	[145,65]
Myeloma cells (U 266)	[72,121,184]
Leukaemia cells	[182,183]
Uterine carcinoma cells	[180]
Amnion-derived cell line (FL)	[145]
Myxoma cells	[180]
Bladder carcinoma cells (T24)	[145]
Epidermoid carcinoma cells	[171]
Astrocytoma cells (U 373)	[145,181]
Glioblastoma cells (SK-MG-4)	[145]
Osteosarcoma cells (MG-63)	[177,161]

human hepatoma cell line HepG2 450 ± 100 high-affinity (K_a 6.7×10^{10} M) and 5000 low-affinity (K_a 2×10^9 M) binding sites were determined [204]. A high-affinity association constant K_a of 77×10^{10} M and about 1500 IL-6 receptors were determined when the binding of 125 I-labelled recombinant human IL-6 to sinusoidal membranes from rat liver was studied (E. Hipp, unpublished work). No evidence for an internalization of 125 I-labelled recombinant human IL-6 by HepG2 cells and by rat hepatocytes *in vivo* was obtained [205]. A similar observation has been described by Aarden *et al.* [206], who found that IL-6 was neither taken up nor consumed by B-cell hybridomas. In contrast, 125 I-labelled recombinant human IL-6 was found to be internalized by CESS cells [202]. Affinity cross-linking using 125 I-labelled recom-

Fig. 4. Cross-linking of 125 I-labelled recombinant human IL-6 to its hepatic receptor

Sinusoidal membranes (SM) from rat liver were incubated with 125 I-labelled recombinant human IL-6 in the presence or absence of recombinant human IL-6 and cross-linked with disuccinimidyl suberate. Arrow denotes the presence of an IL-6 protein-receptor complex.

Fig. 5. Regulation of interleukin-6-receptor mRNA by dexamethasone in HepG2 cells

HepG2 cells were incubated for 18 h with IL-6 (100 units/ml) and dexamethasone (10^{-6} M) as indicated in the Figure. Controls with antibody-neutralized and heat-inactivated IL-6 are included. HeLa cell RNA showed no cross-hybridization with IL-6-receptor cDNA. The probe used in this experiment cross-hybridizes with 28 S rRNA.

binant human IL-6 and sinusoidal membranes from rat, pig or human liver resulted in the detection of a complex between IL-6 and its receptor with an apparent molecular mass of 100 kDa (arrow, Fig. 4). Therefore, a molecular mass of 80 kDa for the hepatic IL-6 receptor protein is suggested (D. Lenz, unpublished work). In the human lymphoplastoid cell line CESS, Coulie *et al.* [202] found three IL-6-binding proteins of about 170, 140 and 90 kDa. Content *et al.* [207] observed in HepG2 cells an increased binding of IL-6 after incubation with dexamethasone. This effect was due to an increase in the number of IL-6 binding sites and not to an increased affinity of IL-6 for its receptor. This is in accordance with data of H. Schoolink (unpublished results), who found increased IL-6-receptor mRNA levels after treatment of HepG2 cells with dexamethasone (Fig. 5). In fetal human hepatocytes in primary culture IL-6-receptor mRNA synthesis was stimulated by dexamethasone, IL-6 or IL-1 [208]. Interestingly, in normal peripheral blood monocytes the synthesis of IL-6-receptor mRNA is strongly reduced by IL-6, IL-1 and lipopolysaccharide [208]. Thus, during inflammation, a switch from monocytes to hepatocytes and activated B-cells as target cells for IL-6 may occur.

Signal transduction

In spite of the low number of hepatic IL-6 receptors, incubation of hepatocytes or hepatoma cells with IL-6 results in a large increase in acute phase protein synthesis, indicating the existence of a very efficient amplification system. Unfortunately, the underlying mechanisms are still unknown. Although, signal transduction by IL-6

was claimed to involve protein kinase C [209], it has been shown by pretreatment of HepG2 cells with phorbol ester that the action of IL-6 was independent of protein kinase C activity [204]. Preliminary experiments from our laboratory gave no evidence for the involvement of protein kinase C, inositol trisphosphate and adenylate cyclase in the IL-6-mediated signal transduction.

INTERLEUKIN-6-RESPONSIVE ELEMENTS IN ACUTE PHASE PROTEIN GENES

To further understand the mechanisms of acute phase protein induction by IL-6, the promoter regions of acute phase proteins have been isolated and ligated to suitable reporter genes. By transfection experiments with deletion mutants it was possible to define *cis*-elements responsible for the induction of acute phase protein gene transcription by IL-6. In the case of the rat α_1 -acid glycoprotein gene an IL-6-responsive enhancer element with a 142 bp sequence located 5300–5150 bp upstream of the transcriptional start site was identified [210]. Two IL-6-responsive elements, between –852 and –777 and between –336 and –165 upstream from the cap site were identified in the rat α_2 -macroglobulin gene [211] (Fig. 6). In the human haptoglobin gene three responsive elements around –157, –111 and –61 have been described [212]. The IL-6-responsive elements of α_1 -acid glycoprotein-, α_2 -macroglobulin and haptoglobin genes contain a common sequence motif (Fig. 7). This consensus sequence is also present in the promoter regions of rat α -, β - and γ -fibrinogen [214], rat T-kininogen [215], human C-reactive protein [216], rat transthyretin [217], murine serum amyloid A [218] and human α_1 -antitrypsin [219] (Fig. 7), although its function in these proteins still remains to be demonstrated. Thus, the sequence motif CTGGGAA in combination with appropriate trans-acting factors plays an important role in the acute phase protein gene regulation. Direct evidence for the involvement of this consensus sequence in IL-6-dependent rat

Fig. 6. Localization of interleukin-6-responsive DNA sequences in the promotor region of the rat α_2 -macroglobulin gene

Deletion mutants of the rat α_2 -macroglobulin promoter region were constructed, fused to the reporter gene chloramphenicol acetyltransferase and transfected into HepG2 cells as described by Kunz *et al.* [211]. Cells were stimulated with recombinant human IL-6 and assayed for chloramphenicol acetyltransferase (CAT) activity.

Acute phase gene	Location	Sequence	Reference
Rat α_1 -acid glycoprotein	–5220	TCTGGGAA	[210]
Rat α_2 -macroglobulin	–171	TCTGGGAA	[211, 213]
Human haptoglobin	–118	ACTGGAAA	[212]
Rat α -fibrinogen	–130	TCTGGGAT	[214]
Rat β -fibrinogen	–151	GCTGGGAA	[214]
Rat γ -fibrinogen	–152	TCTGGGAA	[214]
Rat T-kininogen	–112	GCTGGGTA	[215]
Human C-reactive protein	–77	GTGGGAAA	[216]
Rat transthyretin	–323	ACTGGGAA	[217]
Murine serum amyloid A	–80	AGTGGGAT	[218]
Human α_1 -antitrypsin	–118	ACTGGGAT	[219]

Consensus sequence:

CTGGGAAT

Fig. 7. Consensus sequence in the promoter regions of various acute phase proteins [213]

Fig. 8. Functional assay of an interleukin-6-responsive element of the rat α_2 -macroglobulin promoter

(a) Synthetic oligonucleotide (18-mer) containing the IL-6-responsive element of the rat α_2 -macroglobulin gene. (b) A 95 bp fragment containing the putative IL-6 responsive element (–215 to –121 relative to the cap site) was linked in both orientations (a) either singly (e) or in tandem (f) to the 5'-end of the pUC12 vector containing the SV 40 promoter in front of the chloramphenicol acetyltransferase (CAT) gene (a,d). Alternatively, the 18-mer (a) either singly (e) or in tandem (f) was fused to the chloramphenicol acetyltransferase gene. After transfection into HepG2 cells and subsequent stimulation with IL-6, chloramphenicol acetyltransferase was assayed.

α_2 -macroglobulin induction was obtained from transfection experiments with fusions consisting of the sequence 5'ATCCTTCTGGGAATTCTGG 3' (Fig. 8a) ligated to an SV40 promoter in front of the chloramphenicol acetyltransferase gene. Fig. 8(b) shows that the tandem of two such consensus sequences represents a particularly strong IL-6 responsive element.

As a consequence of the IL-6-induced increase in transcription [210–213,220], elevated mRNA [40,46, 48–51,53,204,217], and acute phase protein [38,41, 47–50,52–55,57,59,61,64,204] concentrations result. Furthermore, IL-6 was found to enhance the secretion of rat α_2 -macroglobulin in rat hepatocyte primary cultures [38].

RECENT DEVELOPMENTS

Baumann *et al.* [19,28] described a hepatocyte-stimulating factor, designated as HSF-III, in conditioned media from squamous carcinoma cells (COLO 16). This factor acts in an additive manner on the same spectrum of acute phase proteins as IL-6 [221]. It is structurally (molecular mass 39 kDa) and immunologically different from IL-6 and acts on a different receptor. Recently, it was shown that HSF-III is identical with leukaemia inhibitory factor (LIF) [222].

CONCLUSION

Due to its pleiotropic action, IL-6 has been intensively studied in many laboratories. It turned out to be an important factor in the immune and in the haematopoietic system and the major mediator in the hepatic acute phase response.

In Fig. 9 we show a simplified scheme of the complex role of IL-6 within the cytokine network that controls acute phase protein synthesis in hepatocytes. Monocytes/macrophages stimulated by bacteria, lipopolysaccharide, viruses, IL-1 and yet unknown factors synthesize and secrete, among other monokines, IL-1, IL-6 and $\text{TNF}\alpha$ (Fig. 9). These three monokines act via specific hepatic receptors. IL-6, however, turned out to be the most potent inducer of acute phase protein synthesis in human hepatocytes. The monokines IL-1 and $\text{TNF}\alpha$ also stimulate fibroblasts, endothelial cells and keratinocytes to synthesize IL-6 and thus amplify its biological effects.

IL-6, IL-1 and $\text{TNF}\alpha$ have been shown to stimulate the release of ACTH from pituitary cells, leading to an increased secretion of glucocorticoids by adrenocortical cells. The glucocorticoids, on one hand, increase the stimulatory effect of monokines on the synthesis of acute phase proteins by hepatocytes, and on the other hand, inhibit the synthesis of monokines.

PERSPECTIVES

Many questions in respect to the mechanisms involved in the acute phase response remain to be answered. Among these is the question of how acute phase states not involving lipopolysaccharide are mediated. Bacterial infections are only one of a large number of possible inflammatory stimuli, suggesting that other endogenous monocyte stimulators must exist. Investigations on the molecular basis of the acute phase response have only begun to elucidate a myriad of fascinating intercellular regulatory events.

Fig. 9. Regulation of hepatic acute phase protein synthesis by inflammatory mediators

Furthermore, it is of great interest to understand the interaction of IL-6 with its hepatic receptor, the regulation of the synthesis of the IL-6-receptor and the pathways of the intracellular signal transduction.

The intracellular mechanism of acute phase protein gene activation after IL-6 stimulation of hepatocytes, in particular a functional analysis of IL-6-responsive element(s), and the isolation and characterization of transcription factors, need to be worked out in greater detail. The fascinating problem of why some proteins behave as strong acute phase reactants in animals and not in man, or vice versa, should be solved. A better knowledge of the pathophysiological mechanisms of the acute phase response will improve the treatment of inflammatory diseases in man.

We thank R. Sittarz and A. Stammschulte for their help with the preparation and Dr. S. Rose-John for the critical reading of this manuscript.

REFERENCES

- Koj, A. (1974) in *Structure and Function of Plasma Proteins* (Allison, A. C., ed.) vol. 1, pp. 73–131, Plenum Press, London, New York
- Kushner, I. (1982) *Ann. NY Acad. Sci.* **389**, 39–49
- Pepys, M. B. & Baltz, M. L. (1983) *Adv. Immunol.* **34**, 141–210
- Schreiber, G. & Howlett, G. (1983) in *Plasma Protein Secretion by the Liver* (Glaumann, H., Peters, T., Jr. & Redman, C., eds.) pp. 423–449, Academic Press, London, New York
- Dinareello, C. A. (1984) *New England J. Med.* **311**, 1413–1418
- Koj, A. (1985) in *The Acute Phase Response to Injury and Infection* (Gordon, A. H. & Koj, A., eds.) vol. 10, pp. 139–232, Elsevier, Amsterdam
- Fey, G. H. & Fuller, G. M. (1987) *Mol. Biol. Med.* **4**, 323–338
- Kushner, I. & Mackiewicz, A. (1987) *Disease Markers* **5**, 1–11
- Kushner, I. (1988) *Methods Enzymol.* **163**, 373–383
- Fahraeus, R. (1921) *Acta. Med. Scand.* **55**, 1
- Sox, H. C., Jr. & Liang, M. H. (1986) *Ann. Intern. Med.* **104**, 515–523
- Abernethy, T. J. & Avery, O. T. (1941) *J. Exp. Med.* **73**, 173–182
- Miller, L. L., Bly, C. G., Watson, M. L. & Bale, W. F. (1951) *J. Exp. Med.* **94**, 431–453
- Kampschmidt, R. F., Upchurch, H. F., Eddington, C. L. & Pulliam, L. A. (1973) *Am. J. Physiol.* **224**, 530–533
- Powanda, M. C., Cockerell, G. L. & Pekarek, R. S. (1973) *Am. J. Physiol.* **225**, 399–401
- Wannemacher, R. W., Jr., Pekarek, R. S., Thompson, W. L., Curnow, R. T., Beall, F. A., Zenser, T. V., DeRubertis, F. R. & Beisel, W. R. (1975) *Endocrinology* **96**, 651–661
- Sipe, J. D., Vogel, S. N., Ryan, J. L., McAdam, K. P. W. J. & Rosenstreich, D. L. (1979) *J. Exp. Med.* **150**, 597–606
- Ritchie, D. G. & Fuller, G. M. (1983) *Ann. NY Acad. Sci.* **408**, 490–502
- Baumann, H., Jahreis, G. P., Sauder, D. N. & Koj, A. (1984) *J. Biol. Chem.* **259**, 7331–7342
- Koj, A., Gaudie, J., Regoeczi, E., Sauder, D. N. & Sweeney, G. D. (1984) *Biochem. J.* **224**, 505–514
- Woloski, B. M. R. N. J. & Fuller, G. M. (1985) *Proc. Natl. Acad. Sci. U.S.A.* **82**, 1443–1447
- Fuller, G. M., Otto, J. M., Woloski, B. M., McGary, C. T. & Adams, M. A. (1985) *J. Cell Biol.* **101**, 1481–1486
- Woloski, B. M. R. N. J., Smith, E. M., Meyer, W. J., Fuller, G. M. & Blalock, J. E. (1985) *Science* **230**, 1035–1037
- Bauer, J., Tran-Thi, T.-A., Northoff, H., Hirsch, F., Schlayer, H.-J., Gerok, W. & Heinrich, P. C. (1986) *Eur. J. Cell. Biol.* **40**, 86–93
- Amrani, D. L., Mauzy-Melitz, D. & Mosesson, M. W. (1986) *Biochem. J.* **238**, 365–371
- Northoff, H., Andus, T., Tran-Thi, T.-A., Bauer, J., Decker, K., Kubanek, B. & Heinrich, P. C. (1987) *Eur. J. Immunol.* **17**, 707–711
- Fuller, G. M., Bunzel, R. J., Woloski, B. M. & Nham, S.-U. (1987) *Biochem. Biophys. Res. Commun.* **144**, 1003–1009
- Baumann, H., Onorato, V., Gaudie, J. & Jahreis, G. P. (1987) *J. Biol. Chem.* **262**, 9756–9768
- Goldmann, N. D. & Liu, T.-Y. (1987) *J. Biol. Chem.* **262**, 2363–2368
- Bauer, J., Birmelin, M., Northoff, G.-H., Northemann, W., Tran-Thi, T.-A., Ueberberg, H., Decker, K. & Heinrich, P. C. (1984) *FEBS Lett.* **177**, 89–94
- Hirata, Y., Ishibashi, H., Kimura, H., Hayashida, K., Nagano, M. & Okubo, H. (1985) *Inflammation* **9**, 201–209
- Kurokawa, S., Ishibashi, H., Hayashida, K., Tsuchiya, Y., Hirata, Y., Sakaki, Y., Okubo, H. & Niho, Y. (1987) *Cell Struct. Funct.* **12**, 35–42
- Ramadori, G., Sipe, J. D., Dinarello, C. A., Mizel, S. B. & Colten, H. R. (1985) *J. Exp. Med.* **162**, 930–942
- Darlington, G. F., Wilson, D. R. & Lachmann, L. B. (1986) *J. Cell. Biol.* **103**, 787–793
- Perlmutter, D. H., Dinarello, C. A., Punsal, P. I. & Colten, H. R. (1986) *J. Clin. Invest.* **78**, 1349–1354
- Koj, A., Kurdowska, A., Magielska-Zero, D., Rokita, H., Sipe, J. D., Dayer, J. M., Demczuk, S. & Gaudie, J. (1987) *Biochem. Int.* **14**, 553–560
- Geiger, T., Andus, T., Klapproth, J., Northoff, H. & Heinrich, P. C. (1988) *J. Biol. Chem.* **263**, 7141–7146
- Andus, T., Geiger, T., Hirano, T., Kishimoto, T., Tran-Thi, T.-A., Decker, K. & Heinrich, P. C. (1988) *Eur. J. Biochem.* **173**, 287–293
- Karin, K., Imbra, R. J., Heguy, A. & Wong, G. (1985) *Mol. Cell Biol.* **5**, 2866–2869
- Andus, T., Geiger, T., Hirano, T., Northoff, H., Ganter, U., Bauer, J., Kishimoto, T. & Heinrich, P. C. (1987) *FEBS Lett.* **221**, 18–22
- Gaudie, J., Richards, C., Harnish, D., Lansdorp, P. & Baumann, H. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 7251–7255
- Billiau, A. (1986) *Nature (London)* **324**, 415
- Sehgal, P. B., May, L. T., Tamm, I. & Vilcek, J. (1987) *Science* **235**, 731–732
- Billiau, A. (1987) *Immunol. Today* **8**, 84–87
- Poupart, P., Vandenabeele, P., Cayphas, S., Van Snick, J., Haegemann, G., Kruys, V., Fiers, W. & Content, J. (1987) *EMBO J.* **6**, 1219–1224
- Andus, T., Geiger, T., Hirano, T., Kishimoto, T. & Heinrich, P. C. (1988) *Eur. J. Immunol.* **18**, 739–746
- Baumann, H. & Müller-Eberhard, U. (1987) *Biochem. Biophys. Res. Commun.* **146**, 1218–1226
- Itoh, N., Matsuda, T., Ohtani, R. & Okamoto, H. (1989) *FEBS Lett.* **244**, 6–10
- Geiger, T., Andus, T., Klapproth, J., Hirano, T., Kishimoto, T. & Heinrich, P. C. (1988) *Eur. J. Immunol.* **18**, 717–721
- Marinkovic, S., Jahreis, G. P., Wong, G. G. & Baumann, H. (1989) *J. Immunol.* **142**, 808–812

51. Ramadori, G., Van Damme, J., Rieder, H. & Meyer zum Büschenfelde, K.-H. (1988) *Eur. J. Immunol.* **18**, 1259–1264
52. Neta, R., Vogel, S. N., Sipe, J. D., Wong, G. G. & Nordan, R. P. (1989) *Lymphokine Res.* **7**, 403–413
53. Baumann, H., Richards, C. & Gauldie, J. (1987) *J. Immunol.* **139**, 4122–4128
54. Ganapathi, M. K., May, L. T., Schultz, D., Brabenec, A., Weinstein, J., Sehgal, P. B. & Kushner, I. (1988) *Biochem. Biophys. Res. Commun.* **157**, 271–277
55. Mackiewicz, A. & Kushner, I. (1989) *Scand. J. Immunol.* **29**, 265–271
56. Daveau, M., Davrinche, C., Julen, H., Hiron, M., Arnaud, P. & Lebreton, J. P. (1988) *FEBS Lett.* **241**, 191–194
57. Koj, A., Gordon, H. H. & Gauldie, J. (1988) *Experientia* **44**, 9–10
58. Perlmutter, D. H., Colten, H. R., Adams, S. P., May, L. T., Sehgal, P. B. & Fallons, R. J. (1989) *J. Biol. Chem.* **264**, 7669–7674
59. Castell, J. V., Gomez-Lechon, M. J., David, M., Hirano, T., Kishimoto, T. & Heinrich, P. C. (1988) *FEBS Lett.* **232**, 347–350
60. Klapproth, J., Castell, J., Geiger, T., Andus, T. & Heinrich, P. C. (1989) *Eur. J. Immunol.*, **19**, 1485–1490
61. Moshage, H. J., Roelofs, H. M. J., Van Pelt, J. F., Hazenberg, B. P. C., Van Leuven, M. A., Limburg, P. C., Aarden, L. A. & Yap, S. H. (1988) *Biochem. Biophys. Res. Commun.* **155**, 112–117
62. Castell, J. V., Gomez-Lechon, M. J., David, M., Andus, T., Geiger, T., Trullenque, R., Fabra, R. & Heinrich, P. C. (1989) *FEBS Lett.* **242**, 237–239
63. Katz, Y., Revel, M. & Strunk, R. C. (1989) *Eur. J. Immunol.* **19**, 983–988
64. Hirano, T., Taga, T., Nakano, N., Yasukawa, K., Kashiwamura, S., Shimizu, K., Nakajima, K., Pyun, K. H. & Kishimoto, T. (1985) *Proc. Natl. Acad. Sci. U.S.A.* **82**, 5490–5494
65. Hirano, T., Yasukawa, K., Harada, H., Taga, T., Watanabe, Y., Matsuda, T., Kashiwamura, S., Nakajima, K., Koyama, K., Iwamatsu, A., Tsunasawa, S., Sakiyama, F., Matsui, H., Takahara, Y., Taniguchi, T. & Kishimoto, T. (1986) *Nature (London)* **324**, 73–76
66. Weissenbach, J., Chernajovsky, Y., Zeevi, M., Shulman, L., Soreq, H., Nir, U., Wallach, D., Perriacaudet, M., Tiollais, P. & Revel, M. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **79**, 2768–2772
67. Sehgal, P. B. & Sagar, A. D. (1980) *Nature (London)* **288**, 95–97
68. Content, J., De Wit, L., Pierard, D., Derynck, R., De Clerque, E. & Fiers, W. (1982) *Proc. Natl. Acad. Sci. U.S.A.* **79**, 2768–2772
69. Haegeman, G., Content, J., Volckaert, G., Derynck, R., Tavernier, J. & Fiers, W. (1986) *Eur. J. Biochem.* **159**, 625–632
70. Van Damme, J., Opdenakker, G., Simpson, R. J., Rubira, M. R., Cayphas, S., Vink, A., Billiau, A. & Van Snick, J. (1987) *J. Exp. Med.* **165**, 914–919
71. Van Damme, J., Cayphas, S., Opdenakker, G., Billiau, A. & Van Snick, J. (1987) *Eur. J. Immunol.* **17**, 1–7
72. Shabo, Y., Lotem, J., Rubinstein, M., Revel, M., Clark, S. C., Wolf, S. F., Kamen, R. & Sachs, L. (1988) *Blood* **72**, 2070–2073
73. Kishimoto, T. & Hirano, T. (1988) *BioEssays* **9**, 11–15
74. Wong, G. G. & Clark, S. C. (1988) *Immunol. Today* **9**, 137–139
75. Yasukawa, K., Hirano, T., Watanabe, Y., Muratani, K., Matsuda, T., Nakai, S. & Kishimoto, T. (1987) *EMBO J.* **6**, 2939–2945
76. Sehgal, P. B., Grieninger, G. & Tosato, G. (eds.) (1989) *Ann. N.Y. Acad. Sci.* **557**
77. Kishimoto, T. (1987) *J. Clin. Immunol.* **7**, 343–355
78. Muraguchi, A., Hirano, T., Tang, B., Matsuda, T., Horii, Y., Nakajima, K. & Kishimoto, T. (1988) *J. Exp. Med.* **167**, 332–344
79. Kishimoto, T. & Hirano, T. (1988) *Annu. Rev. Immunol.* **6**, 485–512
80. Taga, T., Kawanishi, Y., Hardy, R. R., Hirano, T. & Kishimoto, T. (1987) *J. Exp. Med.* **166**, 967–981
81. Takatsuki, F., Okano, A., Suzuki, C., Chieda, R., Takahara, Y., Hirano, T., Kishimoto, T., Hamuro, J. & Akiyama, Y. (1988) *J. Immunol.* **141**, 3072–3077
82. Tosato, G., Gerrard, T. L., Goldman, N. G. & Pike, S. E. (1988) *J. Immunol.* **140**, 4329–4336
83. Van Snick, J., Cayphas, S., Vink, A., Uyttenhove, C., Coulie, P. G., Rubira, M. R. & Simpson, R. J. (1986) *Proc. Natl. Acad. Sci. U.S.A.* **83**, 9679–9683
84. Van Snick, J., Vink, A., Cayphas, S. & Uyttenhove, C. (1987) *J. Exp. Med.* **165**, 641–649
85. Kawano, M., Hirano, T., Matsuda, T., Taga, T., Horii, Y., Iwato, K., Asaoku, H., Tang, B., Tanabe, O., Tanaka, H., Kuramoto, A. & Kishimoto, T. (1988) *Nature (London)* **332**, 83–85
86. Zhang, X. G., Klein, B. & Bataille, R. (1989) *Blood* **74**, 11–13
87. Bergui, L., Schena, M., Gaidano, G., Riva, M. & Caligaris-Cappio, F. (1989) *J. Exp. Med.* **170**, 613–618
88. Aarden, L. A., Lansdorp, P. & De Groot, E. (1985) *Lymphokines* **10**, 175–185
89. Aarden, L. A., Degroot, E. R., Schaap, O. L. & Lansdorp, P. M. (1987) *Eur. J. Immunol.* **17**, 1411–1416
90. Van Damme, J., Cayphas, S., Van Snick, J., Conings, R., Put, W., Lenaerts, J. P., Simpson, R. J. & Billiau, A. (1987) *Eur. J. Biochem.* **168**, 543–550
91. Van Damme, J., Van Beeumen, J., Decock, B., Van Snick, J., De Ley, M. & Billiau, A. (1988) *J. Immunol.* **140**, 1534–1541
92. Simpson, R. J., Moritz, R. L., Rubira, M. R. & Van Snick, J. (1988) *Eur. J. Biochem.* **176**, 187–197
93. Helle, M., Brakenhoff, J. P. J., De Groot, E. R. & Aarden, L. A. (1988) *Eur. J. Immunol.* **18**, 957–959
94. Matsuda, T., Hirano, T. & Kishimoto, T. (1988) *Eur. J. Immunol.* **18**, 951–956
95. Garman, R. D., Jacobs, K. A., Clark, S. C. & Raulet, D. H. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 7629–7633
96. Habetswallner, D., Pelosi, E., Bulgarini, D., Camagna, A., Samoggia, P., Montesoro, E., Gianella, G., Lazzaro, D., Isaachi, G., Testa, U. & Peschle, C. (1988) *Immunology* **65**, 357–364
97. Suda, T., Hodgkin, P., Lee, F. & Zlotnik, A. (1989) *J. Immunol. Methods* **120**, 173–178
98. Uyttenhove, C., Coulie, P. G. & Van Snick, J. (1988) *J. Exp. Med.* **167**, 1417–1427
99. Houssiau, F. A., Coulie, P. G., Olive, D. & Van Snick, J. (1988) *Eur. J. Immunol.* **18**, 653–656
100. Ceuppens, J. L., Baroja, M. L., Lorre, K., Van Damme, J. & Billiau, A. (1988) *J. Immunol.* **141**, 3868–3874
101. Tosato, G. & Pike, S. E. (1988) *J. Immunol.* **141**, 1556–1562
102. Elias, J. A., Trinchieri, G., Beck, J. M., Simon, P. L., Sehgal, P. B., May, L. T. & Kern, J. A. (1989) *J. Immunol.* **142**, 509–514
103. Mizutani, H., May, L. T., Sehgal, P. P. & Kupper, T. S. (1989) *J. Immunol.* **143**, 896–901
104. Le, J., Fredrickson, G., Reis, L. F. L., Diamantstein, T., Hirano, T., Kishimoto, T. & Vilcek, J. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 8643–8647

105. Lotz, M., Jirik, F., Kabouridis, P., Tsoukas, C., Hirano, T., Kishimoto, T. & Carson, D. A. (1988) *J. Exp. Med.* **167**, 1253–1258
106. Takai, Y., Wong, G. C., Clark, S. C., Burakoff, S. J. & Herrmann, S. H. (1988) *J. Immunol.* **140**, 508–512
107. Okada, M., Kitahara, M., Kishimoto, S., Matsuda, T., Hirano, T. & Kishimoto, T. (1988) *J. Immunol.* **141**, 1543–1549
108. Kharazmi, A., Nielsen, H., Rechnitzer, C. & Bendtzen, K. (1989) *Immunol. Lett.* **21**, 177–184
109. Leary, A. G., Ikebuchi, K., Hirai, Y., Wong, G. G., Yang, Y. C., Clark, S. C. & Ogawa, M. (1988) *Blood* **71**, 1759–1763
110. Bot, F. J., Van Eijk, L., Broeders, L., Aarden, L. A. & Löwenberg, B. (1989) *Blood* **73**, 425–437
111. Caracciolo, D., Clark, S. C. & Rovera, G. (1989) *Blood* **73**, 666–670
112. Ikebuchi, K., Wong, G. G., Clark, S. C., Ihle, J. N., Hirai, Y. & Ogawa, M. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 9035–9039
113. Wong, G. G., Ikebuchi, K. L. & Ogawa, M. (1988) *J. Immunol.* **140**, 2040–2044
114. Koike, K., Nakahata, T., Takagi, M., Kobayashi, T., Ishiguro, A., Tsuji, K., Naganuma, K., Okano, A., Akiyama, Y. & Akabane, T. (1988) *J. Exp. Med.* **168**, 879–890
115. Chiu, C.-P., Moulds, C., Coffman, R. L., Rennick, D. & Lee, F. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 7099–7103
116. Ikebuchi, K., Ihle, J. N., Hirai, Y., Wong, G. G., Clark, S. C. & Ogawa, M. (1988) *Blood* **72**, 2007–2014
117. Rennick, D., Jackson, J., Yang, G., Wideman, J., Lee, F. & Hudak, S. (1989) *Blood* **73**, 1828–1835
118. Suzuki, C., Okano, A., Takatsuki, F., Miyasaka, Y., Hirano, T., Kishimoto, T., Ejima, D. & Akiyama, Y. (1989) *Biochem. Biophys. Res. Commun.* **159**, 933–938
119. Ulich, T. R., Del Castillo, J. & Guo, K. (1989) *Blood* **73**, 108–110
120. Hoang, T., Haman, A., Goncalves, O., Wong, G. G. & Clark, S. C. (1988) *Blood* **72**, 823–826
121. Shimizu, S., Yoshioka, R., Hirose, Y., Sugai, S., Tachibana, J. & Konda, S. (1989) *J. Exp. Med.* **169**, 339–344
122. Miyaura, C., Onozaki, K., Akiyama, Y., Taniyama, T., Hirano, T., Kishimoto, T. & Suda, T. (1988) *FEBS Lett.* **234**, 17–21
123. Chiu, C.-P. & Lee, F. (1989) *J. Immunol.* **142**, 1909–1915
124. Shimizu, S., Hirano, T., Yoshioka, R., Sugai, S., Matsuda, T., Taga, T., Kishimoto, T. & Konda, S. (1988) *Blood* **72**, 1826–1828
125. Kohase, M., Henriksen-DeStefano, D., May, L. T., Vliccek, J. & Sehgal, P. B. (1986) *Cell* **45**, 659–666
126. Chen, L., Mory, Y., Zilberstein, A. & Revel, M. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 8037–8041
127. Miyaura, C., Onozaki, K., Akiyama, Y., Taniyama, T., Hirano, T., Kishimoto, T. & Suda, T. (1988) *FEBS Lett.* **234**, 17–21
128. May, L. T., Torcia, G., Cozzolino, F., Ray, A., Tatter, S. B., Santhanam, U., Sehgal, P. B. & Stern, D. (1989) *Biochem. Biophys. Res. Commun.* **159**, 991–998
129. May, L. T., Helfgott, D. C. & Sehgal, P. B. (1986) *Proc. Natl. Acad. Sci. U.S.A.* **83**, 8957–8961
130. Chen, L., Novick, D., Rubinstein, M. & Revel, M. (1988) *FEBS Lett.* **239**, 299–304
131. Ishibashi, T., Kimura, H., Uchida, T., Kariyone, S., Friese, P. & Burstein, S. A. (1989) *Proc. Natl. Acad. Sci. U.S.A.* **86**, 5953–5957
132. Satoh, T., Nakamura, S., Taga, T., Matsuda, T., Hirano, T., Kishimoto, T. & Kaziro, Y. (1988) *Mol. Cell. Biol.* **8**, 3546–3549
133. Woloski, B. M. R. N. J., Smith, E. M., Meyer, W. J., III, Fuller, G. M. & Blalock, J. E. (1985) *Science* **230**, 1035–1037
134. Naitoh, Y., Fukata, J., Tominaga, T., Nakai, Y., Tamai, S., Mori, K. & Imura, H. (1988) *Biochem. Biophys. Res. Commun.* **155**, 1459–1463
135. Zilberstein, A., Ruggieri, R., Korn, J. H. & Revel, M. (1986) *EMBO J.* **5**, 2529–2537
136. Revel, M. & Zilberstein, A. (1987) *Nature (London)* **325**, 581–582
137. Jambou, R. C., Snouwaert, J. N., Abendroth Bishop, G., Stebbins, J. R., Freilinger, J. A. & Fowlkes, D. M. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 9426–9430
138. Hirano, T., Matsuda, T., Hosoi, K., Okano, A., Matsui, H. & Kishimoto, T. (1988) *Immunol. Lett.* **17**, 41–45
139. Poupart, P., De Wit, L. & Content, J. (1984) *Eur. J. Biochem.* **143**, 15–21
140. Reis, L. F. L., Le, J., Hirano, T., Kishimoto, T. & Vilcek, J. (1988) *J. Immunol.* **140**, 1566–1570
141. Sehgal, P. B., Zilberstein, A., Ruggieri, R., May, L. T., Ferguson-Smith, A., Slate, D. L., Revel, M. & Ruddle, F. H. (1986) *Proc. Natl. Acad. Sci. U.S.A.* **83**, 5219–5222
142. Sutherland, G. R., Baker, E., Callen, D. F., Hyland, V. J., Wong, G., Clark, S., Jones, S. S., Eglinton, L. K., Shannon, M. F., Lopez, A. F. *et al.* (1988) *Hum. Genet.* **79**, 335–337
143. Ferguson-Smith, A., Chen, Y. F., Newman, M. S., May, L. T., Ray, A., Sehgal, P. B. & Ruddle, F. H. (1988) *Genomics* **2**, 203
144. Mock, B. A., Nordan, R. P., Justice, M. J., Kozak, C., Jenkins, N. A., Copeland, N. G., Clark, S. C., Wong, G. G. & Rudikoff, S. (1989) *J. Immunol.* **142**, 1372–1376
145. Yasukawa, K., Hirano, T., Watanabe, Y., Muratani, K., Matsuda, T., Nakai, S. & Kishimoto, T. (1987) *EMBO J.* **6**, 2939–2945
146. Tanabe, O., Akira, S., Kamiya, T., Wong, G. G., Hirano, T. & Kishimoto, T. (1988) *J. Immunol.* **141**, 3875–3881
147. Ray, A., Tatter, S. B., May, L. T., Sehgal, P. B. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 6701–6705
148. Brakenhoff, J. P. J., De Groot, E. R., Evers, R. F., Pannekoek, H. & Aarden, L. A. (1987) *J. Immunol.* **139**, 4116–4121
149. Zimmermann, R., Bill, E., Geiger, T., Northoff, H. & Heinrich, P. C. (1988) *Biol. Chem. Hoppe-Seyler* **369**, 950–951
150. May, L. T., Ghrayeb, J., Santhanam, U., Tatter, S. B., Sthoege, Z., Helfgott, D. C., Chiorazzi, N., Griening, G. & Sehgal, P. B. (1988) *J. Biol. Chem.* **263**, 7760–7766
151. Andersson, U. & Matsuda, T. (1989) *Eur. J. Immunol.* **19**, 1157–1160
152. Bauer, J., Ganter, U., Geiger, T., Jacobshagen, U., Hirano, T., Matsuda, T., Kishimoto, T., Andus, T., Acs, G., Gerok, W. & Ciliberto, G. (1988) *Blood* **72**, 1134–1140
153. Navarro, S., Debili, N., Bernaudin, J.-F., Vainchenker, W. & Doly, J. (1989) *J. Immunol.* **142**, 4339–4345
154. Gross, V., Andus, T., Castell, J., Vom Berg, D., Heinrich, P. C. & Gerok, W. (1989) *FEBS Lett.* **247**, 323–326
155. May, L. T., Santhanam, U., Tatter, S. B., Bhardwaj, N., Ghrayeb, J. & Sehgal, P. B. (1988) *Biochem. Biophys. Res. Commun.* **152**, 1144–1150
156. Van Snick, J., Cayphas, S., Szikora, J.-P., Renaud, J.-C., Van Roost, E., Boon, T. & Simpson, R. J. (1988) *Eur. J. Immunol.* **18**, 193–197
157. Simpson, R. J., Moritz, R. L., Van Roost, E. & Van Snick, J. (1988) *Biochem. Biophys. Res. Commun.* **157**, 364–372

158. Nakajima, K., Martinez-Maza, O., Hirano, T., Breen, E. C., Nishanian, P. G., Salazar-Gonzales, J. F., Fahey, J. L. & Kishimoto, T. (1989) *J. Immunol.* **142**, 531–536
159. Lotz, M., Vaughan, J. H. & Carson, D. A. (1988) *Science* **241**, 1218–1221
160. Content, J., De Wit, L., Poupart, P., Opdenakker, G., Van Damme, J. & Billau, A. (1985) *Eur. J. Biochem.* **152**, 253–257
161. Defilippi, P., Poupart, P., Tavernier, J., Fiers, W. & Content, J. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 4557–4561
162. Walther, Z., May, L. T. & Sehgal, P. B. (1988) *J. Immunol.* **140**, 974–977
163. Zhang, Y., Lin, J.-X. & Vilcek, J. (1988) *J. Biol. Chem.* **263**, 6177–6182
164. Zhang, Y., Lin, J.-X., Keung, Y. & Vilcek, J. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 6802–6805
165. Sehgal, P. B., Helfgott, D. C., Santhanam, U., Tatter, S. B., Clarick, R. H., Ghayeb, J. & May, L. T. (1988) *J. Exp. Med.* **167**, 1951–1956
166. Van Damme, J., Schaafsma, M. R., Fibbe, W. E., Falkenburg, J. H. F., Opdenakker, G. & Billiau, A. (1989) *Eur. J. Immunol.* **19**, 163–168
167. Kohase, M., May, L. T., Tamm, I., Vilcek, J. & Sehgal, P. B. (1987) *Mol. Cell Biol.* **7**, 273–280
168. Helfgott, D. C., May, L. T., Sthoeger, Z., Tamm, I. & Sehgal, P. B. (1987) *J. Exp. Med.* **166**, 1300
169. Jirik, F. R., Podor, T. J., Hirano, T., Kishimoto, T., Loskutoff, D. J., Carson, D. A. & Lotz, M. (1989) *J. Immunol.* **142**, 144–147
170. Sironi, M., Breviario, F., Proserpio, P., Biondi, A., Vecchi, A., Van Damme, J., Dejana, E. & Mantovani, A. (1989) *J. Immunol.* **142**, 549–553
171. Kirnbauer, R., Köck, A., Schwarz, T., Urbanski, A., Krutmann, J., Borth, W., Damm, D., Shipley, G., Ansel, J. C. & Luger, T. A. (1989) *J. Immunol.* **142**, 1922–1928
172. Horii, Y., Muraguchi, A., Suematsu, S., Matsuda, T., Yoshizaki, K., Hirano, T. & Kishimoto, T. (1988) *J. Immunol.* **141**, 1529–1535
173. Freeman, G. J., Freedman, A. S., Rabinowe, S. N., Segil, J. M., Horowitz, J., Rosen, K., Whitman, J. F. & Nadler, L. M. (1989) *J. Clin. Invest.* **83**, 1512–1518
174. Fukui, H., Koishihara, Y., Nagamuta, M., Mizutani, Y. & Atsushi, U. (1989) *Immunol. Lett.* **21**, 127–130
175. Guerne, P.-A., Zuraw, B. L., Vaughan, J. H., Carson, D. A. & Lotz, M. (1989) *J. Clin. Invest.* **83**, 585–592
176. Harigai, M., Hara, M., Norioka, K., Kitani, A., Hirose, T., Suzuki, K., Kawakami, M., Masuda, K., Shinmei, M., Kawagoe, M. & Nakamura, H. (1989) *Scand. J. Immunol.* **29**, 289–297
177. Poupart, P., De Wit, L. & Content, J. (1984) *Eur. J. Biochem.* **143**, 15–21
178. Tabibzadeh, S., Santhanam, U., Sehgal, P. B. & May, L. T. (1989) *J. Immunol.* **142**, 3134–3139
179. Frei, K., Malipiero, U. V., Leist, T. P., Zinkernagel, R. M., Schwab, M. E. & Fontana, A. (1989) *Eur. J. Immunol.* **19**, 689–694
180. Hirano, T., Taga, T., Yasukawa, K., Nakajima, K., Nakano, N., Takatsuki, F., Shimizu, M., Murashima, A., Tsunasawa, S., Sakiyama, F. & Kishimoto, T. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 228–231
181. Nishida, T., Nakai, S., Kawakami, T., Aihara, K., Nishino, N. & Hirai, Y. (1989) *FEBS Lett.* **243**, 25–29
182. Biondi, A., Rossi, V., Bassan, R., Barbui, T., Bettoni, S., Sironi, M., Mantovani, A. & Rambaldi, A. (1989) *Blood* **73**, 1279–1284
183. Oster, W., Cicco, N. A., Klein, H., Hirano, T., Kishimoto, T., Lindermann, A., Mertelsmann, R. H. & Herrmann, F. (1989) *J. Clin. Invest.* **84**, 451–457
184. Kishimoto, T., Hirano, T., Taga, T., Yamazaki, K., Matsuda, T., Tang, B., Muraguchi, A., Horii, Y., Kawanishi, Y., Hirata, Y., Yawata, K., Watanabe, Y., Shimizu, M., Heinrich, P. C. & Kawano, M. (1988) *B Cell Develop.* 157–171
185. Coulie, P. C., Cayphas, S., Vink, A., Uyttenhove, C. & Van Snick, J. (1987) *Eur. J. Immunol.* **17**, 1217–1220
186. Fong, Y., Moldawer, L. L., Marano, M., Wei, H., Tatter, S. B., Clarick, R. H., Santhanam, U., Sherris, D., May, L. T., Sehgal, P. B. & Lowry, S. F. (1989) *J. Immunol.* **142**, 2321–2324
187. Jablons, D. M., Mulé, J. J., McIntosh, J. K., Sehgal, P. B., May, L. T., Huang, C. M., Rosenberg, S. A. & Lotze, M. T. (1989) *J. Immunol.* **142**, 1542–1547
188. Brouckaert, P., Spriggs, D. R., Demetri, G., Kufe, D. W. & Fiers, W. (1989) *J. Exp. Med.* **169**, 2257–2262
189. Helfgott, D. C., Tatter, S. B., Santhanam, U., Clarick, R. H., Bhardwaj, N., May, L. T. & Sehgal, P. B. (1989) *J. Immunol.* **142**, 948–953
190. Sehgal, P. B., Walther, Z. & Tamm, I. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 3663–3667
191. Shaw, G. & Kamen, R. (1986) *Cell* **46**, 659–667
192. Cosman, D. (1987) *Immunol. Today* **8**, 16–17
193. Houssiau, F. A., Bukasa, K., Sindic, C. J. M., Van Damme, J. & Van Snick, J. (1988) *Clin. Exp. Immunol.* **71**, 320–323
194. Waage, A., Brandtzaeg, P., Halstensen, A., Kierulf, P. & Espevik, T. (1989) *J. Exp. Med.* **169**, 333–338
195. Frei, K., Leist, P., Meager, A., Gallo, P., Leppert, D., Zinkernagel, R. M. & Fontana, A. (1989) *J. Exp. Med.* **169**, 449–453
196. Houssiau, F. A., Devogelaer, J.-P., Van Damme, J., DeDeuxchaisnes, C. N. & Van Snick, J. (1988) *Arthritis Rheum.* **31**, 784–788
197. Hirano, T., Matsuda, T., Turner, M., Miyasaka, N., Buchan, G., Tang, B., Sato, K., Shimizu, M., Maini, R., Feldmann, M. & Kishimoto, T. (1988) *Eur. J. Immunol.* **18**, 1797–1801
198. Waage, A., Kaufmann, C., Espevik, T. & Husby, G. (1989) *Clin. Immunol. Immunopathol.* **50**, 394–398
199. Castell, J. V., Geiger, T., Gross, V., Andus, T., Walter, E., Hirano, T., Kishimoto, T. & Heinrich, P. C. (1988) *Eur. J. Biochem.* **177**, 357–361
200. Matsuda, T., Hirano, T., Nagasawa, S. & Kishimoto, T. (1989) *J. Immunol.* **142**, 148–152
201. Taga, T., Kawanishi, Y., Hardy, R. R., Hirano, T. & Kishimoto, T. (1989) *J. Exp. Med.* **166**, 967–981
202. Coulie, P. G., Vanhecke, A., Van Damme, J., Cayphas, S., Poupart, P., De Wit, L. & Content, J. (1987) *Eur. J. Immunol.* **17**, 1435–1440
203. Yamasaki, K., Taga, T., Hirata, Y., Yawata, H., Kawanishi, Y., Seed, B., Taniguchi, T., Hirano, T. & Kishimoto, T. (1988) *Science* **242**, 825–828
204. Baumann, H., Isseroff, H., Latimer, J. J. & Jahreis, G. P. (1988) *J. Biol. Chem.* **263**, 17390–17396
205. Castell, J. V., Klapproth, J., Gross, V., Walter, E., Andus, T., Snyers, L., Content, J. & Heinrich, P. C. (1990) *Eur. J. Biochem.*, in the press
206. Aarden, L., Lansdorp, P. & De Groot, E. (1985) in *Lymphokines, Growth and Differentiation of B-cells* (Schreier, M. H. & Smith, K. H., eds.), vol. 10, pp. 175–185, Academic Press, Orlando
207. Content, J., Fontaine, V. & Snyers, L. (1988) *Modulation of IL-6 Receptors in Human Cells*, The New York Academy of Sciences/National Foundation for Cancer Research, December 12–14, 1988

208. Bauer, J., Lengyel, G., Bauer, T. M., Acs, G. & Gerok, W. (1989) *FEBS Lett.* **249**, 27–30
209. Evans, E., Courtois, G. M., Kilian, P. L., Fuller, G. M. & Crabtree, G. R. (1987) *J. Biol. Chem.* **262**, 10850–10854
210. Prowse, K. R. & Baumann, H. (1988) *Mol. Cell. Biol.* **8**, 42–51
211. Kunz, D., Zimmermann, R., Heisig, M. & Heinrich, P. C. (1989) *Nucleic Acids Res.* **17**, 1121–1138
212. Oliviero, S., Morrone, G. & Cortese, R. (1987) *EMBO J.* **6**, 1905–1912
213. Tsuchiya, Y., Hattori, M., Hayashida, K., Ishibashi, H., Okubo, H. & Sakaki, Y. (1987) *Gene* **57**, 73–80
214. Folwkes, D. M., Mullis, N. T., Comeau, C. M. & Crabtree, G. R. (1984) *Proc. Natl. Acad. Sci. U.S.A.* **81**, 2313–2316
215. Fung, W.-P. & Schreiber, G. (1987) *J. Biol. Chem.* **262**, 9298–9308
216. Arcone, R., Gualandi, G. & Ciliberto, G. (1988) *Nucleic Acids Res.* **16**, 3195–3207
217. Fung, W.-P., Thomas, T., Dickson, P. W., Alred, A. R., Milland, J., Dziadek, M., Power, B., Hudson, P. & Schreiber, G. (1988) *J. Biol. Chem.* **263**, 480–488
218. Lowell, C. A., Potter, D. A., Stearman, R. S. & Morrow, J. F. (1986) *J. Biol. Chem.* **261**, 8442–8452
219. Ciliberto, G., Dente, L. & Cortese, R. (1985) *Cell* **41**, 531–540
220. Morrone, G., Ciliberto, G., Oliviero, S., Arcone, R., Dente, L., Content, J. & Cortese, R. (1988) *J. Biol. Chem.* **263**, 12554–12558
221. Baumann, H., Won, K. A. & Jahreis, G. P. (1989) *J. Biol. Chem.* **264**, 8046–8051
222. Baumann, H. & Wong, G. G. (1989) *J. Immunol.* **143**, 1163–1167