Metabolic cleavage of cell-penetrating peptides in contact with epithelial models: human calcitonin (hCT)-derived peptides, Tat(47–57) and penetratin(43–58)

Rachel TRÉHIN^{*1}, Hanne M. NIELSEN^{*†1}, Heinz-Georg JAHNKE^{*‡}, Ulrike KRAUSS[‡], Annette G. BECK-SICKINGER[‡] and Hans P. MERKLE^{*2}

^{*}Drug Formulation & Delivery Group, Department of Chemistry and Applied BioSciences, Swiss Federal Institute of Technology Zurich (ETH Zurich), Winterthurerstrasse 190, CH-8057 Zurich, Switzerland, [†]Department of Pharmaceutics, The Danish University of Pharmaceutical Sciences, 2 Universitetsparken, DK-2100 Copenhagen, Denmark, and [‡]Institute of Biochemistry, University of Leipzig, D-04103 Leipzig, Germany

We assessed the metabolic degradation kinetics and cleavage patterns of some selected CPP (cell-penetrating peptides) after incubation with confluent epithelial models. Synthesis of N-terminal CF [5(6)-carboxyfluorescein]-labelled CPP, namely hCT (human calcitonin)-derived sequences, Tat(47–57) and penetratin(43–58), was through Fmoc (fluoren-9-ylmethoxycarbonyl) chemistry. Metabolic degradation kinetics of the tested CPP in contact with three cell-cultured epithelial models, MDCK (Madin–Darby canine kidney), Calu-3 and TR146, was evaluated by reversed-phase HPLC. Identification of the resulting metabolites of CF-hCT(9–32) was through reversed-phase HPLC fractionation and peak allocation by MALDI–TOF–MS (matrix-assisted laser-desorption ionization–time-of-flight mass spectrometry) or direct MALDI–TOF–MS of incubates. Levels of proteolytic activity varied highly between the investigated epithelial models and the CPP. The Calu-3 model exhibited the highest proteolytic activity.

The patterns of metabolic cleavage of hCT(9–32) were similar in all three models. Initial cleavage of this peptide occurred at the N-terminal domain, possibly by endopeptidase activity yielding both the N- and the C-terminal counterparts. Further metabolic degradation was by aminopeptidase, endopeptidase and/or carboxypeptidase activities. In conclusion, when in contact with epithelial models, the studied CPP were subject to efficient metabolism, a prerequisite of cargo release on the one hand, but with potential for premature cleavage and loss of the cargo as well on the other. The results, particularly on hCT(9–32), may be used as a template to suggest structural modifications towards improved CPP performance.

Key words: cell-penetrating peptides, epithelial delivery, human calcitonin, metabolic cleavage, penetratin, Tat.

INTRODUCTION

The lipid bilayer of the cellular plasma membrane acts as a highly impermeable barrier to most polar molecules, hindering efficiently peptide-, protein- and nucleic acid-based therapeutics to cross this membrane [1,2], and a major hurdle in drug development. Therefore fusion or complexation of such therapeutic agents with CPP (cell-penetrating peptides) is currently one of the strategies proposed to improve their cellular bioavailability. Most of the recognized CPP derive from naturally occurring protein sequences that are endowed with the ability to cross the plasma membrane and even reach the nuclear compartment [3,4]. In the literature, CPP are reported to enter a variety of different cell types [5,6] and enhance the cellular uptake of various macromolecular and even nanoparticulate cargos [7–10]. Depending on their chemical nature, different routes of cellular entry should be expected. CPP of polycationic nature, e.g. penetratin, Tat and oligo-arginine, were shown to share a common uptake mechanism initialized by electrostatic interaction with the negatively charged sulphated proteoglycans and glycoproteins of the extracellular matrix, then triggering endocytosis [11,12]. For transportan and a transportan derivative, a non-endocytic pathway has been concluded, caused by an as yet unspecified perturbation of the plasma membrane [13]. In contrast, several hCT (human calcitonin)-derived CPP

of moderately cationic character were shown to cross the plasma membrane by an endocytic mechanism of still unknown nature [14], but supposedly different from polycationic CPP.

A major obstacle in using peptides as delivery vectors is their rapid clearance when in contact with or passing through the enzymic barriers of epithelia and endothelia [15] and in the bloodstream [16]. Metabolic stability of CPP is thus an important biopharmaceutical factor for their cellular delivery. The peptides should translocate their cargo before they are metabolically cleaved. Nevertheless, in addition to a negative connotation, metabolic cleavage of CPP is also a prerequisite for the release of a chemically ligated cargo after cellular internalization, and has important consequences for the physiological clearance of CPP and their acute and chronic toxicity when used for therapeutic purposes. Major determinants for the proteolytic cleavage of CPP are supposedly their individual chemistries and specific routes of entry. Another one should be the type of cargo that is chemically ligated or physically assembled with the CPP. Depending on the respective combination, different pathways and kinetics of enzymic cleavage may be envisaged.

To our knowledge, interest in the metabolic stability and clearance of CPP in the literature has been limited [13,17]. Elmquist et al. [18] studied the stability of pVEC, a CPP derived from murine vascular endothelial cadherin inside and outside human

Abbreviations used: BCA, bicinchoninic acid; CF, 5(6)-carboxyfluorescein; CPP, cell-penetrating peptides; FCS, foetal calf serum; HBSS, Hanks balanced salt solution; hCT, human calcitonin; MALDI–TOF–MS, matrix-assisted laser-desorption ionization–time-of-flight mass spectrometry; MDCK, Madin–Darby canine kidney; RP-HPLC, reversed-phase HPLC; TEER, transepithelial electrical resistance; TFA, trifluoroacetic acid; TP10, transportan 10.

¹ These authors have contributed equally to this work.

² To whom correspondence should be addressed (email hmerkle@pharma.ethz.ch).

aorta endothelial cells and murine A9 fibroblasts; *pVEC* turned out to degrade rapidly when incubated with the cellular models. By contrast, when replacing all L-amino-acid residues by their D-amino-acid counterparts, *pVEC* was no longer subject to any degradation regardless of media composition and incubation time [17]. In a further study, the authors focused on the metabolic stability of transportan, transportan 10 (TP10), a transportan analogue [19], and penetratin(43–58). The stability of the peptides was in the order of transportan > TP10 > penetratin [13]. More effort has been spent in the area of the metabolic stability of peptide therapeutics themselves. For instance, previous studies in our laboratory demonstrated the cleavage of therapeutic and related model peptides in contact with epithelial models [20–22]. Due to peptidase activity of various types, epithelia act as efficient metabolic barriers to block the epithelial delivery of peptidase-labile peptides.

In a previous study, we demonstrated that the C-terminal fragment of hCT featuring residues 9–32, hCT(9–32), as well as hCT itself were internalized *in vitro* into bovine nasal epithelium [23,24]. In the same model, the C-terminal fragment was also demonstrated to act as a vector for green fluorescent protein [25]. More recently, hCT(9–32) and selected truncated sequences thereof were shown to cross the plasma membrane of cell-cultured MDCK (Madin–Darby canine kidney) cell monolayers following an endocytic mechanism [14]. MDCK cell line is derived from distal renal epithelium and forms a recognized cell culture model to mimic columnar-type absorptive epithelia [26–28]. In addition to the mildly cationic hCT-derived CPP, polycationic sequences such as Tat(47–57), derived from the nuclear transcription-activating protein from the HIV-1 virus, and penetratin (43–58), from the homeodomain of the Antennapedia transcription factor of *Drosophila*, also showed intracellular uptake in MDCK cell monolayers [14]. Furthermore, hCT(9–32), Tat(47–57) and penetratin(43–58) demonstrated either cytoplasmic vesicular or paracellular accumulation [29] in Calu-3 cell layers, derived from a human bronchial submucosal adenocarcinoma used as another columnar-type absorptive model [30], and in TR146 cell multilayers. The TR146 cell line originates from a neck node metastasis of a human buccal carcinoma [31] and serves as a stratified-type epithelial model.

In the present study, we engage in the assessment of the metabolic degradation of hCT(9–32) and the related hCT-derived peptides, Tat(47–57) and penetratin(43–58) in contact with the three confluent epithelial models MDCK, Calu-3 and TR146 [27,32–34]. An objective of the study is to create a basis for potential sequence modifications of hCT(9–32), which could be helpful to improve the metabolic stability of CPP towards proteolytic activity when in contact with or passing airway or buccal epithelia. Our results encourage sequence modifications in the N-terminal segment to stabilize hCT(9–32) against metabolic cleavage and eventually ameliorate its potential as a drug delivery vector.

EXPERIMENTAL

Materials

The MDCK cell line (low resistance, type II) was a gift from the Biopharmacy group of ETH Zurich (Switzerland) [35]. Calu-3 cells were purchased from A.T.C.C. (Rockville, MD, U.S.A.) and TR146 was provided by Imperial Cancer Research Technology (London, U.K.). Cell culture media, L-glutamine, trypsin/EDTA, penicillin, streptomycin and HBSS (Hanks balanced salt solution) were from Life Technologies (Basel, Switzerland). FCS (foetal calf serum) for the cell culture of MDCK was obtained from

Table 1 Amino acid sequences and molecular masses of hCT-derived peptides, Tat(47–57) and penetratin(43–58)

Sequence numbering of hCT-derived peptides corresponds to native hCT.

Peptide	Sequence	Molecular mass* (Da)
	9 12 15 18 21 24 27 31	
hCT(9–32)	LGTYTQDFNKFHTFPQTAIGVGAP	2968.5
hCT(12–32)	YTQDFNKFHTFPQTAIGVGAP	2699.3
hCT(15–32)	DFNKFHTFPQTAIGVGAP	2306.3
hCT(18–32)	KFHTFPQTAIGVGAP	1929.9
Random sequence	FLTAGQNTIQTVPKGTGGHFFADY	2968.5
Tat(47–57)	GYGRKKRRQRRG	2031.9
Penetratin(43–58)	RQIKIWFQNRRMKWKK	2604.7

* The molecular masses of the peptides include the N-terminally ligated CF.

Winiger AG (Wohlen, Switzerland) and for the TR146 cell culture from PAA Laboratories (Linz, Austria). For the Calu-3 model, FetalClone III bovine serum was from HyClone (Logan, UT, U.S.A.). CF [5(6)-carboxyfluorescein], used for N-terminal labelling of peptides, and BSA were obtained from Fluka (Buchs, Switzerland). Vitrogen collagen was from Nutacon (Leimuiden, Belgium). Protein assay kit with BCA (bicinchoninic acid) and all other chemicals were from Sigma (St. Louis, MO, U.S.A.). Cell culture inserts of polyethylene terephthalate (PET; 0.4 µm pore size, 1.6 × 10⁶ pores/cm², 0.9 and 4.2 cm² growth area) and companion plates (six- and 12-well plates) were purchased from Falcon (Becton Dickinson Labware, Franklin Lakes, NJ, U.S.A.); 96-well plates were obtained from TPP (Trasadingen, Switzerland).

Synthesis of hCT fragments, labelling and identification

The sequences of the tested peptides are described in Table 1. Linear hCT fragments were synthesized as described by Rist et al. [36] by automated multiple solid-phase peptide synthesis using a robot system (Syro, MultiSynTech, Bochum, Germany). All fragments were CF-labelled at the N-terminus, whereas the peptide was still bound to the resin with fully protected side chains [37]. The methods were described previously [14]. The peptides were purified by RP-HPLC (reversed-phase HPLC) and characterized by MS.

Synthesis of Tat(47–57) and penetratin(43–58), labelling and identification

Synthesis of N-terminally CF-labelled Tat(47–57) and penetratin(43–58) was performed by solid-phase synthesis on an Applied Biosystems 433A peptide synthesizer (Foster City, CA, U.S.A.) following the Fmoc (fluoren-9-ylmethoxycarbonyl) chemistry method (for sequences see Table 1) at the Institute of Biochemistry (University of Zurich). Two glycine residues were added, one at the N-terminal and one at the C-terminal end of the Tat(47–57) to provide links for fluorochrome binding. In the case of penetratin(43–58), the C-terminal end was amidified to improve its stability. All procedures were as described previously [14].

Cell culture

MDCK cells were grown under standard conditions in minimum essential medium with Earl's salts containing 10% heat-inactivated FCS and 1% penicillin/streptomycin [35]. Calu-3 and TR146 cells were cultured in DMEM, supplemented with 1% penicillin/streptomycin and 10% (v/v) FCS. For metabolism

studies, cells were seeded on PET cell culture inserts in six- (4.2 cm²; MDCK) or 12-well plates (0.9 cm²; Calu-3 and TR146) with a constant density of approx. 2×10^4 , 10^5 and 24×10^3 cells/cm² for MDCK, for Calu-3 and for TR146 respectively. In the case of Calu-3, the cells were plated on to 30 µg/ml collagen-coated filter supports. After Calu-3 cells had attached on the filter overnight, the medium was removed from the apical compartment to allow the monolayer to grow at the air interface. Air-interface conditions have been shown to improve the differentiation of primary cultures of human airway epithelia [38,39]. The three epithelial cell lines were maintained in 5% CO₂ in a humidified atmosphere with 1.5 ml medium in the basal chamber for Calu-3, with 2.5 ml medium in the apical chamber and 3 ml in the basal chamber for MDCK, and with 0.5 ml medium in the apical chamber and 1.5 ml in the basal chamber for TR146.

Confluent MDCK monolayers were formed after 10 days, and culture medium was renewed twice a week. Cells were used between passage numbers 227 and 235. Calu-3 cells grew during 17–19 days and TR146 during 28–30 days. For the latter two cell lines, media were exchanged three times per week. Calu-3 cells were used between passage numbers 36–41 and TR146 between passage numbers 11–15.

BCA protein assay

The BCA assay is based on the formation of a Cu²⁺–protein complex under alkaline conditions followed by reduction of Cu²⁺ to Cu⁺. The reduction is proportional to the quantity of protein present in the sample. BCA forms a purple-blue complex with Cu⁺ providing a basis to monitor the reduction of alkaline Cu²⁺ by proteins. In the present study, the amount of protein was assessed for each cell line and converted to µg/cm². For this purpose, cells were scraped from the filter, suspended in 400 µl HBSS and lysed on ice by a high-intensity ultrasonic processor (Digitane AG, Horgen, Switzerland) for 60 s. After centrifugation, 30 µl of samples were withdrawn and added to 300 µl of BCA assay solution in a 96-well plate. The plate was protected from light and incubated for 2 h at room temperature with mechanical shaking (150 rev./min). The absorbance was read at 570 nm using a microplate reader (Molecular Devices, Sunnyvale, CA, U.S.A.). Standard curves were established using BSA dissolved in HBSS.

Analytical and preparative RP-HPLC of CPP and N-terminal metabolites

By RP-HPLC we studied the following aspects: (i) the chemical stability of selected CF-labelled CPP in buffer, (ii) their metabolic degradation kinetics in contact with MDCK, Calu-3 and TR146 layers, (iii) the quantification of the degradation pattern of CF-labelled hCT(9–32) and its N-terminal metabolites after incubation with MDCK, Calu-3 and TR146. Furthermore, (iv) we applied preparative RP-HPLC to isolate the N-terminal metabolites emerging from hCT(9–32) degradation in contact with Calu-3, to be used for identification by MALDI-TOF-MS (matrix-assisted laser-desorption ionization–time-of-flight mass spectrometry) analysis (see below). Throughout, we used a Merck-Hitachi RP-HPLC (VWR International, Dietikon, Switzerland). Because of the low sensitivity of UV detection, degradation of the peptides was monitored by fluorescence detection of their CF label. Accordingly, detection was limited to intact CF-labelled peptides as well as the resulting N-terminal metabolites carrying the N-terminally ligated fluorophore as model cargo. For hCT-derived peptides, we used a linear gradient mobile phase starting at 10% solvent A consisting of acetonitrile, water and TFA (trifluoroacetic acid) (80:19.99:0.01, by vol.) and 90%

solvent B (10:89.99:0.01, by vol.) to 20% solvent B over 30 min. The flow rate was 1 ml/min. All conditions were identical for analytical and preparative RP-HPLC except for the injected sample volume (analytical, 10 µl; preparative, 400 µl). Fluorescence emission was detected at 517 nm with excitation at 492 nm. A Microsorb-MV100 C18 (150 mm × 4.6 mm; 5 µm) column (Varian, Zug, Switzerland) was used. For Tat(47–57) and penetratin(43–58), we used a linear-gradient mobile phase starting at 90% solvent A (water/TFA, 99.9:0.1, by vol.) and 10% solvent B (acetonitrile/TFA, 99.925:0.075, by vol.) to 100% solvent B over 41 min. A LiChrospher 100 RP-18 column (250 mm × 4 mm, 5 µm; Merck KgaA, Darmstadt, Germany) was used throughout. The flow rate was 1 ml/min and the injected sample volume was 10 µl. Again, fluorescence emission detection was at 517 nm and excitation at 492 nm.

CPP stability in buffer

Selected hCT-derived peptides (40 µM) and 10 µM of Tat(47–57) and penetratin(43–58) were dissolved in HBSS (pH 7.4). The tubes were incubated for 21 days, protected from light at 37 °C with mechanical shaking (150 rev./min). Every second or third day, a sample of 50 µl was taken from each tube and analysed by RP-HPLC to follow the chemical stability of the peptides.

Kinetics of metabolic degradation of CPP in contact with epithelial models

Before investigating metabolic degradation, MDCK cell cultures were equilibrated for 30 min at 37 °C, and Calu-3 and TR146 cells for 15 min at room temperature in HBSS, all at pH 7.4. After equilibration, the TEER (transepithelial electrical resistance) was measured to validate the physical integrity of the epithelial cell layer(s). For measurement, a Millicell-ERS system (Millipore, Bedford, MA, U.S.A.) was used. TEER values (in Ω · cm²) were corrected for the blank filter and normalized for its surface area. The study was performed using the reflection kinetics mode [40], i.e. with the apical side of the monolayer exposed to the peptide solution, whereas the basal side was blocked with a polystyrene well plate acting as an impermeable barrier. Under such conditions, permeation across the cell monolayer is blocked, and the metabolites accumulate in the donor compartment on the apical side. An aliquot of 40 µM CF-labelled hCT(9–32), hCT(12–32), hCT(15–32), hCT(18–32), random sequence and 10 µM Tat(47–57) or penetratin(43–58) were dissolved in HBSS and added to the apical side of the insert. hCT(12–32) was exclusively incubated with Calu-3 monolayers and TR146 multilayers respectively, and hCT(18–32) only with MDCK monolayers. The experiment was conducted at 37 °C, with mechanical shaking (150 rev./min) to minimize the aqueous boundary layer. Samples of 50 µl were taken from the apical side after 10, 20, 30, 40, 50, 60 and 90 min incubation with Calu-3 cells and hourly during 8 h for TR146 and MDCK cells. The kinetics of metabolic degradation of the tested compounds was assessed by RP-HPLC (see above). As a first approximation, the data were evaluated according to pseudo-first-order kinetics. The half-lives, $t_{1/2}$, of the peptides were calculated according to

$$t_{1/2} = \ln 2 / [d(\log C)/dt] \quad (1)$$

where C is the concentration of the peptide and t the time of incubation. To account for the difference in the cross-sectional area of the cell culture inserts, 4.2 cm² with MDCK and 0.9 cm² with both Calu-3 and TR146, the experimental half-lives were normalized by multiplication with a dimensionless factor: 4.2

(MDCK) or 0.9 (Calu-3 and TR146) respectively. This approximation is limited by the assumption of linear kinetics for the full concentration range and excludes enzyme saturation. Also all effects of diffusion kinetics were neglected. For independent, model-free evaluation of the data, we also calculated initial rates of degradation ($\mu\text{M} \cdot \text{cm} \cdot \text{min}^{-1}$) approximated by linear regression of the three initial data pairs of the degradation profiles.

MALDI-TOF-MS

Metabolites emerging from the degradation of CF-labelled hCT(9–32) in contact with epithelial models were identified by MALDI-TOF-MS. Both RP-HPLC-enriched fractions of specific metabolites and samples taken directly from cell culture incubates were analysed. For this purpose, 2 μl of samples containing peptides and/or metabolites were mixed with 3 μl of a saturated matrix solution consisting of 2,5-dihydroxybenzoic acid in acetonitrile/water (2:1, v/v) with 0.1% TFA. This solution (1 μl) was applied to the target plate, allowed to crystallize, de-dusted and analysed with delayed extraction (150 ns) and reflectron mode. Acceleration voltage was at 20 kV. Acquisition mass ranged from approx. 0.5 to 3.5 kDa. Calibration was performed using standard peptides: des-Arg-bradykinin (molecular mass, 904.7 Da), angiotensin 1 (molecular mass, 1297 Da) and Glu-Fibrinopeptide B (molecular mass, 1571.1 Da). All measurements were performed using a MALDI-TOF mass spectrometer (Voyager-DE Elite Biospectrometry workstation, PerSeptive Biosystems, Farmington, MA, U.S.A.).

Identification of metabolites emerging from hCT(9–32) degradation

Metabolites were identified in two ways. Identification of selected N-terminal metabolites of hCT(9–32) was through RP-HPLC fractionation and peak allocation by MALDI-TOF-MS. Briefly, 0.8 ml of 40 μM CF-labelled hCT(9–32) was incubated for 120 min with Calu-3 monolayers following the specifications described for the kinetics study (see above). At 0, 20, 40, 60 and 120 min, 120 μl of samples were taken. The experiment was repeated five times and all samples from 60 and 120 min were pooled for subsequent preparative RP-HPLC (see above). The respective RP-HPLC eluates containing CF-hCT(9–32) and its metabolites were collected using a L-7655 fraction collector (Merck, Darmstadt, Germany), and each fraction was concentrated by freeze drying. Because of fluorescence detection, analysis was limited to N-terminally labelled CF-metabolites. For MS, the freeze-dried fractions were dissolved in 20 μl of HBSS and analysed by MALDI-TOF-MS.

As a second approach, we incubated 0.5 ml of 40 μM CF-labelled hCT(9–32) with Calu-3 and TR146 multilayers, again using the reflection kinetics approach (see above) [40]. Samples were taken from the apical chamber after 10, 20, 50 and 90 min incubation with Calu-3 cells, and after 1, 2, 4, 6, 7 and 8 h incubation with TR146 cells. All samples were directly analysed by MALDI-TOF-MS for identification of N- and C-terminal metabolites. For identification of peptidases involved in the metabolic degradation of hCT(9–32) we used Peptide Cutter, an online software available at the ExPASy Molecular Biology Server (www.expasy.org; Swiss Institute of Bioinformatics, Switzerland).

Quantification of hCT(9–32) and its N-terminal metabolites

A 0.5 ml volume of 40 μM CF-labelled hCT(9–32) was incubated with one of the epithelial models MDCK, Calu-3 or TR146 following the same method as described for the kinetics study (see above). Samples of 50 μl were taken from the apical side after 10, 20, 30, 40, 50, 60 and 90 min incubation with Calu-3

Table 2 Peptide stability in buffer after 21 days

The given values represent mean percentages of intact peptide ($n = 2$).

HCT(9–32)	hCT(12–32)	hCT(15–32)	Random sequence	Tat(47–57)	Penetratin(43–58)
39%	87%	100%	91%	100%	76%

cells, and hourly during 8 h for MDCK and TR146. All samples were analysed by RP-HPLC, and the amount of each N-terminal metabolite was expressed as a percentage of the total fluorescence over time. Allocation of the various RP-HPLC peaks was by MALDI-TOF-MS (see above). Total fluorescence of the CF label was checked to be constant and independent of the lengths of the N-terminal peptide fragments still carrying the CF label [29].

RESULTS

Peptide stability in buffer

For orientation we performed exploratory chemical stability studies of all CF-labelled peptides in HBSS buffer and analysed for intact peptides by RP-HPLC. As demonstrated in Table 2, even after 21 days of incubation in buffer, all peptides, except for hCT(9–32) and penetratin(43–58), showed high percentages of intact peptides, being 87% or higher. Over the same period of time, 76 and 39% of penetratin(43–58) and hCT(9–32) respectively remained chemically intact. Thus all compounds demonstrated reasonable chemical stability in buffer for any of the subsequent metabolic degradation studies of up to 8 h (see below). Moreover, the subsequently observed cleavage of the CPP after incubation with the three epithelial models (see below) was essentially due to proteolytic activity and not the result of chemical degradation.

Kinetics of metabolic degradation of CPP in contact with epithelial models

To evaluate contrasts in the rates of metabolic cleavage between the investigated cell models, we assessed the metabolic degradation of the CF-labelled CPP and the random sequence after incubation with three epithelial models: MDCK, Calu-3 and TR146. Exclusively intact peptides were monitored. Cell layer integrity was checked before each experiment by measuring the TEER values. The measured values were 111 ± 21 ($n = 68$), 362 ± 86 ($n = 54$) and $164 \pm 79 \Omega \cdot \text{cm}^2$ ($n = 58$) for MDCK, for Calu-3 and for TR146 respectively. The values were consistent with previously reported TEER values of the investigated epithelial models [35,39,41–44].

As a rough approximation, the obtained degradation profiles of the CPP and random sequence were evaluated according to pseudo-first-order kinetics. Normalized half-lives as defined in the Experimental section were calculated. Since rate-limiting effects by enzyme saturation and diffusion kinetics were neglected, the significance of the results is empirical. The results are given in Table 3. All compounds incubated in the presence of MDCK cell monolayer demonstrated normalized half-lives ranging from 1421 min for the random sequence to 6065 min for Tat(47–57). Shorter half-lives were observed with the TR146 model showing normalized half-lives in the range 166–601 min. Interestingly, hCT(9–32) was the most stable peptide in the TR146 model.

In contrast, degradation in the presence of Calu-3 was fastest, with normalized half-lives ranging between 15 and 45 min. Tat(47–57) was an exception from this trend by featuring a half-life of 544 min when incubated with Calu-3 (Table 3).

Table 3 Metabolic degradation of hCT-derived peptides, Tat(47–57) and penetratin(43–58) on different epithelial cell models represented as half-life in min

Normalized half-lives are given in parentheses. Results are means $\pm$ S.D. ($n = 3$). n.d., not determined.

Peptides	Half-life (min)		
	MDCK	Calu-3	TR146
hCT(9–32)	397 $\pm$ 56 (1668 $\pm$ 236)	23 $\pm$ 5 (21 $\pm$ 5)	662 $\pm$ 45 (601 $\pm$ 40)
hCT(12–32)	n.d.	17 $\pm$ 2 (15 $\pm$ 2)	420 $\pm$ 50 (382 $\pm$ 45)
hCT(15–32)	n.d.	35 $\pm$ 9 (31 $\pm$ 8)	256 $\pm$ 86 (233 $\pm$ 78)
hCT(18–32)	719 $\pm$ 44 (3019 $\pm$ 187)	n.d.	n.d.
Random sequence	338 $\pm$ 86 (1421 $\pm$ 360)	19 $\pm$ 4 (18 $\pm$ 3)	310 $\pm$ 161 (282 $\pm$ 146)
Tat(47–57)	1444 $\pm$ 408 (6065 $\pm$ 1715)	598 $\pm$ 379 (544 $\pm$ 345)	553 $\pm$ 70 (503 $\pm$ 63)
Penetratin(43–58)	495 $\pm$ 1 (2079 $\pm$ 1)	50 $\pm$ 14 (45 $\pm$ 13)	183 $\pm$ 12 (166 $\pm$ 11)

Table 4 Protein contents and cell densities in MDCK, Calu-3 and TR146 epithelial models

Results are means $\pm$ S.D. ($n = 3$). n.d., not determined.

Cell model	Protein content ($\mu\text{g}/\text{cm}^2$)	Cells/ cm^2
MDCK	269 $\pm$ 19	n.d.
Calu-3	1876 $\pm$ 455	940 329 $\pm$ 51 769
TR146	396 $\pm$ 48	475 309 $\pm$ 26 189

The data may be interpreted with respect to the typical cell densities and protein contents of the three epithelial models. As summarized in Table 4, the number of cells/ cm^2 was 2-fold higher in Calu-3 when compared with TR146. Protein content ($\mu\text{g}/\text{cm}^2$) in Calu-3 was 4–7-fold higher than that of MDCK and TR146, which correlates reasonably well with the higher normalized half-lives found with these models. The high protein content of the Calu-3 model may explain its large enzymic capacity to metabolize peptides.

Model-free evaluation of the data by calculation of the initial rates (in $\mu\text{M} \cdot \text{cm} \cdot \text{min}^{-1}$) of proteolytic degradation of the peptides, as approximated by linear regression of the initial three data pairs of the degradation profiles, led to similar conclusions (results not shown). The Calu-3 model showed the fastest metabolic rates followed by TR146 and MDCK. Moreover, degradation of Tat(47–57) was the slowest among the studied CPP.

Identification of metabolites emerging from hCT(9–32) degradation in contact with Calu-3

We first analysed the occurrence of metabolites of CF-labelled hCT(9–32) after incubation with Calu-3 monolayers. RP-HPLC in combination with fluorescence detection was used. Thus, in addition to intact CF-labelled hCT(9–32), only N-terminal metabolites carrying the fluorescent CF-cargo were detectable, and C-terminal metabolites lacking the CF-cargo excluded. When incubated with Calu-3 monolayers, hCT(9–32) degraded into four main N-terminal metabolites (Figure 1). To identify these metabolites by MS, a pooled sample was fractionated into 13 fractions. RP-HPLC chromatograms of four of them are given in Figure 2 (left panels) and display the respective metabolites. Subsequent peak allocation analysis of the four RP-HPLC fractions by MALDI-TOF-MS allowed to unveil their identity (Figure 2, right panels). For control, pure buffer solution was incubated with Calu-3 instead of peptide solution. MALDI-TOF-MS analysis of this incubate showed only peaks below approx. 0.6 kDa (results not shown). This excludes significant interference of non-CPP-derived compounds with the analysis of CF-labelled hCT(9–32) and its metabolites above 0.6 kDa. The observed m/z

Figure 1 RP-HPLC chromatographs of CF-labelled hCT(9–32) and its N-terminal metabolites after metabolic degradation by Calu-3 layers using the reflection kinetics mode

At zero time, hCT(9–32) was nearly intact and practically represented by a single peak. After 60 min of incubation with Calu-3 cells, hCT(9–32) was almost fully degraded into four main N-terminal CF-labelled metabolites, indicated as metabolites 1–4.

values of the MALDI-TOF-mass spectra correspond to $[M + H]^+$. Furthermore, we typically observed associated $[M + Na]^+$ peaks for each metabolite. As shown in Figure 2, metabolite 1 with a RP-HPLC retention time (Rt) of 13.6 min corresponds to the fragment hCT(9–15). Metabolite 2 (Rt = 14.3 min) was identified as hCT(9–11), metabolite 3 (Rt = 15.6 min) was hCT(9–12) and metabolite 4 (Rt = 18.5 min) was hCT(9–14).

MALDI-TOF-MS analysis was also performed by direct analysis of samples containing N- and C-terminal metabolites emerging from the degradation of hCT(9–32) in contact with Calu-3. From the results obtained after 10, 20, 50 and 90 min, two typical MALDI-TOF-mass spectra are shown in Figure 3. After 20 min, the N-terminal metabolites hCT(9–11), hCT(9–12), hCT(9–15) and their C-terminal counterparts hCT(12–32), hCT(13–32) and hCT(16–32) were detected. Moreover, the fragment hCT(17–32) was monitored without its N-terminal counterpart. After 90 min, additional fragments such as hCT(9–14) and hCT(18–32) were detected, whereas their respective C- and N-terminal counterparts remained undetectable (Figure 3).

Figure 2 Identification of CF-labelled N-terminal metabolites emerging from hCT(9-32) degradation in contact with Calu-3 monolayers

CF-hCT(9-32) was incubated for 120 min with Calu-3 monolayer, using the reflection kinetics mode. Samples (60 and 120 min) were pooled and their N-terminal metabolites fractionated by preparative RP-HPLC. For RP-HPLC peak allocation, fractions were further analysed by MALDI-TOF-MS. Metabolite 1 (Rt = 13.6 min) was hCT(9-15); metabolite 2 (Rt = 14.3 min) was hCT(9-11). Metabolites 3 and 4 (Rt = 15.6 and 18.5 min) respectively were hCT(9-12) and hCT(9-14). For each metabolite, the observed m/z values represent $[M + H]^+$. The associated peaks, observed for each metabolite, represent $[M + Na]^+$.

The fragment hCT(18-32) was monitored after 50 min (results not shown). The fragments hCT(9-11), hCT(12-32) and hCT(13-32) were undetectable after 90 min, suggesting that hCT(12-32) and hCT(13-32) could have been further degraded after 20 min.

Figure 4(A) illustrates a summary of the suggested cleavage pattern of hCT(9-32) in contact with Calu-3. Initial cleavage seems to occur at four different sites, namely between the residues Thr¹¹-Tyr¹², Tyr¹²-Thr¹³ and Asp¹⁵-Phe¹⁶, as demonstrated by the concomitant occurrence of N- and C-terminal counterparts, and between residues Phe¹⁶-Asn¹⁷, documented by the C-terminal fragment. Subsequently, further truncations of initial metabolites

appear between residues Gln¹⁴-Asp¹⁵ and Asn¹⁷-Lys¹⁸, as evidenced by either N- or C-terminal fragments.

Identification of metabolites emerging from hCT(9-32) degradation in contact with MDCK and TR146

When incubated with both MDCK monolayers and TR146 multilayers, CF-labelled hCT(9-32) degraded into a total of eight N-terminal metabolites (Figure 5). Both cell lines yielded roughly identical metabolite patterns, with four metabolites identical with those identified after incubation with Calu-3 (Figure 1).

Figure 3 Typical MALDI-TOF-MS of N- and C-terminal metabolites of CF-labelled hCT(9-32) after metabolic degradation by Calu-3 monolayer using the reflection kinetics mode

Results obtained by direct sampling of cell culture incubates; 20 and 90 min MALDI-TOF-MS spectra shown. For each metabolite, the observed m/z values represent $[M + H]^+$. The associated peaks, observed for each metabolite, represent $[M + Na]^+$. Note the expansion of the 90 min spectrum in the m/z range of approx. 0.7–2 kDa.

Because of the similarity of the cleavage patterns of CF-hCT(9-32) in contact with MDCK and TR146, only TR146 was further considered. Samples were directly analysed by MALDI-TOF-MS. The resulting cleavage pattern is depicted in Figure 4(B). After 2 h of incubation, cleavage sites were found between the residues Thr¹¹-Tyr¹², Tyr¹²-Thr¹³, Gln¹⁴-Asp¹⁵ and Asp¹⁵-Phe¹⁶ as documented by concomitant occurrence of N- and C-terminal counterparts. Between 2 and 8 h of incubation, additional fragments, such as hCT(17-32), hCT(20-32) and the N-terminal metabolite hCT(9-13) were found, but without their respective N- and C-terminal counterparts. A summary of the expected and measured values of the molecular masses corresponding to the different metabolites is given in Table 5.

Quantification of hCT(9-32) and its N-terminal metabolites

Quantitative analysis by RP-HPLC under fluorescence detection yielded typical distribution patterns of CF-hCT(9-32) and its

metabolites as a function of time and depending on the epithelial model (Figure 6). The data are relative and strictly refer to N-terminal fragments still carrying the CF label.

Calu-3. Metabolism kinetics by Calu-3 monolayers was fast. Within 90 min, practically all initial CF-hCT(9-32) was degraded (Figure 6). Four N-terminal metabolites resulted: the major metabolite was hCT(9-12), reaching a total of 53.5% within this time frame. Initially, hCT(9-15) increased in content until 40 min of incubation and then decreased between 60 and 90 min to 8% only. In parallel, hCT(9-14) strongly increased between 60 and 90 min to reach 14%. Overall, the content of hCT(9-11) slightly increased during the experiment.

MDCK. In MDCK, the main N-terminal metabolite was hCT(9-12) reaching a final of 48.5% of the metabolites after 8 h of incubation (Figure 6). During the same time, two other identified N-terminal metabolites, hCT(9-11) and hCT(9-15), reached percentages of 22% each. Metabolite hCT(9-15) decreased slightly in content after 7 h, whereas the fractions of

Figure 4 Scheme of suggested metabolic cleavage sites of CF-labelled hCT(9–32) after incubation with Calu-3 (A) and TR146 (B) layers

Arrowheads indicate cleavage sites in CF-hCT(9–32). Black arrowheads, sites of concomitantly detected N- and C-terminal counterparts; grey arrowheads, only N-terminal fragment detected; open arrowheads, only C-terminal fragment detected. Results indicate similar primary cleavage of CF-hCT(9–32) occurs in the N-terminal domain in both cell models.

hCT(9–11) and hCT(9–12) continued to increase over time. Metabolite hCT(9–14) was present in very low amounts only. Two of the four non-identified N-terminal metabolites, i.e. those with retention times of 15 and 17 min, were clearly monitored during the whole period of the study. The content of the 17-min fragment decreased sharply after 5 h, whereas that of the 15-min fragment was rather constant. The remaining N-terminal metabolites with retention times of 16.3 and 19.5 min were present only in very low amounts.

TR146. When incubated with TR146 multilayers, the fractions of the N-terminal metabolites hCT(9–11), hCT(9–12) and hCT(9–15) increased constantly in a parallel manner over the full course of the study to reach 21, 13 and 18 % respectively, after 8 h (Figure 6). The occurrence of hCT(9–14) was lower than that of the three others and remained approximately constant during the full time frame of 8 h. The 17-min metabolite decreased towards the end of the experiment, whereas the content of the 15-min fragment increased slightly. The 16.3 and 19.5 min metabolites were also detected, but at very low amounts only.

DISCUSSION

Whereas the primary encounter of CPP with membrane-associated proteolytic enzymes, e.g. in the extracellular matrix or in the lipid bilayer, could be similar for all classes of CPP,

their individual pathways of intracellular distribution will decide the type of proteolytic environment the CPP will encounter. For instance, an endocytic pathway may expose CPP-modified therapeutics to late endosomal and lysosomal degradation and, therefore, represent a problematic pathway for drug delivery. The cellular entry of the polycationic CPP penetratin, Tat and oligo-arginine, was identified to follow an endocytic mechanism. Subsequently, the CPP were shown to be either subject to endosomal proteases or leave the endocytic compartment in a rather early stage of endosomal maturation avoiding degradation [45]. Even a translocated cargo may retain its biological activity when transported via an endocytic pathway. For instance, when fused to Tat(47–57) and internalized by endocytosis, there was significant improvement in the biodistribution of active β -glucuronidase imparted by the Tat motif in a mouse model of β -glucuronidase deficiency [46]. Non-endocytic pathways of CPP were also shown to lead to proteolytic cleavage. For example, pVEC, a vascular endothelial cadherin-derived CPP [18], transportan and a transportan derivative [19] showed partial or high biodegradation when incubated with various cell lines [13,17].

In a previous study [14], we demonstrated that truncated sequences of hCT, ranging from hCT(9–32) to hCT(18–32), as well as Tat(47–57) and penetratin(43–58) can translocate the plasma membrane of MDCK monolayers and show punctuated cytoplasmic distribution. In Calu-3 monolayers and TR146 multilayers, hCT(9–32), Tat(47–57) and penetratin(43–58) were subject to either vesicular cytoplasmic or paracellular accumulation [29]. Thus hCT-derived CPP have the potential to carry therapeutic cargos across the plasma membrane of epithelial barriers. The ultimate requirements for a therapeutic agent ligated to a CPP to become active are the following: (i) sufficient metabolic stability of the construct at the site of administration, (ii) subsequent delivery and sufficient release of a functional unit containing the cargo and (iii) efficient clearance from the site without local or systemic toxicity.

To evaluate the potential of hCT-derived peptides, Tat(47–57) and penetratin(43–58) as drug delivery vectors, we investigated their metabolic stability in contact with three epithelial models, MDCK, Calu-3 and TR146. Furthermore, with a special focus on CF-labelled hCT(9–32), we identified typical metabolites and the related patterns of metabolic cleavage. To our knowledge, our investigation is among the first in-depth *in vitro* metabolism studies for CPP, in addition to the work of Langel and co-workers on pVEC [17] and to some extent on transportan, TP10 and penetratin(43–58) [13].

In the presence of Calu-3 monolayers, we observed markedly higher rates of metabolic degradation than with the two other models, MDCK and TR146. Confluent Calu-3 cell cultures are covered with a mucus layer that acts as a protective barrier against microorganisms, toxins and abrasive particulates [32,47]. The total protein content of Calu-3 is significantly higher than that of the two other epithelial models, MDCK and TR146, and correlates well with its observed higher metabolic activity. In contrast, MDCK and TR146 lack the formation of a mucus layer and show weaker proteolytic activity. TR146 originates from a neck-node metastasis of a human buccal carcinoma [31]. Its stratified squamous phenotype is similar to that of the epidermis, which is known for its metabolism of xenobiotics as demonstrated, e.g. by Boderke et al. [48]. Among all tested compounds, Tat(47–57) was the most stable CPP. Even in contact with Calu-3, its metabolic half-life was 544 min, whereas the values of the other CPP ranged from 15 to 45 min only. The metabolic stabilities of the hCT-derived peptides in contact with Calu-3 were poor and exhibited half-lives similar to those of full-length hCT when incubated with rabbit small intestine luminal

Figure 5 Typical RP-HPLC profiles of N-terminal metabolites after metabolic degradation of CF-labelled hCT(9–32) in contact with MDCK and TR146 layers

Study performed under reflection kinetics. At zero time, CF-hCT(9–32) was fully intact and was represented by a single peak. After 8 h of incubation with MDCK and TR146 cells, both cell-culture models yielded practically identical patterns of N-terminal CF-labelled metabolites. Metabolites 1–4 correspond to the four metabolites identified in the Calu-3 model (see Figure 1). The remaining peaks 5–8 indicate non-identified N-terminal CF-labelled metabolites.

Table 5 Measured and expected molecular masses of detected metabolites by MALDI-TOF-MS after incubation in the TR146 model

Measured and expected molecular mass values represent $[M + H]^+$.

Peptides	Measured molecular mass (Da)	Expected molecular mass (Da)
hCT(9–32)	2970.5/2970.9	2968.5
hCT(9–15)	1155.8/1155.4	1155.2
hCT(9–14)	1041.0/1038.1	1040.1
hCT(9–13)	912.5	912.0
hCT(9–12)	811.6/811.6	810.8
hCT(9–11)	648.4/648.3	647.7
hCT(12–32)	2340.2	2338.9
hCT(13–32)	2177.2	2175.7
hCT(15–32)	1947.7	1946.4
hCT(16–32)	1832.7	1831.3
hCT(17–32)	1685.4	1684.1
hCT(18–32)	1570.7	1570.0
hCT(20–32)	1294.8	1294.6

enzymes, trypsin, α -chymotrypsin and elastase [49]. In contrast, when incubated with MDCK, hCT-derived peptides were quite stable.

In Calu-3, hCT(9–32) was initially cleaved at four sites, Thr¹¹-Tyr¹², Tyr¹²-Thr¹³, Asp¹⁵-Phe¹⁶ and Phe¹⁶-Asn¹⁷. The most probable cleavage type at the sites Thr¹¹-Tyr¹², Tyr¹²-Thr¹³, and Asp¹⁵-Phe¹⁶ was by endopeptidases, yielding the occurrence of both the C- and N-terminal counterparts of the native peptide at the same time. Cleavage at Phe¹⁶-Asn¹⁷ may have occurred by aminopeptidase-type cleavage of hCT(16–32) at the N-terminus, because the N-terminal counterpart of hCT(17–32) was not detected. After

50 min incubation, a C-terminal metabolite, hCT(18–32), was also observed, which could have originated from an N-terminal aminopeptidase-type cleavage of hCT(17–32). Fragment hCT(9–14), detected without its C-terminal counterpart, is a probable product of carboxypeptidase-type cleavage of hCT(9–15). This suggestion correlates with the results illustrated in Figure 6, showing an increase in the fraction of hCT(9–14) concomitant to a decrease in hCT(9–15).

In the TR146 model, hCT(9–32) was degraded at similar cleavage sites. However, an additional endopeptidase-type cleavage site was found between residues Gln¹⁴ and Asp¹⁵, since both hCT(9–14) and hCT(15–32) were detected at the same time. In contrast, because of the absence of the C-terminal counterpart of hCT(9–13), this fragment may result from the degradation of longer N-terminal metabolites by carboxypeptidases or endopeptidases. Hypothetically, neither hCT(9–14) nor hCT(9–15) could be the source of hCT(9–13) since both rates did not decrease over time. The longer fragment may correspond to one of the non-identified N-terminal metabolites (see Figure 5). Furthermore, hCT(17–32) may have originated from N-terminal aminopeptidase-type cleavage of hCT(16–32) and hCT(20–32) from endopeptidase-type cleavage of longer C-terminal metabolites.

Finally, for the metabolic cleavage of hCT(9–32) in the MDCK model, we conclude a similar pattern as observed in the TR146 model. This is supported by the identity of the eight metabolites found after incubation in both models (see Figure 5). A particular similarity between MDCK and Calu-3 was found with hCT(9–12), representing the major N-terminal metabolite resulting from predominant cleavage at the site Tyr¹²-Thr¹³.

The various cleavage sites are characteristic for a variety of enzymes. Cleavage between residues Thr¹¹-Tyr¹² probably results

Figure 6 Quantitative evaluation by RP-HPLC of CF-labelled hCT(9-32) and its N-terminal metabolites over time in contact with Calu-3, MDCK and TR146 layers

hCT(9-32) was incubated for up to 8 h with MDCK and TR146, or 90 min with Calu-3. Hourly sampling was carried out with MDCK and TR146, and every 10 (or 30) min with Calu-3. For each time point, the concentration of the various N-terminal metabolites is represented as percentage of normalized total fluorescence. In the MDCK panel, the contributions of hCT(9-14) and peak 19.5 min are too small for graphical distinction.

from chymotrypsin, pepsin or proteinase K activity, whereas cleavage at the linkage $\text{Gln}^{14}\text{-Asp}^{15}$ is indicative of Asp-N endopeptidase activity. Pepsin activity would also correspond to cleavage at $\text{Asp}^{15}\text{-Phe}^{16}$. So far, we have not been able to identify a specific enzyme for the cleavage site at $\text{Tyr}^{12}\text{-Thr}^{13}$. Overall, initial hCT(9-32) degradation is suggested to occur by endopeptidase activity, and it concentrates on the N-terminal domain, followed by cleavage through aminopeptidase, endopeptidase and/or carboxypeptidase activity on the emerging metabolites.

Taking into consideration that hCT(9-32) carries its cargo ligated to the N-terminus, the cargo may be prematurely detached from the CPP before cellular internalization, e.g. in contact with peptidases secreted from the cell layers or peptidases located in the cell membrane. In this way, the cargo may be lost and never reach the cytoplasm. Fast metabolic degradation of CPP in the Calu-3 model might explain the exclusive paracellular staining of Calu-3 cells in the presence of N-terminally labelled CF-hCT(9-32) as observed in a previous study [29]. Conjugation

of the cargo to a more stable site of hCT(9-32) could be an alternative, e.g. to the side chain of Lys^{18} , but needs experimental verification. Comparably, the fluorescence seen in the cytoplasm of MDCK and TR146 models might be the result of the lower metabolic activity of these models. The complete degradation of labelled hCT(9-32) in Calu-3 also explains results from the same study that no intact peptide, but only metabolites, was observed to permeate through this epithelium. The hypothesis is supported by the fact that intact hCT(9-32) was found to permeate the MDCK and TR146 epithelia, yet in very low amounts. In the Calu-3 model, Tat(47-57) was significantly more stable than any of the other tested CPP and was the only CPP to permeate the monolayer partially intact [29]. This further supports the statement that for improved intracellular localization as well as for increased transepithelial permeability, metabolic stability of the peptide is a crucial factor.

In an earlier study, we investigated the metabolic cleavage pattern of full-length hCT in contact with excised nasal mucosa [20]. The initial metabolic cleavage of this peptide was mainly due to chymotryptic and tryptic endopeptidase activity in the midsection of the molecule between residues $\text{Phe}^{16}\text{-Asn}^{17}$, $\text{Lys}^{18}\text{-Phe}^{19}$ and $\text{Phe}^{19}\text{-His}^{20}$, following an endopeptidase activity pattern. Further cleavage was then observed to occur by aminopeptidase activity leading to stepwise truncated C-terminal metabolites. Two additional cleavage sites were located in the N-terminal domain of hCT between amino acids $\text{Met}^8\text{-Leu}^9$ and $\text{Leu}^9\text{-Gly}^{10}$. Degradation of hCT was also studied in contact with excised rabbit small intestine mucosa [49]. The authors showed a similar degradation pattern as that found by Lang et al. [20] in excised bovine nasal mucosa.

We suggest that one of the reasons behind the differences between the degradation patterns of hCT(9-32) versus full-length hCT will probably depend on differences in the steric accessibility for proteolytic enzymes. Indeed, full-length hCT features a disulphide bridge forming a loop-like structure at the N-terminus, between residues Cys^1 and Cys^7 , which may stabilize this region from heavy degradation. Both eel and hCT form a stable loop-forming disulphide bridge at the N-terminus, followed by a stable α -helix in the midsection and a random coil region towards the C-terminus [50]. In contrast, the fragment hCT(9-32) probably will not adopt the same secondary structure since it lacks the respective disulphide-bridged loop at the N-terminus. Therefore the N-terminal domain of hCT(9-32) is suggested to be more easily accessible for proteolytic enzymes.

In conclusion, we demonstrated a higher stability of Tat(47-57) versus hCT-derived CPP and penetratin(43-58) in all three investigated epithelial models, i.e. MDCK, Calu-3 and TR146. Among these three models, Calu-3 featured the highest level of proteolytic activity leading to the rather poor stability of most of the CPP incubated with this model. Initial degradation of hCT(9-32) occurred in its N-terminal domain through endopeptidase activity. The resulting metabolites were then further degraded by aminopeptidases, endopeptidases and/or carboxypeptidases. A selection of metabolites was identical in all three epithelial models, suggesting similarity of the major degradation pathways.

To overcome the enzymic barrier of the epithelial layers, two principal approaches to enhance hCT(9-32) stability can be suggested: (i) the use of specific endopeptidase inhibitors and (ii) the chemical modification of hCT(9-32) in the region of its initial cleavage sites to obtain analogues with higher enzymic stability. Stabilization may be achieved by the replacement of L-amino-acids of hCT(9-32) by their non-physiological D-counterparts, or by N-methylation close to major cleavage sites. Improved stability of hCT-derived CPP will probably exhibit a higher uptake rate.

However, in addition to its negative connotation, it needs to be kept in mind that metabolism of CPP is at the same time a prerequisite for the release of chemically ligated cargos after internalization. We expect the balance between both aspects to require the development of carefully engineered CPP, to avoid premature cleavage and to free the cargo once internalized.

The authors acknowledge Dr R. Brunisholz (ETH Zurich) for his advice on MALDI-TOF-MS, Dr S. R. K. Hoffmann (University of Zurich) for the purification of Tat and penetratin, and Ruth Alder (ETH Zurich) for her help and advice with RP-HPLC. This work was supported by an ETH Zurich grant for R. T., research project number 0-20702-99, and the Commission of the European Union (EU project on Quality of Life and Management of Living Resources, Project No. QLK2-CT-2001-01451). Further, the project was supported by an Alfred Benzon Foundation grant to H. M. N.

REFERENCES

- Gewirtz, A. M., Sokol, D. L. and Ratajczak, M. Z. (1998) Nucleic acid therapeutics: state of the art and future prospects. *Blood* **92**, 712–736
- Juliano, R. L., Alahari, S., Yoo, H., Kole, R. and Cho, M. (1999) Antisense pharmacodynamics: critical issues in the transport and delivery of antisense oligonucleotides. *Pharm. Res.* **16**, 494–502
- Frankel, A. D. and Pabo, C. O. (1988) Cellular uptake of the tat protein from human immunodeficiency virus. *Cell (Cambridge, Mass.)* **55**, 1189–1193
- Mann, D. A. and Frankel, A. D. (1991) Endocytosis and targeting of exogenous HIV-1 Tat protein. *EMBO J.* **10**, 1733–1739
- Schwarze, S. R., Hruska, K. A. and Dowdy, S. F. (2000) Protein transduction: unrestricted delivery into all cells? *Trends Cell Biol.* **10**, 290–295
- Bogoyevitch, M. A., Kendrick, T. S., Ng, D. C. and Barr, R. K. (2002) Taking the cell by stealth or storm? Protein transduction domains (PTDs) as versatile vectors for delivery. *DNA Cell Biol.* **21**, 879–894
- Nagahara, H., Vocero-Akbani, A. M., Snyder, E. L., Ho, A., Latham, D. G., Lissy, N. A., Becker-Hapak, M., Ezhevsky, S. A. and Dowdy, S. F. (1998) Transduction of full-length TAT fusion proteins into mammalian cells: TAT-p27Kip1 induces cell migration. *Nat. Med. (NY)* **4**, 1449–1452
- Torchilin, V. P., Rammohan, R., Weissig, V. and Levchenko, T. S. (2001) TAT peptide on the surface of liposomes affords their efficient intracellular delivery even at low temperature and in the presence of metabolic inhibitors. *Proc. Natl. Acad. Sci. U.S.A.* **98**, 8786–8791
- Astriab-Fisher, A., Sergueev, D., Fisher, M., Shaw, B. R. and Juliano, R. L. (2002) Conjugates of antisense oligonucleotides with the Tat and antennapedia cell-penetrating peptides: effects on cellular uptake, binding to target sequences, and biological actions. *Pharm. Res.* **19**, 744–754
- Koppelhus, U., Awasthi, S. K., Zachar, V., Holst, H. U., Ebbesen, P. and Nielsen, P. E. (2002) Cell-dependent differential cellular uptake of PNA, peptides, and PNA-peptide conjugates. *Antisense Nucleic Acid Drug Dev.* **12**, 51–63
- Richard, J. P., Melikov, K., Vives, E., Ramos, C., Verbeure, B., Gait, M. J., Chernomordik, L. V. and Lebleu, B. (2003) Cell-penetrating peptides: a re-evaluation of the mechanism of cellular uptake. *J. Biol. Chem.* **278**, 585–590
- Lundberg, M., Wikstrom, S. and Johansson, M. (2003) Cell surface adherence and endocytosis of protein transduction domains. *Mol. Ther.* **8**, 143–150
- Lindgren, M. E., Hallbrink, M. M., Elmquist, A. M. and Langel, U. (2004) Passage of cell-penetrating peptides across a human epithelial cell layer *in vitro*. *Biochem. J.* **377**, 69–76
- Tréhin, R., Krauss, U., Muff, R., Meinecke, M., Beck-Sickinger, A. and Merkle, H. P. (2004) Cellular internalization of human calcitonin derived peptides in MDCK monolayers: a comparative study with Tat(47–57) and penetratin(43–58). *Pharm. Res.* **21**, 33–42
- Fix, J. A. (1996) Oral controlled release technology for peptides: status and future prospects. *Pharm. Res.* **13**, 1760–1764
- Garner, P., Sherry, B., Moilanen, S. and Huang, Y. (2001) *In vitro* stability of alpha-helical peptide nucleic acids (alphaPNAs). *Bioorg. Med. Chem. Lett.* **11**, 2315–2317
- Elmquist, A. and Langel, U. (2003) *In vitro* uptake and stability study of pVEC and its All-D analog. *Biol. Chem.* **384**, 387–393
- Elmquist, A., Lindgren, M., Bartfai, T. and Langel, U. (2001) VE-cadherin-derived cell-penetrating peptide, pVEC, with carrier functions. *Exp. Cell Res.* **269**, 237–244
- Soomets, U., Lindgren, M., Gallet, X., Hallbrink, M., Elmquist, A., Balaspiri, L., Zorko, M., Pooga, M., Brasseur, R. and Langel, U. (2000) Deletion analogues of transportan. *Biochim. Biophys. Acta* **1467**, 165–176
- Lang, S. R., Staudenmann, W., James, P., Manz, H. J., Kessler, R., Galli, B., Moser, H. P., Rummelt, A. and Merkle, H. P. (1996) Proteolysis of human calcitonin in excised bovine nasal mucosa: elucidation of the metabolic pathway by liquid secondary ionization mass spectrometry (LSIMS) and matrix assisted laser desorption ionization mass spectrometry (MALDI). *Pharm. Res.* **13**, 1679–1685
- Steinstrasser, I., Koopmann, K. and Merkle, H. P. (1997) Epidermal aminopeptidase activity and metabolism as observed in an organized HaCaT cell sheet model. *J. Pharm. Sci.* **86**, 378–383
- Boderke, P., Bodde, H. E., Ponec, M., Wolf, M. and Merkle, H. P. (1998) Mechanistic and quantitative prediction of aminopeptidase activity in stripped human skin based on the HaCaT cell sheet model. *J. Investig. Dermatol. Symp. Proc.* **3**, 180–184
- Lang, S., Rothen-Rutishauser, B., Perriard, J. C., Schmidt, M. C. and Merkle, H. P. (1998) Permeation and pathways of human calcitonin (hCT) across excised bovine nasal mucosa. *Peptides* **19**, 599–607
- Schmidt, M. C., Rothen-Rutishauser, B., Rist, B., Beck-Sickinger, A., Wunderli-Allenspach, H., Rubas, W., Sadee, W. and Merkle, H. P. (1998) Translocation of human calcitonin in respiratory nasal epithelium is associated with self-assembly in lipid membrane. *Biochemistry* **37**, 16582–16590
- Machova, Z., Muhle, C., Krauss, U., Tréhin, R., Koch, A., Merkle, H. P. and Beck-Sickinger, A. G. (2002) Cellular internalization of enhanced green fluorescent protein ligated to a human calcitonin-based carrier peptide. *ChemBiochem* **3**, 672–677
- Misfeldt, D. S., Hamamoto, S. T. and Pitelka, D. R. (1976) Transepithelial transport in cell culture. *Proc. Natl. Acad. Sci. U.S.A.* **73**, 1212–1216
- Horster, M. F. and Stopp, M. (1986) Transport and metabolic functions in cultured renal tubule cells. *Kidney Int.* **29**, 46–53
- Cho, M. J., Thompson, D. P., Cramer, C. T., Vidmar, T. J. and Scieszka, J. F. (1989) The Madin-Darby canine kidney (MDCK) epithelial cell monolayer as a model cellular transport barrier. *Pharm. Res.* **6**, 71–77
- Tréhin, R., Krauss, U., Beck-Sickinger, A., Merkle, H. P. and Nielsen, H. M. (2004) Cellular uptake but low permeation of human calcitonin derived cell penetrating peptides and Tat(47–57) through well-differentiated epithelial models. *Pharm. Res.* **21**, 1248–1256
- Finkbeiner, W. E., Carrier, S. D. and Teresi, C. E. (1993) Reverse transcription-polymerase chain reaction (RT-PCR) phenotypic analysis of cell cultures of human tracheal epithelium, tracheobronchial glands, and lung carcinomas. *Am. J. Respir. Cell Mol. Biol.* **9**, 547–556
- Rupniak, H. T., Rowlett, C., Lane, E. B., Steele, J. G., Trejdosiewicz, L. K., Laskiewicz, B., Povey, S. and Hill, B. T. (1985) Characteristics of four new human cell lines derived from squamous cell carcinomas of the head and neck. *J. Natl. Cancer Inst.* **75**, 621–635
- Forbes, I. I. (2000) Human airway epithelial cell lines for *in vitro* drug transport and metabolism studies. *Pharm. Sci. Technol. Today* **3**, 18–27
- Nielsen, H. M. and Rassing, M. R. (2000) TR146 cells grown on filters as a model of human buccal epithelium: V. Enzyme activity of the TR146 cell culture model, human buccal epithelium and porcine buccal epithelium, and permeability of leu-enkephalin. *Int. J. Pharm.* **200**, 261–270
- Florea, B. I., Cassara, M. L., Junginger, H. E. and Borchard, G. (2003) Drug transport and metabolism characteristics of the human airway epithelial cell line Calu-3. *J. Control Release* **87**, 131–138
- Rothen-Rutishauser, B., Kramer, S. D., Braun, A., Gunthert, M. and Wunderli-Allenspach, H. (1998) MDCK cell cultures as an epithelial *in vitro* model: cytoskeleton and tight junctions as indicators for the definition of age-related stages by confocal microscopy. *Pharm. Res.* **15**, 964–971
- Rist, B., Entzeroth, M. and Beck-Sickinger, A. G. (1998) From micromolar to nanomolar affinity: a systematic approach to identify the binding site of CGRP at the human calcitonin gene-related peptide 1 receptor. *J. Med. Chem.* **41**, 117–123
- Weber, P. J. A., Bader, J. E., Folkers, G. and Beck-Sickinger, A. G. (1998) A fast and inexpensive method for N-terminal fluorescein-labeling of peptides. *Bioorg. Med. Chem. Lett.* **8**, 597–600
- Yamaya, M., Finkbeiner, W. E., Chun, S. Y. and Widdicombe, J. H. (1992) Differentiated structure and function of cultures from human tracheal epithelium. *Am. J. Physiol.* **262**, 713–724
- Shen, B. Q., Finkbeiner, W. E., Wine, J. J., Mrsny, R. J. and Widdicombe, J. H. (1994) Calu-3: a human airway epithelial cell line that shows cAMP-dependent Cl⁻ secretion. *Am. J. Physiol.* **266**, 493–501
- Lang, S., Langguth, P., Oschmann, R., Traving, B. and Merkle, H. P. (1996) Transport and metabolic pathway of thymocartin (TP4) in excised bovine nasal mucosa. *J. Pharm. Pharmacol.* **48**, 1190–1196
- Richardson, J. C., Scalera, V. and Simmons, N. L. (1981) Identification of two strains of MDCK cells which resemble separate nephron tubule segments. *Biochim. Biophys. Acta* **673**, 26–36
- Loman, S., Radl, J., Jansen, H. M., Out, T. A. and Lutter, R. (1997) Vectorial transcytosis of dimeric IgA by the Calu-3 human lung epithelial cell line: upregulation by IFN- γ . *Am. J. Physiol.* **272**, 951–958

-
- 43 Jacobsen, J., Nielsen, E. B., Brondum-Nielsen, K., Christensen, M. E., Olin, H. B., Tommerup, N. and Rassing, M. R. (1999) Filter-grown TR146 cells as an *in vitro* model of human buccal epithelial permeability. *Eur. J. Oral Sci.* **107**, 138–146
- 44 Nielsen, H. M. and Rassing, M. R. (1999) TR146 cells grown on filters as a model of human buccal epithelium: III. Permeability enhancement by different pH values, different osmolality values, and bile salts. *Int. J. Pharm.* **185**, 215–225
- 45 Fischer, R., Kohler, K., Fotin-Mleczek, M. and Brock, R. (2004) A stepwise dissection of the intracellular fate of cationic cell-penetrating peptides. *J. Biol. Chem.* **279**, 12625–12635
- 46 Xia, H., Mao, Q. and Davidson, B. L. (2001) The HIV Tat protein transduction domain improves the biodistribution of β -glucuronidase expressed from recombinant viral vectors. *Nat. Biotechnol.* **19**, 640–644
- 47 Mathias, N. R., Timoszyk, J., Stetsko, P. I., Megill, J. R., Smith, R. L. and Wall, D. A. (2002) Permeability characteristics of calu-3 human bronchial epithelial cells: *in vitro*–*in vivo* correlation to predict lung absorption in rats. *J. Drug Target* **10**, 31–40
- 48 Boderke, P., Merkle, H. P., Cullander, C., Ponc, M. and Bodde, H. E. (1997) Localization of aminopeptidase activity in freshly excised human skin: direct visualization by confocal laser scanning microscopy. *J. Invest. Dermatol.* **108**, 83–86
- 49 Lu, R. H., Kopeckova, P. and Kopecek, J. (1999) Degradation and aggregation of human calcitonin *in vitro*. *Pharm. Res.* **16**, 359–367
- 50 Motta, A., Andreotti, G., Amodeo, P., Strazzullo, G. and Castiglione Morelli, M. A. (1998) Solution structure of human calcitonin in membrane-mimetic environment: the role of the amphipathic helix. *Proteins* **32**, 314–323
-

Received 12 February 2004/25 May 2004; accepted 11 June 2004

Published as BJ Immediate Publication 11 June 2004, DOI 10.1042/BJ20040238