Figure 2 Time course of quinone production by tyrosinase with t-butylphenol at 0.1 mM

In the assay, concentrations of O_2 and tyrosinase of 0.26 mM and 0.3 μ M, respectively, were used. The quinone accumulation was monitored spectrophotometrically at 400 nm ($\epsilon = 1150 \text{ M}^{-1} \cdot \text{cm}^{-1}$) using a Perkin-Elmer Lambda-2 spectrophotometer interfaced on-line with an Amstrad PC2086 computer. The reaction medium was 0.1 M sodium phosphate buffer, pH 7.0. Temperature was controlled at 25 °C with a Haake D1G circulating bath with a heater/cooler and checked with a Cole-Parmer digital thermometer with a precision of ± 0.1 °C.

tyrosinase and t-butylphenol indicate the appearance of a lag period (Figure 2), and therefore the mechanism proposed in Figure 1(a) is not valid and the steps which must occur in the turnover would be those described in Figure 1(b) without the chemical reactions after the enzymic reaction. This gives rise to the lag period and, when the enzyme concentration increases, this process takes less time, resulting in a shorter lag period, and when the monophenol concentration increases, more diphenol is accumulated, the lag period becoming longer.

José Ramón ROS-MARTÍNEZ,* José Neptuno RODRÍGUEZ-LÓPEZ,*
Ramón Varón CASTELLANOS† and Francisco GARCÍA-CÁNOVAS*‡

*Departamento de Bioquímica y Biología Molecular, Facultad de Biología, Universidad de Murcia, 30071 Murcia, Spain, and †Escuela Universitaria Politécnica, Departamento de Química-Física, Universidad de Castilla-La Mancha, Avda. de España s/n, Campus Universitario, 02071 Albacete, Spain

‡ To whom correspondence should be addressed.

Cabanes, J., García-Cánovas, F., Lozano, J. A. and García-Carmona, F. (1987) *Biochim. Biophys. Acta* **923**, 187–195
Naish-Byfield, S. and Riley, P. A. (1992) *Biochem. J.* **288**, 63–67
Rodríguez-López, J. N., Tudela, J., Varón, R., García-Carmona, F. and García-Cánovas, F. (1992) *J. Biol. Chem.* **267**, 3801–3810

Received 7 April 1993

Sequences of clostridial ferredoxins: determination of the *Clostridium sticklandii* sequence and correction of the *Clostridium acidurici* sequence

Clostridial low-potential 2[4Fe-4S] ferredoxins (Fds) have been among the first isolated (Mortenson et al., 1962) and characterized (Lovenberg et al., 1963) iron-sulphur proteins. These small (approx. 55 residues) molecules have proved amenable to a variety of structural investigations, which have yielded numerous

amino acid sequences (George et al., 1985), a three-dimensional structure (Adman et al., 1976; Backes et al., 1991) and a host of data on the electromagnetic properties and topology of the [4Fe-4S] clusters (Mathews et al., 1974; Moulis et al., 1984).

Recently, these Fds have attracted renewed interest, since they afford useful models, as small and well-defined structures, for the investigation of the influence of the polypeptide chains on the properties of the [4Fe-4S] clusters. Research along these lines may benefit from the availability of molecular variants of 2[4Fe-4S] Fds, which are produced by chemical synthesis of polypeptide chains (Smith et al., 1991) or by using a synthetic gene expressed in *Escherichia coli* (Davasas and Moulis, 1992). Micro-organisms nevertheless remain a valuable source of Fd, the more so since naturally occurring sequences are likely to point out functionally important amino acids. In order to strengthen the basis for such studies, we report here corrections of the sequence of the Fd from *Clostridium acidurici* (Rall et al., 1969), which has long been a benchmark protein in the study of 2[4Fe-4S] Fds (Rabinowitz, 1972), and the newly determined sequence of *Clostridium sticklandii* Fd.

Automated Edman degradation of carboxymethylated *C. acidurici* Fd yielded the complete sequence, with only the three C-terminal residues missing. This sequence displays four changes, as compared with the previously reported one (Rall et al., 1969): D15 has to be replaced by E, D21 by N, Q25 by S, and S28 by D (Figure 1). The first error, in position 15, was initially detected on a 0.18 nm (1.8 Å)-resolution electron-density map derived from X-ray-diffraction data (TranQui et al., 1991) and prompted us to re-sequence the entire protein. Peptides purified from a thermolysin digest of the carboxymethylated ferredoxin allowed the identification of the three C-terminal amino acids (peptide Th-4), and confirmed the four corrections in positions 15, 21 (peptides Th-1 and Th-2), 25 and 28 (peptide Th-3) (Figure 1). The four changes in the sequence do not modify the amino acid analysis of the protein: in both the previously reported sequence (Rall et al., 1969) and the corrected one shown in Figure 1, the four residues come out as 2 Asx, 1 Glx, 1 Ser. This is probably why these sequencing errors have remained undetected in the numerous careful analyses that have been reported in the meantime (Lode et al., 1976).

The evidence for the amino acid sequence of *C. sticklandii* Fd is summarized in Figure 2. The carboxymethylated Fd yielded the sequence down to residue 33, with four undetected residues. The sequences of six overlapping peptides obtained by either chymotrypsin or thermolysin digestion afforded the complete sequence of the protein (Figure 2), which was consistent with amino acid analyses.

The two sequences determined here have been aligned in Figure 3 with those of two other well-characterized clostridial-

	10	20	30	40	50	55
	AYVINEACISGCGACEPECVPVNAISSGGDDRYVIDADTCIDCGACAGVCVPDAPVQA					
Cm-Fd	AYVINEACISGCGACEPECVPVNAISSGGDDRYVIDADTCIDCGACAGVCVPDAPVQA					
Th-1	VINEACISGCGACEPECVPVNA					
Th-2	ISGCGACEPECVPVNA					
Th-3	ISSGGDDRY					
Th-4	VIDADTCIDCGACAGVCVPDAPVQA					

Figure 1 Summary proof of the sequence of *C. acidurici* ferredoxin

C. acidurici (A.T.C.C. 7906, formerly *C. acidurici*; Cato et al., 1986) Fd was purified as described by Gaillard et al. (1986). The final sequence is shown in **bold** characters, and the four corrected residues are doubly underlined. The sequences of the carboxymethylated ferredoxin (Cm-Fd) and of four peptides obtained by thermolysin cleavage of Cm-Fd are shown below. Hyphens stand for unidentified residues. The sequence has been deposited in the EMBL Data Library with accession number P00198.

```

      10      20      30      40      50      55
AYVINDSCISGACEPECPVNAITAGDDKYVIDAATCIDCGACAGVCPVDAQPQE
Cm-Fd  AYVIND·CISGACEPECPVNAIT···DKYVID··········
CT-1   AYVINDSCISGACEPECPVNAITAGDDKY
Th-1    ······VIDAATCIDCGACAGVCPVDAQP·
Th-2    ······ISGACEPECPVNAI
Th-3    ······ITAGDDKY
Th-4    ······VIDAATCIDCGACAG
Th-5    ······VCPVDAQPQE

```

Figure 2 Summary proof of the sequence of *C. sticklandii* ferredoxin

C. sticklandii (A.T.C.C. 12662) Fd was purified as previously described (Sliwkowski and Stadtman, 1987). The final sequence is shown in **bold** characters. The sequences of the carboxymethylated ferredoxin (Cm-Fd), of one peptide obtained by chymotrypsin cleavage (CT-1) and of five peptides obtained by thermolysin cleavage (Th) of Cm-Fd are shown below. Hyphens stand for unidentified residues. The length of the C-terminal peptide (Th-5) was confirmed by liquid-desorption mass spectrometry: the measured monoisotopic mass MH^+ was 1112.4 (calculated = 1112.4933). The sequence has been deposited in the EMBL Data Library with accession number P80168.

```

      10      20      30      40      50      55
Cau  AYVINEACISGACEPECPVNAISSGDDRYVIDADTCIDCGACAGVCPVDAQVQA
Cst  AYVINDSCISGACEPECPVNAITAGDDKYVIDAATCIDCGACAGVCPVDAQPQE
Cpa  AYKIADSCVSGACASCECPVNAISSGDSIFVIDADTCIDCGNCANVCVPGAPVQE
Pas  AYVINDSCIAGACKPECPVNCIQGG·SIYAIADADSCIDCGSASVCPVGAAPNPED

```

Figure 3 Alignment of the sequences of the ferredoxins from *C. acidurici* (Cau), *C. sticklandii* (Cst), *C. pasteurianum* (Cpa) and *Peptostreptococcus asaccharolyticus* (Pas)

The dot in the Pas sequence represents a gap introduced to optimize the alignment.

type ferredoxins: that from *Clostridium pasteurianum*, which has been confirmed by the sequence of the gene (Graves et al., 1985), and that from *Peptostreptococcus asaccharolyticus*, which has been re-sequenced because of inconsistencies between the reported amino acid sequence and the crystallographic data (Backes et al., 1991). More extensive alignments of clostridial Fd sequences are given elsewhere (George et al., 1985). The alignments displayed here confirm that the *P. asaccharolyticus* protein differs from the genuine clostridial ones by the deletion of residue 27, by an additional amino acid at the C-terminus, and by the presence of a cysteine in position 22 (Figure 3). Since the latter protein is the only $2[4Fe-4S]$ Fd for which a high-resolution structure has been solved (Adman et al., 1976; Backes et al., 1991), the availability of another crystallographic model derived from a clostridial Fd would be of great help for the study of these proteins in further detail. The Fd from *C. acidurici*, which has long been known to give the best-quality crystals (Lovenberg et al., 1963; Murthy et al., 1988), is probably the best candidate for this purpose, and its structure is in the process of being elucidated (TranQui et al., 1991). *C. pasteurianum* Fd, which differs from *C. acidurici* Fd in 15 positions (Figure 3), is notoriously difficult to crystallize (Lovenberg et al., 1963; L. Sieker, personal com-

munication). As *C. sticklandii* Fd is most closely related to *C. acidurici* Fd (nine differences), it will be interesting to investigate whether, by its behaviour in crystal growth experiments, it is more similar to *C. acidurici* or to *C. pasteurianum* Fd.

The importance of reliable amino acid sequences is at least twofold in the case of the proteins discussed here. First, they are used as phylogenetic markers (George et al., 1985), and mistakes in a few residues in these small proteins are likely to result in inaccurate evolutionary trees. Second, many properties of these Fds are being investigated in increasing detail, when possible on the basis of high-resolution three-dimensional structures. In such small proteins, most amino acids have important roles in at least one aspect of the structure and function (chain folding, Fe-S cluster ligation, interaction with redox partners, etc.). It is therefore essential that each residue be identified with a high degree of confidence.

We are grateful to Dr. T. C. Stadtman for her kind gift of the *C. sticklandii* strain.

Jacques MEYER,*§ Jean-Marc MOULIS,* Nathalie SCHERRER,†
Jean GAGNON† and Jacques ULRICH‡

*DBMS-Métalloproteines, CENG, 85X, 38041 Grenoble, France, and †Enzymologie Moléculaire and ‡Spectrométrie de Masse des Protéines, Institut de Biologie Structurale, 38027 Grenoble, France

§ To whom correspondence should be addressed.

- Adman, E. T., Sieker, L. C. and Jensen, L. H. (1976) *J. Biol. Chem.* **251**, 3801–3806
- Backes, G., Mino, Y., Loehr, T. M., Meyer, T. E., Cusanovich, M. A., Sweeney, W. V., Adman, E. T. and Sanders-Loehr, J. (1991) *J. Am. Chem. Soc.* **113**, 2055–2064
- Cato, E. P., George, W. L. and Finegold, S. M. (1986) in *Bergey's Manual of Systematic Bacteriology* (Sneath, P. H. A., Mair, N. S., Sharpe, N. E. and Holt, J. G., eds.), pp. 1141–1200, Williams and Wilkins, Baltimore.
- Davashe, V. and Moulis, J.-M. (1992) *Biochem. Biophys. Res. Commun.* **185**, 341–349
- Gaillard, J., Moulis, J.-M., Auric, P. and Meyer, J. (1986) *Biochemistry* **25**, 464–468
- George, D. G., Hunt, L. T., Yeh, L.-S. L. and Barker, W. C. (1985) *J. Mol. Evol.* **22**, 20–31
- Graves, M. C., Mullenbach, G. T. and Rabinowitz, J. C. (1985) *Proc. Natl. Acad. Sci. U.S.A.* **82**, 1653–1657
- Lode, E. T., Murray, C. L. and Rabinowitz, J. C. (1976) *J. Biol. Chem.* **251**, 1675–1682
- Lovenberg, W., Buchanan, B. B. and Rabinowitz, J. C. (1963) *J. Biol. Chem.* **238**, 3899–3913
- Mathews, R., Charlton, S., Sands, R. H. and Palmer, G. (1974) *J. Biol. Chem.* **249**, 4326–4328
- Mortenson, L. E., Valentine, R. C. and Carnahan, J. E. (1962) *Biochem. Biophys. Res. Commun.* **7**, 448–452
- Moulis, J.-M., Meyer, J. and Lutz, M. (1984) *Biochemistry* **23**, 6605–6613
- Murthy, H. M. K., Hendrickson, W. A., Orme-Johnson, W. H., Merritt, E. A. and Phizackerley, R. P. (1988) *J. Biol. Chem.* **263**, 18430–18436
- Rabinowitz, J. C. (1972) *Methods Enzymol.* **24**, 431–446
- Rall, S. C., Bolinger, R. E. and Cole, R. D. (1969) *Biochemistry* **8**, 2486–2496
- Sliwkowski, M. X. and Stadtman, T. C. (1987) *J. Biol. Chem.* **262**, 4899–4904
- Smith, E. T., Feinberg, B. A., Richards, J. H. and Tornich, J. M. (1991) *J. Am. Chem. Soc.* **113**, 688–689
- TranQui, D., Fanchon, E., Vicat, J., Sieker, L. C., Meyer, J., Moulis, J.-M., Gagnon, J. and Duée, E. D. (1991) *Int. Conf. Bioinorganic Chem.* 5th, Oxford, Abstr. D049