GCN2 phosphorylation of eIF2 α activates NF- κ B in response to UV irradiation

Hao-Yuan JIANG¹ and Ronald C. WEK

Department of Biochemistry and Molecular Biology, Indiana University School of Medicine, Indianapolis, IN 46202, U.S.A.

In response to UV irradiation, mammalian cells elicit a gene expression programme designed to repair damage and control cell proliferation and apoptosis. Important members of this stress response include the NF- κ B (nuclear factor- κ B) family. However, the mechanisms by which UV irradiation activates NF- κ B are not well understood. In eukaryotes, a variety of environmental stresses are recognized and remediated by a family of protein kinases that phosphorylate the α subunit of eIF2 (eukaryotic initiation factor-2). In the present study we show that NF- κ B in MEF (murine embryo fibroblast) cells is activated by UV-C and UV-B irradiation through a mechanism requiring eIF2 α phosphorylation. The primary eIF2 α kinase in response to UV is GCN2 (general control non-derepressible-2), with PEK/PERK (pancreatic eIF2 α kinase/RNA-dependent-protein-kinase-like endoplasmic-reticulum kinase) carrying out a secondary function. Our

studies indicate that lowered protein synthesis accompanying eIF2 α phosphorylation, combined with eIF2 α kinase-independent turnover of I κ B α (inhibitor of κ B α), reduces the levels of I κ B α in response to UV irradiation. Release of NF- κ B from the inhibitory I κ B α would facilitate NF- κ B entry into the nucleus and targeted transcriptional control. We also find that loss of GCN2 in MEF cells significantly enhances apoptosis in response to UV exposure similar to that measured in cells deleted for the RelA/p65 subunit of NF- κ B. These results demonstrate that GCN2 is central to recognition of UV stress, and that eIF2 α phosphorylation provides resistance to apoptosis in response to this environmental insult.

Key words: eukaryotic initiation factor-2 (eIF2), general control non-derepressible-2 (GCN2), nuclear factor κ B (NF- κ B), stress response, translational control, UV irradiation.

INTRODUCTION

UV irradiation from the sun is the main causative agent in the formation of non-melanoma skin cancer, and the risk for exposure to harmful UV irradiation is rising with the continued destruction of the protective atmospheric ozone layer. Exposure of mammalian cells to UV irradiation rapidly stimulates many well known transcription factors, such as p53 and NF- κ B (nuclear factor κ B) [1–4]. Whereas activation of p53 is thought to be a direct consequence of nuclear signals triggered by damaged DNA, the molecular mechanisms by which exposure to UV-C and UV-B activate NF- κ B rely on cytoplasmic signals that are generated independently of DNA damage [4–7].

The NF- κ B/Rel family of transcription factors regulate the expression of genes critical for multiple biological processes, including inflammatory reactions, immune responses, cell proliferation and apoptosis [8–11]. The cellular localization of NF- κ B is tightly controlled by a protein family designated I κ Bs (inhibitory κ B), the most prominent and well-studied being I κ B α . Nuclear uptake of NF- κ B is blocked by its tight association with I κ B α . Exposure of cells to a variety of extracellular signals, such as TNF α (tumour necrosis factor α), interleukin-1 or bacterial lipopolysaccharide leads to the rapid phosphorylation of I κ B α at two amino acid residues, namely Ser³² and Ser³⁶, by IKK (I κ B kinase). This targeted I κ B α phosphorylation contributes to its ubiquitin-

ation and, ultimately, proteasome-mediated degradation. Decreased levels of I κ B α lead to the release of NF- κ B, facilitating translocation of this transcription factor into the nucleus to regulate expression of its target genes.

The molecular mechanisms regulating NF- κ B in response to UV exposure are less well understood. Activation of NF- κ B by UV irradiation is slower than cytokine stimulation, taking between 1 and 3 h after UV treatment [12,13]. Whereas ubiquitin-mediated degradation of I κ B α is proposed to be required for full induction of NF- κ B by UV exposure, such degradation appears to occur independently of IKK phosphorylation of I κ B α [13,14]. Recently, Kato et al. [15] suggested that the UV-mediated reduction in I κ B α levels occurs via a mechanism involving CK2 (casein kinase II) phosphorylation of one or more serine residues in the PEST (Pro-Glu-Ser-Thr) domain of I κ B α . CK2 activity is proposed to be UV inducible through a mechanism involving activation of the p38 MAPK (mitogen-activated protein kinase) pathway.

We are interested in the mechanisms by which eukaryotic cells process diverse stress signals to regulate programmes of gene expression designed to prevent or remedy cellular damage. Central to this stress response is a family of protein kinases that phosphorylate the α subunit of eIF2 (eukaryotic initiation factor-2) [16,17]. The eIF2 α kinases includes GCN2 (general control non-derepressible-2), which is activated by amino acid limitation

Abbreviations used: ATF4, activating transcription factor 4; bZIP, basic leucine zipper; CHOP, C/EBP (CCAAT-enhancer-binding protein)-homologous protein; CK2, casein kinase II; CRE, cAMP response element; CREB, cAMP-response-element-binding protein; CreP, constitutive repressor of eIF2 α (eukaryotic initiation factor-2 α) phosphorylation; DMEM, Dulbecco's modified Eagle's medium; EMSA, electrophoretic-mobility-shift assay; ER, endoplasmic reticulum; GADD153 and GADD34, growth arrest and DNA damage inducible gene 153 and 34; GCN2, general control non-derepressible-2; HRI kinase, haem-regulated inhibitory kinase; I κ B, inhibitory κ B; IKK, I κ B kinase; JNK, c-Jun N-terminal kinase; MAPK, mitogen-activated protein kinase; MEF, mouse embryo fibroblast; MG132, N-Cbz (benzyloxycarbonyl)-Leu-Leu-Leu-al; NF- κ B, nuclear factor κ B; OCT1, Octamer-1; PEK, pancreatic eIF2 α kinase; PERK, RNA-dependent-protein-kinase-like ER kinase; PEST, Pro-Glu-Ser-Thr; PKR, protein kinase regulated by RNA; PPP1R15B, phosphorylation/protein phosphatase 1, regulatory (inhibitor) subunit 15b; RSB, reticulocyte standard buffer; SAPK, stress-activated protein kinase; TNF α , tumour necrosis factor α ; TUNEL, terminal deoxynucleotidyltransferase-mediated dUTP-biotin *in situ* nick end labelling; URE, upstream regulatory element.

¹ To whom correspondence should be addressed (email hjiang@iupui.edu).

[18–22], PEK/PERK [pancreatic eIF2 α kinase/RNA-dependent-protein-kinase-like ER (endoplasmic reticulum) kinase], which is important for remedying protein misfolding in the ER [23], HRI kinase (haem-regulated inhibitory kinase), which couples protein synthesis to haem availability in erythroid cells [24] and PKR (protein kinase regulated by RNA), which participates in an antiviral defence pathway induced by interferon [25]. Phosphorylation of eIF2 α during cellular stress impedes the ability of eIF2 to deliver initiator Met-tRNA_i^{Met} to the translation machinery [26]. The resulting reduction in eIF2 activity lowers global protein synthesis concomitant with induced translation of selected mRNAs including ATF4 (activating transcription factor 4), a transcriptional activator of stress remedy genes [18,27,28]. Furthermore, we recently reported that eIF2 α phosphorylation is linked to activation of NF- κ B in response to ER stress and nutritional deprivation by mechanisms independent of IKK phosphorylation of I κ B α [29].

In addition to the stress conditions highlighted above, there have been several reports coupling UV irradiation to eIF2 α phosphorylation or its target transcription activator. Deng et al. [30] reported that UV-C irradiation elicits phosphorylation of eIF2 α by GCN2 via a mechanism that does not require SAPK/JNK (stress-activated protein kinase/c-Jun N-terminal kinase) or p38 MAP kinases. Wu et al. [31] suggested that UV-C treatment induced the eIF2 α kinase PEK. Phosphorylation of eIF2 α by PEK may play a role in cell-cycle arrest after UV irradiation. Exposure of yeast to UV irradiation leads to activation of Gcn4, a bZIP (basic leucine zipper) transcriptional activator related to ATF4, whose expression is induced by GCN2 phosphorylation of eIF2 α during amino acid limitation [32]. Such activation of Gcn4 was proposed to be required for induction of target genes involved in nucleotide biosynthesis, contributing to DNA repair.

Given the important role of eIF2 α phosphorylation as an upstream sensor and regulator in stress-response pathways, and the linkage between eIF2 α kinases and UV exposure, we addressed the question as to whether phosphorylation of eIF2 α facilitates activation of NF- κ B by UV irradiation. We find that mouse embryo fibroblasts (MEFs) deleted for GCN2 or containing a homozygous mutation at the eIF2 α phosphorylation site (Ser⁵¹ → Ala) block activation of NF- κ B. The basis for this reduction is that eIF2 α phosphorylation by GCN2 contributes to significantly lower levels of general translation, including decreased synthesis of I κ B α . Decreased I κ B α translation, along with eIF2 α kinase-independent turnover of I κ B α , contributes to the lowered levels of I κ B α and activation of NF- κ B. These results indicate that, in addition to induced translational expression of regulatory proteins such as ATF4, eIF2 α phosphorylation and the accompanying reduced general translation can regulate stress-gene expression by lowering the levels of key regulatory proteins that are subject to rapid turnover.

EXPERIMENTAL

Cell culture and stress conditions

GCN2^{-/-} and PEK^{-/-}, and their wild-type counterparts, were immortalized by simian-virus-40 large T antigen and have been described previously [29,33,34]. Immortalized S/S MEF cells expressing wild-type eIF2 α , and A/A MEF cells (containing eIF2 α with alanine replacing serine at the position-51 phosphorylation site), were previously described [33]. The RelA/p65^{-/-} MEF cells [35,36] and the immortalized cells were cultured in DMEM (Dulbecco's modified Eagle's medium; BioWhittaker), supplemented with 2 mM glutamine, 1 mM non-essential amino acids, 10% (v/v) fetal-bovine serum, 100 units/ml penicillin, and

100 μ g/ml streptomycin. A/A MEF cells require additional amino acid supplements for growth viability, and therefore the A/A and the wild-type counterpart S/S cells were maintained under these enriched-media conditions. MEF cells were grown to 50–70% confluency and were irradiated at the indicated UV-C dose (254 nm; UV Stratalinker 1800, Stratagene). Alternatively, UV-B irradiation of cells was carried out using a Philips FS-20 UVB lamp that was filtered to eliminate residual UV-C. The intensity of the UV-B exposure was measured for each experiment using an IL1700 radiometer and an SED240 UV-B detection instrument (International Light, Newburyport, MA, U.S.A.) at a distance of 8 cm from the UV-B source to the monolayer of cells. As a control for I κ B α phosphorylation and subsequent degradation, 10 ng/ml TNF α was added to MEF cells in DMEM for 30 min.

EMSAs (electrophoretic-mobility-shift assays)

Nuclear extracts were prepared from MEF cells as described previously [37]. MEF cells were subjected 40 J/m² UV-C irradiation, 1 μ M thapsigargin or 10 ng/ml TNF α , as indicated. To determine the role of ubiquitin-mediated degradation in the activation of NF- κ B, one set of experiments involved the addition of 1 μ M MG132 [*N*-Cbz (benzyloxycarbonyl)-Leu-Leu-Leu-al, a proteasome inhibitor] 30 min prior to exposure to these stress agents. Following the stress treatment, cells were resuspended in 1 ml of cold hypo-osmotic RSB (reticulocyte standard buffer) [10 mM Tris (pH 7.4)/10 mM NaCl/3 mM MgCl₂] supplemented with 0.5% Nonidet P40 and protease inhibitors (100 μ M of PMSF/0.15 μ M aprotinin/1 μ M leupeptin/1 μ M pepstatin). Cell lysis was carried out using a Dounce homogenizer and, following centrifugation (13 000 g, 10 min), the nuclei pellets were resuspended in 2 packed nuclear vol. of extraction buffer C [420 mM KCl/20 mM Hepes (pH 7.9)/1.5 mM MgCl₂/0.2 mM EDTA/20% (v/v) glycerol] supplemented with protease inhibitors. Protein concentrations were determined using the Bradford assay. The sequence of the DNA fragment containing the URE (upstream regulatory element) NF- κ B binding element derived from *c-myc* was 5'-GATCCAAGTCCGGGTTTCCCCAACC-3' [29] and the Octamer-1 (OCT1) binding element was 5'-GATCTGTGCA-ATGCAAATCACTAGAA-3' [37]. Binding reaction mixtures contained ³²P-labelled DNA fragments (20 000–25 000 c.p.m.), 5 μ g of nuclear extract, and 2.5 μ g of poly(dI-dC) as non-specific competitor were added to a solution of 10 mM Hepes, pH 7.9, 4 mM dithiothreitol, 0.5% Triton X-100, 100 mM KCl and 2.5% (v/v) glycerol in a final assay volume of 25 μ l. Binding mixtures were incubated at room temperature for 30 min. DNA/protein complexes were separated by gel electrophoresis and visualized by autoradiography. To address NF- κ B binding specificity, unlabelled competitor URE or CREB (cAMP-response-element-binding protein) DNA fragments were added at the indicated stoichiometry to the binding mixture. The CREB competitor DNA included a previously reported ATF consensus binding sequence, 5'-TGACGTCA-3' [38]. Supershift studies were carried out by including polyclonal antibody specific to p65 (Upstate Biotechnology), p50 (Santa Cruz) or c-Rel (Santa Cruz), as indicated, in the binding mixture. Mixtures were analysed by EMSA and autoradiography.

Preparation of protein lysates and immunoblot analyses

MEF cells that were exposed to UV irradiation, or to no stress, were washed twice with ice-cold PBS and lysed using Breaking Solution [50 mM Tris/HCl (pH 7.9)/150 mM NaCl/1% Nonidet P40/0.1% SDS/100 mM NaF/17.5 mM β -glycerol phosphate/10% glycerol, supplemented with protease inhibitors]. Lysates

were subjected to sonication for 30 s and clarified by centrifugation (13 000 g, 10 min). Protein content was measured by using the Bio-Rad protein quantification kit for detergent lysis, and equal amounts of each sample were subjected to SDS/PAGE. Proteins were transferred to nitrocellulose filters, and the filters were incubated in TBS-T solution [20 mM Tris/HCl (pH 7.9)/150 mM NaCl/0.2 % Tween-20, supplemented with 4 % (w/v) non-fat dried milk] and antibodies that specifically recognize ATF4, ATF3, CHOP [C/EBP (CCAAT-enhancer-binding protein)-homologous protein], p65, I κ B α , actin or eIF2 α . Phosphorylation of eIF2 α and I κ B α were measured by using a polyclonal antibody that recognizes eIF2 α phosphorylated at Ser⁵¹ (Invitrogen, Carlsbad, CA, U.S.A.), or I κ B α phosphorylated at serine-32 (Cell Signaling Technology, Inc., Beverly, MA, U.S.A.). Following three washes in TBS-T, filters were incubated with TBS-T containing secondary antibody conjugated to horseradish peroxidase (Bio-Rad) and spots were visualized using a chemiluminescent substrate. Low- and high-molecular-mass ranges of polypeptide markers (Bio-Rad) were used in the SDS/PAGE analysis to determine the sizes of proteins detected in the immunoblots.

Radiolabelling assays

Translation inhibition was measured as previously described [18]. Briefly, *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells were seeded at 2×10^5 cells per 60 mm-diameter dish and grown to 50 % confluency in DMEM as described above. Cells were washed twice with PBS and incubated in 4 ml of *trans*-labelling medium without methionine or cysteine supplemented with 10 % (w/v) dialysed foetal bovine serum. The MEF cells were then labelled with 500 μ Ci of [³⁵S]Met/Cys Express Labeling Mix (ICN) for 30 min, and washed twice with ice-cold PBS containing non-radiolabelled methionine and cysteine. Cell lysates were prepared using Breaking Solution and clarified by centrifugation. Alternatively, the MEF cells were then subjected to 40 J/m² UV-C, or no stress, and incubated for between 0 and 2 h before transferring to the *trans*-labelling medium for 30 min and the [³⁵S]Met/Cys Labeling Mix for 30 min. Equal amounts of total protein were separated by SDS/PAGE, and radiolabelled protein was visualized by autoradiography. Half-life measurements were carried out by radiolabelling cells under nonstress conditions for 30 min, washing with non-radiolabelled medium, and incubation in medium containing non-radiolabelled methionine and cysteine. MEF cells were then treated with 40 J/m² UV-C, followed by incubation in non-radiolabelled medium for the indicated length of time. Lysates were prepared from the radiolabelled cells, and I κ B α and p65 were immunoprecipitated using protein-specific antibodies. Immunoprecipitated proteins were analysed by SDS/PAGE and visualized by autoradiography. Immunoblots measuring total levels of actin were carried out in parallel to be certain that similar levels of total protein were analysed in each lysate preparation.

Apoptosis assays

GCN2^{-/-} and *RelA/p65*^{-/-} MEF cells, and their wild-type counterparts, were exposed to UV-B or UV-C irradiation and the number of apoptotic cells were visually counted by using the *In Situ* Cell Death Detection Kit (Roche) that uses the TUNEL (terminal deoxynucleotidyltransferase-mediated dUTP-biotin *in situ* nick end labelling) reaction to label fragmented DNA with free 3-ends. Cells were visualized by microscopy, and 100 cells in each field were scored for apoptosis. Mean values $\pm$ S.D. were derived from three independent experiments. ANOVA and a Bonferroni post test (*post hoc* test) of the apoptotic measurements

were carried out by using GraphPad Prism (San Diego, CA, U.S.A.) software. Significant differences were defined as $P < 0.05$ for all tests. Activation of caspases 3 and 8 were measured by immunoblot analysis using polyclonal antibodies specific to each caspase.

RESULTS

Phosphorylation of eIF2 α by GCN2 is induced in response to UV-C and UV-B irradiation

eIF2 α kinases GCN2 and PEK have been suggested to elicit translational control in response to UV irradiation [30,31]. To address this model, we measured eIF2 α phosphorylation in *GCN2*^{-/-} and *PEK*^{-/-} MEF cells and their wild-type counterparts in response to increasing amounts of UV-C and UV-B irradiation. Phosphorylation of eIF2 α , as judged by immunoblot using polyclonal antibody specific to eIF2 α phosphorylated at the regulatory residue Ser⁵¹, occurs with as little as 10 J/m² of UV-C, with a maximum at 40 J/m² (Figure 1A). Similar levels of total eIF2 α were observed between the stressed and non-stressed lysates using antibody that recognizes both phosphorylated and non-phosphorylated forms of eIF2 α . In the case of UV-B exposure, significant eIF2 α phosphorylation was detected between 200 and 400 J/m² (Figure 1B). Induction of eIF2 α phosphorylation was rapid, within 15 min of UV-C exposure of MEF cells, and was sustained following 8 h of UV stress (Figure 1C). Minimal eIF2 α phosphorylation was detected in MEF cells devoid of GCN2 function in response to either UV-C or UV-B irradiation (Figures 1A, B, and C).

Deletion of PEK led to some reduction in eIF2 α phosphorylation in MEF cells exposed to UV-C. Analysis of increasing dosages of UV-C indicated that there was significantly lowered eIF2 α phosphorylation in *PEK*^{-/-} MEF cells treated with 10 or 20 J/m², whereas cells exposed to higher amounts of UV-C (40–160 J/m²) showed induced levels of phosphorylation that were partially decreased compared with wild-type cells (Figure 1D). Phosphorylation of eIF2 α in response to 30 J/m² was modestly delayed with the loss of PEK activity, with eIF2 α phosphorylation in *PEK*^{+/+} being maximally induced between 0.5 and 1 h, and upwards of 3 h in *PEK*^{-/-} cells (Figure 1). No UV induction of eIF2 α phosphorylation was detected using A/A MEF cells, demonstrating the specificity of our immunoblot assay (Figure 1F). We conclude that UV-C or UV-B exposure enhances GCN2 phosphorylation of eIF2 α at Ser⁵¹ in MEF cells, and that PEK functions as a secondary eIF2 α kinase during UV stress.

eIF2 α kinase-targeted genes, ATF4 and CHOP are not induced in response to UV stress

Phosphorylation of eIF2 α induces stress-gene expression through activation of transcription factors ATF4 and NF- κ B [18,27,29]. In the example of ATF4, eIF2 α phosphorylation in response to ER or nutritional stress stimulates ATF4 translation via a mechanism involving translation reinitiation and two open reading frames in the 5'-leader of the ATF4 mRNA [18,28]. Increased levels of ATF4 mRNA are also associated with cellular stress [39]. ATF4 contributes to induction of additional transcription factors, including ATF3 and CHOP/GADD153 (growth arrest and DNA damage inducible gene 153), which together direct a programme of stress-related gene expression [18,33,40]. We measured induction of ATF4 and its target genes in *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells by immunoblot analysis after 40 J/m² UV-C irradiation, an exposure that gave high levels of eIF2 α phosphorylation (Figure 2). No increase in ATF4, ATF3 or CHOP protein levels

Figure 1 Phosphorylation of eIF2α by GCN2 is enhanced in response to UV irradiation

GCN2^{+/+} and *GCN2*^{-/-} MEF cells were subjected to increasing dosages of UV-C (A) or UV-B irradiation (B). After UV treatment, cells were incubated in the culture medium for 3 h and the levels of phosphorylation of eIF2α were measured by immunoblot by using a polyclonal antibody specific to eIF2α phosphorylated at Ser⁵¹. Equal amounts of proteins were analysed in each lane. Levels of total eIF2α were assayed by using an antibody that recognizes both phosphorylated and non-phosphorylated versions of the translation initiation factor. *GCN2*^{+/+} and *GCN2*^{-/-} (C) or *PEK*^{+/+} and *PEK*^{-/-} (E) MEF cells were subjected to 30 J/m² UV-C irradiation, incubated in the culture medium for between 0.25 and 8 h as indicated, and eIF2α phosphorylation was measured by immunoblot. (D) *PEK*^{+/+} and *PEK*^{-/-} MEF cells were subjected to increasing dosages of UV-C, cultured for 3 h, and the amounts of eIF2α phosphorylation were measured by immunoblot. (F) S/S and A/A MEF cells were exposed to 30 J/m² UV-C irradiation, and after between 0.5 and 6 h incubation as indicated, phosphorylated eIF2α and total eIF2α levels were measured by immunoblot. Immunoblots in each panel are representative of three independent experiments.

were detected in either *GCN2*^{+/+} or *GCN2*^{-/-} MEF cells after 8 h of 40 J/m² UV irradiation (Figure 2A). Additionally, there was no increase in the levels of ATF4, ATF3 or CHOP mRNA after 40 J/m² UV-C irradiation in either *GCN2*^{+/+} or *GCN2*^{-/-} MEF

Figure 2 Phosphorylation of eIF2α in response to UV irradiation does not induce expression of ATF4 and its target genes

(A) *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells were exposed to 40 J/m² UV-C or no stress (0 h), and incubated in culture medium for between 0.5 and 8 h, as indicated. Levels of ATF4, CHOP, ATF3 and eIF2α were measured by immunoblot using antibodies specific to each of these proteins. (B) *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells were exposed to increasing dosages of UV-C, as indicated, and then cultured for 6 h prior to immunoblot measurements of ATF4, CHOP, ATF3 and eIF2α. (C) S/S and A/A MEF cells were exposed to increasing dosages of UV-C, cultured for 6 h, and ATF3, phosphorylated eIF2α, and total eIF2α levels were measured by immunoblot analysis. Results shown are representative of three independent experiments.

cells (results not shown). Consistent with previous reports, expression of each of these transcription factors was significantly increased in *GCN2*^{+/+} MEF cells subjected to leucine starvation, and this induction was blocked in *GCN2*^{-/-} cells (Figure 2A) [18,33]. We also measured expression of these eIF2α kinase stress-response genes after exposure from between 1 and 40 J/m² UV irradiation. No expression of ATF4 or CHOP were detected after treatment with these UV-C dosages (Figure 2B). Interestingly, ATF3 expression was significantly increased between 5 and 20 J/m², with minimal expression at 40 J/m² UV-C. This induction was detected in both *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells, as well as in S/S (wild-type eIF2α) and A/A ([Ser⁵¹ → Ala]eIFα) MEF cells, indicating that there are regulatory mechanisms in response to certain stress conditions that can lead to

enhanced expression of ATF3 independent of eIF2 α phosphorylation (Figures 2B and 2C).

Phosphorylation of eIF2 α activates NF- κ B during UV irradiation

To address whether phosphorylation of eIF2 α contributes to NF- κ B activation in MEF cells in response to UV irradiation, we performed an EMSA using radiolabelled DNA containing a NF- κ B binding site. Nuclear lysates were prepared from *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells exposed to 40 J/m² UV-C or no stress. Significant DNA binding was detected in the *GCN2*^{+/+} cells 1 h after the UV exposure, with further increases following 3 and 6 h of UV irradiation (Figure 3A). By contrast, there were only low levels of NF- κ B activity in *GCN2*^{-/-} cells after 6 h of the UV insult. No differences were found in the EMSA using radiolabelled DNA containing OCT1 binding site (Figure 3B). A similar EMSA analysis of *PEK*^{-/-} MEF cells revealed a similar time course for activation of NF- κ B compared with *PEK*^{+/+} cells, although the levels of NF- κ B binding were partially decreased (Figure 3C).

The role of phosphorylation of eIF2 α in the activation of NF- κ B was directly addressed by using MEF cells containing wild-type (S/S) or the mutant form of eIF2 α (A/A). Activation of NF- κ B occurred in the S/S cells exposed to UV, while minimal NF- κ B binding was detected in the A/A cells (Figure 3E). The enhanced DNA binding was specific for NF- κ B, as confirmed by using non-radiolabelled competitor DNA containing an NF- κ B binding site (Figure 3F). By contrast, no competition was found using a non-radiolabelled DNA containing an ATF/CRE (cAMP response element) DNA binding element. The NF- κ B binding was attributed to NF- κ B dimers p65/p50 and p50/p50, as judged by antibody supershift experiments [14] (Figure 3F). These results indicate that eIF2 α phosphorylation by GCN2 is central to activation of NF- κ B in response to UV irradiation.

Decreased levels of I κ B contribute to activation of NF- κ B during UV stress

One mechanism facilitating nuclear localization of NF- κ B involves phosphorylation and proteolysis of I κ B α . We wished to measure I κ B α phosphorylation at Ser³² and I κ B α protein levels in the *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells in response to UV irradiation. As a control, we treated the MEF cells with TNF α , a cytokine that induces IKK phosphorylation of I κ B at Ser³² and Ser³⁶, leading to ubiquitin-mediated proteolysis of I κ B [41,42]. No phosphorylation of eIF2 α was detected after exposure to TNF α (results not shown) [29]. While TNF α induced phosphorylation of I κ B α in both the *GCN2*^{+/+} and *GCN2*^{-/-} cells, as judged by immunoblot using a polyclonal antibody specific to I κ B α phosphorylated at Ser³², no I κ B α phosphorylation was detected in either MEF cell line exposed to UV irradiation (Figure 4A). Total levels of I κ B α were also measured by immunoblot analysis. Consistent with the TNF α -mediated phosphorylation of I κ B, there was a decrease in I κ B α levels in both the *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells after treatment with this cytokine. Interestingly, the I κ B α levels were also rapidly reduced in the *GCN2*^{+/+} MEF cells in response to UV irradiation, with a significant decrease in the amounts of I κ B α within 1 h following UV stress (Figure 4A). By comparison, an appreciable decrease in I κ B α levels was detected in the *GCN2*^{-/-} cells only after 8 h of UV insult. Levels of p65, as well as actin, were similar in *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells during the time course after UV irradiation. A similar reduction in I κ B α levels was measured in UV-treated S/S MEF cells, which was alleviated in A/A cells, indicating that the regulatory function of GCN2 is via phosphorylation of eIF2 α (results not shown).

Figure 3 GCN2 is required for activation of NF- κ B in response to UV exposure

GCN2^{+/+} and *GCN2*^{-/-} (A and B) or *PEK*^{+/+} and *PEK*^{-/-} (C and D) MEF cells were exposed to 40 J/m² UV-C or no stress (0 h). After UV stress, the MEF cells were cultured for the indicated time, and nuclear lysates were utilized in an EMSA with radiolabelled DNA containing a NF- κ B (A and C) or control OCT1 (B and D) binding site. (E) A similar EMSA study was carried out using nuclear lysates prepared from UV-stressed MEF cells containing *eIF2* α with wild-type Ser⁵¹ (S/S) or with alanine replacing serine in the *eIF2* α phosphorylation site at position 51 (A/A). (F) In lanes 2–4, supershift indicates that polyclonal antibodies that specifically recognize NF- κ B subunits, p50, p65, or Rel-C were added to the EMSA binding mixture. None indicates that no antibody was used in the assay (lane 1). To determine the specificity for the NF- κ B binding site in the EMSA, non-radiolabelled DNA containing the NF- κ B site (lanes 5–7) or ATF/CRE (8–10) were added to the EMSA binding mixtures containing nuclear lysates prepared from S/S MEF cells. Competitor DNA was added at 10- or 100-fold molar excess. Arrows indicate DNA complexed with p65/p50 or p50/p50. In each experiment excess free radiolabelled probe was present at the bottom of the panel. Portions of the radiolabelled free probes in (B) and (E) migrated off the bottom of the polyacrylamide gel during the EMSA. EMSA results shown are representative of three independent experiments.

Phosphorylated eIF2 α prevents the exchange of eIF2-GDP for eIF2-GTP that is required for general translation initiation [16,26]. To determine the effects of phosphorylation of eIF2 α during UV irradiation, we measured [³⁵S]methionine/[³⁵S]cysteine incorporation after exposure of *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells exposed to 40 J/m² of UV-C. There was a significant decrease in protein synthesis in *GCN2*^{+/+} cells 1–3 h following UV-C irradiation (Figure 4B). By contrast, *GCN2*^{-/-} MEF cells devoid of

Figure 4 Phosphorylation of eIF2 α facilitates lowered levels of I κ B α in response to UV irradiation

(A) *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells were exposed to 40 J/m² UV-C irradiation or no stress (0), and incubated in the culture medium for up to 8 h as indicated. Alternatively, the MEF cells were incubated with TNF α for 30 min. Equal amounts of proteins were analysed in each lane. Levels of phosphorylated I κ B α , total I κ B α , p65 and actin were measured by immunoblot analysis using polyclonal antibody specific to each protein. (B) Levels of protein synthesis were measured in *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells subjected to 40 J/m² UV-C irradiation or no stress (0). Following the UV stress, cells were incubated in the culture medium for the indicated time. Thirty minutes prior to harvesting, cells were radiolabelled with [³⁵S]Met/Cys. Cells were then washed, and radiolabelled proteins were separated by SDS/PAGE, and visualized by autoradiography. I κ B α and p65 were immunoprecipitated using protein-specific antibodies, subjected to SDS/PAGE, and imaged by autoradiography. To show that equal amounts of proteins were analysed in the electrophoretic analysis, actin was measured by immunoblot analysis using the radioactive lysate preparation. Levels of general protein synthesis, I κ B α and p65 in the *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells are listed relative to 100% synthesis in the non-stressed cells. (C) The turnover of I κ B α was measured by radiolabelling *GCN2*^{+/+} or *GCN2*^{-/-} MEF cells with [³⁵S]methionine/[³⁵S]cysteine for 30 min. Radiolabelled MEF cells were washed, incubated in DMEM supplemented with non-radiolabelled amino acids, and then exposed to 40 J/m² UV-C or no stress (0 min). Following incubation for between 20 and 160 min, I κ B α was immunoprecipitated, separated by SDS/PAGE and visualized by autoradiography. Total levels of actin were measured by immunoblot to insure that equal amounts of lysates were analysed in each experimental lane. Results shown in (A)–(C) are representative of three experiments.

phosphorylated eIF2 α showed minimal reductions in translation following UV stress. These results are similar to that reported by Deng et al. [30], who found that loss of GCN2 activity alleviated the translation repression associated with UV exposure. We extended our analysis to the synthesis of I κ B α and p65 by using immunoprecipitation followed by SDS/PAGE and autoradiography to visualize the radiolabelled proteins. We found that translation of I κ B α was significantly decreased upon eIF2 α phosphorylation in the *GCN2*^{+/+} MEF cells as compared with minimal change in the UV-stressed *GCN2*^{-/-} cells (Figure 4B). Interestingly, synthesis of p65 was more resistant to the decrease in translation initiation, with significant p65 expression in the *GCN2*^{+/+} cells after 1.5 h of the UV treatment. In the *GCN2*^{-/-} MEF cells subjected to UV, p65 synthesis was decreased to about half the levels measured for non-stressed cells (Figure 4B).

We next compared the turnover of I κ B α in *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells after exposure to UV. Using pulse-chase experiments we found that there was a rapid decay of [³⁵S]-methionine/[³⁵S]cysteine-labelled I κ B α in both *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells after UV stress (Figure 4C). The UV-triggered turnover of I κ B α had been reported previously and is disrupted by proteasome inhibitors, suggesting that ubiquitin-assisted degradation was taking place [13]. These results suggest that turnover in response to UV irradiation occurs independently of GCN2 phosphorylation of eIF2 α . We conclude that decreased synthesis of the repressing I κ B α is the central contribution of eIF2 α phosphorylation in the activation of NF- κ B in response to UV irradiation. Decreased translation of I κ B α , combined with turnover of I κ B α , results in detachment from NF- κ B. Release of NF- κ B from I κ B α would lead to translocation of the transcription factor into the nucleus and expression of NF- κ B-targeted genes.

Different mechanisms activate NF- κ B in response to UV and ER stress

We had shown previously that activation of NF- κ B in response to ER stress involved the release of I κ B α from NF- κ B without measurable reductions in I κ B α levels [29]. Thus it appears that the mechanisms by which eIF2 α phosphorylation modulates NF- κ B activity can vary depending on the cellular stress condition and the participation of additional stress-response pathways. We wished to formally delineate the role of ubiquitin-mediated degradation of I κ B α in the activation of NF- κ B between the UV and ER stress conditions. Wild-type MEF cells were pretreated with MG132, followed by exposure to UV-C, thapsigargin (a standard ER stress agent), TNF α or no stress. Inhibition of ubiquitin-mediated degradation significantly decreased activation of NF- κ B in response to UV or TNF α exposure, but did not diminish NF- κ B induction in response to ER stress (Figure 5). These results support the notion that there are differences between the mechanisms activating NF- κ B during ER stress and in response to UV or TNF α treatments. Ubiquitin-mediated turnover of I κ B α , which would contribute to lowered levels of this inhibitory protein, does not appear to be central to modulation of NF- κ B activity in response to ER stress.

Loss of GCN2 eIF2 α kinase enhances apoptosis in response to UV irradiation

NF- κ B regulates the transcription of genes involved in stress remediation and apoptosis. We wished to determine whether impaired activation of NF- κ B in *GCN2*^{-/-} MEF cells affected their viability upon treatment with UV irradiation. *GCN2*^{+/+} and *GCN2*^{-/-} MEF cells were exposed to UV-C irradiation and the number of apoptotic cells was counted by using the TUNEL assay. While less than 10% of *GCN2*^{+/+} MEF cells elicited apoptosis

Figure 5 Phosphorylation of eIF2 α in response to UV or ER stress activates NF- κ B by different mechanisms

Wild-type MEF cells were pretreated with MG132 as indicated, followed by exposure to UV-C (lanes 4 and 5), the ER stress agent thapsigargin (TG, lanes 6 and 7), TNF α (lanes 8 and 9), or no stress agent (None, lanes 2 and 3). Nuclear lysates were prepared from these MEF cells and included in an EMSA with radiolabelled DNA containing a binding site for NF- κ B (top panel) or OCT1 (bottom panel). Free probe (FP) indicates that only radiolabelled DNA was included in the EMSA. Arrows indicate the protein–DNA complexes. EMSA results shown are representative of three independent experiments.

after 8 h of the UV insult, almost 80% of cells devoid of GCN2 eIF2 α kinase activity underwent programmed cell death ($P < 0.0001$) (Figure 6A). UV-triggered apoptosis has been shown to be facilitated by cleavage and activation of a cascade of caspase proteases [43]. Cleavage of caspase 3 and 8 were measured by immunoblot and found to be significantly enhanced in the *GCN2*^{-/-} MEF cells compared with the wild-type cells (Figure 6B). Cleavage and activation of caspase 8, an early event in the caspase cascade, was at a maximum 3 h after UV irradiation. By comparison, cleavage of caspase 3 was detected at low levels 3 h after the UV insult, with maximum levels of cleaved protein following 6 and 8 h of the stress. These results indicate that phosphorylation of eIF2 α by GCN2 facilitates activation of caspases in response to UV irradiation, contributing to enhanced apoptosis.

Loss of RelA/p65 was reported to enhance apoptosis in response to UV irradiation [44,45]. We found that whereas the pattern of induced eIF2 α phosphorylation was similar in *RelA/p65*^{+/+} and *RelA/p65*^{-/-} MEF cells in response to UV exposure, the *RelA/p65*^{-/-} cells preferentially elicited cleavage of caspase 3 and 8 (Figure 6C). Furthermore, about 50% of the *RelA/p65*^{-/-} MEF cells underwent apoptosis 8 h after the UV treatment, while only 10% of the wild-type cells were apoptotic. These results are consistent with the idea that activation of NF- κ B and its target genes is one important reason for GCN2 protection against UV-induced apoptosis.

DISCUSSION

Translation repression by eIF2 α phosphorylation activates NF- κ B and reduces apoptosis in response to UV irradiation

Lowered levels of the inhibitory protein I κ B α are central to mechanisms activating NF- κ B in response to UV irradiation. We

Figure 6 Loss of GCN2 activity enhances apoptosis in response to UV exposure

GCN2^{-/-} and *RelA/p65*^{-/-} MEF cells, and their wild-type counterparts, were exposed to 30 J/m² UV-C irradiation or no stress (0), and incubated in the culture medium for up to 8 h as indicated. (A) Apoptosis was monitored by using the TUNEL assay to detect fragmented DNA. Wild-type *GCN2*^{+/+} cells and *RelA/p65*^{+/+} gave similar results, and the *GCN2*^{+/+} apoptotic measurements are presented. Means values $\pm$ S.D. were derived from three independent experiments. Two-way ANOVA and *post hoc* analyses indicated that there were significant differences in apoptosis between genotypes ($P < 0.001$) and with time ($P < 0.001$) as indicated by the asterisks. (B) Cleavage and activation of caspase (CASP) 3 and 8 were monitored by immunoblot analysis using polyclonal antibody specific to each caspase. Arrowheads indicate the non-cleaved caspase 8 (upper band), and cleaved/activated caspase 8 (two lower bands). The polyclonal antibody used in the caspase 3 immunoblot recognizes only the cleaved, activated caspase 3 polypeptide highlighted by the arrowhead. (C) *RelA/p65*^{+/+} and *RelA/p65*^{-/-} MEF cells were subjected to 30 J/m² UV-C irradiation or no stress (0), and cultured for between 0.5 and 8 h as indicated. Levels of phosphorylated eIF2 α , total eIF2 α , actin and cleaved and activated caspase 3 and 8 were measured by immunoblot analysis. Arrowheads indicate the uncleaved and cleaved/activated versions of caspase 8, and activated caspase 3, as described for (B). Immunoblot experiments shown in (B) and (C) are representative of three independent experiments.

show that reduced general protein synthesis in response to UV irradiation is the result of eIF2 α phosphorylation by GCN2, and this lowered translation facilitates decreased levels of I κ B α (Figures 1 and 4). Translation of I κ B α mRNA appears to be very sensitive to eIF2 α phosphorylation, with a notable reduction in I κ B α synthesis as compared with RelA/p65 translation (Figure 4B). Lowered synthesis of I κ B α , coupled with the rapid turnover of this inhibitory protein by eIF2 α kinase-independent mechanisms, is thought to trigger the nuclear translocation of NF- κ B and activation of its target genes (Figures 3 and 4). Loss of GCN2 or RelA/p65 increases activation of capases 3 and 8 and enhances apoptosis in response to UV irradiation, suggesting that GCN2 modulation of NF- κ B activity is important for signalling programmed cell death (Figure 6).

The basic features of this model are supported by a recent report by Wu et al. [46] that appeared after the completion of the present study. Wu and colleagues found that eIF2 α phosphorylation contributed to decreased I κ B α levels in response to UV-C exposure, and that PEK/PERK was the eIF2 α kinase facilitating this regulation, as judged by analysis of MC-7 cells transfected with a dominant-negative version of PEK. We analysed *PEK*^{-/-} MEF cells for eIF2 α phosphorylation in response to UV stress, and found a partial reduction compared with wild-type cells (Figure 1). These results suggest that GCN2 is the primary eIF2 α kinase, with a near elimination of eIF2 α phosphorylation over a range of UV-C exposures, and PEK may carry out a secondary role. The results obtained in the present study is also consistent with previous reports that conclude that I κ B α turnover in response to UV stress is independent of IKK phosphorylation of I κ B α N-terminal residue Ser³² (Figure 4) [13,14].

In addition to NF- κ B, eIF2 α phosphophorylation can enhance expression of ATF4 and its target genes. However, there was minimal expression of the ATF4 response pathway after UV exposure (Figure 2), suggesting that the ATF4/CHOP pathway is not required for the anti-apoptotic function of GCN2 in response to UV irradiation. MAP kinase cascades anchored by JNK and p38 are also induced by UV exposure and contribute to expression of genes regulating cellular stress responses, including cell-cycle control and apoptosis [47]. In the example of p38, induction of this MAP kinase by UV irradiation is proposed to be critical for CK2-directed ubiquitination and degradation of I κ B α via the proteasome [15]. Deng et al. [30] found that GCN2 is not required for the phosphorylation of JNK and p38 MAP kinases. These results suggest that GCN2 is not a sensor of UV stress for multiple stress pathways. Furthermore, eIF2 α phosphorylation appears to be effectively induced upon UV exposure in the absence of either JNK or p38 MAP kinase activities, further supporting a delineation between these stress pathways [30]. Combined with the earlier reports, our study suggests that while GCN2-deficient MEF cells are prevented from inducing NF- κ B, and its protective function against UV-directed apoptosis, these cells retain the pro-apoptotic signals of the JNK MAP kinase pathway.

Phosphorylation of eIF2 α triggers changes in the levels of regulatory proteins

Our results indicate that there are at least two fundamental processes by which eIF2 α phosphorylation can regulate stress response pathways. GCN2 activation of NF- κ B in response to UV irradiation illustrates the model whereby a decrease in general translation can preferentially lower the levels of regulatory proteins subject to rapid turnover. In the example of the inhibitory I κ B α protein, this reduction would lead to activation of the NF- κ B transcriptional regulator in response to UV irradiation. Another example is CReP/PPP1R15B [constitutive repressor of eIF2 α

phosphorylation/protein phosphatase 1, regulatory (inhibitor) subunit 15b], which functions in the dephosphorylation of eIF2 α under basal conditions in the absence of stress. In response to translation inhibition, there is reduced synthesis of the labile CReP that contributes to elevated levels of eIF2 α phosphorylation [48]. Such a mechanism could also be adapted to the lowered synthesis of an activator protein subject to a short half-life. In this case, such protein turnover would lead to repression of a stress-responsive pathway.

The second fundamental process by which eIF2 α phosphorylation can control stress response pathways involves preferential translation of specific mRNAs. There are several examples of such gene-specific translation control, including *ATF4*, *CAT-1* (cationic amino acid transporter 1) and *C/EBP α* [16,18,28,49,50]. Phosphorylation of eIF2 α leads to preferential translation of ATF4, triggering expression of a large number of genes involved in metabolism, antioxidation and apoptosis [27]. It is noteworthy that exposure to UV irradiation does not enhance ATF4 levels, despite significant induction of eIF2 α phosphorylation (Figure 2). Furthermore, expression of CHOP was not induced in response to UV-C, consistent with an earlier study [51]. This result would suggest that eIF2 α phosphorylation alone is not sufficient for increased ATF4 expression under certain stress conditions. The underlying difference between ER and nutritional stresses, which induce ATF4 translation, and UV irradiation, may involve lowered ATF4 mRNA that accompanies this UV treatment of MEF cells. In addition to UV exposure, we previously reported that ER stress in *PEK*^{-/-} MEF cells does not elicit expression of ATF4, or its target genes *ATF3* and *CHOP* [33]. Initially this may not appear to be significant, given that PEK is the primary eIF2 α kinase that is activated when misfolded proteins accumulate in the ER. However, it was found that *PEK*^{-/-} MEF cells treated for extended periods of time with the ER stress agent thapsigargin display robust eIF2 α phosphorylation that is comparable with that shown by *PEK*^{+/+} cells. This delayed eIF2 α phosphorylation was nearly absent in MEF cells with a combined deletion of *PEK* and *GCN2*, suggesting that the secondary eIF2 α kinase activated by ER stress is GCN2. These two examples, whereby significant eIF2 α phosphorylation is not linked to induced levels of ATF4 protein, suggest that there are one or more mechanisms that can thwart ATF4 expression. Such mechanisms would provide for versatility in the regulatory proteins induced by eIF2 α phosphorylation. Therefore, the programme of gene expression can be tailored to a given stress condition.

The transcriptional regulator ATF3 is induced in response to diverse environmental stress conditions [52,53]. Previously, we found that eIF2 α phosphorylation, and the subsequent expression of ATF4, was required for increased ATF3 levels in response to amino acid limitation or ER stress [33]. ATF3 was found to be important for expression of CHOP during amino acid limitation, and for induction of GADD34, an eIF2 α protein phosphatase regulatory subunit involved in feedback control, in response to either nutrient or ER stress. However, upon exposure of MEF cells to between 5 and 20 J/m² UV-C, significant ATF3 expression was measured independently of GCN2 activity (Figure 2). This observation indicates that there are multiple mechanisms that enhance ATF3 expression in response to different stresses, and that ATF4 induced by eIF2 α phosphorylation is required for only a subset of these stress conditions.

eIF2 α kinases activate NF- κ B in response to diverse stress conditions

Previously we reported that eIF2 α phosphorylation is required for the activation of NF- κ B in MEF cells exposed to ER stress

or leucine starvation [29]. Characterization of the mechanisms regulating NF- κ B during ER stress indicated that whereas I κ B α was released from NF- κ B, there was no significant reduction in the steady-state levels of I κ B α . Supporting this conclusion, the addition of the proteasome inhibitor MG132 along with ER stress did not block activation of NF- κ B in MEF cells (Figure 5) [29]. This is a distinguishing difference compared with UV stress, whereby proteasome inhibitors block activation of NF- κ B (Figure 5) [13].

An explanation for how eIF2 α phosphorylation can contribute to different mechanisms of NF- κ B regulation involves the synergism between eIF2 kinases and other stress-response pathways. In the example of UV irradiation, decreased translation by GCN2 phosphorylation of eIF2 α is proposed to combine with p38-MAP-kinase-directed turnover of I κ B α to significantly lower the steady-state levels of this repressor protein, facilitating NF- κ B translocation into the nucleus and direct gene expression. Disruption of one stress pathway, loss of GCN2 or deletion of the *p38 α* gene, blocks activation of NF- κ B in response to UV irradiation (Figure 3) [15]. Decreased translation accompanying PEK phosphorylation of eIF2 α is suggested to prevent further synthesis of secreted proteins that would further overload the folding capacity of the ER [23,54]. Using pulse-chase style experiments, we have also found lowered I κ B α synthesis during ER stress, but no measurable lowering of the steady-state levels of I κ B α ([29]; H. Y. Jiang, unpublished work). There does not appear to be a directed turnover of I κ B α during ER stress as has been described for UV irradiation. We do not yet understand the mechanism by which eIF2 α phosphorylation contributes to release of I κ B α from NF- κ B in response to ER stress. Elaborating on our discussion above, in the case of activation of NF- κ B by ER stress, lowered translation may combine with preferential synthesis of a regulatory protein, such as an NF- κ B subunit, or alternatively targeted degradation of a regulator distinct from I κ B α .

We thank Hong Chang for technical assistance, Dan Spandau and Jana Narasimhan for helpful discussions, and Douglas Cavener (Pennsylvania State University, University Park, PA, U.S.A.), Barbara McGrath (Pennsylvania State University, University Park, PA, U.S.A.), Gail Sonenshein (Boston University, Boston, MA, U.S.A.), Randal Kaufman (University of Michigan, Ann Arbor, MI, U.S.A.), and Harikrishna Nakshatri for sharing cell lines and reagents (where no address is given, the people cited are in our Department). We also acknowledge the Biochemistry Biotechnology Facility at Indiana University for technical support. This study was supported in part by grants R01GM49164 and R01GM64350 (to R. C. W.) from the National Institutes of Health, and by a Biomedicine Research Grant from Indiana University School of Medicine (to H.-Y.J.).

REFERENCES

- Bender, K., Blattner, C., Knebel, A., Iordanov, M., Herrlich, P. and Rahmsdorf, H. J. (1997) UV-induced signal transduction. *J. Photochem. Photobiol. B Biol.* **37**, 1–17
- Dent, P., Yacoub, A., Fisher, P. B., Hagan, M. P. and Grant, S. (2003) MAPK pathways in radiation responses. *Oncogene* **22**, 5885–5896
- Herrlich, P., Blattner, C., Knebel, A., Bender, K. and Rahmsdorf, H. J. (1997) Nuclear and non-nuclear targets of genotoxic agents in the induction of gene expression. Shared principles in yeast, rodents, man and plants. *Biol. Chem.* **378**, 1217–1229
- Vousden, K. H. (2002) Activation of p53 tumor suppressor protein. *Biochim. Biophys. Acta* **1602**, 47–59
- Devary, Y., Rosette, C., DiDonato, J. A. and Karin, M. (1993) NF- κ B activation by ultraviolet light not dependent on a nuclear signal. *Science* **261**, 1442–1445
- Jayaraman, J. and Privet, C. (1995) Activation of p53 sequence-specific DNA binding by short single strands of DNA requires the p53 C-terminus. *Cell* **81**, 1021–1029
- Liu, Z.-G., Baskaran, R., Lea-Chou, E. T., Wood, L. D., Chen, Y., Karin, M. and Wang, J. Y. J. (1996) Three distinct signalling responses by murine fibroblasts to genotoxic stress. *Nature (London)* **384**, 273–276
- Barkett, M. and Gilmore, T. D. (1999) Control of apoptosis by Rel/NF- κ B transcription factors. *Oncogene* **18**, 6910–6924
- Baud, V. and Karin, M. (2001) Signal transduction by tumor necrosis factor and its relatives. *Trends Cell Biol.* **11**, 372–377
- Chen, F. E. and Ghosh, G. (1999) Regulation of DNA binding by Rel/NF- κ B transcription factors: structural views. *Oncogene* **18**, 6845–6852
- Li, Q. and Verma, I. M. (2002) NF- κ B regulation in the immune system. *Nat. Rev. Immunol.* **2**, 725–734
- Huang, T. T., Feinsberg, S. L., Suryanarayanan, S. and Miyamoto, S. (2002) The zinc finger domain of NEMO is selectively required for NF- κ B activation by UV radiation and topoisomerase inhibitors. *Mol. Cell. Biol.* **22**, 1513–1525
- Li, N. and Karin, M. (1998) Ionizing radiation and short wavelength UV activate NF- κ B through two distinct mechanisms. *Proc. Natl. Acad. Sci. U.S.A.* **95**, 13012–13017
- Bender, K., Gottlicher, M., Whiteside, S., Rahmsdorf, H. J. and Herrlich, P. (1998) Sequential DNA damage-independent and -dependent activation of NF- κ B by UV. *EMBO J.* **17**, 5170–5181
- Kato, T., Delhase, M., Hoffmann, A. and Karin, M. (2003) CK2 is a C-terminal I κ B kinase responsible for NF- κ B activation during the UV response. *Mol. Cell* **12**, 829–839
- Dever, T. E. (2002) Gene-specific regulation by general translation factors. *Cell* **108**, 545–556
- Wek, R. C. (1994) eIF-2 kinases: regulators of general and gene-specific translation initiation. *Trends Biochem. Sci.* **19**, 491–496
- Harding, H. P., Novoa, I., Zhang, Y., Zeng, H., Wek, R., Schapira, M. and Ron, D. (2000) Regulated translation initiation controls stress-induced gene expression in mammalian cells. *Mol. Cell* **6**, 1099–1108
- Hinnebusch, A. G. (1997) Translational regulation of yeast GCN4. A window on factors that control initiator-tRNA binding to the ribosome. *J. Biol. Chem.* **272**, 21661–21664
- Wek, S. A., Zhu, S. and Wek, R. C. (1995) The histidyl-tRNA synthetase-related sequence in eIF2 α protein kinase GCN2 interacts with tRNA and is required for activation in response to starvation for different amino acids. *Mol. Cell. Biol.* **15**, 4497–4506
- Zhang, P., McGrath, B. C., Reinert, J., Olsen, D. S., Lei, L., Gill, S., Wek, S. A., Vattam, K. M., Wek, R. C., Kimball, S. R. et al. (2002) The GCN2 eIF2 α kinase is required for adaptation to amino acid deprivation in mice. *Mol. Cell. Biol.* **22**, 6681–6688
- Anthony, T. G., McDaniel, B. J., McGrath, B. C., Cavener, D. R., McNurlan, M. A. and Wek, R. C. (2004) Preservation of liver mass during dietary leucine starvation occurs at the expense of skeletal muscle mass in GCN2 knockout mice. *J. Biol. Chem.* **279**, 36553–36561
- Harding, H. P., Calfon, M., Urano, F., Novoa, I. and Ron, D. (2002) Transcriptional and translational control in the mammalian unfolded protein response. *Annu. Rev. Cell Dev. Biol.* **18**, 575–599
- Chen, J.-J. (2000) Heme-regulated eIF2 α kinase. In *Translational Control of Gene Expression* (Sonenberg, N., Hershey, J. W. B. and Mathews, M., eds.), pp. 529–546, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Kaufman, R. J. (2000) Double-stranded RNA-activated protein kinase. In *Translational Control of Gene Expression* (Sonenberg, N., Hershey, J. W. B. and Mathews, M., eds.), pp. 503–528, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Hershey, J. W. B. and Merrick, W. C. (2000) Pathway and mechanism of initiation of protein synthesis. In *Translational Control of Gene Expression* (Sonenberg, N., Hershey, J. W. B. and Mathews, M., eds.), pp. 33–88, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Harding, H. P., Zhang, Y., Zeng, H., Novoa, I., Lu, P. D., Calfon, M., Sadri, N., Yun, C., Popko, B., Paules, R., Stojdl, D. F. et al. (2003) An integrated stress response regulates amino acid metabolism and resistance to oxidative stress. *Mol. Cell* **11**, 619–633
- Vattam, K. M. and Wek, R. C. (2004) Reinitiation involving upstream open reading frames regulates ATF4 mRNA translation in mammalian cells. *Proc. Natl. Acad. Sci. U.S.A.* **101**, 11269–11274
- Jiang, H. Y., Wek, S. A., McGrath, B. C., Scheuner, D., Kaufmann, R. J., Cavener, D. R. and Wek, R. C. (2003) Phosphorylation of the α subunit of eukaryotic initiation factor 2 is required for activation of NF- κ B in response to diverse cellular stress. *Mol. Cell. Biol.* **23**, 5651–5663
- Deng, J., Harding, H., Raught, B., Gingras, A., Berlanga, J., Scheuner, D., Kaufman, R., Ron, D. and Sonenberg, N. (2002) Activation of GCN2 in UV-irradiated cells inhibits translation. *Curr. Biol.* **12**, 1279–1286
- Wu, S., Hu, Y., Wang, J. L., Chatterjee, M., Shi, Y. and Kaufman, R. J. (2002) Ultraviolet light inhibits translation through activation of the unfolded protein response kinase PERK in the lumen of the endoplasmic reticulum. *J. Biol. Chem.* **277**, 18077–18083
- Engelberg, D., Klein, C., Marinetto, H., Struhl, K. and Karin, M. (1994) The UV response involving the RAS signalling pathway and AP-1 transcription factors is conserved between yeast and mammals. *Cell* **77**, 381–390
- Jiang, H. Y., Wek, S. A., McGrath, B. C., Lu, D., Hai, T., Harding, H. P., Wang, X., Ron, D., Cavener, D. R. and Wek, R. C. (2004) Activating transcription factor 3 (ATF3) is integral to the eIF2 kinase stress response. *Mol. Cell. Biol.* **24**, 1365–1377

- 34 Scheuner, D., Song, B., McEwen, E., Liu, C., Laybutt, R., Gillespie, P., Saunders, T., Bonner-Weir, S. and Kaufman, R. J. (2001) Translational control is required for the unfolded protein response and *in vivo* glucose homeostasis. *Mol. Cell* **7**, 1165–1176
- 35 Beg, A. A. and Baltimore, D. (1996) An essential role for NF- κ B in preventing TNF- α -induced cell death. *Science* **274**, 782–784
- 36 Nozaki, S., Sledge, G. W. and Nakshatri, H. (2001) Repression of GADD153/CHOP by NF- κ B: A possible cellular defense against endoplasmic reticulum stress-induced cell death. *Oncogene* **20**, 2178–2185
- 37 Jiang, H. Y., Petrovas, C. and Sonenshein, G. E. (2002) RelB-p50 NF- κ B complexes are selectively induced by cytomegalovirus immediate-early protein 1: differential regulation of Bcl-x(L) promoter activity by NF- κ B family members. *J. Virol.* **76**, 5737–5747
- 38 Vinson, C. R., Hai, T. and Boyd, S. M. (1993) Dimerization specificity of the leucine zipper-containing bZIP motif on DNA binding: prediction and rational design. *Genes Dev.* **7**, 1047–1058
- 39 Siu, F., Blain, P. J., LeBlanc-Chaffin, R., Chen, H. and Kilberg, M. S. (2002) ATF4 is a mediator of the nutrient-sensing response pathway that activates the human asparagine synthetase gene. *J. Biol. Chem.* **277**, 24120–24127
- 40 Ma, Y., Brewer, J. W., Diehl, J. A. and Hendershot, L. M. (2002) Two distinct stress signaling pathways converge upon the CHOP promoter during the mammalian unfolded protein response. *J. Mol. Biol.* **318**, 1351–1365
- 41 Ghosh, S., May, M. J. and Kopp, E. B. (1998) NF- κ B and rel proteins: Evolutionarily conserved mediators of immune responses. *Annu. Rev. Immunol.* **16**, 225–260
- 42 Karin, M. and Ben-Neriah, Y. (2000) Phosphorylation meets ubiquitination: The control of NF- κ B activity. *Annu. Rev. Immunol.* **18**, 621–663
- 43 Kaina, B. (2003) DNA damage-triggered apoptosis: critical role of DNA repair, double-strand breaks, cell proliferation and signaling. *Biochem. Pharmacol.* **66**, 1547–1554
- 44 Karin, M. and Lin, A. (2002) NF- κ B at the crossroads of life and death. *Nat. Immunol.* **3**, 221–227
- 45 Wang, C. Y., Mayo, M. W. and Baldwin, A. S. (1996) TNF- and cancer-therapy-induced apoptosis: Potentiation by inhibition by NF- κ B. *Science* **274**, 784–787
- 46 Wu, S., Tan, M., Hu, Y., Wang, J.-L., Scheuner, D. and Kaufman, R. J. (2004) Ultraviolet light activates NF- κ B through translation inhibition of I κ B α synthesis. *J. Biol. Chem.* **279**, 34898–34902
- 47 Karin, M. and Hunter, T. (1995) Transcriptional control by protein phosphorylation: signal transmission from the cell surface to the nucleus. *Curr. Biol.* **5**, 747–757
- 48 Jousse, C., Oyadomari, S., Novoa, I., Lu, P. D., Zhang, H., Harding, H. P. and Ron, D. (2003) Inhibition of a constitutive translation initiation factor 2 α phosphatase, CReP, promotes survival of stressed cells. *J. Cell Biol.* **163**, 767–775
- 49 Calkhoven, F. C., Muller, C. and Leutz, A. (2000) Translational control of C/EBP α and C/EBP β isoform expression. *Genes Dev.* **14**, 1920–1932
- 50 Yaman, I., Fernandez, J., Liu, H., Caprara, M., Komar, A. A., Koromilas, A. E., Zhou, L., Snider, M. D., Scheuner, D., Kaufman, R. J. and Hatzoglou, M. (2003) The zipper model of translational control: a small upstream ORF is the switch that controls structural remodeling of an mRNA leader. *Cell* **113**, 519–531
- 51 Schmitt-Ney, M. and Habener, J. F. (2000) CHOP/GADD153 gene expression response to cellular stresses inhibited by prior exposure to ultraviolet light wavelength band c (UVC). *J. Biol. Chem.* **275**, 40839–40845
- 52 Hai, T. and Hartman, M. G. (2001) The molecular biology and nomenclature of the activating transcription factor/cAMP responsive element binding family of transcription factors: Activating transcription factor proteins and homeostasis. *Gene* **273**, 1–11
- 53 Hai, T., Wolfgang, C. D., Marsee, D. K., Allen, A. E. and Sivaprasad, U. (1999) ATF3 and stress responses. *Gene Expression* **7**, 321–335
- 54 Harding, H. P., Zhang, Y. and Ron, D. (1999) Protein translation and folding are coupled by an endoplasmic-reticulum-resident kinase. *Nature (London)* **397**, 271–274

Received 8 July 2004/31 August 2004; accepted 8 September 2004

Published as BJ Immediate Publication 8 September 2004, DOI 10.1042/BJ20041164