REVIEW ARTICLE

The transcriptional regulation of the preproenkephalin gene

Gary WEISINGER

Faculty of Medicine, Technion, Haifa, Israel

INTRODUCTION

The expression of eukaryotic genes results from the interplay of a multitude of signal-transduction pathways with an array of *cis*-acting genetic elements, affecting gene transcription at different levels. These *cis*-acting DNA-response elements are able to reversibly enhance, repress or silence transcription in response to regulatory factors that bind to them [1–5]. Once these *trans*-acting DNA-binding factors (usually proteins) bind to their appropriate *cis*-acting DNA-response element, direct or indirect protein–protein interaction(s) between the resultant DNA-factor complex(s) and the gene's transcription apparatus then allow the consequent RNA expression [3–5]. With nearly a decade of intensive research within this field, the specifics of how combinations of extracellular signals are transduced as biochemical messages, into a particular pattern of *in vivo* gene expression, are still not well understood. To address these questions, we and others have focused on the neuropeptide transmitter gene preproenkephalin (ppEnk) and hence the current review is to highlight the current status of ppEnk RNA regulation.

The use and abuse of exogenous opiate drugs (e.g. opium) for cultural, medicinal, military (e.g. the opium wars between Britain and China in the 1800s, see ref. [6]) and social reasons has warranted a steady interest in how they affect human behaviour. More recently, researchers have recognized that endogenous opiates may also exist, which bind to the 'natural' opiate receptors. Hughes and his co-workers [7] were the first to describe two endogenous substances from pig brain, which act as agonists at opiate-receptor sites. Identical endogenous opiates were independently isolated from calf brain soon after [8]. These pentapeptides [7] are called [Leu]enkephalin (Tyr-Gly-Gly-Phe-Leu) and [Met]enkephalin (Tyr-Gly-Gly-Phe-Met) (see Table 1). Since then, approximately 25 different endogenous opiate peptides, isolated from the mammalian nervous system, have been identified [9]. They are all derived from three distinct genes, encoding the precursors for pro-opiomelanocortin [10], pro-enkephalin [11–16] and prodynorphin [17]. Prior to the cloning of these genes it was widely assumed that the enkephalins were derived from both the preprodynorphin as well as the ppEnk genes. The existence of β -endorphin and dynorphin peptides,

which resemble the enkephalins, was the reason for this confusion [18–20]. These matters were ultimately resolved, not only by the cloning of these three genes, but also by the recognition of differences in distribution of these peptides in the pituitary and the brain [21]. The result of much of this work within the central and peripheral nervous systems has been to establish the importance of enkephalin as a neurotransmitter within the stimulus–secretion–synthesis coupling model of regulatory neurobiology [5,22,23].

THE STRUCTURE OF THE ppEnk GENE

(I) ppEnk primary structure

The primary structure for the ppEnk gene was determined for a number of mammalian (human, hamster, bovine and rat) and amphibian (African clawed toad) species (see Table 2). The rat and toad (*Xenopus laevis*) ppEnk genes are each made up of three equivalent exons (Table 2; [24–28a]), while the human and hamster genes have four transcribed exons (Figures 1 and 2; [12,14,16,28–30]). It seems that the untranslated first two exons of the human ppEnk gene are located within, or homologous with, the 5' and 3' ends of the first rat ppEnk exon respectively (see Figure 2). Additionally, the human ppEnk gene has been localized to chromosome 8, at q23–q24 [9]. There is, as yet, no information about ppEnk's chromosome localization in other species. As we have incomplete information for bovine ppEnk, exon comparisons, except for the enkephalin-peptide-coding exon, are not yet possible (Table 2).

The first exon of rat ppEnk is 192 nucleotides in length if measured from its most 5' transcribed base (Figure 1; [25,31,32]). Exon 1 is separated from exon 2 by a 511-base intervening sequence, intron A (Figure 1; [33]). The translated prohormone begins at the fourth nucleotide of exon 2 and encodes a 24-amino-acid signal peptide followed by 33 amino acids of pro-enkephalin within this 141-nucleotide exon (Figure 1; [25,27]). Between the second and third rat ppEnk exons are approximately 3500 nucleotides of intron B [25]. Exon 3, although 978 nucleotides in length, only codes for 223 amino acids of pro-enkephalin [25]. The last 310 nucleotides of this exon, like the first exon, are untranslated [25,27].

The human ppEnk gene shows many similarities, as well as some significant differences, with the rat ppEnk gene. While the translated portion of the human and rat ppEnk genes are very similar (Figure 2) the first rat exon is divided into a 70-base exon I and a 56-base exon II by a small 87-base intron A in the human ppEnk gene (Figure 2; [12,29]). It is as if a small transposable element inserted itself within the first original exon of this gene (see Figure 2). This remarkable difference between the putative untranslated regulatory 5' ends of the human and rat ppEnk genes may indicate the origin of some of the tissue (basal) and treatment (stress) regulatory differences in the expression of this gene between these two species. However, much still needs to be

Table 1 The enkephalin peptides

Peptide	Sequence	Copies/proenkephalin
Octapeptide	Tyr-Gly-Gly-Phe-Met-Arg-Gly-Leu	1
Heptapeptide	Tyr-Gly-Gly-Phe-Met-Arg-Phe	1
[Met]Enkephalin	Tyr-Gly-Gly-Phe-Met	4
[Leu]Enkephalin	Tyr-Gly-Gly-Phe-Leu	1

Table 2 Cloned ppEnk cDNAs and genes

Species	cDNA/gene	Number of exons	Enkephalin peptides		Reference
			[Met]Enkephalin	[Leu]Enkephalin	
Human	cDNA/gene	4	6	1	Comb et al. [12]
Human	cDNA	4	6	1	Comb et al. [29]
Human	Gene	4	6	1	Legon et al. [14], Noda et al. [16]
Hamster	Gene	4	6	1	Zhu et al. [30]
Bovine	cDNA	2(?)	6	1	Noda et al. [15], Gubler et al. [13]
Rat	cDNA	3	6	1	Yoshikawa et al. [27]
Rat	Gene	3	6	1	Rosen et al. [25]
Toad	Gene	3	7	0	Wong et al. [26], Martens and Herbert [24]

Figure 1 Proenkephalin expression from the rat ppEnk gene

The translated prohormone begins at the fourth nucleotide of exon 2 and encodes a 24-amino-acid signal peptide followed by 33 amino acids of proenkephalin. While 223 amino acids of proenkephalin come from exon 3, the last 310 nucleotides of this exon, like the first exon, are untranslated [25,27,28].

done to understand the regulation of this gene, before these subtleties can be properly understood.

Rosen and co-workers [25] performed a comparison between the four species variants of the ppEnk gene, presented in Table 2. Wong et al. [26] repeated a portion of this comparison with more complete *Xenopus* data and Zhu et al. [30] did the same for hamster. The overall degree of homology between the coding regions of these genes was very high (see Figure 2 for data concerning three of these species: human, rat and *Xenopus*, with percentage sequence similarity compared with the rat sequence in brackets under each exon). The highest level of sequence similarity is focused around the enkephalin-peptide-coding sequences

[25,26]. This high degree of sequence conservation from toad to human may indicate the general importance of the enkephalin peptide. A 5' upstream region of 100 bases is highly conserved (98%) between the human and hamster genes [30] and therefore may hold some regulatory sequences not in common with the other species. One major difference between the *Xenopus* proenkephalin gene and the mammalian form of the ppEnk gene is that the toad gene codes for seven [Met]enkephalin and no [Leu]enkephalin peptides, while all of the mammalian genes code for six [Met]enkephalin and one [Leu]enkephalin peptides [24,26]. Again, any functional significance to this difference has still to be determined.

Figure 2 Comparison of the rat, human and toad ppEnk genes

The homology comparisons were against the equivalent region of the rat gene and are represented in parentheses as the percentage under each non-rat exon. They were performed by the author using the NUCALN program (Lipman, D. J. and Wilbur, W. J., Mathematics Research Branch, Building 31, NIADDK, National Institutes of Health, Bethesda, MD, U.S.A.). It should be noted that the first two exons of the human gene are very similar to the 5' and 3' ends of the rat exon 1. It is as if a transposable element jumped, in recent evolutionary history, into the centre of rat ppEnk first exon to form the two human exons.

(ii) ppEnk RNA initiation

Transcription can be regulated at a number of levels; recently we and others have focused on the definition of ppEnk RNA initiation. The rat gene, for example, has four closely spaced somatic transcriptional start sites [31,32,34,35], which seems also to be the case for the human form of this gene (see discussion of this in ref. [31]). By Northern-blot analysis these different rat ppEnk mRNA species are not apparent, as the four start sites are maximally 42 bases apart within a 1.4–1.5 kb mature mRNA (see Table 2 and references therein; [31,32,35]). The same pattern of somatic ppEnk RNA initiation has been validated for the non-neuronal, highly ppEnk-expressing rat heart [36,37]. Additionally, the *Xenopus laevis* (toad) ppEnk gene has at least two major somatic transcriptional start sites, separated by 96 nucleotides (Figure 2; [26]). The more distal of these start sites is 28 bases downstream from a TATA box [26].

Another species of ppEnk mRNA has been described, which is found in rat spermatogenic tissue [32,33,38–41]. This ppEnk mRNA is approximately 1700 nucleotides in length and is the dominant form found in testis and epididymis [33,39,41]. This increase in mRNA size was determined to be due to an alternate set of closely spaced multiple start sites, located within intron A of ppEnk (Figures 1 and 2; [39]). At least five of these RNA start sites are located within a 20-nucleotide region [40]. Alternatively, Borsook et al. [42] claim that germ-cell expression of human ppEnk mRNA is from the somatic 'normal' start site(s) when studied in transgenic mice. In the rat, both the multiple germ-cell start sites and the proximal two somatic start sites, do not have

any 5' TATA box control elements, but GC-rich sequences are found upstream of their respective start sites [31,32,39]. This is not apparent for the distal two somatic start sites, which are four nucleotides apart and do have an appropriate double TATA box element [31]. Moreover, the expression of the germ-cell transcripts in the epididymis seems to be regulated by a newly identified spermatozoa-associated factor [38]. The somatic start sites have also been found to be uniquely tissue- and treatment (stress)-regulated, which will be discussed further below [31,32,34–37].

(iii) Other mechanisms of ppEnk modulation

RNA expression can additionally be modulated at other levels; for example, alternate splicing of the hnRNA, to form different mRNA species, as well as by RNA elongation attenuation, as has been described for some oncogenes (see [43], for example). RNA stability can also regulate the half-life of any RNA species (see the example of the *c-myc* proto-oncogene; [43]). Garrett et al. [33] tried to implicate a mechanism of alternate splicing to explain the 1700-nucleotide testes and epididymus ppEnk mRNA. This was later disputed and disproven by Kilpatrick et al. [39], who showed that the different ppEnk mRNA species was due to an alternate set of germ-cell transcriptional start sites (see Figure 1, intron A).

Another incongruity has recently been recognized by us, which may imply a novel additional transcriptional level [32,35; G. Weisinger, J. D. DeCristofaro and E. F. LaGamma, unpublished work). When studying ppEnk mRNA expression within the nervous system, using either exon 1 or exon 3 probes

by Northern, nuclease-protection or primer-extension analysis, a signal disparity becomes obvious. This is not due to probe size or other probe-associated artifacts, as different probes or oligomers derived from exon 1 or exon 3 sequences give the same pattern. An example of this is the apparent inducibility of exon 1 striatal ppEnk mRNA sequences (1.4 kb), which followed whole animal cholinergic drug treatments ([32,35]; G. Weisinger, J. D. DeCristofaro and E. F. LaGamma, unpublished work). We usually measured a 10–15-fold induction. Alternatively, on using exon 3 probes in the same experiment no induction was seen ([35]; G. Weisinger, J. D. DeCristofaro and E. F. LaGamma, unpublished work). A possible explanation could involve the importation of a similarly sized foreign first exon on induction, to replace the normal ppEnk exon 1. A variation of this notion was recently described by Brooks and co-workers [44]. Using *in situ* hybridization techniques they identified a novel population of cells in the reticular thalamic nucleus and basal forebrain. These cells contained ppEnk RNA transcripts that were initiated within intron A and did not contain ppEnk exon 2 sequences [44]. If a similar mechanism is possible in the striatum our incongruous observations above become explainable. More experimentation is needed before any conclusions can be reached.

DeCristofaro and co-workers [35] have recently identified an inducible smaller-molecular-mass adrenomedullary ppEnk mRNA species. This species appears as a major species in adrenomedullary mRNA following a course of whole animal oxotremorine treatment [35]. This smaller mRNA species is unlikely to be degraded RNA, due to its definition and reproducibility. Additionally, it may be related to similarly sized ppEnk mRNA species reported in the central nervous system [27,45] and rat ovary [46]. These smaller mRNA species require further characterization.

PREPROENKEPHALIN mRNA HAS A WIDE TISSUE DISTRIBUTION IN MAMMALS

(I) Distribution

Enkephalin peptides are found widely throughout the nervous system. Access to ppEnk DNA/RNA probes, derived from the available cloned sequences (see Tables 1 and 2), as well as to anti-enkephalin antibodies, has allowed investigators to determine the relative abundances of ppEnk RNA and its peptides in discrete regions of the unperturbed nervous system. Figure 3 shows the relative average composite tissue distribution of ppEnk RNA in discrete structures from the unperturbed rat nervous and sexual systems. The data in this Figure was compiled from eight separate studies [27,32,47–52].

The adult rat striatum has the greatest content of enkephalin-derived peptides of any brain region (Figure 3; [27,32,47–54]). This high level of striatal ppEnk mRNA expression is apparently dependent upon cerebral cortical afferents [55]. Within the rat nervous system the thalamus and the adrenal medulla have the lowest basal levels of ppEnk mRNA expression [32,50,52]. Although there are only low levels of steady-state ppEnk mRNA in the basal rat adrenal medulla, the stores of enkephalin peptide have been localized to adrenaline-containing chromaffin cells [56–60]. Moreover, quantities of the proenkephalin prohormone have also been identified in the soluble portion of both rat and bovine adrenal chromaffin cells [56,59], and it seems to be developmentally regulated [58,60].

It is interesting that the tissue with the third highest levels of ppEnk mRNA, presented in Figure 3, is the testis, a non-neuronal tissue [32,33,39,61]. In this case, Garrett and co-workers [33] claim that most of the ppEnk mRNA present in the testes is not translated and processed into the pentapeptide. A detailed

Figure 3 Relative tissue distribution of ppEnk mRNA

Here is the relative composite mean tissue distribution of ppEnk mRNA in discrete structures from unperturbed rat nervous and sexual systems. The data in this Figure were compiled from eight separate studies [27,32,47–52]. The abbreviations for the structures are as follows: St, striatum; OT, olfactory tubular; Te, testes; NA, nucleus accumbens; OB, olfactory bulb; Hy, hypothalamus; Pn, pineal; CS, cervical spine; Ep, epididymus; Po, pons and medulla oblongate; FC, frontal cortex; Mi, midbrain; Pi, pituitary; Ce, cerebellum; Hi, hippocampus; AM, adrenal medulla; Th, thalamus; Ov, ovary. As a frame of reference the 100% value used for striatum here, represents a mean of 34 pg of ppEnk mRNA/ μ g of total RNA [48,50,52].

relative ppEnk tissue expression pattern has not been performed for other species to date, but a limited study comparing Sprague-Dawley rats and Syrian hamsters is in the literature [50]. This study was particularly interesting due to the finding of comparatively high ppEnk expression in the hamster adrenal medulla (123% of that found in hamster striatum). The other structures compared in this study, hippocampus, hypothalamus and striatum showed relative ppEnk levels comparable with the rat (Figure 3). Pittius et al. [62] found a similar rank order in bovine tissues, where ppEnk mRNA levels are 20–400 times higher in the adrenal medulla than in the central nervous system.

In another study, which used the *in situ* hybridization approach to study ppEnk mRNA expression in sheep [63], the highest densities of ppEnk-expressing cells were found in the nucleus accumbens, caudate-putamen (striatum), olfactory tubercle, the central nucleus of the amygdala, the paraventricular nucleus of the hypothalamus, the suprachiasmatic nucleus and the gigantocellular division of the medullary reticular formation. This widespread distribution and relative expression of ppEnk RNA in sheep was in agreement with similar data for rats (see Figure 3; [63]). Additionally, as can be seen in Figure 3, the rat pituitary does not, as a whole, express much ppEnk RNA. The reason for this could be the sublocalization of expression, as was found for the sheep pituitary [63]. In sheep, a dense ppEnk signal was observed in the intermediate lobe of the pituitary and not at all in the anterior or neural lobes [63]. In the other studies contributing to Figure 3, the pituitary was dissected and analysed as a whole, hence averaging a strong signal in a few cells over a whole structure.

(II) ppEnk expression within tissues or brain structures

(a) Nervous system structures

A few reports have tried to define further the exact subregion, or cells, expressing ppEnk within a few of the structures shown in

Figure 3. Senba and co-workers [64] used *in situ* hybridization to show a moderate number of ppEnk mRNA-expressing neurons in the glomerular layer of the rat olfactory bulb. Moreover, in the granule-cell layer, numerous small ppEnk RNA-expressing cells were also found [64]. In the rat cerebellum, Spruce and co-workers [65] localized ppEnk expression to Golgi cells, macro-neurons with deep cerebellar nuclei and a population of Purkinje cells.

In the central nervous system the ppEnk expression is found both in astrocytes [65–70] and nerve cells (fibres, terminals and cell bodies) [65,69–72], but not other neuronal support cells [68]. This pattern of enkephalin expression in the brain, with their interaction with brain opioid receptors is consistent with the notion that, in this context, the enkephalins are neurotransmitters and neuromodulators [73].

(b) Non-nervous-system structures

In addition to finding ppEnk expression within the nervous system, there is an increasing number of reports of non-nervous-tissue expression. One study, focusing on ppEnk expression in the epididymis, found a very dense local expression in the initial segment of the epididymis [38]. The authors claimed that while ppEnk RNA expression was limited to a small region of this structure, it comprised as much as 5% of total cellular RNA there [38]. In the testes, the 1700-nucleotide germ-cell mRNA species can be found in spermatocytes, round spermatids and condensed spermatids [74]. Of these three types of germ cells, the highest ppEnk levels are found in the round spermatids [74]. Zinn and colleagues [40] suggest that as ppEnk products are synthesized and stored in acrosomes during spermatogenesis and the gene products are released from the mature sperm after the acrosome reaction, ppEnk gene products may have unique functions during fertilization [40,75].

In the rat ovary, where very low levels of steady-state ppEnk mRNA were found (compared with striatum, see Figure 3), following careful investigation, significant levels of ppEnk RNA were found in the theca cells of the preovulatory follicles [76]. It seems, as for the pituitary, that whole-tissue RNA analysis may not be subtle enough to detect specific RNA expression. Abundant ppEnk expression in the rat oviduct, which was restricted to the secretory and ciliated epithelium of the mucosa, was also found by Rosen et al. [76]. Basal ppEnk expression was also studied in the rat uterus and identified in the deep glandular layer of the endometrium. This expression was dramatically induced upon pregnancy in the same uterine cells, primarily at the site of implantation [42,76]. Borsook et al. [42] therefore concluded that ppEnk played some type of autocrine/paracrine role within the female reproductive system.

High level ppEnk RNA expression has also been reported for the rat ventricular heart muscle [36,37,77,78]. Similarity between multiple RNA start site usage between heart and brain ppEnk was pointed out [31,32,36]. Springhorn and Claycomb [36,78] speculated that the enkephalins may be involved, at different levels, in the differentiation of both neonatal and adult heart cells. Some other reports have identified ppEnk RNA expression not only in rat heart, brain and testes, but also in articular chondrocytes, bone, developing skin, fibroblasts, intestine (neonate), liver (neonate), lung, kidney and skeletal muscle (neonate) [73,79–85]. The authors of four of these reports suggested linking proenkephalin expression to mesodermal lineage tissue growth and/or differentiation. They came to their conclusion using sensitive *in situ* hybridization techniques. Furthermore, the authors of one of these reports [85] hypothesized that enkephalin peptides may have a dual function in mesodermal

tissue; first, in the control of tissue regeneration, as well as in the control of pain. Kew and Kilpatrick [82], having only used the cruder Northern analysis approach, claimed multiple embryonic origins of the ppEnk-expressing tissues. More specifically, liver and lung arise from embryonic endoderm, skeletal muscle, heart and kidney from mesoderm and brain from ectoderm. Skin's origin is from both ectoderm (epidermis) and mesoderm (dermis), and intestine has both endodermal and mesodermal origins [82].

White blood cells can also express significant levels of ppEnk mRNA [86–97]. Murine CD-4-positive [91,92], rat [95,97] and human [86,90] T-cells together with monocytes [86] and macrophages [98] express this ppEnk mRNA. T-cells seem to express not only the 1400-nucleotide species of ppEnk mRNA, but also the spermatogenic 1700-nucleotide form [91]. In the case of T-cells the smaller RNA form predominates, while in testis and epididymis the larger species is more obvious [33,39,91]. Kuis et al. [86] report that the ppEnk prohormone is processed into [Met]enkephalin in monocytes but not in T-cells. Hence, in T-cells only the prohormone is found [86]. As thymocytes do not process the prohormone to the pentapeptide, Linner et al. [91,92] suggest a thymocyte growth and differentiation role for proenkephalin. Furthermore, the ppEnk mRNA expression, in this case, is regulated by Interleukin-1B [92]. Alternatively, as monocytes do produce and secrete [Met]enkephalin, the authors suggest that the enkephalins contribute to the regulation of both local and systemic aspects of the host's immune response [86].

The wide distribution of the ppEnk gene products throughout mammalian tissue may indicate that they serve more than one function. As suggested above these putative functions range from the traditional neurotransmission and neuromodulation to growth and differentiation promotion of particular cell types. A number of investigators have begun to show a correlation between ppEnk gene expression and cell division for various cell types [38,73,74,79,84,87–89]. These various cell types include different neuroendocrine cells [87–89], articular chondrocytes [79], spermatocytes [38,74] and embryonic fibroblasts [73,84]. Gustin and co-workers [87] have even suggested that endogenous enkephalins may be an integral component of different brain tumours. On a different track, Kuis and colleagues [86] have advanced the possible involvement of [Met]enkephalin in the immune response. This last novel suggestion may indicate enkephalin as a candidate second messenger for mammalian neural-immune interactions and will be further discussed later.

PREPROENKEPHALIN mRNA REGULATION

(i) In development

In general the opioid peptide precursor proenkephalin is produced by a variety of rat tissues, during early post-natal development. In most tissues the steady-state levels of ppEnk gene expression decrease as the animal (i.e. rat) becomes adult. Three exceptions to this pattern are brain, heart and epididymis, where levels of ppEnk mRNA increase as the animal ages [38,53,78,82,99].

Adult rat striatum has the highest content of ppEnk mRNA (see Figure 3) and enkephalin-derived peptides in the brain. Schwartz and Simantov [53] undertook to carefully map the developmental expression of rat striatal ppEnk mRNA, and they found a biphasic developmental profile of ppEnk expression, with an initial peak 2 days after birth, a drop to embryonic levels by post-natal day 7 and a second increase to adult levels over the second to fourth weeks following birth [53]. This developmental biphasic ppEnk mRNA profile was not mirrored by striatal enkephalin peptides [53]. The enkephalin peptide increased steadily during the first post-natal week of life, which was

followed by a more gradual rate of increase [100]. Rosen and Polakiewicz [99] performed a similar but wider developmental study, focusing not only on striatum, but also hypothalamus and pontin. Although the biphasic developmental pattern mentioned above was seen in all three of these brain structures, adult striatal ppEnk mRNA levels were considerably less in the latter study when compared with the study by Schwartz and Simantov [53]. This difference may reflect the different rat species used in each of these studies.

Another rat tissue that expresses more ppEnk mRNA in the adult than the neonate is the heart [36,37,78,82]. The developmental profile of ppEnk mRNA in this tissue starts with a relatively low level of expression for the first 2 weeks after birth, which decreases even further over the next 2 weeks [78]. By 3 months of age ppEnk mRNA was maximally expressed [78]. In the same study ppEnk RNA, in cultured primary rat ventricular cardiac muscle cells, was shown to be very responsive to cyclic AMP stimulation. As rat cardiac muscle cells cease cell division by 3 weeks of age [101–103] and at this same time (3 weeks of age) functional adrenergic innervation of the heart occurs [102,103], Springhorn and Claycomb [78] speculate that there is a correlation between functional adrenergic innervation of the heart and the subsequent cyclic AMP modulation of ppEnk mRNA.

The pattern of ppEnk gene expression in the epididymis during development is localized to the principal cells of the initial segment [38,42]. This 1700-nucleotide mRNA species is strongly induced at about 44 days after birth [38]. This developmental age is associated with the initial entry of spermatozoa into the epididymal lumen. It has been claimed that this 1700-base-long ppEnk species is translated to a much lesser extent than the 1450-base transcript [41]. Recently, the DNA regulatory element responsible for this translational blockage has been defined, and is currently being characterized [41]. Additionally, upon hypophysectomy, a 60-fold decrease in epididymal ppEnk mRNA levels occurs, which can be prevented by testosterone replacement [38]. Garrett et al. [38] speculate that this developmental increase in ppEnk mRNA is due to a spermatozoa-associated factor, hence it is also responsible for this epididymal ppEnk mRNA developmental profile.

(ii) Trans-synaptic regulation of adrenal ppEnk

In the first two post-natal weeks of the rat's life, adrenal catecholamine pathway maturation, adrenal innervation, functional neurotransmitter release and adrenocortical responsiveness occur [104–113]. Moreover, catecholamine biosynthetic pathways can be augmented by trans-synaptic impulse activity by the end of the first post-natal week [105,107–115]. We, and others, have shown that this sympathoadrenal innervation of the splanchnic nerve to the adrenal, even though anatomically in place prior to birth, only becomes functional, for ppEnk pathways, by the end of the first week of life [108,109,112,114,116].

Once functional innervation of the adrenal medulla occurs, a mechanistic paradox, regarding trans-synaptic activity to the adrenal medulla and enkephalin biosynthesis, becomes apparent. It appears that enkephalin biosynthesis is both positively and negatively regulated by trans-synaptic stimulation [31,32,35,50,108,115–120]. That is, steady-state levels of rat ppEnk mRNA, proenkephalin and enkephalin peptide are induced by decreased trans-synaptic activity, by cholinergic-nicotinic receptor blockage, denervation or adrenal explantation [108,114–118,120,121]. Conversely, by increasing trans-synaptic activity through hypoglycaemic shock, moderate cold stress or treatment with cholinergic agonists, both rat enkephalin mRNA

and peptide content rise [31,32,35,112,116–118,122,123]. Quite a number of hypotheses have been suggested to explain this paradox, with little supportive data (discussed in LaGamma et al. [108]). The exception, however, is the substantial body of evidence indicating that many of the relevant receptor-mediated signal-transduction pathways converge together on the regulatory region of the ppEnk promoter (discussed below, in the next section) [29,31,32,61,114,122,124–134]. Hence, when an environmental signal is conveyed to the genome, a ppEnk-specific combination of multiple *trans*-acting nuclear factors are regulated through these final common second-messenger pathways. This will be discussed in detail below.

Levels of hamster and bovine adrenal medullary ppEnk mRNA are many times higher than for the rat [30,50,62,135]. In fact adrenal medulla ppEnk mRNA expression in both bovine and hamster is greater than for their respective striata [30,50,62]. In the case of the hamster, a decrease in trans-synaptic activity leads to a drop in ppEnk mRNA and peptide levels (i.e. 54% after 2 days and 90% by 11 days) [50,135]. As this is the exact opposite effect to that found in the rat, a biochemical comparison of ppEnk expression in both of these species' adrenal medulla would be most revealing.

Reserpine, an anti-hypertensive drug, was originally known for its inhibition of catecholamines uptake into adrenal chromaffin granules [136]. Immediately after treatment, a reserpine-induced depletion of co-localized catecholamines and enkephalin peptides occurs, which is followed by a reflex stimulation of trans-synaptic activity [137–140]. Moreover, reserpine treatment of bovine chromaffin cells, as well as hamster adrenal medulla, produces a dose-dependent increase of both tyrosine hydroxylase and ppEnk mRNA species [135,141–143]. The case of the rat is different, in that reserpine treatment leads to an increase in the adrenal medullary tyrosine hydroxylase mRNA, and co-localized ppEnk mRNA decreases [144–146]. Here an apparent replacement of vesicular enkephalin pentapeptide stores was due to processing of soluble proenkephalin prohormone [56,59]. Hence, there are different types of increased trans-synaptic activity for the rat, one that increases ppEnk mRNA expression (hypoglycaemia, cold stress and cholinergic treatments, see above for discussion) and the other (reserpine treatment) that leads to a down-regulation of ppEnk. This suggests that other factors must work in concert to change the effect of increased trans-synaptic activity on ppEnk expression.

(iii) Glucocorticoids and ppEnk

Glucocorticoids regulate the expression of various neurotransmitters in the central nervous system [49,147–150]. The initially controversial suggestion that glucocorticoids can regulate neuropeptide genes directly, by activating glucocorticoid receptors, is now well accepted [151]. These activated receptors bind to specific DNA sequences associated with the regulated gene and in turn regulate its transcription [151]. ppEnk has one and perhaps more such elements [108,129,148,152]. This element has since been generally defined and called the glucocorticoid response element (GRE, see below) [151]. It has now become apparent that rat forebrain regions, which express ppEnk mRNA, also express corticosteroid receptors [153–155]. Moreover, Ahima and colleagues [147] reported the requirement of glucocorticoids for the maintenance of basal ppEnk mRNA expression in most regions of the rat forebrain [121,147]. In addition, Chao and McEwen [148] and others, found that these hormones are necessary for ppEnk mRNA-modulated expression in the brain striatum [148], explanted adrenals [107,121] or in rat adrenal medullae *in vivo* [107]. If one compares the effect of glucocorticoid

treatments on ppEnk mRNA expression in adrenalectomized and sham-operated rats, one finds considerable regional heterogeneity in response [147]. Ahima and colleagues [147] conclude that, while ppEnk expression may be only minimally affected by circulating glucocorticoids in the unperturbed rat, during stress (where blood glucocorticoid levels can show dramatic changes) these hormones play a major role in any ppEnk mRNA response. This is particularly likely in the striatum, the hippocampus [147] and the hypothalamus [150].

THE BIOCHEMISTRY OF SIGNAL TRANSDUCTION TO THE ppEnk GENE

One of the major challenges confronting modern biochemists, cell biologists and physiologists is the delineation of the molecular mechanisms that cells, tissues and whole animals use to adapt their respective phenotypes in response to environmental signals. Over the last decade, biochemical studies have been essential in the elucidation and the molecular characterization of different classes of transcription factors (see reviews: [1,3,4,22,43,156–159]). The interaction and regulation of these classes of transcription factors with *cis*-acting DNA elements are important in permitting the precision and nuance of gene regulation required for complex functional systems, such as the nervous system. In the following discussion, the focus will be on those signal-transduction-related *trans*-acting factors that interact with ppEnk *cis*-acting DNA regulatory sequences.

(I) The cyclic AMP response elements (CREs) and ppEnk

A family of nuclear factor (NF-1) proteins, together with the transcription factors, AP-1, AP-2 and AP-4, have been shown to modulate ppEnk mRNA transcription in response to changes in cyclic AMP levels [125–127,132,133,160–162]. These data were obtained using *in vitro* cell lines together with transfection studies. A 35-bp region of DNA, now known to contain CREs, is closely linked to the human ppEnk promoter and positioned upstream to it (–70 to –105 bp). This CRE region has been shown to act as a second-messenger-inducible *cis*-acting DNA enhancer, which functions in an orientation- and position-independent manner (Table 3; [125–128]). Since its discovery, it has been found to regulate ppEnk transcription not only by cyclic AMP, but also by phorbol ester (an activator of protein kinase C), Ca^{2+} [126–128] and *in vitro* by membrane depolarization-dependent intracellular pathways (Table 3; [133]). Point mutational analysis of this CRE region has shown that it is made up of two different DNA elements, CRE1 and CRE2 (see Figure 4). CRE1 is separated from CRE2 by 5 bp or about a 180° rotation around the DNA [126] and they act cooperatively to regulate ppEnk transcription [126,127,133]. The upstream CRE1 element has little enhancer activity by itself, but greatly augments cyclic AMP inducibility if combined with the more downstream CRE2 [163]. These CREs

Figure 4 Summary of comparative ppEnk promoter footprint data

The top diagram shows most of the superimposed binding factor positions with respect to the four ppEnk somatic transcriptional start sites and the cyclic AMP enhancers (CRE1 and CRE2). The data in the second and third diagrams (rat striatum and adrenal medulla) are summarized from LaGamma and Weisinger [131] and LaGamma et al. [122] respectively. The data presented in the bottom diagram are from Rattner et al. [186]. Closed arrowheads represent the main ppEnk start sites used in that tissue, while the open arrowheads are weak RNA start sites. In the bottom diagram the horizontal arrowhead indicates that the tissue-specific start site pattern has yet to be determined. The red boxes represent tissue-specific ppEnk promoter footprints.

Figure 5 Summary of upstream ppEnk DNAase I hypersensitivity sites and some other DNA functional element locations

Tissue distribution of regions of active chromatin (DNAase I hypersensitivity) associated with the rat ppEnk gene, are shown relative to the principal rat ppEnk transcriptional start site, in untreated animals [187]. Superimposed on this are the approximate locations of positive and negative regulating glucocorticoid response elements [3,151,192–194] as well as a putative Z-DNA-forming sequence [164].

are well conserved between species, with a 1 bp difference between human and rat sequences [25,131,164]. This relatively high, local sequence identity over the CREs is particularly interesting, as most of the sequences immediately 5' to the CREs become extremely unrelated. Indeed, a putative Z-DNA structure, over 150 bp in length, can be found in this upstream ppEnk region only in the rat sequence (see Figure 5) and not the human (compare Durkin et al. [164] and Comb et al. [29]). Another group has suggested that this type of sequence could also form a triplex DNA structure [165].

CRE2 (TGCGTCA), whose binding site closely resembles the phorbol ester response element (i.e. TRE = TGACTCA), also

Table 3 Signal transduction of ppEnk RNA

Regulator	Element	Effect on ppEnk RNA
(1) Cyclic AMP, phorbol esters, Ca^{2+}	CRE	Increased
(2) Nicotine, muscarinic agonists	CRE/CARE	Increased (striatum)
	CARE	Increased (adrenal medulla)
(3) Glucocorticoid hormones	GRE (+) and CRE	Increased
	GRE (–)/?	Decreased
(4) NF- κ B, ConA+PMA	BETA2	Increased (lymphocytes)

has the consensus CRE motif contained within it (CGTCA) [126,164]. Using affinity-purified AP-1 proteins, or antibodies to them, high-affinity binding of AP-1 complex proteins to the CRE2 site has also been demonstrated (the same AP-1 proteins bind with a lower affinity to a CRE1 oligonucleotide) [126,127,131,162]. The AP-1 family of proteins are composed of many proteins in the Fos, FRA and Jun families of transcription factor [5]. They act through binding to the CREs as, at the very least, a heterodimer [132]. Kobierski and colleagues [162] demonstrated that two members of the Jun sub-API family can either activate (D-Jun) or inhibit (B-Jun) CRE2-driven transcription through competitive binding. It also seems that CRE1 (TGGCGTA) has other, higher-affinity binding factors (CREB and EnkTF-1) [164,166,167]. A recent report demonstrated the importance of c-fos co-expression on ppEnk inducibility in bovine chromaffin cell cultures [168]. The investigators compared the relative contribution of c-fos, c-jun and CREB in a particular ppEnk induction model [168]. Moreover, they claimed that in their model the CRE2 DNA element was more critical than the CRE1 element [168]. Another report claims that both CRE1 and CRE2 elements are equally important in ppEnk induction by discussing the putative secondary hairpin structure CRE1 can make with CRE2, which is necessary for its enhancer activity [169].

Hyman et al. [128] report an AP-2 element, located slightly downstream of CRE2, which acts synergistically with it (see Figure 4). When an *HpaII* restriction site within this AP-2 site was CpG methylated *in vitro*, not only was AP-2 binding inhibited, but ppEnk expression was also prevented on vector transfection into a glioma cell line [170]. Both of these groups indicate that only appropriate factor binding to both of the CREs as well as the AP-2 elements will lead to the maximal response of the ppEnk 'CRE' enhancer from cyclic AMP and phorbol ester treatments [128,170].

AP-4 is another transcription factor which has been found to bind over the CRE2 locus [127,163]. Even though its transcriptional role in regulating the ppEnk gene is still unclear, AP-4 has been found necessary for simian virus 40 (SV40) enhancer regulation [171]. It seems that a synergism must occur between AP-1 and AP-4 to allow both heterologous and homogeneous SV40 enhancer action [171]. This may also be the case for ppEnk due to the overlapping binding sites of these two proteins (see Figure 4). This possibility remains to be validated.

Considerable *in vitro* data exist, which document the functional importance of the ppEnk CREs and AP-2 *cis*-acting elements in controlling the human ppEnk promoter. These results were determined using multiple established cell lines (discussed above). Unfortunately, there are few studies to verify any *in vivo* significance of these regulatory genetic elements. The few studies in the literature fall into two types, the transgenic animal approach [40,42,172,173] and a transplantation paradigm [174,176].

Transgenic studies of mice, in which specific segments of the human ppEnk promoter are introduced stably into the genome, have recently focused on a 193-nucleotide fragment of upstream sequence within the human ppEnk promoter region (or a larger 5' region but only for germ-cell tissue [42]). This sequence contains both of the CRE elements as well as the AP-2-binding site. Data from these transgenic animal experiments have demonstrated that enough information is coded in this 193 bp promoter to mimic basal ppEnk expression in most rodent tissues, the primary exception being the almost non-expression of this chimeric construct in the brain striatum [40,172,173,177]. Moreover, this limited promoter region, which contains the whole CRE enhancer, confers trans-synaptic ppEnk up-regulation in

three well-studied models. The first of these is the up-regulation of striatal ppEnk mRNA following haloperidol treatments in the whole animal [178–181]. Secondly, adjuvant injection of mouse footpads has been shown to induce ppEnk expression in the spinal cord [182]. Finally, ppEnk gene expression is regulated by CREB in the paraventricular nucleus in response to hypertonic saline stress [183]. All of these model paradigms may therefore involve the CRE enhancer sequences.

At least one major and well-documented level of synaptic regulation was absent from these transgenic mice, which involves the inducibility of ppEnk mRNA in the hippocampus following seizure [172]. The transcription factors, which recognize the sequences in this 193 bp ppEnk promoter fragment, were up-regulated in the hippocampus following seizures, yet these constructs could not be induced. Hence, as this construct did not contain all the necessary sequences to allow ppEnk induction, it follows that another *cis*-acting seizure element must exist either upstream or downstream of this 193 bp ppEnk fragment.

An alternate method of defining functional *in vivo cis*-acting elements is through a transplantation procedure we recently developed [174,175]. In this case, stably, genetically modified primary astrocytes were stereotactically transplanted into a syngeneic host animal. In our initial experiments we inserted a human ppEnk promoter–chloramphenicol acetyltransferase reporter vector into primary rat striatal astrocytes in culture and then transplanted them into a syngeneic rat striatum [174–176]. Here, as opposed to the traditional transgenic experiments, we managed to reproduce striatal levels of ppEnk RNA expression. Additionally, we demonstrated that endogenous striatal dopamine levels contribute to the high basal levels of striatal ppEnk RNA expression [176].

(ii) The cholinergic–nicotinic pathways

Cholinergic receptor modulation of transmitter genes is an important maturational factor affecting transmitter choice in the developing rat autonomic nervous system [106]. These same signals serve a parallel function in the mature animal, where they maintain the transmitter phenotype, as well as play a role in modulating transmitter phenotypic plasticity [185]. We have reported that there is a synergistic effect of combined cholinergic nicotinic and muscarinic receptor agonist treatments, at the level of steady-state ppEnk mRNA, in the adrenal medulla (Table 3; [32,35,122]). In this case the mixed cholinergic nicotinic and muscarinic agonists induced adrenal medullary ppEnk mRNA more than 100-fold [32,35,122]. Additionally, we also demonstrated a 10–15-fold induction of ppEnk exon 1 RNA sequences from the same animals in the brain striatum (Table 3; [32,131]). It seems that different levels of transcription are activated between these two tissues in the same animal for the same gene. Hence, it would not be surprising if a different mosaic of transcription factors are involved in each tissue.

To functionally define the *cis*-acting element(s) *in vivo*, transgenic or transplantation paradigms, as discussed in the last section, could prove very useful. Unfortunately, these methods have yet to be applied successfully to this problem. Another approach, that of studying the composition of transcription factors in ppEnk-induced versus basal adrenal medulla [122] or striatum [131,184], has provided significant data. By means of gel shift analysis and DMS interference footprinting methodologies LaGamma and colleagues [122] characterized the unique pattern of adrenal medullary ppEnk promoter DNA-binding factors from basal or combined nicotine plus oxotremorine-treated animals (see Figure 4). It is quite apparent that the pattern of factor binding in the adrenal medulla is different to that found in

the striatum (see Figure 4; [108,122,130,131]). Interestingly, CREB, Fos and Jun family proteins were clearly footprintable over human or rat CRE elements using striatal extracts and specific antibodies [131,184]. Moreover, the striatal extracts from the ppEnk-induced animals gave even stronger and wider footprints over the CRE enhancer region than the basal extracts [131]. The identities of the Fos and Jun proteins were confirmed in the same paper [131], using other methods. Alternatively, nuclear extracts prepared from rat adrenal medulla failed to footprint any factors over the CRE enhancer region for basal or ppEnk-induced animals (see Figure 4; [122]). These data are consistent with the notion that the ppEnk RNA induction measured in cholinergic agonist-treated animal basal ganglia (striata) may be a direct result of CRE enhancer activity. On the other hand, the different type and stronger ppEnk mRNA induction seen in the adrenal medulla may be the result of factor binding to an as yet undescribed 'cholinergic agonist response element(s)' (i.e. 'CARE' element). It is also apparent from the footprint data, that this 'CARE' element does not reside in this region of the ppEnk promoter [122] but may overlap with other regions [186].

As the above 'CARE' element(s) were not able to be assigned to the above region of the ppEnk promoter, DNAaseI hypersensitivity (hs) assays were employed to identify other regions of active chromatin associated with the rat ppEnk gene [187]. These investigators found hs sites -200, -1800, -2500 and -3000 bp upstream of the principal rat ppEnk transcriptional start site in untreated animals (Figure 5; [187]). While the -1800, -2500 and -3000 bp hs sites were constitutively expressed in the liver, striatum and adrenal medulla with no apparent tissue specificity, the -200 bp hs site was most intense in the unstimulated adrenal medulla and less intense in the striatum (Figure 5; [187]). This -200 bp hs site was most intense slightly upstream of the CRE enhancer and covered a region that included this enhancer region. It was not visible in rat liver. When the basal state tissues were compared with those from combined nicotine plus oxotremorine-treated animals, no differences of DNAaseI hypersensitivity were seen [187]. These authors conclude that the relative intensity of the -200 bp hs site between basal tissues could indicate a chromatin structure which is poised for increased transcription. In other words, the tissue with the strongest -200 bp hs site, the adrenal medulla, could lead to the greatest ppEnk induction (> 100-fold) after mixed cholinergic treatments to the whole animal [32,35], while the striatum and its less intense -200 bp hs site could only induce a much smaller effect. In the liver, where there is no -200 bp hs site there is no ppEnk inducibility [187].

Data from Joshi and Sabol [129], suggest the presence of another enhancer between -2700 and -1050 bp upstream of the rat ppEnk RNA start sites (discussed in the next section). Hence, two of the upstream hs sites (-1800 and -2500) may be associated with this other putative genetic element. The fact that the three farthest upstream hs sites are constitutive in the tissues studied may only reflect the necessity for constitutively open chromatin at those sites. If this is so then one could imagine the open passage of a variety of transcription factors to a particular site on the DNA. This may be similar to the Fos to FRA transition discussed by others [5].

(iii) The glucocorticoid pathways

Glucocorticoid modulation of ppEnk in the central and peripheral nervous systems [49,107,121,123,147,148], as well as in some non-nervous systems [91,186,188-191], is well recognized (discussed above). The molecular mechanisms of these gluco-

corticoid effects have yet to be well characterized. It seems that in hormonally regulated eukaryotic cells, glucocorticoid-activated cytosolic receptors translocate to the nucleus and bind to a particular 15 bp GRE [(T/C)GGT(A/T)CA(A/C)(A/T)-TGT(T/C)CT]. This DNA-activated receptor binding thereby enhances or otherwise regulates gene expression [3,151,192-194].

Joshi and Sabol [129] found that rat ppEnk transcription is regulated by the co-operative action of glucocorticoids and cyclic AMP (Table 3). That is, glucocorticoids potentiate ppEnk transcription by directly interacting with *cis*-acting element(s) other than the CREs. Beato [151] provides evidence that steroid-glucocorticoid receptor complexes usually bind to a positive GRE within 3000 bp of a specific gene's transcriptional start site to have its effect (Figure 5). Another report demonstrated that this was also possible within the first intron of the GRE-regulated gene (Figure 5; [195]). In the rat ppEnk's case, both 5' as well as intron A GREs may be present and functional (Figure 5; [107,129]).

CRE function has been demonstrated not only from rat sequences confined between -53 to -103 upstream of the start of ppEnk transcription (see Figure 4; E1), but also a functional GRE was found between -103 to -148 5' of the principal rat start site, E1 [129]. In this case, this 5' GRE had a inhibitory effect, and no sequences within this -103 to -148 bp region seemed to match the consensus GRE sequence (Figure 5). Additionally, as cycloheximide (a protein synthesis inhibitor) did not have an effect on ppEnk induction in this system, this 5' GRE is apparently activated by direct binding of the steroid-receptor complex [129]. There may be different GRE functional *cis*-acting elements for positive and negative action, where necessary. Hence, this 5' ppEnk-negative GRE is perhaps a new sequence that still allows activated-receptor binding. On the other hand, within the rat ppEnk's intron A three putative GRE-related sequences similar to the original GRE (see above) have been identified (Figure 5; [107,129]) as follows: GTTCCTcatTGTCCT (at +196 to +210 w.r.t. the E1 start site, 12/15 match), ATGGCAtccTGTCCT (at +356 to +342, on the minus strand, a 12/15 match) and ATGGCAtccTGTTCT (at +399 to +385, on the minus strand, a 12/15 match).

In these cases, functional studies may show positive GRE activity from one or more of these intron A elements. Moreover, it has been suggested that the ppEnk gene may be regulated by both positive as well as negative GREs, which do not function alone, but within the context of neighbouring *cis*-acting elements [129].

(iv) Lymphocytic induction pathways

ppEnk has been shown to be expressed in both T [91,97,98,186,196], as well as B lymphocytes [94,188]. Upon activation with concanavalin A (ConA) plus phorbol myristate acetate (PMA), T lymphocytes accumulate high levels of enkephalin peptide as well as steady-state ppEnk mRNA (Table 3; [186]). A ubiquitous transcription factor, NF- κ B was found to bind to a new *cis*-acting upstream element 'BETA2' (GGGGACGTCCCC; see Figure 4, Table 3). The induction of NF- κ B, in T lymphocytes, by ConA plus PMA treatment, paralleled the activation of the ppEnk promoter (Table 3; [186]). Point mutation experiments confirmed the functional importance of this ppEnk BETA2 element for lymphocytic ppEnk promoter induction [186]. Two studies [186,197] pointed out another BETA element (BETA1), further upstream of BETA2 (see Figure 4). This BETA1 element was found not to bind the NF- κ B or to be functionally involved in lymphocytic ppEnk induction [186]. Rattner and colleagues [186] conclude that this BETA2 element

should be considered a new *cis*-acting ppEnk-activating element, independent of those previously described.

Another activating *cis*-acting DNA region has been reported upstream of the rat ppEnk promoter, which is active in T lymphocytes [196]. This activating element is located approximately 150 bp upstream of the ppEnk E1 start site (Figure 4) and is in the same region as the 5' negative GRE (discussed above, Figure 5). It functions by allowing the HTLV1 Tax₁ protein to bind to it, thereby transactivating the ppEnk promoter in T cells as well as a glioma cell line [196]. Joshi and Dave [196] advance the notion that this type of transactivation may provide the bi-directional communication between the nervous and immune systems, which has been suspected in HTLV1-related diseases. This bi-directional flow of information between immune cells and the neuroendocrine systems is further supported by new data from Rosen's group [98]. They noted that ppEnk gene expression in macrophages could be induced *in vivo* by lipopolysaccharide administration and further enhanced by the presence of adrenaline. They proposed [98] that this type of ppEnk regulation reflects both ascending and descending routes of intersystemic interactions. They concluded that both of these routes of communication are components of a complete homeostatic circuit of communication [98].

Steroid hormones have also been shown to inhibit enkephalin peptide, prohormone and mRNA biosynthesis in B lymphocytes [188]. This indicates that the 5' negative GRE, discussed earlier, may also be active in B lymphocytes. In this case no DNA binding or functional studies have been reported, thus the identity and position of the B-cell negative GRE remains to be identified. We do know that new protein synthesis is not required for early ppEnk mRNA biosynthesis, but is required to prevent its superinduction [188]. This indicates, that for this cell type there is a transcriptional repressor maintaining 'normal' basal ppEnk mRNA levels. Thus the mechanism of B-cell ppEnk promoter activation could be by down-regulation of this putative transcription repressor factor.

ppEnk gene product biosynthesis in lymphocytes may, as discussed earlier, play a role not only in cell growth and differentiation but also in cell immunity itself. Moreover, the enkephalin peptide that both T and B lymphocytes secrete, could itself be the signal which can potentially allow immune cell modulation of the nervous system during stress and disease.

SUMMARY AND FUTURE DIRECTIONS

In this article, the characterization of ppEnk expression at the DNA, RNA and the genetic control element level has been discussed. These three issues although well studied still have many open questions to investigate. Examples of these outstanding issues include the characterization of unusual steady-state ppEnk RNA transcripts (see Figure 1, lane 4 of [35], for example). Further studies are required to characterize and define the genetic elements that have been described in this review as well as others that may surface in the future. Until now only the CREs, and to a lesser extent the BETA2 elements, have been defined and characterized. For the others (e.g. GRE(s), CARE, upstream enhancer [114], TAX transactivating element and so on) we still do not know not only the DNA sequence binding requirements but also information on what binds to them and how they function. A detailed analysis of 'the natural experiments of nature', that examines species sequence differences at the functional level by transfection and/or *in vitro* transcription experiments is still necessary. Furthermore, we know that there are differences in the expression of ppEnk in particular tissues between different species [e.g. bovine (very high levels) versus rat

(very low levels) basal adrenal medulla]. Whether this is due to differences in the gene's sequence or the protein and cellular background is still very much an open question.

A more direct approach to the study of ppEnk transcriptional regulation, which has been minimally invoked up to now, is the nuclear run-on approach. In this type of experiment we not only assess the ability of any particular mitogen or stressor to affect *de novo* ppEnk RNA synthesis, but also are able to study the transcriptional level invoked and the type of transcriptional products produced. Only a few studies have been published looking at these types of questions [198–200]. These run-on transcription studies have demonstrated that nicotine and histamine induction of ppEnk RNA is an early event in chromaffin cells [198]. The mRNA produced in these experiments was shown to be unaffected by α -amanitin and hence was very stable [198]. The other study indicated that ppEnk RNA induction by cyclic AMP and phorbol esters is independent of protein synthesis in cultured striatal neurons [199,200]. More recently, we have found unusual small sense and antisense ppEnk-related RNA molecules, that may play a role in the tissue- and treatment-specific induction of this gene in rats (G. Weisinger, J. D. DeCristofaro and E. F. LaGamma, unpublished work). In short, it seems that run-on transcription studies will reveal a wealth of information regarding ppEnk transcriptional mechanisms.

The pentapeptide enkephalin molecules have been known for their neurotransmitter function for two decades. As discussed in this review, other roles for ppEnk gene products, ranging from neuromodulation and tissue regeneration to cell proliferation and differentiation, are now being suggested [36,40,42,73,74,78,81,85,91]. Two reports tie ppEnk gene expression to both local and systemic aspects of host's immune response [86,196]. One of these claim that a proenkephalin gene product may be the bi-directional signal between the nervous and immune systems [196]. In any case it is very clear that ppEnk expression is not confined to the nervous system and hence the further definition of alternate roles for ppEnk gene products is an important avenue for future research.

The ppEnk gene and gene products are well conserved from toad to man. The gene products are found throughout the body and can serve numerous functions from neurotransmitter and neuromodulator, to growth and differentiation factors. With such a large variety of functions one would imagine multiple levels of gene regulation at both tissue-specific and function-specific levels. Not only has this been found, but also the regulation of this gene, as complex as it is, should be viewed as a model for complex eukaryotic gene regulation. Some genes may only require a simple developmental on and/or off signal, as is the case for β -globin in red blood cells. This is clearly not the case for ppEnk, the regulation of which must be so finely tuned that it needs not only multiple transcriptional start sites, putative control of RNA elongation attenuation, the possibility of alternate splicing, and RNA stability, but also multiple *cis*-acting elements each responding to a mixture of independent receptor-mediated signal-transduction pathways. As is evident from much early ppEnk work *in vitro*, many treatments can induce the ppEnk promoter in most cells in a Petri dish (see [124–128]). It was thought that, due to the difficulties of working *in vivo*, this approach was the only way to pick apart the gene. Since then, however, many new discoveries have revealed that ppEnk regulation *in vivo* has an even more complex nature. Following both *in vivo* and *in vitro* studies, at least six types of *cis*-acting ppEnk DNA regulatory elements have now been identified (i.e. CRE enhancer, CARE element(s), positive as well as negative GREs, BETA2 element and Tax₁ transactivation element). It seems that this gene, like many others, will turn out to be

controlled by a mosaic of positive and negative control elements. Hence, the DNA context, as well as the activated signal-transduction pathways, will together dictate ppEnk gene output. We have come a long way in understanding the regulation of this opiate gene, but we still have a long way to go.

I would like to thank Professors Edmund F. LaGamma, Haim Rosen, Rabi Simantov and Oren Zinder for their critical review of this manuscript.

REFERENCES

- Johnson, P. F. and McKnight, S. L. (1989) *Annu. Rev. Biochem.* **58**, 799–839
- Jones, N. C., Rigby, P. W. J. and Ziff, E. B. (1988) *Genes Dev.* **2**, 371–378
- Miner, J. N. and Yamamoto, K. R. (1991) *Trends Biol. Sci.* **16**, 423–430
- Mitchell, P. J. and Tjian, R. (1989) *Science* **245**, 271–378
- Morgan, J. I. and Curran, T. (1991) *Annu. Rev. Neurosci.* **14**, 421–451
- Robertson, J. R. (1980) 1699–1976: China, Scribner Press
- Hughes, J., Smith, T. W., Kosterlitz, H. W., Fothergill, L. A., Morgan, B. A. and Morris, H. R. (1975) *Nature (London)* **258**, 577–579
- Simantov, R. and Snyder, S. H. (1976) *Proc. Natl. Acad. Sci. U.S.A.* **73**, 2515–2519
- Litt, M., Buroker, N. E., Kondole, S., Douglass, J., Liston, D., Sheehy, R. and Magenist, R. E. (1988) *Am. J. Hum. Genet.* **42**, 327–334
- Nakanishi, S., Inoue, A., Kita, T., Nakamura, M., Chang, A. C. Y., Cohen, S. N. and Numa, S. (1979) *Nature (London)* **278**, 423–427
- Comb, M., Herbert, E. and Crea, R. (1982) *Proc. Natl. Acad. Sci. U.S.A.* **78**, 360–364
- Comb, M., Seeburg, J., Adelman, J., Eiden, L. and Herbert, E. (1982) *Nature (London)* **295**, 663–666
- Gubler, U., Seeburg, P., Hoffmann, B. J., Gage, L. P. and Udenfriend, S. (1982) *Nature (London)* **295**, 206–208
- Legon, S., Glover, D. M., Hughes, J., Lowry, P. J., Rigby, P. W. J. and Watson, C. J. (1982) *Nucleic Acids Res.* **10**, 7905–7918
- Noda, M., Furutani, Y., Takahashi, H., Toyosato, M., Hirose, T., Inayama, S., Nakanishi, S. and Numa, S. (1982) *Nature (London)* **295**, 202–206
- Noda, M., Teranishi, Y., Takahashi, H., Toyosato, M., Notake, M., Nakanishi, S. and Numa, S. (1982) *Nature* **297**, 431–434
- Kakidani, H., Furutani, Y., Takahashi, H., Noda, M., Morimoto, Y., Hirose, T., Asai, M., Inayama, S., Nakanishi, S. and Numa, S. (1982) *Nature (London)* **298**, 245–249
- Bradbury, A. F., Smyth, D. G., Snell, C. R., Birdsall, N. J. M. and Hulme, E. C. (1976) *Nature (London)* **260**, 793–795
- Goldstein, A., Tachibana, S., Lowney, L. I., Hunkapiller, M. and Hood, L. (1979) *Proc. Natl. Acad. Sci. U.S.A.* **76**, 6666–6670
- Li, C. H. and Chung, D. (1976) *Proc. Natl. Acad. Sci. U.S.A.* **73**, 1145–1148
- Krieger, D. T., Liotta, A. S., Brownstein, M. J. and Zimmerman, E. A. (1980) *Recent Prog. Horm. Res.* **36**, 277–344
- He, X. and Rosenfeld, M. G. (1991) *Neuron* **7**, 183–196
- Laufer, R. and Changeux, J. P. (1989) *Mol. Neurobiol.* **3**, 1–53
- Martens, G. J. M. and Herbert, E. (1984) *Nature (London)* **310**, 251–254
- Rosen, H., Douglass, J. and Herbert, E. (1984) *J. Biol. Chem.* **259**, 14309–14313
- Wong, M., Rius, R. A. and Loh, Y. P. (1991) *Mol. Brain Res.* **11**, 197–205
- Yoshikawa, K., Williams, C. and Sabol, S. L. (1984) *J. Biol. Chem.* **259**, 14301–14308
- Herz, A. (1993) Opioids I, chapt. 14, pp. 308–346, Springer-Verlag, Tokyo
- Herz, A. (1993) Opioids I, chapt. 18, pp. 423–447, Springer-Verlag, Tokyo
- Comb, M., Rosen, H., Seeburg, P., Adelman, J. and Herbert, E. (1983) *DNA* **2**, 213–229
- Zhu, Y. S., Robertson, H. D. and Inturrisi, C. E. (1994) *DNA Cell Biol.* **13**, 25–35
- Weisinger, G., DeCristofaro, J. D. and LaGamma, E. F. (1990) *J. Biol. Chem.* **265**, 17389–17392
- Weisinger, G., DeCristofaro, J. D. and LaGamma, E. F. (1992) *J. Biol. Chem.* **267**, 4508–4512
- Garrett, J. E., Collard, M. W. and Douglass, J. C. (1989) *Mol. Cell. Biol.* **9**, 4381–4389
- Weisinger, G., DeCristofaro, J. D. and LaGamma, E. F. (1992) *Protocols in Molecular Biology (13); Protocols in Molecular Neurobiology* (Longstaff, A. and Revest, P., eds.), ch. 8, pp. 115–138, Humana Press, NJ
- DeCristofaro, J. D., Weisinger, G. and LaGamma, E. F. (1993) *Mol. Brain Res.* **18**, 133–140
- Springhorn, J. P. and Claycomb, W. C. (1992) *Am. J. Physiol.* **263**, H1560–H1566
- Rao, S. M. and Howells, R. D. (1992) *Regul. Peptides* **40**, 397–408
- Garrett, J. E., Garrett, S. H. and Douglass, J. (1990) *Mol. Endocrinol.* **4**, 108–118
- Kilpatrick, D. L., Zinn, S. A., Fitzgerald, M., Higuchi, H., Sabol, S. L. and Meyerhardt, J. (1990) *Mol. Cell. Biol.* **10**, 3717–3726
- Zinn, S. A., Ebert, K. M., Mehta, N. D., Joshi, J. and Kilpatrick, D. L. (1991) *J. Biol. Chem.* **266**, 23850–23855
- Rao, S. M. and Howells, R. D. (1993) *J. Biol. Chem.* **268**, 22164–22169
- Borsook, D., Rosen, H., Collard, M., Dressler, H., Gerrup, K., Comb, M. J. and Hyman, S. E. (1992) *Mol. Endocrinol.* **6**, 1502–1512
- Marcu, K. B., Bossone, S. A. and Patel, A. J. (1992) *Annu. Rev. Biochem.* **61**, 809–860
- Brooks, P. J., Funabashi, T., Kleopoulous, S. P., Mobbs, C. V. and Pfaff, D. (1993) *Mol. Brain Res.* **19**, 22–30
- Monstein, H.-J. and Geijer, T. (1988) *Biosci. Rep.* **8**, 225–261
- Kilpatrick, D. L., Howells, R. D., Noe, M., Bailey, L. C. and Udenfriend, S. (1985) *Proc. Natl. Acad. Sci. U.S.A.* **82**, 7467–7469
- Aloyo, V. J. (1991) *Neuroendocrinology* **54**, 594–598
- Branch, A. D., Unterwald, E. M., Lee, S. E. and Kreek, M. J. (1992) *Mol. Brain Res.* **14**, 231–238
- Chao, H. M. and McEwen, B. S. (1990) *Endocrinology* **126**, 3124–3130
- Franklin, S. O., Yoburn, B. C., Zhu, Y.-S., Branch, A. D., Robertson, H. D. and Inturrisi, C. E. (1991) *Mol. Brain Res.* **10**, 241–250
- Iadarda, M. J., Naranjo, J. R., Duchemin, A. M. and Quach, T. T. (1989) *Peptides* **10**, 687–692
- Zhu, Y.-S., Branch, A. D., Robertson, H. D., Huang, T. H., Franklin, S. O. and Inturrisi, C. E. (1992) *Mol. Brain Res.* **12**, 173–180
- Schwartz, J. P. and Simantov, R. (1988) *Dev. Brain Res.* **40**, 311–314
- Yang, H.-Y. T., Hong, J. S. and Costa, E. (1977) *Neuropharmacology* **16**, 303–307
- Uhl, G. R., Navia, B. and Douglas, J. (1988) *J. Neurosci.* **8**, 4755–4764
- Fischer-Colbrie, R., Diez-Guerra, J., Emson, P. C. and Winkler, H. (1986) *Neuroscience* **18**, 167–174
- Livett, B. G., Day, R., Elde, R. P. and Howe, P. R. C. (1982) *Neuroscience* **7**, 1323–1332
- Wan, D. C.-C., Marley, P. D. and Livett, B. G. (1991) *Mol. Brain Res.* **9**, 135–142
- Wilson, S. P. (1987) *J. Neurochem.* **49**, 1550–1556
- McMillian, M. K., Hudson, P. M., Lee, D. Y., Thai, L., Hung, G. H. and Hong, J. S. (1993) *Dev. Brain Res.* **71**, 75–80
- Yoshikawa, K., Maruyama, K., Aizawa, T. and Yamamoto, A. (1989) *FEBS Lett.* **246**, 193–196
- Pittius, C. W., Kley, N., Loeffler, J. P. and Holtt, V. (1985) *EMBO J.* **4**, 1257–1260
- Matthews, S. G., Heavens, R. P. and Sirinathsinghji, D. J. S. (1992) *Mol. Brain Res.* **12**, 349–355
- Senba, E., Simmons, D. M. and Swanson, L. W. (1990) *Neuroscience* **38**, 629–641
- Spruce, B. A., Curtis, R., Wilkins, G. P. and Glover, D. M. (1990) *EMBO J.* **9**, 1787–1795
- Batter, D. K. and Kessler, J. A. (1991) *Mol. Brain Res.* **11**, 65–69
- Hauser, K. F., Osborne, J. G., Stiene-Martin, A. and Melner, M. H. (1990) *Brain Res.* **522**, 347–353
- Melner, M. H., Low, K. G., Allen, R. G., Nielsen, C. P., Young, S. L. and Saneto, R. P. (1990) *EMBO J.* **9**, 791–796
- Villijn, M. H., Vaysse, P. J., Zukin, R. S. and Kessler, J. A. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 6551–6555
- Shinoda, H., Marini, A. M., Cosi, C. and Schwartz, J. P. (1989) *Science* **245**, 415–417
- Hokfelt, T., Elde, R., Johansson, O., Terenius, L. and Stein, L. (1977) *Neurosci. Lett.* **5**, 25–31
- Sar, M., Stumpf, W. E., Miller, R. J., Chang, K.-J. and Cuatrecasas, P. (1978) *J. Comp. Neurol.* **182**, 17–38
- Rosen, H., Polakiewicz, R. D. and Simantov, R. (1990) *Biochem. Biophys. Res. Commun.* **171**, 722–728
- Kilpatrick, D. L., Borland, K. and Jin, D. F. (1987) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 5695–5699
- Kew, D., Muffly, K. E. and Kilpatrick, D. L. (1990) *Proc. Natl. Acad. Sci. U.S.A.* **87**, 1943–1947
- Rosen, H., Itin, A., Schiff, R. and Keshett, E. (1990) *Mol. Endocrinol.* **4**, 146–154
- Howells, R. D., Kilpatrick, D. L., Bailey, C., Noe, M. and Udenfriend, S. (1986) *Proc. Natl. Acad. Sci. U.S.A.* **83**, 1960–1963
- Springhorn, J. P. and Claycomb, W. C. (1989) *Biochem. J.* **258**, 73–78
- Villiger, R. M. and Lotz, M. (1992) *EMBO J.* **11**, 135–143
- Rosen, H., Polakiewicz, R. D., Benzakine, S. and Bar-Shavit, Z. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **88**, 3705–3709
- Keshet, E., Polakiewicz, R. D., Itin, A., Ornoy, A. and Rosen, H. (1989) *EMBO J.* **8**, 2917–2923
- Kew, D. and Kilpatrick, D. L. (1990) *Mol. Endocrinol.* **4**, 337–340
- Polakiewicz, R. D. and Rosen, H. (1990) *Mol. Cell. Biol.* **10**, 736–742
- Polakiewicz, R. D., Behar, O. Z., Comb, M. J. and Rosen, H. (1992) *Mol. Endocrinol.* **6**, 399–408
- Rosen, H. and Bar-Shavit, Z. (1994) *J. Cell. Biochem.* **55**, 334–339
- Kuis, W., Villiger, P. M., Leser, H.-G. and Lotz, M. (1991) *Am. Soc. Clin. Invest.* **88**, 817–824

- 87 Gustin, T., Bachelot, T., Verna, J.-M. G., Molin, L. F., Brunet, J.-F. M., Berger, F. R. and Benabid, A. L. (1993) *Cancer Res.* **53**, 4715–4719
- 88 Zagon, I. S., Isayama, T. and McLaughlin, P. J. (1994) *Mol. Brain Res.* **21**, 85–98
- 89 Zagon, I. S. and McLaughlin, P. J. (1991) *Brain Res.* **542**, 318–323
- 90 Kwon, B. S. and Weissmann, S. M. (1989) *Proc. Natl. Acad. Sci. U.S.A.* **86**, 1963–1967
- 91 Linner, K. M., Beyer, S. and Sharp, B. M. (1991) *Endocrinology* **128**, 717–724
- 92 Linner, K. M., Nicol, S. E. and Sharp, B. M. (1993) *J. Pharmacol. Exp. Ther.* **267**, 1566–1572
- 93 Padros, M. R., Vindrola, O., Zunszain, P., Fainboin, L., Finkelman, S. and Nahmod, V. E. (1989) *Life Sci.* **45**, 1805–1811
- 94 Rosen, H., Behar, O., Abramsky, O. and Ovadia, H. (1989) *J. Immunol.* **143**, 3703–3707
- 95 Roth, K. A., Lorenz, R. G., Unanue, R. A. and Weaver, C. T. (1989) *FASEB J.* **3**, 2401–2407
- 96 Sibinga, N. E. S. and Goldstein, A. (1988) *Annu. Rev. Immunol.* **6**, 219–249
- 97 Zurawski, G., Benedik, M., Komb, B. J., Abrams, J. S., Zurawski, S. M. and Lee, F. D. (1986) *Science* **232**, 772–775
- 98 Behar, O., Ovadia, H., Polakiewicz, R. D. and Rosen, H. (1994) *Endocrinology* **134**, 475–481
- 99 Rosen, H. and Polakiewicz, R. (1989) *Dev. Brain Res.* **46**, 123–129
- 100 Bayon, A., Shoemaker, W. J., Bloom, F. E., Mauss, A. and Guillemin, R. (1979) *Brain Res.* **179**, 93–101
- 101 Claycomb, W. C. (1975) *J. Biol. Chem.* **250**, 3229–3235
- 102 Claycomb, W. C. (1976) *J. Biol. Chem.* **251**, 6082–6089
- 103 Claycomb, W. C. (1977) *Biochem. J.* **168**, 599–601
- 104 Anderson, D. J. and Axel, R. (1986) *Cell* **47**, 1079–1090
- 105 Carmichael, S. W. (1986) *The Adrenal Medulla*. vol. 4, Cambridge University Press, Cambridge
- 106 Hamill, R. W. and LaGamma, E. F. (1988) *Autonomic Failure* (Bannister, R., ed.), pp. 23–43, Oxford University Press, Oxford
- 107 LaGamma, E. F., White, J. D., McKelvey, J. F. and Black, I. B. (1987) *Ann. N. Y. Acad. Sci.* **493**, 270–272
- 108 LaGamma, E. F., DeCristofaro, J. D., Agarwal, B. L. and Weisinger, G. (1989) *Int. J. Dev. Neurosci.* **7**, 499–511
- 109 Lau, C., Ross, L. L., Whitmore, W. L. and Slotkin, T. A. (1987) *Neuroscience* **22**, 1067–1075
- 110 Lau, C., Franklin, M., McCarthy, L., Pylipiw, A. and Ross, L. L. (1988) *Dev. Brain Res.* **44**, 109–117
- 111 Lillien, L. E. and Claude, P. (1985) *Nature (London)* **317**, 632–634
- 112 Slotkin, T. A. (1985) *Developmental Neurobiology of the Autonomic Nervous System*, pp. 69–133, Humana Press, NJ
- 113 Unger, A. and Phillips, J. H. (1983) *Physiol. Rev.* **63**, 787–843
- 114 LaGamma, E. F., Agarwal, B. L. and DeCristofaro, J. D. (1992) *Mol. Brain Res.* **13**, 187–197
- 115 LaGamma, E. F. and DeCristofaro, J. D. (1989) *Molecular Biology of Stress* (Zinder, O. and Brenitz, S., eds.), pp. 153–165, UCLA Press, Los Angeles, CA
- 116 Kanamatsu, K., Unsworth, C. D., Diliberto, E. J., Jr., Viveros, O. H. and Hong, J. S. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 9245–9249
- 117 DeCristofaro, J. D. and LaGamma, E. F. (1990) *Neuroscience* **35**, 203–210
- 118 Eiden, L. E. (1987) *Cell. Mol. Neurobiol.* **7**, 339–352
- 119 Fischer-Colbrie, R., Iacangelo, A. and Eiden, L. E. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 3240–3244
- 120 Yoburn, B. C., Franklin, S. O., Calvano, S. E. and Inturrisi, C. E. (1987) *Life Sci.* **49**, 2495–2503
- 121 Inturrisi, C. E., Branch, A. D., Robertson, H. D., Howells, R. D., Franklin, S. O., Shapiro, J. R., Calvano, S. E. and Yoburn, B. C. (1988) *Mol. Endocrinol.* **2**, 633–640
- 122 LaGamma, E. F., DeCristofaro, J. D. and Weisinger, G. (1991) *Mol. Cell. Neurosci.* **2**, 517–525
- 123 Stachowiak, M. K., Lee, P. H. K., Rigual, R. J., Viveros, O. H. and Hong, J. S. (1988) *Mol. Brain Res.* **3**, 263–274
- 124 Comb, M., Liston, D., Martin, M., Rosen, H. and Herbert, E. (1985) *EMBO J.* **4**, 3115–3122
- 125 Comb, M., Birnberg, N. C., Seasholtz, A., Herbert, E. and Goodman, H. M. (1986) *Nature (London)* **323**, 353–356
- 126 Comb, M., Mermod, N., Hyman, S. E., Pearlberg, J., Ross, M. E. and Goodman, H. M. (1988) *EMBO J.* **7**, 3793–3805
- 127 Hyman, S. E., Comb, M., Lin, Y.-S., Pearlberg, J., Green, M. R. and Goodman, H. M. (1988) *Mol. Cell. Biol.* **8**, 4225–4233
- 128 Hyman, S. E., Comb, M., Pearlberg, J. and Goodman, H. M. (1989) *Mol. Cell. Biol.* **9**, 321–324
- 129 Joshi, J. and Sabol, S. L. (1991) *Mol. Endocrinol.* **5**, 1069–1080
- 130 LaGamma, E. F., Goldstein, N. K., Snyder, J. B. and Weisinger, G. (1989) *Mol. Brain Res.* **5**, 131–140
- 131 LaGamma, E. F. and Weisinger, G. (1991) *Mol. Cell. Neurosci.* **2**, 427–439
- 132 Sonnenberg, J. L., Rauscher, F. J., Morgan, J. I. and Curran, T. (1989) *Science* **246**, 1622–1625
- 133 Van Nguyen, T., Kobierski, L., Comb, M. and Hyman, S. E. (1990) *J. Neurosci.* **10**, 2825–2833
- 134 Weisinger, G. and LaGamma, E. F. (1992) *Protocols in Molecular Biology* (13); *Protocols in Molecular Neurobiology* (Longstaff, A. and Revest, P., eds.), ch. 9, pp. 139–154, Humana Press, NJ
- 135 Franklin, S. O., Zhu, Y.-S., Yoburn, B. C. and Inturrisi, C. E. (1991) *Mol. Pharmacol.* **40**, 515–522
- 136 Kirshner, N. (1962) *J. Biol. Chem.* **237**, 2311–2317
- 137 Mueller, R. A., Thoenen, H. and Axelrod, J. (1969) *J. Pharmacol. Exp. Ther.* **169**, 74–79
- 138 Sietzen, M., Schober, M., Fischer-Colbrie, R., Scherman, D., Sperk, G. and Winkler, H. (1987) *Neuroscience* **22**, 131–139
- 139 Thoenen, H., Mueller, R. A. and Axelrod, J. (1969) *J. Pharmacol. Exp. Ther.* **169**, 249–254
- 140 Viveros, O. H. and Wilson, S. P. (1983) *J. Auton. Nerv. Sys.* **7**, 41–58
- 141 Eiden, L. E., Giraud, P., Affolter, H.-U., Herbert, E. and Hotchkiss, ?. (1984) *Proc. Natl. Acad. Sci. U.S.A.* **81**, 3949–3953
- 142 Stachowiak, M. K., Hong, J. S. and Viveros, O. H. (1990) *Brain Res.* **510**, 277–288
- 143 Waschek, J. A., Pruss, R. M., Siegel, R. E., Eiden, E. E., Bader, M.-F. and Aunis, D. (1987) *Ann. N. Y. Acad. Sci.* **493**, 308–323
- 144 Biguet, N. F., Buda, M., Lamouroux, A., Samolyk, D. and Mallet, J. (1986) *EMBO J.* **5**, 287–291
- 145 Mocchetti, I., Guidotti, A., Schwartz, J. P. and Costa, E. (1985) *J. Neurosci.* **5**, 3379–3385
- 146 Tank, A. W., Lewis, E. J., Chikaraishi, D. M. and Weiner, N. (1985) *J. Neurochem.* **45**, 1030–1033
- 147 Ahima, R. S., Garcia, M. M. and Harlan, R. E. (1992) *Mol. Brain Res.* **16**, 119–127
- 148 Chao, H. M. and McEwen, B. S. (1991) *Mol. Brain Res.* **9**, 307–311
- 149 Harlan, R. E. (1988) *Mol. Neurobiol.* **2**, 183–200
- 150 Lightman, S. L. and Young, W. S. (1989) *Proc. Natl. Acad. Sci. U.S.A.* **86**, 4306–4310
- 151 Beato, M. (1989) *Cell* **56**, 335–344
- 152 LaGamma, E. F. and Adler, J. E. (1987) *Mol. Brain Res.* **2**, 125–130
- 153 Ahima, R. S. and Harlan, R. E. (1990) *Neuroscience* **39**, 579–604
- 154 Ahima, R. E., Krozowski, Z. and Harlan, R. (1991) *J. Comp. Neurol.* **313**, 522–538
- 155 Harlan, R. E., Shivers, B. D., Romano, G. J., Howells, R. D. and Pfaff, D. (1987) *J. Comp. Neurol.* **258**, 159–184
- 156 Orkin, S. H. (1990) *Cell* **63**, 665–672
- 157 Roeder, R. G. (1991) *Trends Biochem. Sci.* **16**, 402–408
- 158 Rosenfeld, M. G. (1991) *Genes Dev.* **5**, 897–907
- 159 Weintraub, H., Davis, R., Tapscott, S., Thayer, M., Krause, M., Benezra, R., Blackwell, T. K., Turner, D., Rupp, R., Hollenberg, S., Zhuang, Y. and Lassar, A. (1991) *Science* **251**, 761–766
- 160 Chu, H.-M., Fischer, W. H., Osborne, T. F. and Comb, M. J. (1991) *Nucleic Acids Res.* **19**, 2721–2728
- 161 Folkesson, R., Monstein, H.-J., Geiger, T. and Terenius, L. (1989) *Mol. Brain Res.* **5**, 211–217
- 162 Kobierski, L. A., Chu, H.-M., Tan, Y. and Comb, M. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **88**, 10222–10226
- 163 Goodman, R. H. (1990) *Annu. Rev. Neurosci.* **13**, 111–127
- 164 Durkin, R. C., Weisinger, G., Holloway, M. P. and LaGamma, E. F. (1992) *Biochim. Biophys. Acta* **1131**, 349–351
- 165 Baran, N., Lapidot, A. and Manor, H. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **88**, 507–511
- 166 Montminy, M. R. and Bilezikian, L. M. (1987) *Nature (London)* **328**, 175–178
- 167 Borsook, D., Konradi, C., Falkowski, D., Comb, M. and Hyman, S. E. (1994) *Mol. Endocrinol.* **8**, 240–248
- 168 Bacher, B., Wang, X. and Holtt, V. (1994) *Regul. Peptides (Suppl. 1)*, S171
- 169 McMurray, C. T., Wilson, W. D. and Douglass, J. D. (1991) *Proc. Natl. Acad. Sci. U.S.A.* **88**, 666–670
- 170 Comb, M. J. and Goodman, R. H. (1991) *Nucleic Acids Res.* **18**, 3975–3982
- 171 Mermod, N., Williams, T. J. and Tjian, R. (1988) *Nature (London)* **332**, 557–561
- 172 Donovan, D. M., Takemura, M., O'Hara, B. F., Brannock, M. T. and Uhl, G. R. (1992) *Proc. Natl. Acad. Sci. U.S.A.* **89**, 2345–2349
- 173 Takemura, M., Donovan, D. M. and Uhl, G. R. (1992) *Mol. Brain Res.* **13**, 207–212
- 174 LaGamma, E. F., Weisinger, G., Lenn, N. J. and Strecker, R. E. (1993) *Cell Transplantation* **2**, 207–214
- 175 LaGamma, E. F., Weisinger, G., Lenn, N. J. and Strecker, R. E. (1992) *Neurol. Transplant. Plast.* **3**, 244–245

- 176 LaGamma, E. F., Strecker, R. E., Lenn, N. J., DeCristofaro, J. D. and Weisinger, G. (1995) *Exp. Neurol.*, in the press
- 177 Borsook, D., Falkowski, O., Burstein, R., Strassmann, A., Konradi, C., Dauber, A., Comb, M. and Hyman, S. E. (1994) *Mol. Endocrinol.* **8**, 116–125
- 178 Schwartz, J. P., Simantov, R., Ryan, J. P. and Uhl, G. P. (1988) *Progress in Endocrinology* (Imura, H., Shizume, K. and Yoshida, S., eds.), pp. 1125–1129, Elsevier, New York
- 179 Petrack, B., Emmett, M. R., Rao, T. S., Kim, H. S. and Wood, P. L. (1990) *Life Sci.* **46**, 687–691
- 180 Mocchetti, I., Schwartz, J. P. and Costa, E. (1985) *Mol. Pharmacol.* **28**, 86–91
- 181 Morris, B. J., Holt, V. and Herz, A. (1988) *Neuroscience* **25**, 525–532
- 182 Iadarola, M. J., Douglass, J., Civelli, O. and Naramjo, J. P. (1988) *Brain Res.* **455**, 205–212
- 183 Borsook, D., Konradi, C., Falkowski, O., Comb, M. and Hyman, S. E. (1994) *Mol. Endocrinol.* **8**, 240–248
- 184 Konradi, C., Kobilski, L. A., Nguyen, T. V., Heckers, S. and Hyman, S. E. (1993) *Proc. Natl. Acad. Sci. U.S.A.* **90**, 7005–7009
- 185 Black, I. B., Adler, J. E., Dreyfus, C. F., Friedman, W. F., LaGamma, E. F. and Roach, A. H. (1987) *Science* **243**, 1263–1268
- 186 Rattner, A., Korner, M., Rosen, H., Baeuerle, P. A. and Citri, Y. (1991) *Mol. Cell. Biol.* **11**, 1017–1022
- 187 Holloway, M. P. and LaGamma, E. F. (1992) *J. Biol. Chem.* **267**, 19819–19823
- 188 Behar, O. Z., Ovadia, H., Polakiewicz, R. D., Abramsky, O. and Rosen, H. (1991) *Endocrinology* **129**, 649–655
- 189 Gubler, U. (1987) *Molecular Cloning of Hormone Genes* (Habener, J. F., ed.), pp. 229–276, Humana Press, Clifton, NJ
- 190 Yoshikawa, K. and Sabol, S. L. (1986) *Biochem. Biophys. Res. Commun.* **139**, 1–10
- 191 Yoshikawa, K. and Sabol, S. L. (1986) *Mol. Brain Res.* **1**, 75–83
- 192 Karin, M., Haslinger, A., Holtgreve, H., Richards, R. I., Krauter, P., Westphal, H. M. and Beato, M. (1984) *Nature (London)* **308**, 513–519
- 193 Toohey, M. G., Morley, K. L. and Peterson, D. O. (1986) *Mol. Cell. Biol.* **6**, 4526–4538
- 194 Yamamoto, K. R. (1985) *Annu. Rev. Genet.* **19**, 209–252
- 195 Slater, E. P., Rabenau, O., Karin, M., Baxter, J. D. and Beato, M. (1985) *Mol. Cell. Biol.* **5**, 2984–2992
- 196 Joshi, J. and Dave, H. P. G. (1992) *Proc. Natl. Acad. Sci. U.S.A.* **89**, 1006–1010
- 197 Korner, M., Rattner, A., Mauxion, F., Sen, R. and Citri, Y. (1989) *Neuron* **3**, 563–572
- 198 Favin, L. J., Kley, N. and Hoell, V. (1990) *J. Biol. Chem.* **265**, 19116–19121
- 199 Giraud, P., Kowalski, C., Barthel, F., Becquet, D., Renard, M., Grino, M., Boudouresque, F. and Loeffler, J. F. (1991) *Neuroscience* **43**, 67–79
- 200 Giraud, P., Rowalski, C., Barthel, F., Becquet, D., Renard, M., Grino, M., Boudouresque, F. and Loeffler, J. R. (1995) *Neuroscience*, in the press