Rat mammary-gland transferrin: nucleotide sequence, phylogenetic analysis and glycan structure

Héctor ESCRIVÁ,*§ Annick PIERCE,† Bernadette CODDEVILLE,† Fernando GONZÁLEZ,‡ Monique BENAÏSSA,† Didier LÉGER,† Jean-Michel WIERUSZESKI,† Geneviève SPIK† and Mercè PAMBLANCO*||

*Departament de Bioquímica i Biologia Molecular and ‡Departament de Genètica i Servei de Bioinformàtica, Facultat de Ciències Biològiques, Universitat de València, 46100 Burjassot, Spain and †Laboratoire de Chimie Biologique, Unité Mixte de Recherche 111 du Centre National de la Recherche Scientifique, Université des Sciences et Technologies de Lille, 59655 Villeneuve d'Ascq Cedex, France

The complete cDNA for rat mammary-gland transferrin (Tf) has been sequenced and also the native protein isolated from milk in order to analyse the structure of the main glycan variants present. A lactating-rat mammary-gland cDNA library in λ gt10 was screened with a partial cDNA copy of rat liver Tf and subsequently rescreened with 5' fragments of the longest clones. This produced a 2275 bp insert coding for an open reading frame of 695 amino acid residues. This includes a 19-amino acid signal sequence and the mature protein containing 676 amino acids and one N-glycosylation site in the C-terminal domain at residue 490. Phylogenetic analysis was carried out using 14 translated Tf nucleotide sequences, and the derived evolutionary tree shows that at least three gene duplication events have occurred during Tf evolution, one of which generated the N- and C-terminal

domains and occurred before separation of arthropods and chordates. The two halves of human melanotransferrin are more similar to each other than to any other sequence, which contrasts with the pattern shown by the remaining sequences. Native rat milk Tf is separated into four bands on native PAGE that differ only in their sialic acid content: one biantennary glycan is present containing either no sialic acid residues or up to three. The complete structures of the two major variants were determined by methylation, m.s. and 400 MHz ^1H -n.m.r. spectroscopy. They contain either one or two neuraminic acid residues ($\alpha 2 \rightarrow 6$)-linked to galactose in conventional biantennary *N*-acetyl-lactosamine-type glycans. Most contain fucose ($\alpha 1 \rightarrow 6$)-linked to the terminal non-reducing *N*-acetylglucosamine.

INTRODUCTION

Transport of iron between sites of absorption, storage and utilization and its delivery to all cells in the organism was the first function ascribed to transferrin (Tf), but this protein is also essential for the growth and differentiation of a variety of cells and plays a significant role in bacteriostasis (for references, see [1]). It belongs to a family of two-sided iron-binding glycoproteins [2] consisting of a single polypeptide chain with M_r values in the range 76000–81000. It is widely distributed in physiological fluids and cells of invertebrates and vertebrates (for reviews, see [1–4]), and is synthesized mainly in the liver, but lower amounts are also produced in other organs, such as the testes, brain and mammary gland [5].

Tf gene expression is regulated in a tissue-specific fashion by a diversity of factors such as iron, vitamins and hormones (reviewed in [6]). In the mammary gland, the expression of the Tf gene is modulated during the reproductive cycle. In rats, in contrast with mice and rabbits, Tf concentrations in milk and mammary-gland Tf mRNA levels vary biphasically, increasing up to parturition and then decreasing to undetectable levels at mid-lactation before increasing again in late lactation. However, the role of Tf synthesized locally in the mammary gland is unknown. It may be involved in cell growth and differentiation in the mouse mammary gland [7] and may have a function in the involution of the mammary gland because at this period it is a major cytosolic protein of the mammary-gland epithelial cells [8]. Other studies have suggested that production of Tf may be one of the

mechanisms through which oestrogens favour mammary-gland development in rats [9].

Several sequences of Tfs have been resolved by cDNA cloning [10–23], and sequences of partial cDNA clones from rat [24–26], mouse [20] and bovine [27] Tf have also been reported. The internal homology found between the N- and C-terminal halves of vertebrate Tfs supports the notion that these proteins probably evolved by intragenic duplication from one ancestor, and analysis of the genomic organization of human Tf and chicken egg white Tf (ovotransferrin) indicates that this common ancestor itself derived from a primordial gene by internal duplication [28]. Few studies have been designed to gain a better understanding of the evolution of the Tf gene family [2,4,16,29].

Tfs have variable carbohydrate content (0–2 glycans per molecule) and glycan structures (bi-, tri-, tetra- or penta-antennary glycans with or without one fucose residue), although the carbohydrate moiety is always of the N-linked asparagine type. A comparative study of the glycan primary structures of serum and egg white Tfs and lactotransferrins from several species led to the conclusion that Tf glycans are specific for each Tf and, for a given Tf, specific to the species (reviewed in [30]).

In the present work we describe the determination of the complete rat Tf sequence, which led to a comparison of this sequence with all known complete Tf sequences, allowing us to construct phylogenetic trees of this protein family. This paper also deals with the purification of rat Tf isolated from milk as well as the structural analysis of the glycan moiety of the two major glycovariants. We compare their structures with those

Abbreviation used: Tf, transferrin.

§ Present address: Centre d'Immunologie et de Biologie Parasitaire, U167 de l'INSERM, Institut Pasteur, Lille, France.

|| To whom correspondence should be addressed.

The novel nucleotide sequence data published here will appear in the EMBL/GenBank/DBJ Nucleotide Sequence Databases under the accession number X77158.

already described for rat serum Tf [31] in order to relate Tf glycan structure to its site of biosynthesis.

EXPERIMENTAL

RNA extraction and preparation

Lactating mammary gland was removed and after being washed with cold 0.9% NaCl, frozen by clamping in liquid nitrogen, ground in a mortar and kept at -80°C until analysis. Total cellular RNA was extracted from frozen tissues essentially as described [32]. Poly(A)-rich RNA was prepared by oligo(dT)-cellulose affinity chromatography.

Construction of a cDNA library

A λ gt10 cDNA library was constructed from lactating-rat mammary-gland poly(A)-rich RNA using a commercial kit (Pharmacia) and following the procedure of the manufacturer. Duplicate Hybond N membranes (Amersham, Bucks., U.K.) were probed with a 688 bp rat serum Tf cDNA [26] (kindly provided by Dr. Stallard, University of Washington State at Pullman, WA, U.S.A.) ^{32}P -labelled using a commercial random-priming kit (Boehringer, Mannheim, Germany). Extensive washes were carried out and the dried membranes were exposed overnight. Positive clones were rescreened until considered pure. The rat mammary-gland cDNA library was also screened against a lactotransferrin probe (a gift from Dr. Rado [33]).

Sequence analysis

cDNA inserts isolated from clones as described [34] and after digestion of λ DNA with *NotI* or *EcoRI* and GeneClean II (Bio 101 Inc., La Jolla, CA, U.S.A.) purification (according to the manufacturer's protocol) were subcloned into pBS(SK) for further analysis. Plasmid DNA was prepared as described [34]. Nucleotide sequences of the appropriate fragments subcloned into M13 phage vectors were determined by the dideoxychain-termination method using modified T7 DNA polymerase (Sequenase), in accordance with the supplier's instructions, and labelling with $[\alpha\text{-}^{35}\text{S}]\text{thio}[\text{dCTP}]$. The orientation of the cDNA cloned in M13 vectors was tested as described previously [35]. Sequences of both strands were obtained for the sequence presented in this paper.

PCR analysis

Primers used to amplify Tf cDNA were d(TGYCTDGCDDGT-NCCDGAYAA) (Y = C or T; D = A, G or T; N = A, C, G or T) based on the N-terminal polypeptide sequence [25] and the nucleotide sequence d(GGATAATCCAGCCTGCAGAC) corresponding to the 5' region of our clone. The reagents used for PCR were from Promega (Madison, WI, U.S.A.). Amplification reaction mixtures consisted of 100 pmol of primers, $1 \times$ PCR buffer (10 mM Tris/HCl, pH 8.3, 50 mM KCl and 1.5 mM MgCl_2), 0.25 mM dNTPs (final concentration) and 1 ng of the DNA amplified from the library. *Taq* DNA polymerase (2.5 units) was added to a final volume of 100 μl , and 100 μl of mineral oil was overlayed. All samples were heated at 95°C for 4 min. Cycling parameters for PCR were denaturation at 92°C for 1 min with annealing at 60°C for 1 min and extension at 72°C for 1 min. A total of 40 cycles was performed for all PCR reactions on a programmed Intelligent Heating Block IHB 2024 (Beckman).

Phylogenetic analysis of the Tf family

Complete nucleotide sequences of Tfs were obtained from the GenBank Database (release 76.0) and are listed in Table 1. Only the nucleotides coding for the mature peptide were considered in the analysis. Translated sequences were aligned by means of the hierarchical clustering algorithm as implemented in the CLUSTAL V program [36]. This alignment strategy consists of the progressive alignment of groups of sequences according to the branching order in a hypothetical phylogenetic tree. Pairwise alignments are performed using the dynamic programming method of Needleman and Wunsch [36], which guarantees an optimal alignment for two sequences. Sets of sequences are then aligned using the average score at each position [36]. Two different alignments were used for the phylogenetic derivations in the present study. First, all the sequences were aligned as described previously. Second, given that the N- and C-termini of Tfs have arisen by duplication of an ancestral sequence, and that substantial identity is still present between both halves, each sequence was divided into its N- and C-termini (see Table 1 for the exact position used in this division for each sequence) and the resulting 28 sequences were aligned using the same procedure as above. Both amino acid alignments were 'back-translated' into the corresponding nucleotide alignments, which were then used for phylogenetic analysis. Distances between each pair of

Table 1 Complete nucleotide sequences of Tfs used in the phylogenetic analysis

C-term, first nucleotide of the C-terminal half considered in the alignments.

Protein	Organism	Locus	Accession no.	Begin	End	C-term	Reference
Tf	Cockroach	BLBTRANS	L05340	58	2187	1114	10
	<i>Manduca sexta</i>	MOTTRNFE	M62802	79	2067	1120	11
	<i>Xenopus laevis</i>	XLTRSFER	X54530	93	2183	1077	12
	Hen	GGCONR	X02009	134	2191	1154	13
	Rabbit	OCTRNFNM	X58533	79	2106	1090	14
	Pig	SSTF	X12386	3	2090	1026	15
	Horse	HRSTFRA	M69020	82	2139	1102	16
	Rat	RATTF	X77158	75	2102	1074	This report
	Human	HUMTF	M12530	88	2124	1093	17
	Pig	PIGPLF	M92089	337	2337	1345	18
LactoTf	Cow	BTACTRA	X57084	90	2156	1107	19
	Mouse	MUSULT	J03298	61	2124	1075	20
	Human	HSLTFR	X52941	52	2127	1072	21
MelanoTf	Human	HUMP971	M12154	118	2274	1141	22

sequences were determined for both nucleotide alignments using the MEGA program [37]. Among the different nucleotide distances available, the Tajima–Nei distance was chosen for further analysis according to the recommendations of Nei [38]; as most pairwise distances between sequences lie in the range 0.3–1.0 substitutions per site, there is no strong transition–transversion bias and the sequences differ significantly from the equidistribution of the four nucleotides. Several methods [distance methods with UPGMA (unweighted pair group method using arithmetic averages) and the neighbour-joining algorithm for clustering and the maximum parsimony method] were used to reconstruct the phylogenetic trees of both alignments of the sequences. Parsimony and distance methods for the reconstruction of the trees were completed with bootstrap analysis [39] to set confidence limits on the branching points. These methods were employed as implemented in the program MEGA [37] and several programs in the PHYLIP package [40]. The matrices obtained with the Tajima–Nei distance measure were used with the neighbour-joining method [41]. The number of synonymous and non-synonymous substitutions between pairs of sequences was estimated using the method proposed by Nei and Gojobori as implemented in the program MEGA.

Milk sample collection

Milk samples were obtained from 15 adult Wistar rats on different days of lactation after anaesthetization with sodium pentobarbital (50 mg/kg) and intraperitoneal injection of oxytocin (2 international units) to stimulate milk flow. The samples collected were stored at -80°C until use.

Fractionation of milk and separation of the Tf glycovariants

Thawed milk samples were pooled, diluted 1:1 with distilled water and delipidated by centrifugation at 34000 *g* for 30 min at 4°C . Whey obtained by centrifugation at 34000 *g* for 60 min at 30°C and at pH 4.6 was dialysed against water and lyophilized. It was dissolved in 0.22 M sodium acetate at pH 7.0, fully iron-saturated by adding a solution of FeCl_3 in 0.1 M trisodium citrate/ NaHCO_3 , pH 8.2 (1.5 mg of Fe/mg of protein), and loaded on to a column (1.8 cm $\times$ 18 cm) of SP-Sephadex C-50 (Pharmacia, Uppsala, Sweden). Elution was performed with the same solution at a flow rate of 26 ml/h. Fractions absorbing at 465 nm were pooled, dialysed against water and lyophilized. This pool of fractions was fully iron-saturated and filtered on a 0.22 μm -pore-size Millipore filter before being applied in 50 mM Tris/HCl, pH 8.6, to a Mono Q (HR 5/5) column QAE (Pharmacia FPLC system) [42] and eluted with a linear gradient of 0–1 M NaCl in the same buffer at a flow rate of 1 ml/min. Each fraction collected from several identical runs was pooled, concentrated to 2 ml first on an Immersible CX-10 filter (Millipore Corp., Bedford, MA, U.S.A.) and then to 0.5 ml on a Centricon 30 filter (Amicon, Pulli, Switzerland) and finally desalted on a Pharmacia Phast Desalting column (Sephadex G-25) using the f.p.l.c. system.

Characterization of the Tf glycovariants

Lyophilized rat milk Tf and separated Tf glycovariants were identified by Western blotting from a native 10–15% gradient polyacrylamide gel (Pharmacia Phast System Separation). Immunovisualization was with a rabbit antibody directed against rat serum Tf (dilution 1:2000) (Flobio SA, Courbevoie, France) and horseradish peroxidase-conjugated goat anti-rabbit IgG (dilution 1:1000) (Diagnostic Pasteur, Marnes La Coquette, France). The peroxidase activity bound to nitrocellulose was

developed in 0.04% diaminobenzidine in Tris-buffered saline containing 0.01 M H_2O_2 . Rat milk Tf was desialylated on a *Clostridium perfringens* type III-A-immobilized neuraminidase column (Sigma, St. Louis, MO, U.S.A.) at 37°C by recycling over 24 h [43].

G.l.c.

Oligosaccharide alditols from the glycovariants of rat milk Tf were prepared by hydrazinolysis, N-reacetylation and reduction [44]. The resulting mixture was desalted on a Bio-Gel P-2 column (0.2 cm $\times$ 40 cm) eluted with distilled water. Oligosaccharide alditols were monitored by u.v. absorbance at 206 nm. The molar compositions of the monosaccharides were determined after methanolysis [44] and g.l.c. of the trimethylsilylated methyl glycosides on a capillary CP sil SCB column (0.25 mm $\times$ 25 m) [44].

Identification of m.s. products

The oligosaccharide alditols were permethylated by the method of Hakomori modified by Paz-Parente et al. [44] using the lithium methanesulphinyll carbanion reagent. The methylated and acetylated methyl glycosides were identified by g.l.c./m.s. [44] with a Ribermag R 10-10 mass spectrometer (Riber, Rueil-Malmaison, France) coupled to the data system Sydar 121.

N.m.r. analysis

The oligosaccharide alditols were repeatedly dissolved in $^2\text{H}_2\text{O}$ at room temperature and at pD 7 with intermediate freeze-drying [45]. The deuterium-exchanged oligosaccharide alditols were submitted to ^1H -n.m.r. spectroscopy performed at 400 MHz on a Bruker AM-400 WB spectrometer operating in the pulsed Fourier-Transform mode and equipped with a Bruker Aspect 3000 computer, at a probe temperature of 27°C (Centre Commun de Mesures, Université de Lille Flandres-Artois, France). Chemical shifts (δ) were expressed as p.p.m. downfield from the sodium 4,4-dimethyl-4-silapentane-1-sulphonate, but were actually measured by reference to internal acetone ($\delta = 2.225$ p.p.m.) with an accuracy of 0.002 p.p.m.

RESULTS

Rat Tf cDNA screening and sequencing

From 10^5 colonies of the initial mammary-gland cDNA library (1.5×10^7 independent clones), screened with rat liver Tf cDNA

Figure 1 Restriction sites used for sequencing the cDNA of Tf isolated from rat mammary gland

The non-coding region is indicated by stippling inside the cDNA. The arrows indicate the direction and length of the sequence determinations of each subcloned DNA fragment. Fragments A [26] and B are the probes used to screen the initial library. Fragment C is the probe used to screen the amplified library assayed by PCR. E, *EcoRI*; St, *StuI*; Ps, *PstI*; H, *HaeIII*; P, *PvuII*; B, *BamHI*; S, *SmaI*; A, *AclI*.

1	CCACACACACCGAGAGGATGAGGTTGCTGTGGGTGCCCTGCTGGCTTGTGCCGCCCTGGGACTGTGT	
	-1 +1 M R F A V G A L L A C A A L G L C	17
69	CTGGCTGTCCTGACAAAACGGTCAAATGGTGGCGAGTGTCTGAGCATGAGAACACCAAGTGATCAGT	
	L A V P D K T V K W C A V S E H E N T K C I S	40
138	TTCCGTGACACATGAAAACCGTCTTCAGCTGATGGCCCCCGCTTGCCCTGTGTGAAGAAAACCTCC	
	F R D H M K T V L P A D G P R L P C V K K T S	63
207	TATCAAGATTGCATCAAGGCCATTCTGGAGGTGAAGCTGATGCCATTACCTTGGATGGGGGTGGGTG	
	Y Q D C I K A I S G G E A D A I T L D G G W V	86
276	TACGATGCAGGCCTGACTCCCAACAACTGAAGCCTGTGGCAGCAGAGTTTATGGATCACTTGAACAT	
	Y D A G L T P N N L K P V A A E F Y G S L E H	109
345	CGACAGCCCACTACTTGGCTGTGGCCGTGGTGAAGAGGGAACAGACTTCCAGCTGAACCCAGTCCAG	
	R Q T H Y L A V A V V K K G T D F Q L N Q L Q	132
414	GGCAAGAAGTCTGCCACACTGGCCCTGGGCAAGTGTGCAGGCTGGATTATCCCCATTGGCTTACTTTTC	
	G K K S C H T G L G R S A G W I I P I G L C F	155
483	TGTAACCTGGCAGAGCCCCGCAAGCCTCTTGAGAAAGCTGTGGCCAGTTTCTTCTCGGGCAGTTGTGTC	
	C N L P E P R K P L E K A V A S F F S G S C V	178
552	CCCTGTGCAGATCCAGTGGCCTTCCCCAGCTGTGTCAACTGTGTCCAGGCTGTGGCTGTCTCCCGACT	
	P C A D P V A F P Q L C P G C G C S P T	201
621	CAACCGTTCTTGGCTACGTAGGCGCCTTCAAGTGTCTGAGAGATGGAGGTGGAGATGTGGCCTTTGTC	
	Q P F F G Y V G A F K C L R D G G G D V A F V	224
690	AAGCATACAACCATATTTGAGGTCTTGCCACAGAAGCTGCAGGGATCAATATGAGCTGTCTGCTT	
	K H T T I F E V L P Q K A D R D Q Y E L L C L	247
759	GACAATACCCGCAAGCCAGTGGATCAGTATGAGGACTGCTACCTAGCCCGGATCCCTTCTCATGTGTT	
	D N T R K P V D Q Y E D C Y L A R I P S H A V	270
828	GTGGCTCGAAATGGAGATGGCAAAGAGGACTTGATCTGGGAGATCCTCAAAGTGGCTCAGGAACACTTT	
	V A R N G D G K E D L I W E I L K V A Q E H F	293
897	GGCAAAGGCAATCAAAGACTTCCAACCTGTTCGGCTCTCCTCTTGGGAAAGACCTGCTGTTTAAGGAT	
	G K G K S K D F Q L F G S P L G K D L L F K D	316
966	TCTCGCTTGGGCTGTTACGTGCCCCCAAGGATGAGCTACAGGCTGTACCTCGGCCACAGCTATGTCAC	
	S R F G L L R A P K D G L Q A V P R P Q L C H	339
1035	TGCCATTCGAAATCAGCGGAAGCTGTCCGGATGCCATCGACAGCGCGCCAGTGAAATGGTGTGCACATG	
	C H S K S A G S C P D A I D S A P V K W C A L	362
1104	AGTCAACAAGAGAGAGCCAAGTGTGATGAGTGGAGCTCACAGGCAATGGGCAGATAGAGTGTGATCA	
	S H Q E R A K C D E W S V T G N G Q I E C E S	385
1173	GCAGAGAGCACTGAGGACTGCATTGACAAGATTGTGTAATGGAGAAGCAGATGCCATGAGCTTGGATGGA	
	A E S T E D C I D K I V N G E A D A M S L D G	408
1242	GGTCATGCCTACATAGCAGGCCAGTGTGGACTGTGCCCGTCATGGCAGAGAACTATGATATCTCTTCG	
	G H A Y I A G Q C G L V P V M A E N Y D I S S	431
1311	TGTACAAACCCACAATCAGATGTCTTTCTTAAAGGGTATTATGCCGTGGCTGTGGTGAAGGCATCAGAC	
	C T N P Q S D V F P K G Y Y A V A V V K A S D	454
1380	TCCAGCATCAACTGGAACAACCTGAAAGGCAAGATCCTGCCATACTGGAGTAGACAGAACCAGCCGGC	
	S S I N W N N L K G K K S C H T G V D R T A G	477
1449	TGGAACATCCCTATGGCCTGTGTTCAGCAGGATCAACCACTGCAAGTTCGATGAATTTTTCAGTCAA	
	W N I P M G L L F S R I N H C K F D E F F S Q	500
1518	GGCTGTGCTCCTGGCTATAAGAAGAAATCCACCCTCTGTGACCTGTGTATTGGCCCAGCAAAATGTGCT	
	G C A P G Y K K N* S T L C D L C I G P A K C A	523
1587	CCGAACAACAGAGAGGGATATAATGGTTATACAGGGCTTTCCAGTGCCTCGTTGAGAAGGGAGACGTA	
	P N N R E G Y N G Y T G A F Q C L V E K G D V	546
1656	GCCTTTGTGAAGCACCAGACTGTCTGGAAAACACGAACGAAAGAACTGCTGCATGGGCTAAGGAT	
	A F V K H Q T V L E N T N G K N T A A W A K D	569
1725	CTGAAGCAGGAAGACTTCCAGCTGTGTGCCPDGTGTTACCAAGAAGCCTGTAACCGAGTTCCGCCACC	
	L K Q E D F Q L L C P D G T K K P V T E F A T	592
1794	TGCCACCTGGCCCAAGCTCCAACCATGTTGTGGTCTCACGAAAAGAGAAGGCAGCCCGGTTAGCACT	
	C H L A Q A P N H V V V S R K E K A A R V S T	615
1863	GTGCTGACTGCCCAGAAGGATTTATTTTGGAAAGGTGACAAGGACTGCACTGGCAATTTCTGTTTGTTC	
	V L T A Q K D L F W K G D K D C T G N F C L F	638
1932	CGGTCTTCCACCAAGGACCTTCTGTTTCAGAGATGACACCAAGTGTTTGACTAACTTCCAGAAGGTACC	
	R S S T K D L L F R D D T K C L T K L P E G T	661
2001	ACATATGAAGAGTACTTAGGAGCAGAGTACTTGAAGCTGTTGGAACATAAGGAAGTGTTCACACTCA	
	T Y E E Y L G A E Y L Q A V G N I R K C S T S	684
2070	CGACTCTAGAAGCCTGCACTTTCCACAAAAGTTAAATCCAAGAGGTGGGTGCCACTGTGGTGGAGGA	
	R L L E A C T F H K S	
2139	GGATGCCCCCGTGATCCATGGGCTTCTCCTGGCCTCCATGCCCTGAGCGGCTGGGGCTAACTGTGTCCG	
2208	TCTTCACTGCTGTGTGTACACATACACAGAGCACAAAATAAATAAATAAATAAATAAATAAATAAATAA	

Figure 2 Nucleotide sequence of rat Tf cDNA and its deduced amino acid sequence

The nucleotide and amino acid numbers are indicated to the left and right of the sequence respectively. The probable signal peptide, the consensus N-linked glycosylation site and the possible polyadenylation signals are underlined.

[26], 250 clones were determined to be true positives, confirming high representation of the milk Tf messenger. This rat mammary-gland cDNA library was also screened against a lactotransferrin

probe but no positive clones were found, indicating that lactotransferrin is either not expressed by rat mammary gland or is expressed at an undetectable level.

Figure 3 Phylogenetic tree of the Tf family produced using whole sequences (a) and both halves of each sequence (b)

Sequence names are given in Table 1. The trees were constructed by the neighbour-joining method on the matrix of Tajima–Nei distances between the complete nucleotide alignment of whole sequences (not shown). The number at each node represents the percentage of its appearance in 2000 replicates of the bootstrap test performed. A branching point is considered to be highly significant when it appears in at least 95% of the replicates. In (b) alignment considered the N- and C-termini of each sequence as separate operational taxonomic units. This alignment is available on request.

*Eco*RI endonuclease hydrolysis of the longest (1838 bp) Tf cDNA insert produced two fragments of 1105 and 733 bp. The nucleotide sequence of the two restriction fragments was determined. By using the 5' 1105 bp *Eco*–*Eco* fragment (B) for screening, we obtained an insert of 2248 bp containing the coding sequence of the mature protein but lacking some of the signal peptide amino acids. Repeated attempts to obtain clones covering the complete 5' end of the Tf cDNA from the library were unsuccessful. Therefore we decided to use PCR to amplify cDNA segments corresponding to the 5' region of the gene. We obtained a PCR-amplified product of approx. 500 bp. This size, in addition to the 1838 bp of the previously sequenced clone, corresponded exactly to the complete size of other previously described Tf cDNAs. Screening of the amplified library (4×10^4) with fragment C (the 5' 258 bp *Eco*–*Sma*I fragment obtained from the 2248 bp insert) as a probe produced eight independent clones. One of the isolated inserts was 2275 bp long. In Figure 1 all the restriction sites used in the sequencing of the complete rat Tf cDNA are shown. This cDNA (Figure 2) has an overall open reading frame of 695 amino acid residues. By comparing our N-terminal amino acid sequence with the partial one previously described [25], the first amino acid (valine) can be located at position 20. The putative upstream signal sequence containing the ATG start codon is 19 amino acids long. These results

established that the mature rat Tf protein is composed of 676 amino acids and contains only one potential glycosylation site according to the presence of the tripeptide code sequence Asn-Xaa-Thr/Ser for N-glycosidically linked glycans. This site is located in the C-terminal domain at residue 490. Taking into account the amino acid and carbohydrate composition of the glycan moiety, the calculated M_r of rat monosialylated Tf variant from mammary tissue is 75928.

Evolutionary analysis of the Tfs

Translated sequences of Tfs obtained from the GenBank database were aligned as described in the Experimental section. As a result of the alignment (results not shown but available on request), all the relevant positions for the phylogenetic analyses are coincident with previously published alignments [29], despite the inclusion of new sequences and the different alignment method employed. The tree obtained using the complete sequences is shown in Figure 3(a). This tree was obtained by the neighbour-joining method applied to the Tajima–Nei distance [41]. The same tree was obtained using other methods of reconstruction, although the confidence interval limits varied slightly between the different methods, as expected given the intrinsic randomness of the

Table 2 Average number of estimated synonymous and non-synonymous substitutions from pairwise comparisons of the two halves of each sequence

Comparison	Synonymous	Non-synonymous
Global average	0.673	0.406
Between N-termini	0.624	0.369
Between C-termini	0.633	0.340
Between N- and C-termini	0.714	0.455
Between N- and C-termini (intraspecific)	0.695	0.443

bootstrap method [40]. The first notable feature is that, in all the trees, insect Tfs are clear outgroups for the remaining sequences. This tree also shows that at least three duplication events have occurred during the evolution of Tfs. An initial duplication occurred before the separation of arthropods and chordates, as both insect Tfs have duplicated N- and C-termini. A second duplication occurred in the branch leading to vertebrates before the emergence of land animals. This duplication gave rise to human melanotransferrin, which can be observed to be more ancient than any other vertebrate Tf. The third duplication took place before the appearance of mammals, and lactotransferrins are the resulting products.

The phylogenetic tree shown in Figure 3(b), derived using the two halves of all the sequences, also shows the general pattern described above with the presence of three gene duplications. The tree derived from whole sequences (Figure 3a) is perfectly paralleled by that derived from the two protein ends (Figure 3b), with one notable exception. The two halves of human melanotransferrin are closer to each other than to any other sequence, which contrasts with the pattern shown by the remaining sequences. This fact is also supported by a comparison of the intraspecific rate of divergence between the two halves of each sequence (results not shown).

Further analyses have been performed to determine whether the higher homology previously described between N- and C-termini of Tfs is mainly due to selective pressure for identical mutations to occur in both halves of the molecule [29] or to other processes that could homogenize the sequence in both halves (intragenic recombination, for instance). The proportion of synonymous and non-synonymous substitutions for the 28 half-sequences has been computed. The estimated number of synonymous substitutions is very close to the saturation level of 0.75 for most pairwise comparisons. In contrast, the estimated proportion of non-synonymous changes is lower, without noticeable differences between N- and C-termini (Table 2). Both synonymous (0.605) and non-synonymous (0.350) substitution rates between the two halves of melanotransferrin are significantly lower than the corresponding rates in the other sequences (0.695 and 0.443 on average respectively).

Characterization of rat milk Tf

Rat milk Tf, which was resolved into four bands by native PAGE (Figure 4, lane 1), migrated on SDS/PAGE with an apparent M_r of about 80000 and was identified as rat Tf by Western blotting and antibody detection (results not shown). As Tf was resolved as a major band after neuraminidase treatment of the milk Tf fraction (Figure 4, lane 2), differences between lanes 2 and 1 suggested that the four bands differ at least in the number of sialic acid residues. Comparison of these Tf glycovariants with those obtained from rat serum Tf (Figure 4, lane 3), which contain one trisialylated biantennary glycan [31], suggests the

Figure 4 Native PAGE of rat milk Tf fractions using Phast Gel (gradient 10–15%) and Coomassie Blue staining

Lane 1, rat milk Tf fraction eluted from the SP-Sephadex column; lane 2, neuraminidase-treated rat milk Tf eluted from the SP-Sephadex column; lane 3, rat serum Tf glycovariant rTF-1 [31].

Table 3 Molar carbohydrate compositions of rmTf glycovariants and oligosaccharide alditols released by reductive alkaline cleavage of the glycan–protein linkage

The molar ratios were calculated on the basis of three mannose residues.

Compound	Molar carbohydrate composition					
	NeuAc	Fuc	Gal	Man	GlcNAc	GlcNAc-ol
rmTf-1 protein	0.1	0.8	1.9	3.0	3.6	0.0
rmTf-2 protein	0.7	0.8	1.6	3.0	3.5	0.0
rmTf-2 oligosaccharide alditol	0.6	0.8	2.1	3.0	3.3	0.6
rmTf-3 protein	1.9	0.6	1.7	3.0	3.6	0.0
rmTf-3 oligosaccharide alditol	1.7	0.7	2.2	3.0	3.4	0.4

Table 4 Molar ratios of the monosaccharide methyl ethers present in the methanolysate of the permethylated oligosaccharide alditols released by hydrazinolysis from rmTf glycovariants

Molar ratios were calculated on the basis of one residue of 2,4-Me₂-Man per mol of oligosaccharide alditol.

Monosaccharide methyl ether	Molar ratio	
	rmTf-2	rmTf-3
2,3,4-Me ₃ -Fuc	0.5	0.6
2,3,4,6-Me ₄ -Gal	0.8	0.0
2,3,4-Me ₃ -Gal	0.9	1.9
3,4,6-Me ₃ -Man	1.8	1.9
2,4-Me ₂ -Man	1.0	1.0
3,6-Me ₂ -GlcNAcMe	2.4	2.6
1,3,5-Me ₃ -GlcNAcMe-ol	0.4	0.5
4,7,8,9-Me ₄ -NeuAcMe	0.6	1.8

Table 5 ^1H -n.m.r. chemical shifts in structural reporter-group protons of the constituent monosaccharides of the two rmTf glycovariants

Chemical shifts are in p.p.m. relative to internal acetone at $\delta = 2.225$ at 25 °C. Compounds are represented by short-hand symbolic notations as follows: ■, Gal; ●, GlcNAc; ◆, Man; ○, NeuAc; □, Fuc. For the numbering system of the residues, see the text. n.d., Not determined.

Reporter group	Residue	Chemical shift (p.p.m.)	
		rmTf-2	rmTf-3
H-1	GlcNAc-2	n.d.	n.d.
	Man-3	n.d.	n.d.
	Man-4	5.139	5.136
	Man-4'	4.928	4.943
	GlcNAc-5	4.608	4.606
	GlcNAc-5'	4.585	4.606
	Gal-6	4.449	4.445
	Gal-6'	4.472	4.445
	α -D-Fuc(1→6)	4.903	4.901
H-2	Man-3	4.257	4.250
	Man-4	4.198	4.197
	Man-4'	4.114	4.117
H-3 _{ax}	α -D-NeuAc(2→6)	1.717	1.718
H-3 _{eq}	α -D-NeuAc(2→6)	2.673	2.674
CH ₃	α -D-Fuc(1→6)	1.225	1.224
N-Ac	GlcNAc-1-ol	2.056	2.057
	GlcNAc-2(-/+Fuc)	2.080/2.093	2.092
	GlcNAc-5	2.072	2.070
	GlcNAc-5'	2.048	2.066
	α -D-NeuAc(2→6)	2.031	2.031

Figure 5 Structure of the rmTf glycovariants rmTf-2 and rmTf-3

presence in rat milk Tf of one glycan containing none, one, two or three neuraminic acid residues. The mono- and bi-sialylated glycan forms were in similar proportions and were the most abundant of the four glycovariants detected. The glycovariants were separated into three fractions from the Mono Q column when elution was carried out with an NaCl concentration gradient from 0 to 50 mM. From 1 ml of milk, 50 µg of rat milk (rm)Tf-1, 900 µg of rmTf-2 and 430 µg of rmTf-3 were obtained. The rmTf-4 glycovariant was present in amounts too small to be isolated.

Structures of rat Tf glycan

The carbohydrate contents of the rat milk Tf glycovariants rmTf-1, rmTf-2 and rmTf-3 were determined to be 2.5, 2.6 and 2.8 % respectively. The molar carbohydrate content of the glycovariants given in Table 3 shows that the proportions of sialic acid, galactose, mannose and fucose of the oligosaccharide alditols of two of the variants (rmTf-2 and rmTf-3) are similar to those found in native protein by methanolysis and g.l.c., indicating that no degradation of the monosaccharides occurred during the alkaline treatment. The major difference between the three purified glycovariants is the number of neuraminic acid residues, in agreement with the results obtained by native PAGE.

Results shown in Table 4 indicate that the glycan in the two oligosaccharide alditols possesses a common trimannosyl-*N,N'*-diacetylchitobiose core and the results are consistent with the presence of a biantennary structure of the *N*-acetyl-lactosamine type that is fucosylated.

The complete oligosaccharide structure was determined only on rmTf-2 and rmTf-3 which were purified to homogeneity and in sufficient amounts for ¹H-n.m.r. spectroscopy. The typical spectral features match the structural-reporter groups of the classical structure reported for several glycoproteins [45]. On the basis of the n.m.r. data compiled in Table 5 we can deduce the structures shown in Figure 5 for the rmTf-2 and rmTf-3 fractions.

DISCUSSION

In the present study, we describe the first isolation and characterization of a cDNA clone encoding the entire rat Tf and the structural analysis of the variant forms of the glycans of rat Tf isolated from milk.

Tf is a major milk whey protein in some species (rabbit and rat) but in man it is virtually undetectable in milk; in contrast, lactotransferrin is present at a high concentration in human milk but is undetectable in rat milk. Mouse milk, however, contains a 3:1 mixture of Tf and lactotransferrin [31]. The biological significance of the high ratio of Tf to lactotransferrin in mouse milk is not well understood, and neither is the differential expression of Tf and lactotransferrin genes in different species. It would be of interest to know whether the lactotransferrin gene is expressed in rat tissues and, if not, whether it is functional or even present in the rat genome.

The sequenced rat Tf cDNA is 2275 bp long with a 5' non-coding flanking region of 17 nucleotides and a 3' non-coding sequence of 173 nucleotides. The remaining 2085 bases code for the rat Tf preprotein, including an upstream signal sequence of 19 amino acids, and the secreted protein comprising 676 amino acids, three residues less than that of man [17] and rabbit [14] Tfs. The 178-amino acid sequence from residue 499 to the C-terminal residue 676 of mammary-gland Tf is identical with the partial sequence described for rat liver Tf [26], except for three residues located at positions 669 (Glu → Asp), 674 (His → Thr) and 675 (Lys → Ala) and for short sequences in the 3' flanking region

all of which are located in the last exon as compared with human Tf gene organization [28,46]. The sequence reported here shows, particularly in this zone, homologous residues to other Tfs. Given that polymorphism is known to occur in the rat Tf gene [47], the almost perfect identity between liver and mammary-gland Tf suggests that the rat Tf gene exists as a single copy.

The rat Tf signal sequence ends with the -1 amino acid residue alanine and obeys the -1, -3 rule [Ala (-1), Cys (-3)] for signal-cleavage sites. The signal peptide alignment of different Tfs has, except for *Manduca sexta* Tf, chicken ovotransferrin and human melanotransferrin, a well conserved *c* region (results not shown). In these latter three Tfs, the -3 and -1 positions are occupied, as usual, by small neutral residues. The sequence similarity in the *h* region consists of the conservation of the -11 or the -12 leucine residue, depending on the class of Tf. The rat Tf analysed by us contains a signal sequence of 19 amino acid residues, as do other Tfs. However, other authors [24] have found for a rat recombinant Tf a presequence of 20 amino acids with a supplementary lysine located after the first methionine residue.

As stated earlier, rat Tf also exhibits extensive identity (approx. 70%) with Tf from other species. In particular, our cDNA sequence showed 76% identity with that of human serum Tf. Analysis of 14 complete nucleotide sequences coding for mature Tfs has shown that at least three gene duplication events have occurred during the evolution of these sequences. These duplications have been proposed previously [11,28-29] on the basis of analyses of complete and partial amino acid sequences. The present analysis was performed using alignments of complete nucleotide sequences. The inferred phylogenetic Tf trees are in accord with fossil records. The same tree topology can be obtained using either half of Tfs, but when both halves are used simultaneously for the reconstruction, the halves of human melanotransferrin are observed to be closer to each other than to any other sequence. This implies that the divergence between each half is substantially lower than the variation relative to sequences that diverged several million years before, as the original duplication happened before the diversification of arthropods. This same phenomenon can be observed in Tfs from the remaining vertebrates, as the estimated number of substitutions per site is higher between the two halves in insects than in vertebrates. Consequently, some mechanism must have been acting on vertebrates such that the two halves of their Tfs are more similar to each other than would be expected.

At present, it is not possible to ascertain whether this similarity is due to selective pressure or to some other mechanism, such as intragenic recombination or gene conversion, that could also homogenize the sequence in both halves. Gene duplications are usually accompanied by an acceleration of the evolutionary rate [48,49], explained either by relaxed selection on the duplicated sequence or by an increased selection for the newly acquired function. In this case, it has been shown that human melanotransferrin has evolved at a lower rate than the remaining Tfs. As the evolutionary rates of Tfs in insects and the remaining ones in vertebrates are approximately equal, it seems clear that the pace of evolution of melanotransferrin has slowed relative to the others. This can be explained only by increased selection pressure on the new sequence. Two different kinds of constraints are often invoked to account for increased selection pressure acting on a sequence: structural and functional. As far as we know, there is no known function that can be ascribed to human melanotransferrin in normal cells, as its presence has only been detected in significant amounts of melanomas [22]. Therefore it is difficult to imagine a functional constraint that would slow the rate of evolution in this sequence. On the other hand, the location of

melanotransferrin on the cell surface may make a structural constraint more likely, although whether this is really due to selection of to other mechanisms of sequence homogenization still remains uncertain.

Rat Tf possesses only one potential glycosylation site for N-glycosidically linked glycans located at asparagine-490 in the C-terminal lobe. The presence of only one glycan/molecule is in agreement with the carbohydrate content of this Tf (about 3%) and confirms that, in all known Tfs from different species, apart from man and one horse glycovariant, there is only one glycan per molecule [30].

In order to determine whether Tf glycosylation is tissue-specific, we compared our results with those obtained previously for rat serum Tf [31]. The structures of the major molecular glycovariants of rat milk Tf (Figure 5) are of the biantennary N-acetyl-lactosamine type, containing one or two sialic acid residues and zero or one residue of fucose. The fully sialylated form of the glycan is one of the several forms of glycan isolated from rat serum Tf that has been previously characterized [31]. Rat serum Tf also possesses different forms of a biantennary trisialylated glycan with the $\{\beta\text{-Gal-(1}\rightarrow\text{3)}[\alpha\text{-Neu-5-Ac(2}\rightarrow\text{6)}]\text{GlcNAc}(\beta\text{1)}\}$ sequence present on the serum rTF-1 and rTF-2 glycovariants. The other difference between rat Tfs isolated from milk and serum is the presence or absence of an $(\alpha\text{1}\rightarrow\text{6})$ fucose residue in trace amounts in the serum protein. Fucosylation occurs in milk proteins but practically never in the corresponding serum proteins, as previously shown for human lactoferrin [50,51] and mouse Tf [52], leading to the conclusion that fucosylation is tissue-specific. Expression of fucosyltransferase activity might also be dependent on cell culture conditions [42,53]. A knowledge of the factors that induce fucosylation may be important in the understanding of the mechanisms of regulation of glycan biosynthesis. In order to determine whether glycans are markers of evolution, some authors [30] undertook a comparative study of the glycan primary structures of Tfs from several different species. Our results also led to the conclusion that Tf glycans are specific for each Tf and, for a given Tf, specific to the species.

This work was supported by grants from the *Institució valenciana d'estudis i investigació* (Code 735 and 883) and by the Laboratoire de Chimie Biologique (UMR no. 111 du CNRS), Université des Sciences et Technologies de Lille (Director Professor A. Verbert). The *Acciones integradas hispano-francesas* (244 área 03, during 1989; and 178 área 04, during 1990) have contributed enormously to this work. We thank Santiago Elena and Celia Buades of the Genetics Department of our University for help in the initial stages of the phylogenetic analysis. We also thank Dr. R. J. Pierce for critical reading of the manuscript.

REFERENCES

- De Jong, G., van Dijk, J. P. and van Eijk, H. G. (1990) *Clin. Chim. Acta* **190**, 1–46
- Bowman, B. H., Yang, F. and Adrian, G. S. (1988) *Adv. Genet.* **25**, 1–38
- Aisen, P. and Listowsky, I. (1980) *Annu. Rev. Biochem.* **49**, 357–393
- Montreuil, J., Mazurier, J., Legrand, D. and Spik, G. (1985) in *Proteins of Iron Storage and Transport* (Spik, G., Montreuil, J., Crichton, R. R. and Mazurier, J., eds.), pp. 25–38, Elsevier, Amsterdam
- Levin, M. J., Tuil, D., Uzan, G., Dreyfus, J. C. and Kahn, A. (1984) *Biochem. Biophys. Res. Commun.* **122**, 212–217
- Zakin, M. M. (1992) *FASEB J.* **6**, 3253–3258
- Lee, E. Y. H., Barcellos-Hoff, M. H., Chen, L. H., Parry, G. and Bissell, M. J. (1987) *In Vitro Cell Dev. Biol.* **23**, 221–226
- Keon, B. H. and Kweenan, T. W. (1993) *Protoplasma* **172**, 43–48
- Escalante, R., Houdebine, L. M. and Pamblanco, M. (1993) *J. Mol. Endocrinol.* **11**, 151–159
- Jamroz, R. C., Gasdaska, J. R., Bradfield, J. Y. and Law, J. H. (1993) *Proc. Natl. Acad. Sci. U.S.A.* **90**, 1320–1324
- Bartfeld, N. S. and Law, J. H. (1990) *J. Biol. Chem.* **265**, 21684–21691
- Moskaitis, J. E., Pastori, R. L. and Schoenberg, D. R. (1990) *Nucleic Acids Res.* **18**, 6135–6135
- Jeltsch, J. M. and Chambon, P. (1982) *Eur. J. Biochem.* **122**, 291–295
- Banfield, D. K., Chow, B. K. C., Funk, W. D., Robertson, K. A., Umelas, T. M., Woodworth, R. C. and MacGillivray, R. T. A. (1991) *Biochim. Biophys. Acta* **1089**, 262–265
- Baldwin, G. and Weinstock, J. (1988) *Nucleic Acids Res.* **16**, 8720
- Carpenter, M. A. and Broad, T. E. (1993) *Biochim. Biophys. Acta* **1173**, 230–232
- Yang, F., Lum, J. B., McGill, J. R. et al. (1984) *Proc. Natl. Acad. Sci. U.S.A.* **81**, 2752–2756
- Lydon, J. P., O'Malley, B. R., Saucedo, O., Lee, T., Headon, D. R. and Conneely, O. M. (1992) *Biochim. Biophys. Acta* **1132**, 97–99
- Pierce, A., Colavizza, D., Benaissa, M. et al. (1991) *Eur. J. Biochem.* **196**, 177–184
- Pentecost, B. T. and Teng, C. T. (1987) *J. Biol. Chem.* **262**, 10134–10139
- Powell, M. J. and Ogden, J. E. (1990) *Nucleic Acids Res.* **18**, 4013–4013
- Rose, T. M., Plowman, G. D., Teplow, D. B., Dreyer, W. J., Hellstrom, K. E. and Brown, J. P. (1986) *Proc. Natl. Acad. Sci. U.S.A.* **83**, 1261–1263
- Mead, P. E. and Tweedie, J. W. (1990) *Nucleic Acids Res.* **18**, 7167–7167
- Schreiber, G., Dryburgh, H., Millership, A. et al. (1979) *J. Biol. Chem.* **254**, 12013–12019
- Aldred, A. R., Howlett, G. J. and Schreiber, G. (1984) *Biochem. Biophys. Res. Commun.* **122**, 960–965
- Huggenwik, J. I., Idzerda, R. L., Haywood, L., Lee, D. C., McKnight, G. S. and Griswold, M. D. (1987) *Endocrinology* **120**, 332–340
- Gilmont, R. R., Coulter, G. H., Sylvester, S. R. and Griswold, M. D. (1990) *Biol. Reprod.* **43**, 139–150
- Park, I., Schaeffer, E., Sidoli, A., Baralle, F. E., Cohen, G. N. and Zakin, M. M. (1985) *Proc. Natl. Acad. Sci. U.S.A.* **84**, 1769–1773
- Baldwin, G. S. (1993) *Comp. Biochem. Physiol. B* **106**, 203–218
- Spik, G., Coddeville, B. and Montreuil, J. (1988) *Biochimie* **70**, 1459–1468
- Spik, G., Coddeville, B., Strecker, G. et al. (1991) *Eur. J. Biochem.* **195**, 397–405
- Chomczynski, P. and Sacchi, N. (1987) *Anal. Biochem.* **162**, 156–159
- Rado, T. A., Wei, W. and Benz, E. J. (1987) *Blood* **70**, 989–993
- Sambrook, J., Fritsch, E. F. and Maniatis, T. (1989) *Molecular Cloning. A Laboratory Manual*, 2nd edn., Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY
- Gardner, R. C., Howarth, H. J., Hahn, P., Brown-Luedi, M., Shepherd, R. J. and Messing, J. (1981) *Nucleic Acids Res.* **9**, 2871–2888
- Higgins, D. G., Bleasby, A. J. and Fuchs, R. (1992) *Comput. Appl. Biosci.* **8**, 189–191
- Kumar, S., Tamura, K. and Nei, M. (1993) *MEGA: Molecular Evolutionary Genetics Analysis*, version 1.01, Pennsylvania State University, Philadelphia
- Nei, M. (1991) in *Recent Advances in Phylogenetics Analysis of DNA Sequences* (Miyamoto, M. M. and Cracraft, J. L., eds.), pp. 90–128, Oxford University Press, Oxford
- Efron, B. (1982) *The Jackknife, the Bootstrap and Other Resampling Plans*, Society for Industrial and Applied Mathematics, Philadelphia
- Felsenstein, J. (1993) *Phylogenetic Inference Package (PHYLP)*, version 3.5, University of Washington, Seattle
- Saitou, N. and Nei, M. (1987) *Mol. Biol. Evol.* **4**, 406–425
- Campion, B., Léger, D., Wieruszski, J. M., Montreuil, J. and Spik, G. (1989) *Eur. J. Biochem.* **184**, 405–413
- Corfield, A. P., Beau, J. M. and Schauer, R. (1978) *Hoppe-Seyler's Z. Physiol. Chem.* **359**, 1335–1342
- Montreuil, J., Bouquelet, S., Debray, H. et al. (1994) in *Carbohydrate Analysis, A Practical Approach* (Chaplin, M. F. and Kennedy, J. F., eds.), pp. 181–293, IRL Press, Oxford
- Vliegenthart, J. F. G., Dorland, L. and van Halbeek, H. (1983) *Adv. Carbohydr. Chem. Biochem.* **41**, 209–374
- Schaeffer, E., Lucero, M. A., Jeltsch, J. M. et al. (1987) *Gene* **56**, 109–116
- Nagabuchi, M., Kawamoto, Y., Nishikawa, T. and Nishimura, M. (1993) *Biochem. Genet.* **31**, 147–154
- Li, W.-H. (1985) in *Population Genetics and Molecular Evolution* (Ohta, T. and Aoki, K., eds.), pp. 333–352, Japan Scientific Societies Press, Tokyo
- Ohta, T. (1993) *Genetics* **134**, 1271–1276
- Spik, G., Strecker, G., Fournet, B. et al. (1982) *Eur. J. Biochem.* **121**, 413–419
- Derisbourg, P., Wieruszski, J. M., Montreuil, J. and Spik, G. (1990) *Biochem. J.* **269**, 821–825
- Leclercq, Y., Sawatzki, G., Wieruszski, J. M., Montreuil, J. and Spik, G. (1987) *Biochem. J.* **247**, 571–578
- Jacquinet, P.-M., Léger, D., Wieruszski, J.-M., Coddeville, B., Montreuil, J. and Spik, G. (1994) *Glycobiology* **4**, 617–624