BIOCHEMICAL LETTERS

Clarification of the nucleotide sequence at the 5'-end of the cDNA for rat liver carnitine palmitoyltransferase II

Controversy has arisen surrounding the structure of the cDNA corresponding to rat liver carnitine palmitoyltransferase II (CPT II). The problem stems from reports by Brady et al. [1,2] in which it was suggested that our published sequence [3] lacked a considerable stretch of nucleotides at the 5'-end, and thus predicted a protein lacking the N-terminus of the natural product. We have previously countered this argument on a variety of grounds [3–6]. Particularly puzzling to us was that in [2] Brady et al. had apparently isolated a rat CPT II genomic clone that differed substantially from our cDNA in terms of its nucleotide sequence 5' to the ATG codon that we had assigned as the translation start site (Figure 1). In support of the authenticity of this material, Brady et al. [2] noted that it contained a nucleotide sequence (B-1 in Figure 1b) encoding an octapeptide which they believe to represent the true N-terminus of the mature protein. For these reasons we had to entertain the possibility that our cDNA was, after all, short at its 5' end or, alternatively, that rat liver contains CPT II transcripts corresponding to both of the sequences in question (Figures 1a and 1b). Accordingly, poly(A)⁺ RNA was isolated from livers of normal rats and of animals treated with diethylhexylphthalate (DEHP; an inducer of peroxisomal and mitochondrial enzymes [8]). From this, first-strand

cDNA was synthesized by using reverse transcriptase and oligo(dT) as the primer. With cDNA generated from poly(A)⁺ RNA of DEHP-treated rats, a PCR reaction was carried out using as primers oligonucleotides W-1 and BW-1(c) (the complement of BW-1), corresponding to portions of the 5'-untranslated and coding regions, respectively, of the sequence found by Woeltje et al. [3] (Figure 1a). The product was subsequently cloned into M13 and its sequence determined in both orientations. Its 5' end was found to be identical with that of our CPT II cDNA, containing the translation stop signal (TAG) 22 bp upstream of a methionine codon ([3] and Figure 1a).

First-strand cDNA was also synthesized by using as primer an oligonucleotide complementary to region BW-2 in Figure 1, i.e. BW-2(c). After subsequent addition of oligonucleotide W-1 or B-1 (Figure 1) to act as the 5' primer, the mixture was subjected to PCR. A positive control using oligonucleotides W-1 and BW-2(c) as primers and our original CPT II cDNA as template was included. As seen in Figure 2(a) (lane 7) the control gave rise to a DNA fragment of the expected size (~ 0.6 kb). PCR products were also generated with primer pair B-1/BW-2(c) (lanes 3 and 4) or W-1/BW-2(c) (lanes 5 and 6). However, unlike the situation with W-1/BW-2(c), where a 0.6 kb band was readily detected as the major product [signal intensity being greater with poly(A)⁺ RNA from DEHP-treated rats, as expected], the DNA fragments amplified by using primers B-1/BW-2(c) were much smaller in size (compare lanes 5 and 6 with lanes 3 and 4). Moreover, a

Figure 1 Comparison of postulated 5' ends for rat liver CPT II

(a) cDNA sequence determined by Woeltje et al. [3]; (b) composite genomic and cDNA sequence determined by Brady et al. [2,7]. Nucleotide numbering assigns +1 to the A of the translation-initiating codon in sequence (a). The more darkly shaded regions represent published sequence that is identical in both constructs. The more lightly shaded region in sequence (b) has been stated to be the same as that in sequence (a) [1]. Segments of each sequence on which oligonucleotides used for PCR reactions and Southern-blot analyses were based are underlined. They carry the prefix W, B or BW, depending on whether they relate to the Woeltje, Brady or both sequences. Putative transcription start sites in the genomic clone were assigned to nucleotides -127 and -97 [2]. Stop codons are denoted by the asterisks.

Figure 2 PCR products generated from the 5' end of CPT II cDNA

Poly(A)⁺ RNA was isolated from either normal (N) or DEHP-treated (D) rat liver and used for first-strand synthesis of cDNA as described in the text. P represents a positive control using the original CPT II cDNA of Woeltje et al. [3]. PCR reactions were performed by using as the 5' primer oligonucleotide W-1 or B-1, unique to sequences a and b respectively (Figure 1); oligonucleotide BW-2(c), common to both sequences, served as the 3' primer. (a) Ethidium bromide-stained agarose gel containing the PCR products. (b) Southern blot of the agarose gel probed with ³²P-5'-end-labelled oligonucleotide BW-3(c) (Figure 1). 'c' denotes oligonucleotides complementary to those underlined in Figure 1.

Southern blot of the gel, using ³²P-labelled oligonucleotide BW-3(c) as the probe (Figure 2b), revealed strong hybridization in lanes 5–7, with no signal detectable in lanes 3 and 4. Since oligonucleotide BW-3(c) is specific for rat liver CPT II cDNA, this result indicated that the PCR product obtained with primer combination W-1/BW-2(c) represented an authentic DNA copy of the corresponding region of CPT II mRNA. By contrast, the material generated from primers B-1 and BW-2(c) did not derive from the CPT II mRNA. Moreover, both oligonucleotide W-1(c) and a 1.4 kb internal fragment of our CPT II cDNA [3] hybridized strongly on Northern blots to a 2.5 kb transcript in poly(A)⁺ RNA from livers of normal and DEHP-treated rats (results not shown). We have repeatedly failed to obtain PCR products, or to identify CPT II transcripts on Northern blots, using oligo-

nucleotides upstream of nucleotide –3 in the Brady et al. [2] sequence (Figure 1b).

These results indicate that the cDNA isolated by Woeltje et al. [3] is indeed full-length and that it represents the major, if not the sole, transcript for CPT II in rat liver. As to the origin or identity of the genomic PCR product isolated by Brady et al. [2], we can offer no explanation at present. Although the possibility exists that it represents an authentic region of the rat genome, several considerations invite scepticism. First, the clone was obtained through the use of PCR under conditions where susceptibility to non-specific product formation is very high, namely the use of long oligonucleotides (> 30 bases) with a low annealing temperature (50 °C). Second, the fragment spanning bp –345 to –285 of the genomic clone shares 88% identity with the polylinker region of the vector M13mp18, frequently used for sequencing purposes. Third, from S1-nuclease analysis, two regions of the genomic clone were postulated as potential start sites for transcription [2]. However, one of these sites lies 4 bp internal to sequence said to be contained in a rat liver CPT II cDNA obtained previously by the same authors [2,8]. One is therefore led to suspect that the material presented in [2] arose from a cloning artifact. Furthermore, earlier reports in which a degenerate oligonucleotide corresponding to region B1 of Figure 1(b) was used in studies of CPT II gene regulation [7,9] become questionable.

This work was supported by grants from the National Institutes of Health (DK18573), the Chilton Foundation and Sandoz Pharmaceuticals.

Brian C. WEIS, Daniel W. FOSTER and J. Denis McGARRY

Departments of Internal Medicine and Biochemistry and the Gifford Laboratories for Diabetes Research, University of Texas Southwestern Medical Center, Dallas, TX 75235, U.S.A.

- 1 Brady, P. S., Liu, J. S., Park, E. A., Hansen, R. W. and Brady, L. J. (1991) *FASEB J.* **5**, A817
- 2 Brady, P. S., Park, E. A., Liu, J., Hansen, R. W. and Brady, L. J. (1992) *Biochem. J.* **286**, 779–783
- 3 Woeltje, K. F., Esser, V., Weis, B. C., Sen, A., Cox, W. F., McPhaul, M. J., Slaughter, C. A., Foster, D. W. and McGarry, J. D. (1990) *J. Biol. Chem.* **265**, 10720–10725
- 4 Brown, N. F., Esser, V., Gonzalez, A. D., Evans, C. T., Slaughter, C. A., Foster, D. W. and McGarry, J. D. (1991) *J. Biol. Chem.* **266**, 15446–15449
- 5 McGarry, J. D., Sen, A., Brown, N. F., Esser, V., Weis, B. C. and Foster, D. W. (1992) in *Current Concepts in Carnitine Research* (Carter, A. L., ed.), pp. 137–151, CRC Press, Boca Raton, FL
- 6 Brown, N. F., Sen, A., Soltis, D. A., Jones, B., Foster, D. W. and McGarry, J. D. (1993) *Biochem. J.* **294**, 79–86
- 7 Brady, P. S., Feng, Y. and Brady, L. J. (1988) *J. Nutr.* **118**, 1128–1136
- 8 Reubsat, F. A. G., Veerkamp, J. H., Dirven, H. A. A. M., Bruckwilder, M. L. P., Hashimoto, T., Trijbels, J. M. F. and Monnens, L. A. H. (1990) *Biochim. Biophys. Acta* **1047**, 264–270
- 9 Brady, P. S. and Brady, L. J. (1989) *Biochem. J.* **258**, 677–682