BIOCHEMICAL JOURNAL LETTERS

Still degrading after all these years

In recent years, the mechanisms by which peptides are degraded or processed have received increasing attention by a number of groups. In most cases, there is a considerable gap between understanding the actions of purified peptidases and their role *in vivo*. This has been most dramatically illustrated for substance P and related tachykinins. For example, as many as half a dozen different enzymes have been proposed to act physiologically on substance P [1]. We purified one such enzyme, termed 'substance P-degrading enzyme' in 1980–1981 from human brain [2], and indicated some unusual and highly distinctive features of its action on substance P: notably a three-point cleavage pattern, cleaving between Gln⁶–Phe⁷, Phe⁷–Phe⁸ and Phe⁸–Gly⁹, sensitivity to metallopeptidase inhibitors (e.g. EDTA and *o*-phenanthroline), but not phosphoramidon, and preferential recognition of substance P as a substrate. We also put forward evidence, based on the unique cleavage pattern and inhibitor sensitivity, that this enzyme was rate-limiting in initiating the inactivation of the biological activity of substance P in brain [1,2]. This work and its conclusions were subsequently criticized by Turner, Kenny and colleagues based on their work with porcine kidney and brain membranes [3], and later human brain membranes [4], and an impression has developed that a phosphoramidon-insensitive substance P-cleaving endopeptidase with the precise cleavage pattern we described does not exist.

Accordingly, we are both pleased but somewhat surprised to find in the recent article by Stephenson & Kenny [5] that all of the key features of this enzyme have been redescribed. In view of their frequent earlier references to our work in a different context, relationship of 'substance P-degrading enzyme' to endopeptidase 24.11 [3–6], it was disappointing to note that our original description, plus the confirmatory reports from at least four other groups [7–10], have not been referenced here, even though they have clearly influenced this work. The issue is not whether 'substance P-degrading enzyme' and 'endopeptidase 2' are identical, or whether our original operational name should be replaced by a systematic name, but rather that the potential role of 'substance P-degrading enzyme' has been neglected because of dis-

proportionate emphasis on the enzymic cleavages and inhibitor sensitivity attributed to endopeptidase 24.11 in pig membranes [3], which as Stephenson & Kenny illustrate in their Fig. 3 [5], may be an idiosyncratic circumstance amongst mammalian species. Thus, we should like to restate the original hypothesis that the 'substance P-degrading enzyme', potentially a family of related enzymes, is a major system initiating the biological inactivation of substance P, and, importantly, perhaps other peptides. This last point is made to emphasize that we have never characterized the enzyme as completely specific for substance P, although this incorrect interpretation has been put forward in reviews [6], but rather identified the enzyme by a trivial name which indicated the assay upon which its purification was based. Indeed, in our original description, we noted that angiotensin I, bradykinin, and LHRH were alternative substrates, as confirmed by Stephenson & Kenny [5]. Nevertheless, the distribution of the 'substance P-degrading enzyme' in mammalian spinal cord, as determined by immunocytochemistry [11], closely parallels the distribution of immunoreactive substance P, suggesting the association of this enzyme with substance P remains the most viable possibility for one of its physiological functions. With hindsight, the trivial name 'substance P-degrading enzyme' has undoubtedly contributed to confusion, and thus we would propose that the concerned groups conjointly introduce a generic name, and come to agreement as a matter of urgency as to how many such distinct enzymes can conform to the following description: phosphoramidon-insensitive neutral metalloendopeptidase preferentially hydrolysing substance P in at least two of the three indicated cleavage sites. By this description, we believe that 'endopeptidase 2' [5], and enzymes identified by the groups of Bauer [7], Selinger [8], Ishii [9], and Nyberg & Terenius [10] would be included. We have summarized the physical and enzymic comparison to 'substance P-degrading enzyme' [1,2,11] to the enzyme from [5] in Table 1.

Lesley PROBERT and Michael R. HANLEY

MRC Molecular Neurobiology Unit, MRC Centre, Cambridge CB2 2QH, U.K.

Table 1. Comparison of 'substance P-degrading enzyme' and 'kidney endopeptidase 2'

Enzyme	Membrane-associated	$10^{-3} \times$ Subunit M_r	K_m for substance P (μM)	Phosphoramidon-insensitive, EDTA-sensitive	pH optimum
Substance P-degrading enzyme	Yes	70	29	Yes	Neutral
Kidney endopeptidase 2	Yes	74–80	86	Yes	Neutral