The adipocyte $G_o\alpha$ -immunoreactive polypeptide is different from the α subunit of the brain G_o protein

Bruno ROUOT,*§ Josiane CARRETTE,* Max LAFONTAN,† Phuong LAN TRAN,‡ Jean-Alain FEHRENTZ,* Joël BOCKAERT* and Madeleine TOUTANT*

*Centre CNRS-INSERM de Pharmacologie-Endocrinologie, F-34094 Montpellier Cedex 2, †INSERM U 317, Institut de Physiologie, Rue Magendie, F-31400 Toulouse, and ‡Laboratoire d'Immunogénétique Moléculaire, Rue de l'Ecole de Médecine, F-75006 Paris, France

Rat adipose tissue possesses two *Bordetella pertussis* toxin (PTX) substrates and, in the same 39–41 kDa molecular mass range, positive immunoreactivity has also been reported with antibodies against the α subunit of G_o , the major brain GTP-binding protein (G-protein). In this study, the presence of the brain $G_o\alpha$ subunit at 39 kDa in adipocytes was reassessed, since direct correspondence between PTX substrates and $G_o\alpha$ immunoreactivity has not yet been clearly established. On resolutive SDS/polyacrylamide-gel electrophoresis, the PTX substrates of human adipocytes were compared with the three PTX substrates found in brain. No ADP-ribosylated substrate at the level of the 39 kDa brain $G_o\alpha$ could be detected in adipocyte membranes. Immunoblotting of human adipocyte membranes stained with our anti- $G_o\alpha$ antibodies confirmed the presence of a positive immunoreactivity in this tissue, but the apparent molecular mass of the immunoreactive polypeptide in adipocytes was higher than that found in nervous tissues. Taken together, these results indicate that the brain $G_o\alpha$ subunit is not present in adipose tissue. They also suggest the existence of a G-protein in adipocytes which is immunologically related to $G_o\alpha$ but having a slightly higher molecular mass.

INTRODUCTION

Guanine-nucleotide-binding regulatory proteins (G-proteins) are a family of receptor-coupled signal-transducing proteins that regulate a variety of second messenger systems and ion channels [1,2]. Diversity in this heterotrimeric G-protein family mainly comes from the α subunit of G-proteins. On the α subunits of various proteins, the *Bordetella pertussis* toxin (PTX) catalyses an ADP ribosylation which abolishes the effects transduced by the PTX-sensitive G-proteins. Based on electrophoretic mobility, three different groups of PTX substrates can be distinguished. The group which comigrates with 41 kDa molecular mass proteins is now assumed to be the polypeptide corresponding to the α subunit sequence termed α_{11} [3]. The α subunits of the major brain GTP-binding protein G_o ($G_o\alpha$) migrate on SDS/polyacrylamide-gel electrophoresis (SDS/PAGE) with a mobility corresponding to a 39 kDa protein. Recently, the presence of PTX substrates with a migration corresponding to a 40 kDa protein has been demonstrated in many tissues [4] and the substrates have been purified from brain [5] and neutrophils [6]. This 40 kDa PTX substrate seems to correspond, in the human leukaemic cell line HL-60, to the sequence termed α_{12} , although α_{13} was also identified in these cells [7].

Adipocytes were the first peripheral tissue cells in which two PTX substrates were simultaneously detected [8]. The polypeptide with the slower electrophoretic mobility was assigned to be at the 41 kDa level, while the other was reported either at the 40 kDa [8] or the 39 kDa

level [9,10]. Immunological studies with anti- $G_o\alpha$ antibodies indicated the existence of a cross-reactivity in rat adipocytes [10,11]. It was thus suggested that the low-molecular-mass PTX substrate in adipocytes was identical to the brain $G_o\alpha$ subunit [10]. In a previous study we also showed that our anti- $G_o\alpha$ antibodies exhibited a positive immunoreactivity in human fat-cell membranes [12]. These previous observations appear contradictory to recent studies indicating that the presence of the G_o protein seems limited to brain tissues and neuroendocrine cells such as chromaffin cells or those of the anterior pituitary and the islets of Langerhans [13,14]. In the present study, we investigated the nature of the $G_o\alpha$ immunoreactive polypeptide present in adipocytes.

EXPERIMENTAL

Isolation of adipocytes and preparation of membranes

Human abdominal subcutaneous adipose tissue samples were obtained from premenopausal women (25–40 years old) undergoing elective surgical lipectomy. Adipocytes were isolated by collagenase digestion of adipose fragments according to the method of Rodbell [15]. After lysis and defatting by centrifugation, plasma membranes were separated from mitochondria and other cell components of particulate fractions by centrifugation in the presence of a self-forming gradient of Percoll at 10000 g for 15 min in a fixed-angle rotor, as previously described [16]. The plasma membrane fraction was washed twice to remove Percoll. The membrane preparation was stored

Abbreviations used: G-proteins, guanine-nucleotide-binding proteins; PTX, *Bordetella pertussis* toxin; PAGE, polyacrylamide-gel electrophoresis.
§ To whom all correspondence should be addressed.

at -80°C . Some control experiments were performed with dog and rat adipocyte membranes prepared in a similar manner.

Preparation of brain PTX substrates and tissue membranes

Purification of G-proteins from bovine cerebral cortex was achieved as indicated by Sternweis & Robishaw [17]. The mixture of the three PTX substrates obtained is designated as 'purified brain G-proteins' throughout the paper. Enriched and pure G_{α} subunits were obtained as previously described [12]. Pure G_{α} was used to raise polyclonal antibodies in rabbit. Crude membranes from various tissues were prepared as described [12].

Preparation of HL-60 cell homogenates

HL-60 human promyelocytic leukaemia cells were grown in RPMI 1640 medium supplemented with 15% heat-inactivated fetal calf serum, 2 mM-glutamine, 10 mM-Hepes, 100 units of penicillin/ml and 100 μg of streptomycin/ml at 37°C in a humidified atmosphere containing 5% CO_2 . Cells were seeded twice weekly at a density of 3×10^5 cells/ml. Cells were washed in a solution of Dulbecco's phosphate-buffered saline (Gibco), resuspended in 20 mM-Tris/HCl, pH 8/1 mM-EDTA/1 mM-dithiothreitol (DTT) and subjected to three cycles of freezing and thawing. The resulting homogenate was stored at -30°C until use.

PTX-catalysed ADP-ribosylation

Brain PTX substrates (14 μg of protein), adipocyte membranes (85 μg), HL-60 homogenate (90 μg), and 25–100 μg of various tissue membranes were incubated for 60 min with [^{32}P]NAD $^{+}$ (New England Nuclear) in the presence of preactivated pertussis toxin (List Biological Laboratories, Campbell, CA, U.S.A.), as previously described [12]. The reaction was stopped by adding SDS (2%) with 100 μg of bovine serum albumin/ml, and proteins were precipitated with trichloroacetic acid. After centrifugation, the pellets were washed twice with diethyl ether and treated as described below before electrophoresis.

SDS/PAGE and immunoblotting

Prior to electrophoresis, samples were subjected to *N*-ethylmaleimide treatment and electrophoresis as described previously [18] for autoradiography or immunoblotting.

Immunoblots were carried out on the same gel, and after electrotransfer of the proteins on to nitrocellulose, the sheet was reacted as previously described [12] with the affinity-purified antibodies against the G_{α} subunit. Antibodies against the C-terminal peptide common to G_{i1} and G_{i2} , KNNLKDCGLF, were also used after purification with Affi-gel 10 (Bio-Rad) linked to the LKDCGLF peptide. Immunostaining was achieved by incubation with ^{125}I -labelled Protein A (10^5 c.p.m./ml) followed by autoradiography of the nitrocellulose sheet. Only the relevant part of the autoradiograms or immunoblots are shown in the Figures.

Protein determination

Proteins were measured by the method of Lowry *et al.* [19], with bovine serum albumin as the standard.

Fig. 1. Comparison by SDS/PAGE of the PTX substrates from adipocytes with those from other tissues

Samples from various tissues were ADP-ribosylated in the presence of [^{32}P]NAD and PTX, as described in the Experimental section, before being loaded. Samples were: lanes 1 and 4, purified calf-brain PTX substrates (0.14 μg); lane 2, human adipocyte membranes (13 μg); lane 3, membranes from HL-60 cells (25 μg).

RESULTS

The three bovine brain PTX substrates first described by Neer *et al.* [20] with apparent molecular masses of 39, 40 and 41 kDa, can be clearly discriminated under our resolvable SDS/PAGE conditions (Fig. 1, lanes 1 and 4). The [^{32}P]ADP-ribosylated PTX substrates of human adipocytes (lane 2), when compared with the three substrates of brain (lanes 1 and 4) and with the 40 kDa substrate of the leukaemic cell line HL-60 (lane 3), clearly matched the two brain substrates of lower mobility, assumed to be G_{i1} and G_{i2} .

The lack of the 39 kDa PTX substrate in adipocytes suggested that the G_{α} immunoreactivity found by Rapijko *et al.* [10] and by our group [12] was due to a subunit which is different from the brain G_{α} polypeptide. Assessment of the electrophoretic mobility of the human adipocyte G_{α} immunoreactive polypeptide was then performed on a gel of increased length (Fig. 2). The mobility of the G_{α} -related polypeptide (lanes 2 and 8) was compared with those of G_i and G_{α} subunits detected either after [^{32}P]ADP ribosylation (lanes 3, 6, 9 and 12) or with specific antibodies in tissues such as rat brain cortex (lanes 1 and 7), human platelets (lanes 5 and 11) and purified brain G-proteins (lanes 4 and 10). The autoradiograms of the [^{32}P]ADP-ribosylated substrates of adipose (lanes 3 and 9) and brain (lanes 6 and 12) tissues, obtained after electrophoresis and electrotransfer on to nitrocellulose sheets, are shown in Figs. 2(a) and 2(b).

The same nitrocellulose sheets were then incubated either with anti- G_{α} antibodies (part a) or with antibodies directed against the common C-terminal sequence of the G_{i1} and G_{i2} (part b). Therefore, after staining with ^{125}I -Protein A, autoradiography of the nitrocellulose sheets depicted both [^{32}P]ADP-ribosylated substrates and immunoreactivities with anti- G_{α} (part c) or anti- $G_{i1,2}$ (part d) antibodies. According to our previous data [12,21], anti- G_{α} antibodies recognized a single protein in brain cortical membranes (Fig. 2c, lane 1) and in the purified brain G-protein preparation (Fig. 2c, lane 4). A clear G_{α} immunoreactivity was detected in adipocyte mem-

Fig. 2. Comparison of electrophoretic mobilities of $[^{32}\text{P}]$ ADP-ribosylated PTX substrates with $G_o\alpha$ and $G_{11,2}\alpha$ immunoreactivity polypeptides of adipocytes and other tissues

This two-step experiment was performed on a single electrophoresis gel loaded with identical samples on both sides. Lanes 1 and 7, rat-brain cortex membranes (20 μg); lanes 2 and 8, human adipocyte membranes (120 μg); lanes 3 and 9, $[^{32}\text{P}]$ ADP-ribosylated adipocyte membranes (10 μg); lanes 4 and 10, purified calf-brain G-proteins (0.5 μg and 0.9 μg respectively); lanes 5 and 11, human platelets (125 μg); lanes 6 and 12, $[^{32}\text{P}]$ ADP-ribosylated brain PTX substrates (0.1 μg). After electrophoresis, as described in the Experimental section, proteins were transferred on to a nitrocellulose sheet which was subjected to a first autoradiography (parts *a* and *b*) in order to assess the relative position of the ADP-ribosylated substrates. Thereafter, (*a*) and (*b*) were incubated with anti- $G_o\alpha$ and anti- $G_{11,2}\alpha$ antibodies respectively and, after staining with protein A, subjected to another autoradiography. Thus, the second autoradiograms (*c* and *d*) reveal both $[^{32}\text{P}]$ ADP-ribosylated substrates and immunoreactive polypeptides.

branes (Fig. 2*c*, lane 2). However, assuming the presence of an identical G_o protein in brain and in adipocytes, the level of this protein in Percoll-purified adipocyte membranes would only represent 0.06% of the protein content, which is substantially lower than the 1% found in brain cortex. Note the slower mobility of ADP-ribosylated substrates compared with the non-ADP-ribosylated ones, for example compare the $G_o\alpha$ immunoreactivity of lanes 1 and 4 (Fig. 2) with the ADP-ribosylated $G_o\alpha$ which is the lower band in lane 6. A similar observation could be made with $F_{11,2}$ immunoreactivities (lane 8) and the corresponding ADP-ribosylated substrates of adipocytes (lane 9). Interestingly, Fig. 2(*c*) revealed that the $G_o\alpha$ -immunoreactive adipocyte substrate (Fig. 2*c*, lane 2) migrated between the two ADP-ribosylated PTX substrates from adipocytes (Fig. 2*c*, lane 3). No immunoreactivity was detected with the anti- $G_o\alpha$ antibodies in human platelets (Fig. 2*c*, lane 5). In Fig. 2(*d*), very little $G_{11,2}\alpha$ immunoreactivity is revealed in rat cortical membranes compared with that of $G_o\alpha$ (Fig. 2*c*, lane 1) but a higher level of both proteins was found in adipocytes (lane 8) and in the purified brain G-protein preparation (lane 10). The immunoreactivity present in human platelets corresponded essentially to the 40 kDa PTX substrate, the G_α subunit of the higher electrophoretic migration generally considered as G_{12} [7]. Comparison of parts (*c*) and (*d*) of Fig. 2 indicated that the $G_o\alpha$ immunoreactivity in adipocytes migrated at a level similar to the higher molecular mass $G_{11,2}$ immunoreactive polypeptide of adipocytes; that is, the $G_{11}\alpha$ subunit.

The electrophoretic mobilities of adipocyte $G_o\alpha$ immunoreactivity was not species-dependent, since the human and rat adipose tissues both exhibited a positive immunoreactivity at a similar level which was different from that of the bovine brain $G_o\alpha$ subunit (results not shown).

DISCUSSION

Comparison of the ADP-ribosylated PTX substrates of adipocytes with those of brain revealed that no labelling was detectable at the $G_o\alpha$ level (39 kDa) in human fat cells. Instead, the two adipocyte PTX substrates corresponded to higher molecular masses, depicted here as the 40 and 41 kDa PTX substrates. Taking into account the fact that ADP-ribosylated substrates migrate at a slower rate than unmodified α subunits, one can assume that the two ^{32}P -labelled bands after ADP-ribosylation in adipocytes derived from two immunoreactive polypeptides revealed with the anti- $G_{11,2}$ C-terminal antibodies. Although the 39 kDa $G_o\alpha$ PTX substrate was absent, a cross-reactivity with our anti- $G_o\alpha$ antibodies was seen in adipose tissue at a level which closely corresponds to the immunoreactivity of the $G_{11}\alpha$ subunit (Fig. 2).

A plausible explanation to account for the $G_o\alpha$ immunoreactivity in adipose tissue would be to assume a cross-reactivity of our anti- $G_o\alpha$ antibodies with the G_α subunit. However, the antibodies towards the 39 kDa substrate only recognized the lower 39 kDa subunit in the mixture of purified brain G-proteins, despite the presence of two G_α subunits in this preparation (Fig. 2). Furthermore, a substantial amount of the 40 kDa substrate in human platelets was not detected by $G_o\alpha$ antibodies. These observations seem to discount a possible cross-reactivity of our antibodies with $G_{11}\alpha$ and $G_{12}\alpha$ subunits. Theoretically, it could also be that our $G_o\alpha$ antibodies recognized the $G_{13}\alpha$ subunit which is possibly present in adipocytes. However, under our experimental electrophoresis conditions (Fig. 1), the PTX substrates of HL-60, reported to possess G_{12} and $G_{13}\alpha$ subunits, had different mobilities to that of the $G_{11}\alpha$ subunit, which migrates at the same level as the $G_o\alpha$ immunoreactive polypeptide. Therefore, a cross-reactivity of $G_{13}\alpha$ with our $G_o\alpha$ antibodies is unlikely.

It should be recalled that our observation is not unique and that Rapiejko *et al.* [10], using antibodies prepared by Northup against $G_o\alpha$, also found immunoreactivity in fat-cell membranes. The fact that antibodies against the peptide sequence close to the N-terminal of $G_o\alpha$ did not detect any immunoreactivity in adipose tissue [22] indicates that these antibodies were directed against other epitopes than our antibodies and those of Northup. This discrepancy indicates that the total sequence of the brain $G_o\alpha$ is not present in adipocytes and suggests specific $G_o\alpha$ -like post-translational modifications in adipose tissue [22,23].

The presence of a positive immunoreactivity towards anti- $G_o\alpha$ antibodies is not however restricted to adipocytes. Our antibodies have also identified a G-protein α subunit in skeletal muscle transverse tubules which had a higher apparent molecular mass than the brain G_o protein [18]. Similarly, Tanfin *et al.* [24] reported the presence of a novel form of G_o in the rat myometrium, based on analysis with antibodies against specific peptides of various G_α subunits. These data, together with our

findings with adipocytes, strongly suggest the existence in various tissues of G-proteins which have not yet been fully characterized, and which may differ from the well-established G-proteins, either by their sequences or by the differences in their post-translational modifications.

We thank P. Brabet and V. Homburger for purifying the G_{α} antibodies, C. Bunoust, M. Passama and J. P. Toutant for their photographs of gels and autoradiographies and Y. Audigier for stimulating discussions. The secretarial assistance of A. Turner-Madeuf and P. Costagliola is greatly appreciated.

REFERENCES

- Gilman A. G. (1987) *Annu. Rev. Biochem.* **56**, 615–649
- Neer, E. J. & Clapham, D. E. (1988) *Nature (London)* **333**, 129–134.
- Jones, D. T. & Reed, R. R. (1987) *J. Biol. Chem.* **262**, 14241–14249
- Brabet, P., Pantaloni, C., Rouot, B., Toutant, M., Garcia-Sainz, A., Bockaert, J. & Homburger, V. (1988) *Biochem. Biophys. Res. Commun.* **152**, 1185–1192
- Katada, T., Oinuma, M., Kusakabe, K. & Ui, M. (1987) *FEBS Lett.* **213**, 353–358
- Gierschik, P., Sidiropoulos, D., Spiegel, A. & Jakobs, K. (1987) *Eur. J. Biochem.* **165**, 185–194
- Murphy, P. M., Eide, B., Goldsmith, P., Brann, M., Gierschik, P. & Spiegel, A. (1987) *FEBS Lett.* **221**, 81–86
- Malbon, C. C., Rapiejko, P. J. & Garcia-Sainz, J. A. (1984) *FEBS Lett.* **176**, 301–306
- Owens, J. R., Frame, L. T., Ui, M. & Cooper, D. M. F. (1985) *J. Biol. Chem.* **260**, 15946–15952
- Rapiejko, P. J., Northup, J. K., Evans, T., Brown, J. E. & Malbon, C. C. (1986) *Biochem. J.* **240**, 35–40
- Ros, M., Northup, J. K. & Malbon, C. C. (1988) *J. Biol. Chem.* **263**, 4362–4368
- Homburger, V., Brabet, P., Audigier, Y., Pantaloni, C., Bockaert, J. & Rouot, B. (1987) *Mol. Pharmacol.* **31**, 313–319
- Terashima, T., Katada, T., Oinuma, M., Inoue, Y. & Ui, M. (1987) *Brain Res.* **417**, 190–194
- Asano, T., Semba, R., Kamiya, N., Ogasawara, N. & Kato, K. (1988) *J. Neurochem.* **50**, 1164–1169
- Rodbell, M. (1964) *J. Biol. Chem.* **239**, 375–380
- Belsham, G. J., Denton, R. M. & Tanner, M. J. A. (1980) *Biochem. J.* **192**, 457–467
- Sternweis, P. C. & Robishaw, J. D. (1984) *J. Biol. Chem.* **259**, 13806–13813
- Toutant, M., Barhanin, J., Bockaert, J. & Rouot, B. (1988) *Biochem. J.* **254**, 405–409
- Lowry, O., Rosebrough, N., Farr, A. & Randall, I. (1951) *J. Biol. Chem.* **193**, 265–275
- Neer, E. J., Lok, J. M. & Wolf, L. (1984) *J. Biol. Chem.* **259**, 14222–14229
- Brabet, P., Dumuis, A., Sebben, M., Pantaloni, C., Bockaert, J. & Homburger, V. (1988) *J. Neurosci.* **8**, 701–708
- Chang, K. J., Pugh, W., Blanchard, S. G., McDermed, J. & Tam, J. P. (1988) *Proc. Natl. Acad. Sci. U.S.A.* **85**, 4929–4933
- Hinsch, K.-D., Rosenthal, W., Spicher, K., Binder, T., Gausepohl, H., Frank, R., Schultz, G. & Joost, H. G. (1988) *FEBS Lett.* **238**, 191–196
- Tanfin, Z., Milligan, G., Goureau, O., Unson, C. & Harbon, S. (1988) *Biol. Cell* **64**, 10a

Received 3 January 1989/6 March 1989; accepted 15 March 1989