Adrenergic regulation of formation of inositol phosphates in rat submandibular acini

Claire DOUGHNEY, Robert L. DORMER and Margaret A. McPHERSON*

Department of Medical Biochemistry, University of Wales College of Medicine, Heath Park, Cardiff CF4 4XN, U.K.

Formation of inositol phosphates in response to adrenergic secretagogues was studied in rat submandibular acini labelled with *myo*-[2-³H]inositol. Noradrenaline rapidly (within 5 s) increased radioactivity incorporated into inositol 1,4,5-trisphosphate and inositol tetrakisphosphate, with less rapid (within 1 min) increases in inositol 1,3,4-trisphosphate being observed. Inositol polyphosphate formation was less sensitive to noradrenaline than was stimulation of mucin secretion and was mediated by stimulation of α - but not β -adrenergic receptors. The β -adrenergic agonist isoproterenol, which is a potent stimulator of mucin secretion [McPherson & Dormer (1984) *Biochem. J.* **224**, 473–481] did not increase formation of inositol mono-, bis- or polyphosphates during a 15 min incubation. The results suggest that inositol phosphates do not mediate β -adrenergic stimulation of mucin secretion in rat submandibular acini. In addition, rat submandibular acinar cells contain a Ca^{2+} pool which can be mobilized by isoproterenol [McPherson & Dormer (1984) *Biochem. J.* **224**, 473–481], without involvement of inositol polyphosphates as second messengers.

INTRODUCTION

The rat submandibular gland is seromucous, consisting predominantly of mucous acinar cells (Scott & Pease, 1959) which synthesize and secrete high- M_r mucins. The glands are innervated by autonomic nerves and sympathetic neurotransmitters are potent stimulators of mucin secretion *in vivo* (Bogart, 1975; Martinez & Cassity, 1982) and *in vitro* (Bogart & Picarelli, 1978; Quissell & Barzen, 1980; McPherson & Dormer, 1984a).

Our studies using isolated acini preparations (McPherson & Dormer, 1984a) have shown that adrenergic agonists act directly on the acinar cells causing increased mucin secretion primarily by stimulation of β -adrenergic receptors. β -Adrenergic agonists increase cyclic AMP concentrations (Quissell *et al.*, 1981) and cell Ca^{2+} mobilization (McPherson & Dormer, 1984a); however, the mechanism by which mucin secretion is triggered is not clear.

In a variety of secretory cells, stimulation by agonists results in phosphatidylinositol phosphate breakdown and formation of water-soluble inositol phosphates (Berridge & Irvine, 1984; Michell *et al.*, 1984). In exocrine pancreatic (Streb *et al.*, 1983) and parotid (Aub & Putney, 1985) acinar cells, $\text{Ins}(1,4,5)\text{P}_3$ has been proposed as a second messenger mediating cell Ca^{2+} mobilization and the triggering of enzyme secretion in response to cholinergic stimulation.

In the present study we have investigated whether adrenergic neurotransmitters increase formation of inositol phosphates in rat submandibular acini and whether these metabolites are involved in mediating increases in mucin secretion and Ca^{2+} mobilization observed in response to β -adrenergic stimulation (McPherson & Dormer, 1984a). A preliminary account of some of this work has been reported (Doughney *et al.*, 1986).

EXPERIMENTAL

Materials

myo-[2-³H]inositol (16.3 Ci/mmol) and [³H] $\text{Ins}(1,4,5)\text{P}_3$ (1 Ci/mmol) were purchased from Amersham International. AG 1-X8 formate form ion exchange resin was from Bio-Rad, U.K.

The following were from Sigma: soybean trypsin inhibitor (type 1-S), hyaluronidase (type 1), L-isoproterenol (isoprenaline) hydrochloride, L-noradrenaline (dissolved initially in 0.1 M-HCl), DL-propranolol. Phenolamine mesylate [2-*N*-(3-hydroxyphenyl)-*p*-toluidinomethyl-2-imidazoline mesylate; Rogitine] was from Ciba, Horsham, Sussex, U.K. Collagenase was from Millipore (Worthington type II). Bovine serum albumin (fraction V) was from Miles Laboratories, Stoke Poges, Bucks., U.K. and Eagle's minimal medium (MEM) essential amino acid supplement was from Gibco. Scintillation fluid was Liquiscint (National Diagnostics, Somerville, NJ, U.S.A.). All other reagents, from BDH, were of the highest purity available.

Preparation of submandibular acini

Submandibular glands were obtained from two male Wistar rats (250–280 g) that had been starved overnight and killed by stunning and decapitation. Acini were prepared as described (McPherson *et al.*, 1983; McPherson & Dormer, 1984a). Briefly, glands were injected with collagenase solution [Ca^{2+} - and Mg^{2+} -free Hank's balanced salt solution (Gibco) containing collagenase (Worthington II) (64 units/ml) and hyaluronidase (500 units/ml)], cut into fragments and incubated for 60 min at 37 °C with shaking at 120 cycles/min. Tissue was mechanically dispersed by pipetting; acini were filtered through cheese cloth and purified by centrifugation through KHB, pH 7.4, containing 4% (w/v) bovine serum albumin.

Abbreviations used: KHB, Krebs–Henseleit bicarbonate buffer; InsP , InsP_2 , InsP_3 and InsP_4 , inositol mono-, bis-, tris- and tetrakis-phosphates respectively; $\text{Ins}(1,4,5)\text{P}_3$ and $\text{Ins}(1,3,4)\text{P}_3$, isomers of inositol trisphosphate where positions of phosphate groups assigned.

* To whom correspondence and reprint requests should be addressed.

Fig. 1. Separation of $\text{Ins}(1,4,5)\text{P}_3$ and $\text{Ins}(1,3,4)\text{P}_3$ by anion-exchange h.p.l.c.

Duplicate, neutralized HClO_4 extracts of submandibular acini, stimulated for 15 min by $100 \mu\text{M}$ -noradrenaline, were chromatographed on a Partisil SAX-10 column as described in the Experimental section. ●, Extract with no addition; □, extract + 20000 d.p.m. of D-myo-[2- ^3H]Ins(1,4,5) P_3 (Amersham International); ---, ammonium formate gradient.

Labelling with [^3H]inositol

Acini (approx. $3 \times 10^6/\text{ml}$) were washed and incubated in 3 ml of KHB, pH 7.4, containing Eagle's minimal medium (MEM) essential amino acids, 1.2 mM-glucose, soybean trypsin inhibitor (0.01%), 2% (w/v) bovine serum albumin, 10 mM-inosine, 0.4 mM-adenosine and myo-[2- ^3H]inositol ($5\text{--}33 \mu\text{Ci}/\text{ml}$) for 60 min at 37°C with shaking at 60 cycles/min.

They were then centrifuged and resuspended in KHB with 1% (w/v) bovine serum albumin + 10 mM-LiCl for 10 min before addition of agonists for the times stated in the Figure legends. [^3H]inositol phosphates were extracted by precipitating acini with a final concentration of 3.6% (w/v) HClO_4 on ice for 20 min and centrifuging (10 s in an Eppendorf Microfuge). Supernatants were removed and either adjusted to pH 8–9 by addition of an equal volume of 0.5 M-KOH/9 mM-sodium tetraborate/1.9 mM-EDTA/3.8 mM-NaOH for anion-exchange chromatography on AG 1-X8 formate resin or neutralized with 1 M- K_2HPO_4 for anion-exchange h.p.l.c.

Separation of [^3H]inositol phosphates

Anion-exchange chromatography on AG 1-X8 columns. The same procedure as that described by Berridge *et al.* (1983) was used. Extracts (equivalent to 1.5–4.1 mg of acinar cell protein) were applied to 1 ml columns of AG 1-X8 (formate form) and eluted sequentially with: (a) 20 ml of water, (b) 10 ml of 5 mM-disodium tetraborate/60 mM-sodium formate, (c) 10 ml of 0.2 M-ammonium formate in 0.1 M-formic acid, (d) 14 ml of 0.4 M-ammonium formate in 0.1 M-formic acid, (e) 10 ml of 1.0 M-ammonium formate in 0.1 M-formic acid; 2 ml fractions were collected of which 1 ml was assayed for ^3H by liquid-scintillation counting (Packard Tri-Carb 4430 counter). Counting efficiency was approx. 60%; quench corrections were carried out by use of an external

standard. A representative elution profile is shown in Doughney *et al.* (1986). The fraction eluted with 1.0 M-ammonium formate (e) contains InsP_3 and InsP_4 (Batty *et al.*, 1985). Our results show that this fraction, which is designated InsP_3 , contains 79–87% InsP_3 and is a mixture of $\text{Ins}(1,4,5)\text{P}_3$ and $\text{Ins}(1,3,4)\text{P}_3$ (Doughney *et al.*, 1986; Fig. 3).

Separation of [^3H]Ins P_3 and [^3H]Ins P_4 . Extracts (equivalent to 0.79–2.04 mg of acinar cell protein) were applied to a Zorbax SAX anion-exchange h.p.l.c. column (25 cm $\times$ 4.6 mm, packed by Dupont Instruments). The column was equilibrated with water and eluted at a flow rate of 1.25 ml/min. Water was eluted for 6 min, followed by a linear gradient to 1.1 M-ammonium formate (adjusted to pH 3.7 with orthophosphoric acid) over the next 24 min. After 5 min at 1.1 M-formate, the eluant was returned to water. Fractions (1.25 ml) were collected and 1 ml was counted for ^3H as described above. The separation method was checked by use of [^3H]Ins P_3 and [^3H]Ins P_4 standards prepared from submandibular acini stimulated for 15 min with $100 \mu\text{M}$ -noradrenaline. Extracts were subjected to AG 1-X8 anion exchange chromatography as described above, except that after elution of InsP_2 (d), InsP_3 was eluted with 0.8 M-ammonium formate in 0.1 M-formic acid and InsP_4 was eluted with 1.0 M-ammonium formate in 0.1 M-formic acid as described by Batty *et al.* (1985). The labelled fractions were lyophilized, redissolved in 1.5 ml of water, spiked with nucleotide markers (ATP, ADP and AMP; 50 nmol each) and applied to the Zorbax SAX column. The mean elution times for ATP, [^3H]Ins P_3 and [^3H]Ins P_4 were 18.3 ± 0.06 min ($n = 37$); 17.2 ± 0.08 min ($n = 36$) and 22.1 ± 0.12 min ($n = 36$) respectively. In addition, when duplicate crude extracts were run in the presence or absence of [^3H]Ins(1,4,5) P_3 (Amersham), this standard eluted at the same time as [^3H]Ins P_3 . The InsP_4 (see

Table 1. Formation of [³H]inositol phosphates in response to adrenergic agonists and antagonists in rat submandibular acini

Acini were labelled with [³H]inositol, incubated under the conditions shown for 15 min at 37 °C and [³H]inositol phosphates were separated on AG1-X8 formate as described in the Experimental section and as shown in Doughney *et al.* (1986). Results are means $\pm$ S.E.M. for the number of observations (in parentheses) from at least five experiments. Significance was determined by unpaired Student's *t*-test: **P* < 0.001, ***P* < 0.02 for difference from control; †*P* < 0.01 for difference from Ins P formation in the presence of noradrenaline alone; ††*P* < 0.001 for difference from Ins P_3 formation in the presence of noradrenaline alone.

Additions	Radioactivity (d.p.m./flask)		
	Ins P	Ins P_2	Ins P_3
None	30246 $\pm$ 2271 (10)	10380 $\pm$ 1309 (10)	491 $\pm$ 48 (11)
Isoproterenol (10 μ M)	30945 $\pm$ 4726 (5)	11627 $\pm$ 2451 (5)	572 $\pm$ 76 (6)
Noradrenaline (10 μ M)	38917 $\pm$ 2375 (11)**	13981 $\pm$ 1448 (11)	1590 $\pm$ 172 (11)*
Noradrenaline (10 μ M) + propranolol (30 μ M)	38090 $\pm$ 1622 (5)**	10467 $\pm$ 1245 (5)	1426 $\pm$ 209 (5)*
Noradrenaline (10 μ M) + phentolamine (30 μ M)	30231 $\pm$ 1392 (5)†	9482 $\pm$ 211 (5)	409 $\pm$ 23 (5)††

Fig. 3) fraction has been further characterized by Batty *et al.* (1985), who suggested that it is most likely to be Ins(1,3,4,5) P_4 .

Separation of Ins(1,4,5) P_3 and Ins(1,3,4) P_3 . Neutralized extracts (equivalent to 0.74–1.26 mg of acinar cell protein) were applied to a Partisil SAX 10 anion-exchange h.p.l.c. column (25 cm $\times$ 4.6 mm, packed by Technicol) equilibrated with water and eluted at a flow rate of 1.25 ml/min. A linear gradient to 0.66 M-ammonium formate (adjusted to pH 3.7 with orthophosphoric acid) was eluted in 5 min and this concentration was held for a further 2 min. As shown in Fig. 1, [³H]Ins(1,3,4) P_3 and [³H]Ins(1,4,5) P_3 were then eluted by a linear gradient up to 1.1 M-ammonium formate/phosphoric acid over 12 min. Fractions (1.25 ml) were collected until just before elution of ATP, when 0.25 ml fractions were collected. Separation of the isomers was checked using ATP as a marker co-eluting with Ins(1,3,4) P_3 (Irvine *et al.*, 1985), and [³H]Ins(1,4,5) P_3 (Amersham). The mean elution times for ATP, [³H]Ins(1,3,4) P_3 and [³H]Ins(1,4,5) P_3 were 12.3 $\pm$ 0.03 min (*n* = 30), 12.5 $\pm$ 0.04 min (*n* = 31) and 14.0 $\pm$ 0.07 min (*n* = 31) respectively.

RESULTS

Actions of adrenergic agonists and antagonists on formation of [³H]inositol phosphates

Actions of isoproterenol and noradrenaline on formation of radiolabelled inositol phosphates in rat submandibular acini are shown in Table 1. Inositol phosphates, Ins P , Ins P_2 and Ins P_3 were separated using AG 1-X8 formate as described in the Experimental section. Data from at least five experiments, which showed similar elution profiles to that shown in Doughney *et al.* (1986), were combined.

Table 1 shows that isoproterenol at a concentration (10 μ M) which gives maximum stimulation of mucin release in rat submandibular acini (McPherson & Dormer, 1984a) did not significantly increase formation of Ins P , Ins P_2 or Ins P_3 during 15 min incubation. By contrast, noradrenaline (10 μ M) caused a small but significant increase in [³H]Ins P formation and a marked (approx. 3-fold) increase in [³H]Ins P_3 formation after 15 min. At this time approx. 70% of the radioactivity of the Ins P_3 fraction was in Ins(1,3,4) P_3 and 17% in

Ins(1,4,5) P_3 . Stimulation of Ins P_3 formation by noradrenaline was not inhibited by the β -adrenergic blocker propranolol at a concentration (30 μ M) which markedly inhibits noradrenaline stimulation of mucin release (McPherson & Dormer, 1984a). The α -adrenergic blocker phentolamine, by contrast, completely inhibited increases in Ins P and Ins P_3 formation in response to noradrenaline (Table 1) and does not affect noradrenaline stimulation of mucin release (McPherson & Dormer, 1984a).

Dose-dependence of [³H]Ins P_3 formation in response to noradrenaline

Fig. 2 shows the dose-dependence of noradrenaline stimulation of formation of inositol phosphates, expressed as a percentage of control values. Formation of [³H]Ins P_3 was less sensitive to noradrenaline than was stimulation of mucin secretion. Thus, 10 μ M-noradrenaline gave maximum stimulation of mucin release but did not maximally stimulate Ins P_3 formation. Moreover, 1 μ M-noradrenaline increased mucin secretion to approx. 67% of maximum, whereas no detectable increase in Ins P_3 was observed (Fig. 2).

Time-course of formation of Ins(1,4,5) P_3 , Ins(1,3,4) P_3 and Ins P_4 in response to noradrenaline

The time course of formation of Ins(1,4,5) P_3 , Ins(1,3,4) P_3 and Ins P_4 in response to noradrenaline is shown in Fig. 3. Noradrenaline (100 μ M) caused rapid (within 5 s) increases in formation of Ins(1,4,5) P_3 and Ins P_4 . Increased formation of Ins(1,3,4) P_3 was less rapid with no significant increase being observed 10 s after stimulation. After 1 min, Ins(1,3,4) P_3 formation was increased markedly and continued to rise during 15 min stimulation with noradrenaline (Fig. 3), at which time radioactivity incorporated into Ins(1,3,4) P_3 was much greater than that incorporated into the other inositol polyphosphates formed. Isoproterenol (10 μ M) did not significantly increase formation of inositol polyphosphates after 15 min incubation (Table 1) when increases in calcium mobilization in response to this agonist are seen (McPherson & Dormer, 1984a). To determine whether a transient formation of an inositol polyphosphate could give rise to this increased calcium mobilization, we have examined the actions of isoproterenol at shorter times of incubation. In two experiments in which Ins P_3 and Ins P_4 were separated by Zorbax SAX h.p.l.c.,

Fig. 2. Dose-dependence of noradrenaline stimulation of inositol phosphate formation

Acini, prelabelled with [^3H]inositol ($5\text{ }\mu\text{Ci/ml}$) for 60 min, were incubated for 15 min in the presence of various concentrations of noradrenaline. [^3H]inositol phosphates were separated on AG1-X8 formate as described in the Experimental section. The trace showing dose-dependence of mucin release in response to noradrenaline (----) is from McPherson & Dormer (1984a) and is drawn in for comparison. $\circ$, [^3H]InsP₃; $\bullet$, [^3H]InsP₂; $\blacktriangle$, [^3H]InsP. Results, which are means $\pm$ S.E.M. for three experiments, are expressed as a percentage of radioactivity in inositol phosphates from unstimulated acini (range: 13820–53746 d.p.m./flask for InsP, 6136–21436 d.p.m./flask for InsP₂ and 270–766 d.p.m./flask for InsP₃).

incorporation of radioactivity into [^3H]InsP₃ was 98% of control after 10 s and 86% of control after 1 min incubation with isoproterenol ($10\text{ }\mu\text{M}$). [^3H]InsP₄ formation was 89% of control after 10 s and 85% of control after 1 min incubation with isoproterenol ($10\text{ }\mu\text{M}$). Separation of the isomers Ins(1,4,5)P₃ and Ins(1,3,4)P₄ (using Partisil SAX h.p.l.c.) revealed that isoproterenol ($10\text{ }\mu\text{M}$) also did not alter the ratio of these isomers after a short (1 min) incubation time (results not shown). Thus, no increases in formation of either Ins(1,4,5)P₃, Ins(1,3,4)P₃ or InsP₄ were observed in response to isoproterenol ($10\text{ }\mu\text{M}$).

DISCUSSION

In rat submandibular acini, we have previously shown that isoproterenol and noradrenaline increase mucin secretion by stimulation of β - but not α -adrenergic receptors (McPherson & Dormer, 1984a). In the present study our results show that β -adrenergic stimulation of acini induced by either isoproterenol or noradrenaline in the presence of an α -adrenergic blocker was ineffective in increasing formation of inositol mono-, bis- or polyphosphates.

The α -adrenergic action of noradrenaline caused marked increases in inositol polyphosphate formation after 15 min, which mainly reflected increases in Ins(1,3,4)P₃. The result is similar to that in rat parotid glands in which Irvine *et al.* (1984) first demonstrated the presence of two InsP₃ isomers and showed that after 15 min incubation with carbamoylcholine, Ins(1,3,4)P₃ was the predominant isomer formed. Ins(1,4,5)P₃ and InsP₄ were produced rapidly (5 s) in response to noradrenaline in submandibular acini with a lag before Ins(1,3,4)P₃ formation increased. This pattern is similar

to that of cholinergic stimulation in cerebral cortex (Batty *et al.*, 1985) and exocrine pancreas (C. Doughney, R. L. Dormer & M. A. McPherson, unpublished). It is consistent with initial formation of Ins(1,4,5)P₃, phosphorylation to Ins(1,3,4,5)P₄ and subsequent dephosphorylation to Ins(1,3,4)P₃, as has been suggested for other tissues (Batty *et al.*, 1985; Irvine *et al.*, 1986). In rat submandibular acini slightly more radioactivity was incorporated into Ins(1,3,4)P₃ than into Ins(1,4,5)P₃ under control conditions. This is similar to exocrine pancreas (C. Doughney, R. L. Dormer & M. A. McPherson, unpublished work) but different from parotid in which slightly more radioactivity was seen in Ins(1,4,5)P₃ (Irvine *et al.*, 1985) and markedly different from cerebral cortex in which approx. 96% of radioactivity incorporated into InsP₃ was into Ins(1,4,5)P₃. This may reflect differences in metabolism of inositol phosphates and does not rule out the possibility that a separate pathway of Ins(1,3,4)P₃ formation might exist in some tissues. Our results suggest that increased formation of inositol polyphosphates in response to noradrenaline is not involved in triggering mucin secretion. Ins(1,4,5)P₃ and InsP₄ are formed rapidly enough to mediate α -adrenergic stimulation of K⁺ efflux induced by noradrenaline (Quissell & Barzen, 1980) suggesting a possible role in control of secretion of submandibular fluid. Similarly, in rat parotid glands α -adrenergic stimulation increases fluid secretion (Batzri *et al.*, 1973; Schramm & Selinger, 1975) and formation of inositol phosphates (Aub & Putney, 1985), whereas β -adrenergic stimulation increases only amylase release (Schramm & Selinger, 1975), with no effect on inositide turnover (Aub & Putney, 1985).

Ins(1,4,5)P₃ has been implicated in causing cell Ca²⁺ mobilization in response to carbamoylcholine in rat parotid glands (Aub & Putney, 1985) and exocrine

Fig. 3. Time course of formation of $\text{Ins}(1,4,5)\text{P}_3$, InsP_4 and $\text{Ins}(1,3,4)\text{P}_3$ in submandibular acini stimulated with noradrenaline

Acini prelabelled with [^3H]inositol ($33 \mu\text{Ci}/\text{ml}$ for 60 min) were incubated for various times in the presence or absence of $100 \mu\text{M}$ -noradrenaline and neutralized HClO_4 extracts were subjected to anion-exchange h.p.l.c. as described in the Experimental section. Results are means $\pm$ S.E.M. from three experiments, with duplicate incubations in each experiment. $\bullet$, [^3H]Ins(1,3,4) P_3 ; $\blacksquare$, [^3H]Ins(1,4,5) P_3 ; $\blacktriangle$, [^3H]Ins P_4 .

pancreas (Streb *et al.*, 1983). In rat pancreatic acini there is good evidence that carbamoylcholine mobilizes Ca^{2+} from the rough endoplasmic reticulum, giving rise to increases in cytosolic free Ca^{2+} concentration which trigger amylase secretion (Richardson & Dormer, 1984; Richardson *et al.*, 1984). Isoproterenol mobilizes Ca^{2+} and stimulates protein secretion in submandibular (McPherson & Dormer, 1984a; McPherson *et al.*, 1985) and parotid (Mangos *et al.*, 1975; Kanagasuntheram & Randle, 1976; McPherson & Hales, 1978; Butcher, 1980) acinar cells but does not change InsP_3 formation (this paper; Aub & Putney, 1985). In submandibular acini our results indicate that isoproterenol mobilizes Ca^{2+} from a slowly exchanging intracellular pool (McPherson & Dormer, 1984a). Thus, in these cells, where there is evidence that Ca^{2+} is a modulator rather than a primary trigger for secretion (Quissell *et al.*, 1981; McPherson & Dormer, 1984a,b), β -adrenergic agonists can mobilize an intracellular Ca^{2+} pool without involvement of an

inositol polyphosphate as a second messenger. Furthermore, increased formation of metabolites of phospholipase C or phospholipase A_2 (Bradbury & McPherson, 1985) do not mediate β -adrenergic secretory responses in submandibular acinar cells. The role of rapid increases in formation of inositol polyphosphates in response to the physiological neurotransmitter noradrenaline remains to be elucidated.

This work was supported by the Cystic Fibrosis Research Trust, U.K.

REFERENCES

- Aub, D. L. & Putney, J. W. (1985) *Biochem. J.* **225**, 263–266
- Batty, I. R., Nahorski, S. R. & Irvine, R. F. (1985) *Biochem. J.* **232**, 211–215
- Batzri, S., Selinger, Z., Schramm, M. & Robinovitch, M. R. (1973) *J. Biol. Chem.* **248**, 361–368
- Berridge, M. J. & Irvine, R. F. (1984) *Nature (London)* **312**, 315–321
- Berridge, M. J., Dawson, R. M. C., Downes, C. P., Heslop, J. P. & Irvine, R. F. (1983) *Biochem. J.* **212**, 473–482
- Bogart, B. I. (1975) *J. Ultrastruct. Res.* **52**, 139–155
- Bogart, B. I. & Picarelli, J. (1978) *Am. J. Physiol.* **235**, C256–C268
- Bradbury, N. A. & McPherson, M. A. (1985) *Biochem. Soc. Trans.* **13**, 1174–1175
- Butcher, F. R. (1980) *Biochim. Biophys. Acta* **630**, 254–260
- Doughney, C., Dormer, R. L. & McPherson, M. A. (1986) *Biochem. Soc. Trans.* **14**, 1025–1026
- Irvine, R. F., Letcher, A. J., Lander, D. J. & Downes, C. P. (1984) *Biochem. J.* **223**, 237–243
- Irvine, R. F., Anggard, E. E., Letcher, A. J. & Downes, C. P. (1985) *Biochem. J.* **229**, 505–511
- Irvine, R. F., Letcher, A. J., Heslop, J. P. & Berridge, M. J. (1986) *Nature (London)* **320**, 631–634
- Kanagasuntheram, P. & Randle, P. J. (1976) *Biochem. J.* **160**, 547–564
- Mangos, J. A., McSherry, N. R., Barber, T., Arvanitakis, S. N. & Wagner, V. (1975) *Am. J. Physiol.* **299**, 560–565
- Martinez, J. R. & Cassity, N. (1982) *Arch. Oral Biol.* **27**, 159–166
- McPherson, M. A. & Dormer, R. L. (1984a) *Biochem. J.* **224**, 473–481
- McPherson, M. A. & Dormer, R. L. (1984b) *Biochem. Soc. Trans.* **12**, 1091–1092
- McPherson, M. A. & Hales, C. N. (1978) *Biochem. J.* **176**, 855–863
- McPherson, M. A., Dodge, J. A. & Goodchild, M. C. (1983) *Clin. Chim. Acta* **135**, 181–188
- McPherson, M. A., Dormer, R. L., Dodge, J. A. & Goodchild, M. C. (1985) *Clin. Chim. Acta* **148**, 229–237
- Michell, R. H., Hawkins, P. T., Palmer, S. & Kirk, C. J. (1984) in *Calcium Regulation in Biological Systems* (Ebashi, S., *et al.*, eds.), pp. 85–103, Academic Press, New York
- Quissell, D. O. & Barzen, K. A. (1980) *Am. J. Physiol.* **238**, C99–C106
- Quissell, D. O., Barzen, K. A. & Lafferty, J. L. (1981) *Am. J. Physiol.* **241**, C76–C85
- Richardson, A. E. & Dormer, R. L. (1984) *Biochem. J.* **219**, 679–685
- Richardson, A. E., Brown, G. R. & Dormer, R. L. (1984) *Biochem. Soc. Trans.* **12**, 1066–1067
- Schramm, M. & Selinger, Z. (1975) *J. Cyclic Nucleotide Res.* **1**, 181–192
- Scott, B. L. & Pease, D. C. (1959) *Am. J. Anat.* **104**, 115–162
- Streb, H., Irvine, R. F., Berridge, M. J. & Schulz, I. (1983) *Nature (London)* **306**, 67–69