REVIEW ARTICLE

Carbohydrates as antigenic determinants of glycoproteins

Ten FEIZI and Robert A. CHILDS

Applied Immunochemistry Research Group, Medical Research Council, Clinical Research Centre, Watford Road, Harrow, Middlesex HA1 3UJ, U.K.

Introduction

Antibodies are being used increasingly as reagents in biochemical and biological studies of glycoproteins, for example, in the search for glycoproteins with important roles in embryonic development, cell differentiation and oncogenesis, and as a means of monitoring the biosynthesis and subcellular location of specific glycoproteins such as enzymes and receptors. Crucial for the proper application of antibodies in such experiments is knowledge of the determinants they recognize. A glycoprotein may contain many antigenic determinants, each consisting of up to seven monosaccharides or amino acids in consecutive sequence (Kabat, 1976) or, in the case of polypeptides, in non-continuous sequence brought together by chain folding (Lerner, 1984; Walter, 1986). Unravelling the specificities contained in conventional (polyclonal) antisera raised against glycoproteins often presents problems due to the mixture of antibodies present with different combining specificities. It is often difficult to distinguish specificities directed against peptide determinants from those directed against carbohydrate structures on a glycoprotein. Monoclonal antibodies have the advantage in this respect since each is a single antibody population with a narrow specificity. As a result of work with naturally-occurring monoclonal autoantibodies and the widespread use of hybridoma-derived antibodies, there has been an increased awareness of carbohydrates as antigenic determinants of glycoproteins, since it has become apparent that many differentiation antigens of normal and neoplastic cells are oligosaccharide determinants of glycoproteins and glycolipids rather than peptide determinants (Feizi, 1985, 1987; Hakomori, 1985). Thus, carbohydrate antigens which were previously the interest of specialist immunochemists are now of wide relevance to both biochemists and cell biologists.

We do not intend here to make an exhaustive review of antigens found to date on glycoprotein oligosaccharides; rather we shall discuss some examples of whole cells (haematopoietic and embryonic) and specific glycoproteins (a receptor and a glycosyltransferase) where carbohydrate structures have been found to be major antigenic determinants. We shall also discuss approaches to the study of antigenicity (reactivity with antibodies) and immunogenicity (ability to elicit antibodies) of the oligosaccharides of glycoproteins and the opportunities now available for exploiting antibodies with defined carbohydrate specificities as probes of the structure, tissue distribution and function of glycoprotein oligosaccharides. We also include in this article a hypothesis which considers the biological relevance and possible inter-relationships of three areas of knowledge: (a) the occurrence of the same carbohydrate antigens/structures on diverse and unrelated glycoproteins, (b) marked

changes in glycoprotein oligosaccharide antigens/structures during embryonic development, differentiation and oncogenesis and (c) the occurrence of an array of carbohydrate-binding proteins (lectins) in mammalian cells.

Glycoprotein oligosaccharides as major antigens of whole cells

Antibodies have been used extensively to monitor changes in antigenicity at the cell surface during embryogenesis and cellular differentiation. Two cell types which have been studied extensively are cultured teratocarcinoma cell lines which can be chemically induced to undergo changes that mimic early embryonic events, and a promyelocytic cell line HL60 which can be induced to differentiate along either the granulocytic or monocytic pathways.

Work with syngeneic polyclonal antisera to the mouse teratocarcinoma cell line, F9, showed that early embryonic cells express an antigen, termed F9 antigen (Jacob, 1979). The structures of the F9 antigenic determinant(s) were not established; however, it was shown that undifferentiated teratocarcinoma cells and embryonic cells are rich in glycoproteins with long carbohydrate chains of poly-*N*-acetyl-lactosamine type which react with antibodies to F9 antigen (Muramatsu *et al.*, 1979). Work with monoclonal antibodies has shown that several carbohydrate antigens which change dramatically at successive stages of differentiation are borne on poly-*N*-acetyl-lactosamine chains of poly-disperse, high-*M_r* glycoproteins (Childs *et al.*, 1983). Among these antigens are the I antigenic determinants associated with branched poly-*N*-acetyl-lactosamine structures (Feizi *et al.*, 1971; Watanabe *et al.*, 1979) which are expressed throughout the pre-implantation stage (Kapadia *et al.*, 1981), SSEA-1 antigen associated with α 1-3 fucosylated *N*-acetyl-lactosamine (Gooi *et al.*, 1981) which first appears at the 8-cell stage of the mouse embryo when compaction takes place (Solter & Knowles, 1978) and the i antigen associated with unbranched poly-*N*-acetyl-lactosamine structures (Niemann *et al.*, 1978) which first appears at the onset of differentiation of the primary endoderm (Kapadia *et al.*, 1981), as shown in Fig. 1. These antigens diminish or disappear at later stages of development and become restricted to only certain cell types in adult tissues.

An important aspect of changes in expression of these oligosaccharide antigens is the masking effect of additional glycosylations (Feizi, 1981*a,b*) or other modifications such as sulphation (Mehmet *et al.*, 1986; Tang *et al.*, 1986). Thus, just as an antigen may be formed by each new substitution, the antigenicity of the underlying structure may be masked. Each sugar linkage is formed by a specific glycosyltransferase; thus,

Antibodies	Structures recognized	Immunofluorescence of embryonic cells		
		1-4-cell stage	8-cell stage	Primary endoderm
anti-I Ma	Gal β 1-4GlcNAc β 1-6Gal/GalNAc	+	+	+
anti-I Step	Gal β 1-4GlcNAc β 1-6Gal β 1-4GlcNAc	+	+	+
anti-SSEA-1	Gal β 1-4GlcNAc β 1-3Gal β 1-4GlcNAc... 1,3 Fuca	-	+	+
anti-i Den	Gal β 1-4GlcNAc β 1-3Gal β 1-4GlcNAc β 1-3Gal β 1-4GlcNAc...	-	-	+

Fig. 1. Changes in carbohydrate structures on the surface of mouse embryonic cells deduced from the sequential changes in their reactivities with carbohydrate sequence-specific monoclonal antibodies, anti-I Ma, anti-I Step, anti-SSEA-1 and anti-i Den

+, -, Reaction or lack of reaction, respectively, by immunofluorescence.

depending on the relative activities of these enzymes and the availabilities of their donor and acceptor substrates, the proportions of the various oligosaccharide structures and hence the antigenicities will vary. Conversely, incomplete biosynthesis or the actions of glycosidases may reveal cryptic antigens which, from the immunologist's point of view, behave as 'jumping' antigens, i.e. antigens which 'disappear' and 'reappear' at different stages of cell differentiation or cell transformation (Feizi, 1981b). This is important to bear in mind since the appearance of new antigens at the cell surface may not be synonymous with the biosynthesis of new membrane proteins, and loss of antigenicity may not mean lack of biosynthesis of the relevant oligosaccharides.

There are a number of analogies between the embryo/teratocarcinoma cells (Martin, 1980) and promyelocytic cells (Gahmberg *et al.*, 1979). A high proportion of monoclonal antibodies to the promyelocytic leukaemia cell line, HL60, recognize an antigen related to SSEA-1 (reviewed by Feizi, 1984, and Magnani, 1984). Here, also, diffusely migrating cell surface glycoproteins of high M_r (greater than 100 000) are major carriers of this antigen detected both by immunoblotting of total cell lysates and immunoprecipitation of radiolabelled cell surface glycoproteins (Feizi, 1983; Urdal *et al.*, 1983; Tetteroo *et al.*, 1984; Childs & Feizi, 1987). In addition, the i antigenic determinant, associated with linear poly-*N*-acetyl-lactosamine structures (i Den) is borne on these glycoproteins (Childs & Feizi, 1987, and Fig. 2). There is a lack of reactivity with anti-I Step (Fig. 2) which recognizes long, branched poly-*N*-acetyl-lactosamine structures (cf. Fig. 1). With anti-I Ma, which recognizes the short branch-point *N*-acetyl-lactosamine sequence, there is a strong reactivity with a different set of glycoproteins of average M_r 170 000. Thus, two major glycoprotein families are discerned on the surface of HL60 cells with respect to their sugar chain immunoreactivities, the first having linear poly-*N*-acetyl-lactosamine backbones and corresponding in structure to the *N*-glycosidically linked oligosaccharides (Mizoguchi *et al.*, 1984; Spooner *et al.*, 1984) and the second having short, branched backbones of the type occurring among the *O*-glycosidically linked oligosaccharides of the major sialoglycoprotein termed leukosialin (Carlsson *et al.*, 1986).

Among the peripheral blood cells of man, the SSEA-1

determinant is almost exclusively expressed on granulocytes (reviewed by Feizi, 1984, and Magnani, 1984). HL60 cells that have been induced to differentiate along the granulocyte pathway also express SSEA-1 (Tetteroo *et al.*, 1984). In comparison with undifferentiated HL60 cells, the total amount of glycoproteins carrying the SSEA-1 determinant is reduced, since immunoreactive glycoproteins are barely detectable by immunoblotting of total cell lysates (R. A. Childs, unpublished work). However, by immunoprecipitation of radioiodinated cell surface glycoproteins of granulocytes, components migrating at 250, 230, 150, 67 and 52 kDa are detectable (Tetteroo *et al.*, 1984). Biochemical studies have confirmed that α 1-3 fucosylated linear poly-*N*-acetyl-lactosamine structures persist in granulocytes, albeit in

Fig. 2. Expression of I, i, and SSEA-1 antigenic determinants on cell surface glycoproteins on HL60 cells

After treatment with neuraminidase and radiolabelling by galactose oxidase/ NaB^3H_4 (Gahmberg & Hakomori, 1973) the cells were lysed with the zwitterionic detergent Empigen, immune precipitates were obtained and subjected to SDS/polyacrylamide gel electrophoresis and autoradiography as described previously (Childs *et al.*, 1983; Childs & Feizi, 1987). Labelled bands appear white. Positions of M_r markers are indicated to the left. (R. A. Childs, unpublished work.)

smaller amounts than in promyelocytic (HL60) cells (Spooncer *et al.*, 1984; Mizoguchi *et al.*, 1984).

On monocytes, the SSEA-1 and i antigens cannot be detected, but are revealed after treatment of the cells with neuraminidase (Thorpe & Feizi, 1984). Reduced expression of SSEA-1 was also noted on HL60 cells which were induced to differentiate into monocytes, and the antigen could be revealed by neuraminidase treatment (Tetteroo *et al.*, 1984). This antigen is also revealed on acute lymphoblastic leukaemia cells after neuraminidase treatment (Stockinger *et al.*, 1984; Knapp *et al.*, 1984). Thus, antibodies to 3-fucosyl-*N*-acetyl-lactosamine have been useful as reagents in studies of the glycosylation changes during differentiation not only in mouse embryonic cells but also in human leucocytes.

There are species differences in the expression of several carbohydrate antigens. For example, human but not mouse granulocytes express SSEA-1 antigen (Knowles *et al.*, 1982; Thorpe & Feizi, 1984). Moreover, the SSEA-1 determinant is restricted to very few tissues in the adult mouse (Fox *et al.*, 1983) but this structure is widely distributed in human tissues and secreted glycoproteins (reviewed by Gooi *et al.*, 1981). In addition, there are cell-type differences in the relative proportions of glycoproteins and glycolipids that carry this antigen. For example, the SSEA-1, I and i antigens are expressed predominantly on glycoproteins of embryonal carcinoma cells (Childs *et al.*, 1983), but in myeloid cells they are expressed both on glycoproteins and glycolipids (Urdal *et al.*, 1983; Uemura *et al.*, 1985). Information so far available suggests that the Y antigen (2',3-difucosyl-*N*-acetyl-lactosamine) is predominantly expressed on glycolipids of myeloid cells since it was readily detected by immunostaining of glycolipids derived from granulocytes (Uemura *et al.*, 1985), but was barely detectable on glycoproteins of HL 60 cells by immunoblotting of total cell lysates (R. A. Childs, unpublished work).

These are just two differentiation systems in which glycoprotein oligosaccharides feature as major antigenic determinants of whole cells. They may be added to a growing list of carbohydrate antigens on cell surfaces headed by the ABO, Lewis and P blood group antigens of erythrocyte membranes (Watkins, 1980; Kabat, 1982) and the glycoprotein antigens of the normal and neoplastic epithelia of the gastrointestinal tract (Feizi *et al.*, 1984).

Oligosaccharides as major antigens of glycoproteins

The immunochemical approach is an important part of biochemical and biological studies of glycoprotein hormones, receptors and enzymes. It has long been assumed that the predominant specificities of antibodies raised against cell-bound and free glycoproteins are directed toward the protein moiety. We wish to challenge this assumption and cite here two examples of glycoproteins where carbohydrate structures have been shown to be important immunogenic components: these are the receptor for epidermal growth factor and the glycoprotein enzyme galactosyltransferase of human milk. In particular, we wish to discuss two aspects of this immunogenicity: firstly, that the carbohydrate antigenic determinants may be the basis of antigenic cross-reactions among glycoproteins that are distinct both functionally and at the level of their polypeptide structures, and, secondly, that important information

can be derived about the carbohydrate structures of glycoproteins by elucidating the specificities of antibodies to glycoproteins.

A431 cells are especially rich in receptors for epidermal growth factor. Numerous hybridoma antibodies have been raised to this receptor glycoprotein by immunizing mice with the cell membranes or the isolated receptor. As part of a programme of studies of carbohydrate antigens on biologically active glycoproteins, we have looked for carbohydrate specificities among some 20 hybridoma antibodies to the receptor glycoprotein raised in different laboratories and found evidence for blood-group-related carbohydrate specificities in no less than 15 (Gooi *et al.*, 1983b, 1985a,b). The immunogenicity of the carbohydrate moiety of this receptor glycoprotein is endorsed by the work of others (Parker *et al.*, 1984; Le Pendu *et al.*, 1985).

These observations prompted direct analyses of the carbohydrate antigens on the purified EGF receptor glycoprotein, using, in addition to the 'anti-receptor' antibodies, a panel of monoclonal antibodies directed at various members of the blood group family of oligosaccharide antigens (Childs *et al.*, 1984). In Fig. 3 are shown the structures of antigens detected on the backbone and peripheral domains of the carbohydrate chains of the receptor glycoprotein. These include several Type 1 (Gal β 1-3GlcNAc) and Type 2 (Gal β 1-4GlcNAc) based structures such as branched SSEA-1 and 2',3-difucosyl-*N*-acetyl-lactosamine (Y antigen structure) on poly-*N*-acetyl-lactosamine backbones, variants of blood group A antigen, and blood group Lewis^a also on poly-*N*-acetyl-lactosamine backbones. These structures are apparently on oligosaccharides that are *N*-glycosidically linked to the receptor protein, since no evidence has been obtained for the presence of *O*-linked structures on the epidermal growth factor receptor from this cell line (Mayes & Waterfield, 1984; Childs *et al.*, 1984; Cummings *et al.*, 1985).

Monoclonal antibodies have also been used in immunoblotting experiments (Childs *et al.*, 1986) to investigate the carbohydrate antigens of the glycoprotein enzyme, galactosyltransferase, purified in soluble form from human milk. The structures deduced to be present on the enzyme isolated from a presumed group A secretor and a group O non-secretor are summarized in Fig. 4. More recent studies using oligosaccharides released from galactosyltransferase indicate that the Le^a and SSEA-1 antigens occur both on *O*- and *N*-glycosidically linked chains of the enzyme glycoprotein (Tang & Feizi, 1987). These observations, in turn, prompted a study of blood-group-related carbohydrate specificities among rabbit antisera and affinity-purified antibodies to galactosyltransferase, and revealed that various antibody preparations contain anti-blood group A, anti-blood group HLe^b or anti-Le^a and anti-SSEA-1 related specificities as minor but non-trivial populations (Childs *et al.*, 1986). The importance of these specificities lies in the fact that they account for the immunofluorescence of sections of formalin-fixed, paraffin-embedded epithelial cells of the human gastrointestinal tract, immunofluorescence which has previously been assumed to be due exclusively to reactivity with galactosyltransferase. Absorption of these antibodies with reference blood group substances abolished the immunofluorescence both at the plasma membranes and the Golgi regions of these epithelial cell sections. Thus carbo-

Antibodies	Structures recognized	Designation of antigen
M39	Gal β 1-4GlcNAc	
3CIB12	Gal β 1-4GlcNAc β 1 1,3 Fuca	Branched SSEA-1 or X
C14	Gal β 1-4GlcNAc β 1-3Gal β 1-3/4GlcNAc 1,2 1,3 FucaFuca	Blood group Le ^b -related
TL5	GalNAc α 1-3Gal β 1-3/4GlcNAc 1,2 Fuca	Blood group A
G49	GalNAc α 1-3Gal β 1-3/4GlcNAc 1,2 1,4/3 Fuca Fuca	Blood group ALe ^{b/y}
CF4	Gal β 1-3GlcNAc β 1-3Gal β 1-3/4GlcNAc 1,4 Fuca	Blood group Le ^a -related

Fig. 3. Carbohydrate sequences on the backbone and peripheral regions of oligosaccharides of the epidermal growth factor deduced from immunoblotting experiments with six monoclonal antibodies

The occurrence of the C14 (Brown *et al.*, 1983) and CF4 (Young *et al.*, 1983) antigenic determinants on oligosaccharide backbones of poly-*N*-acetyl-lactosamine type was deduced from the loss of reactivities with the two antibodies after treatment with endo- β -galactosidase (adapted from Childs *et al.*, 1984).

Antibodies	Structures recognized	Designation of antigen	Source of galactosyltransferase	
			Blood group A secretor	Blood group O non-secretor
anti-SSEA-1	Gal β 1-4GlcNAc 1,3 Fuca	SSEA-1 or X antigen	+	+
CF4	Gal β 1-3GlcNAc 1,4 Fuca	Blood group Le ^a	+	+
G49	GalNAc α 1-3Gal β 1-3/4GlcNAc 1,2 1,4/3 Fuca Fuca	Blood group ALe ^{b/y}	+	—

Fig. 4. Carbohydrate sequences on human milk galactosyltransferase purified from a presumed blood group A secretor and a group O non-secretor, deduced from immunoblotting of the purified glycoproteins with monoclonal antibodies anti-SSEA-1, CF4 and G49 (Childs *et al.*, 1986)

hydrate specificities are a feature, not only among hybridoma antibodies but also among conventional antisera raised against glycoproteins. When such antisera are used for immunohistochemical studies, different glycoproteins known to express blood group antigens, e.g. mucins (Feizi *et al.*, 1984) and intestinal hydrolases (Triadou, 1984) as well as glycolipids are likely to react in addition to galactosyltransferase. As shown in Figs. 2 and 3 the epidermal growth factor receptor and galactosyltransferase cross-react with a monoclonal antibody (G49) with anti-blood-group-A-related specificity which was originally raised against the epidermal growth factor receptor. These two glycoproteins also cross-react when tested with the anti-Le^a antibody, CF4 (Figs. 2 and 3).

Antigenicities of blood-group-related carbohydrate structures of glycoproteins can be readily and convincingly demonstrated (Childs *et al.*, 1986) due to the availability of glycoproteins and tissues from individuals of known blood group and secretor status. However, as discussed in the following section, the conclusive demonstration of other types of carbohydrate specificities can be difficult, but their importance should not be underestimated. The generality of this problem is

reinforced by evidence that rabbit antibodies to two other enzyme glycoproteins, the acid hydrolases of *Dictyostelium discoideum* (Freeze *et al.*, 1984) and to a fungal glucosyltransferase (Pazur *et al.*, 1986) are predominantly directed at carbohydrate determinants of these glycoproteins. Our experience indicates that antisera to glycoproteins should not be used for subcellular localization studies unless the presence of carbohydrate specificities has been ruled out, or any carbohydrate specificities present are proven to be unique to the glycoproteins in question.

These are just a few examples of the significance of glycoprotein oligosaccharides as antigens in the field of biochemistry. Other striking examples of a combined immunochemical/biochemical approach to the study of glycoprotein oligosaccharides are the discovery of the novel pentasaccharide sequence:

(blood group Sd^a determinant) on the *N*-linked chains of a well-known glycoprotein, the Tamm-Horsfall glyco-

protein (Donald *et al.*, 1984), the pentasaccharide structure:

among the *O*-linked chains of glycophorin A from Cad antigen-positive erythrocytes (Blanchard *et al.*, 1983), and the identification of chondroitin sulphate-related sequences as antigens of nerve adhesion molecules (Chou *et al.*, 1985; Kruse *et al.*, 1985) and as target antigens of autoantibodies associated with polyneuropathy (Kabat *et al.*, 1984). Another unexpected observation has been the detection by immunochemical means of a high- M_r component rich in *i* and sialosyl-*i* sequences in preparations of human α_1 -acid glycoprotein (R. A. Childs, K. Schmid & T. Feizi, cited by Childs & Feizi, 1987). Purified preparations of this glycoprotein express *i* antigen activity which is enhanced after desialylation. Affinity chromatography of the asialoglycoprotein on an anti-*i* column has shown that the *i* activity is associated with a minor component in preparations of α_1 -acid glycoprotein, representing approx. 1% of the carbohydrate moiety; immunoblotting experiments of the unretained and the specifically eluted *i*-active fraction after SDS/polyacrylamide-gel electrophoresis have revealed that the *i*-antigen activity is not associated with α_1 -acid glycoprotein (M_r 43000), but it is borne on diffusely migrating high- M_r glycoproteins (M_r > 100000) that do not immunostain with antibodies to α_1 -acid glycoprotein. We have raised the possibility that some of the biological properties of α_1 -acid glycoprotein may reside in this hitherto unsuspected component. The immunochemical approach has also lead to the finding of several fuco-oligosaccharide structures of the blood-group-oncodevelopmental family in human seminal fluid glycoproteins (Hanisch *et al.*, 1985).

Although all glycoprotein oligosaccharide sequences are potential antigens, the antigenic determinants characterized thus far largely involve the backbone and peripheral domains common to *O*- and *N*-linked chains (Feizi *et al.*, 1984) and the core regions of *O*-linked chains (Rahman & Longenecker, 1982; Steuden *et al.*, 1985; Hirohashi *et al.*, 1985). Since neoglycolipid constructs are of proven immunogenicity (Lai & Kabat, 1985), it should be possible to raise monoclonal antibodies to desired neoglycolipid constructs (Tang & Feizi, 1987) or other oligosaccharide structures, such as various core-region variants of *N*-linked oligosaccharides (Kobata, 1984) and the distinctive oligosaccharide structures that occur on lysosomal enzymes (Kornfeld & Kornfeld, 1985), pituitary hormones (Baenziger, 1985) and tumour derived glycoproteins (Kobata & Yamashita, 1984; Yamamoto *et al.*, 1985). Such antibodies would be invaluable for mapping the distribution of specific structures on purified glycoproteins and within cells.

The problem of determining whether antibodies have carbohydrate specificities

There is as yet no simple way of knowing if an antibody is directed at the carbohydrate or peptide moiety of a glycoprotein with which it reacts. Reduction or abolition of immunoreactivity of whole cells or purified glycoproteins following treatment with proteinases does not necessarily reflect anti-protein specificity; rather, it may reflect a loss of immunoreactive glyco-

peptides from the cell surface or the loss of multivalent expression of a carbohydrate determinant. Retention of antigenicity following heat treatment (100 °C for 5 min) in the presence of SDS as, for example, under the conditions one may use for immunoblotting, would favour a carbohydrate rather than protein epitope. However, there are examples where peptide determinants have survived such treatments, for example, the determinant on the β -galactoside-binding protein of bovine tissues recognized by a hybridoma antibody 36/8 (Carding *et al.*, 1984). Traditionally, susceptibility to mild periodate treatment is considered indicative of the carbohydrate nature of antigens. A mild periodate oxidation procedure was standardized recently for destroying monosaccharides at the non-reducing termini of antigenically-active carbohydrate chains (Woodward *et al.*, 1985). Using this procedure, the fucose-dependent Le^a antigen was destroyed without altering the protein epitopes on bovine serum albumin. As discussed in that report, carbohydrate determinants located on non-terminal domains of carbohydrate chains are unaffected by this procedure. We have observed that under the conditions used by Woodward *et al.* (1985) the linear poly-*N*-acetyl-lactosamine determinant recognised by an anti-*i* antibody (Tho) and the branched poly-*N*-acetyl-lactosamine determinant recognised by anti-I (Step) that are expressed on meconium glycoproteins and sheep gastric mucins, respectively, are not destroyed, whereas SSEA-1, *i* (Den), on meconium, and I (Ma), on sheep gastric mucins, are destroyed (P. W. Tang & T. Feizi, unpublished work). Both the anti-*i* Tho (Tang *et al.*, 1986) and anti-I Step (Gardas, 1982; Uemura *et al.*, 1983) can react with internal domains of poly-*N*-acetyl-lactosamine chains and do not require terminal galactose residues to be present. In contrast, anti-SSEA-1 has a strict requirement for α 1-3-linked terminal fucose (Gooi *et al.*, 1981; Hounsell *et al.*, 1981) and anti-I Ma and anti-*i* Den require terminal galactose residues (Feizi *et al.*, 1971; Tang *et al.*, 1986). Moreover, there remains a question mark as to whether peptide determinants are always resistant to mild periodate treatment, since the Rh(D) antigen of erythrocytes, now thought to be of polypeptide nature (Moore *et al.*, 1982), was susceptible to the mildest periodate oxidation conditions (Moskowitz & Treffers, 1950; Morgan & Watkins, 1951).

Other ways of carbohydrate elimination, e.g. treatments of whole cells or glycoproteins with exo- or endo-glycosidases or biosynthesis in the presence of *N*-glycosylation inhibitors, e.g. tunicamycin (Struck & Lennarz, 1980), may be used, but a lack of effect may mean that the wrong enzymes have been used or that the oligosaccharides are *O*- rather than *N*-glycosidically linked. A loss or reduction of antigenicity after chemical deglycosylation (Mort & Lampert, 1977) would favour a carbohydrate specificity, providing the peptide moiety is unaffected, but complete deglycosylation by chemical means may be difficult to achieve (Tang & Feizi, 1987).

From all of the aforesaid, it is clear that there is a need for procedures that enable direct antigenic analyses of glycoprotein-derived oligosaccharides. With this in mind, a micro-immunoassay procedure was designed recently (Tang *et al.*, 1985; Tang & Feizi, 1987) whereby oligosaccharides that are *O*- and *N*-glycosidically linked to protein can be sequentially released and their antigenicities assayed as lipid conjugates (neoglycolipids) in the absence of the peptide moiety. The procedure

Fig. 5. Schematic representation of an *N*-glycosidically-linked glycoprotein oligosaccharide of poly-*N*-acetyl-lactosamine type showing reactivities with monoclonal antibodies and susceptibilities to endo- β -galactosidase

The oligosaccharide depicted is a hypothetical structure of the type presumed to occur on embryonic glycoproteins (Muramatsu *et al.*, 1978; Childs *et al.*, 1983; Pennington *et al.*, 1985). To the left are shown unsubstituted poly-*N*-acetyl-lactosamine backbones, with short branches reactive with anti-I Ma (Feizi *et al.*, 1971) and M18 (Gooi *et al.*, 1983c) antibodies (Ma⁺, M18⁺); long branched sequences reactive with anti-I Step (Step⁺) and unbranched sequences reactive with anti-i Den (Den⁺). As shown on the right, these determinants are masked (Den⁻, Ma⁻, M18⁻) in the presence of terminal sialic acid residues but are revealed by sialidase (S⁺); the M18 determinant is masked by terminal α -galactosyl residues and is revealed by α -galactosidase (G⁺). Also shown are the α -galactosyl termini reactive with an anti-blood group B antibody (B⁺), designated NB10/8B4 (Voak *et al.*, 1983, Pennington *et al.*, 1985) and the mono- and di-fucosyl structures reactive with anti-SSEA-1 (Gooi *et al.*, 1981) and C14 antibodies (Brown *et al.*, 1983) respectively. Internal β -galactosyl linkages on linear poly-*N*-acetyl-lactosamine domains that are susceptible to digestion with endo- β -galactosidase (Fukuda *et al.*, 1979) are indicated by the short black arrows. Internal β -galactosyl linkages at branch-points, which are resistant to endo- β -galactosidase (Scudder *et al.*, 1984) are indicated by the white arrows. The inset depicts the I(Ma), M18, SSEA-1 and C14 antigenic determinants, carried on an *N*-glycosidically linked glycoprotein oligosaccharide with short backbone sequences, which are resistant to digestion with endo- β -galactosidase. For convenience, the various possible structures are depicted as occurring on a single complex (*N*-glycosidically linked) carbohydrate chain. However, it is not yet known whether, for example, the branched (I type) and linear (i type) backbone structures exist on the same carbohydrate chains (even though they may occur on the same high- M_r glycoproteins) or whether the fucose-containing antigens, SSEA-1 and C14, and the α -galactosyl and sialosyl termini occur on the same carbohydrate chains. Moreover, it has not yet been established whether any of the carbohydrate chains of embryonic cells have core regions of *O*-glycosidic type. Symbols: ●, *N*-acetylglucosamine- β ; ○, mannose- α or β ; □, galactose- β ; ⊗, galactose- α ; △, fucose- α ; □, sialic acid- α . (J. Pennington & T. Feizi in Pennington, 1985).

should be adaptable not only for initial screening to determine whether antibodies are directed at carbohydrate antigens on glycoproteins, but also for eventually identifying the antigenic oligosaccharides among heterogeneous mixtures released from glycoproteins.

Apart from their roles as 'free-standing' antigenic determinants on glycoproteins, oligosaccharides have marked effects on the antigenicities of peptide moieties, for example with the blood group M and N antigens on human sialoglycoproteins (Lisowska, 1981; Waśniowska *et al.*, 1985) and certain viral glycoprotein antigens (Alexander & Elder, 1984). It has been suggested that the immunogenicities of peptide moieties of certain viral envelope glycoproteins may also be influenced by their oligosaccharides (Elder *et al.*, 1986).

Antibodies as probes of structure, tissue distribution and function of oligosaccharides

Elucidation of the structures of oligosaccharide antigens involves processes that are labour-intensive by comparison with protein and nucleic acid sequencing techniques which are now largely automated. Moreover, conclusive structural assignments for novel antigenic sequences requires rigorously purified oligosaccharides for analysis by g.l.c.-m.s. and n.m.r. as well as antigenic analyses (Hounsell, 1987). But the reward that follows from using such defined oligosaccharides to characterize the binding specificity of antibodies is the availability of sequence-specific antibodies which have tremendous potential as probes of oligosaccharide structure. We

predict that micro-immunosequencing procedures using well-characterized monoclonal antibodies in conjunction with lectins (Finne *et al.*, 1980; Cummings & Kornfeld, 1982) and exo- and endo-glycosidases (Pennington *et al.*, 1985; Scudder *et al.*, 1986; Tang *et al.*, 1986) may before long become the order of the day, particularly with glycoprotein oligosaccharides of the poly-*N*-acetyl-lactosamine series which are usually so heterogeneous and hard to purify that sequence information can at best be given as averages (Mizuguchi *et al.*, 1984; Spooncer *et al.*, 1984).

Fig. 5 shows a hypothetical carbohydrate chain with poly-*N*-acetyl-lactosamine backbone sequences characteristic of mouse embryonic cells. Various antigenic determinants recognized by monoclonal antibodies and the predicted cleavage sites using endo- β -galactosidase, and the changes in antigenicity after treatment with sialidase or α -galactosidase, are indicated.

Immunohistochemistry with monoclonal antibodies specific for structural variants among heterogeneous glycoprotein oligosaccharides holds promise for visualizing the distribution and topography of the individual structures (Pennington *et al.*, 1985; Mollicone *et al.*, 1985), and should serve as a guide for the design of appropriate experiments to test the functions of each structure. For example, it was following on from the immunohistochemical experiments described above which revealed the distribution of the I, i and SSEA-1 antigens on the mouse embryo that meaningful investigations were performed of the involvement of poly-*N*-acetyl-lactosamine sequences and their α 1-3 fucosylated derivatives in the adhesion (compaction) of embryonic cells. The results have implicated carbohydrate sequences of the poly-*N*-acetyl-lactosamine series in this first cell-to-cell interaction event in the embryo (Bird & Kimber, 1984; Fenderson *et al.*, 1984; Rastan *et al.*, 1985).

From observations made thus far with these antibodies to the blood group family of oligosaccharide structures, it is clear that a diverse array of macromolecules are decorated with these structures. These include mucins, intestinal hydrolases, galactosyltransferase and the receptor for epidermal growth factor to name but a few, as well as glycosphingolipids. The observation that only selected carbohydrate antigens among those on A431 cells are represented on the receptor for epidermal growth factor (Childs *et al.*, 1984) suggests that glycoproteins in a given cell may be divided into subsets based on their content of oligosaccharide antigens/structures. We have speculated previously that oligosaccharide structures shared between the epidermal growth factor receptor, other glycoproteins and glycolipids may be the basis of interactions mediated by specific carbohydrate-binding proteins (endogenous lectins) through which the receptor glycoprotein might be linked to other extra- and intra-cellular signalling systems (Feizi & Childs, 1985). These thoughts were stimulated by reports that two of the hybridoma antibodies to the epidermal growth factor receptor, TL5 and 29.1, which were found to have blood group A-related specificities (Gooi *et al.*, 1983b, 1985a) can behave as growth factor agonists: antibody TL5 was reported to stimulate DNA synthesis in fibroblasts when bound to these cells and cross-linked by anti-mouse antibodies (Schreiber *et al.*, 1983). Both antibodies TL5 and 29.1 were reported to enhance self-phosphorylation of the partially purified epidermal growth factor receptor

and to increase the affinity of the receptor for the growth factor (Yarden & Schlessinger, 1985). Taken together, these observations may be important clues to the functions of the blood group antigens. It would be an exciting development if endogenous lectins complementary to the blood group isoantigens could be demonstrated in individuals of different blood groups. There is already evidence for reactivities with blood-group oligosaccharides among a family of soluble mammalian lectins (S. R. Carding, R. A. Childs & T. Feizi, cited by Scudder *et al.*, 1982; Leffler & Barondes, 1986).

Other observations that are consistent with the involvement of carbohydrate-recognizing proteins in growth regulation or cell activation include the modulation of the activities of growth factor receptors (Bremer *et al.*, 1984, 1986) and of protein kinase C (Kim *et al.*, 1986; Fujiki *et al.*, 1986) by specific glycolipids, the lack of signal transduction by glycoprotein hormones after they have been deglycosylated (Bahl & Kalyan, 1983; Sairam & Bhargavi, 1985), and the inhibition of signal transduction by glycopeptides and oligosaccharides (Calvo & Ryan, 1985).

Our original hypothesis (Feizi & Childs, 1985) can now be extended and one can envisage a network (Fig. 6) based on glycoconjugates (glycoproteins, proteoglycans and glycolipids) linked through specific lectins which serve as an adjustable connecting system between various response pathways and hence tune the cells response to various stimuli. For example, through such interactions, down- or up-regulation and recycling of various receptor systems may be co-ordinated. Control of the network would be two-fold: activity of the appropriate lectins and expression of the correct carbohydrate chains. Among the lectins in this scheme we include (Feizi, 1985) the carbohydrate-binding proteins of infective agents. The lectin activities may be regulated not only by the absolute amounts of the proteins, but by short range factors such as cation levels (Goldstein & Hayes, 1978; Barondes, 1984; Lis & Sharon, 1986). Changes have been observed in activities of lectins, or levels of proteins antigenically related to lectins, during embryonic development (Barondes, 1984) and in stimulated and transformed cells (Carding *et al.*, 1985; Raz & Lotan, 1986; Gabius *et al.*, 1986). A most striking example of uncoupling of functional lectin-oligosaccharide complexes as a result of structural defects in glycoprotein oligosaccharides is I cell disease, where there is a failure of correct routing (by mannose 6-phosphate binding proteins) of newly synthesized acid hydrolases to lysosomes due to a lack of phosphorylation of mannose residues of these enzyme glycoproteins (Sly & Fisher, 1982; Kornfeld & Kornfeld, 1985). Apart from mediation of intracellular events by interaction of lectins with glycoprotein oligosaccharide variants, other possible effects of variable glycosylation include impairment of fibroblast adhesion (Edwards *et al.*, 1976; Stojanovic *et al.*, 1983) and changes in metastatic potential of tumour cells (Finne, 1983). Moreover, glycosyltransferase activities (and by inference glycoprotein oligosaccharide structures) may be modulated by endogenous lectins (Scudder *et al.*, 1982).

Concluding remarks

In this article we have discussed the wide relevance of the antigenicity and immunogenicity of glycoprotein oligosaccharides to the fields of biochemistry and cell

Fig. 6. Diagram of a cell membrane depicting a hypothetical network based on specific oligosaccharide-lectin interactions as a means of coupling different glycoprotein receptors

In (a) glycoproteins with like carbohydrate structures would be linked by appropriate lectins, e.g. glycoprotein A with B (network I) and glycoprotein B with a glycolipid and with glycoprotein C (network II). In addition, linkage of glycoconjugates with different carbohydrate structures would be achieved through intermediaries involving lectins of different binding specificities, e.g. glycoprotein A, with glycoprotein C via networks I and II. In (b) network I would be interrupted either due to the absence of lectin I or a lack of its activity resulting from a deficiency of essential factors such as calcium ions. Networks II and III would be unaffected. In (c) both networks I and II would be unlinked due to inappropriate glycosylation of glycoproteins A and B, although the relevant lectin activities are present; the result would be an uncoupling of glycoproteins A, B and C. Lectin I is here represented as a membrane-associated protein which varies in activity or the amounts available, and lectins II and III as soluble proteins; lectin III has an oligosaccharide specificity distinct from the other two, and network III may be unaffected by conditions which disrupt networks I and II. Lectin-sugar networks need not be confined to plasma membranes but might operate in intracellular membranes and even in the cytosol. Symbols: ●, ◀, ■, ○, monosaccharides linked in various ways; ∩, ceramide moiety of a glycolipid which shares a terminal oligosaccharide sequence with glycoproteins B and C.

biology. We have highlighted examples of single carbohydrate antigens on plasma membrane glycoproteins which behave as distinctive markers of differentiation stages. We have seen how antigenic cross-reactions between different glycoproteins may be mediated by shared oligosaccharide sequences. We now raise the possibility that interactions *in vivo* between different receptor systems may be mediated by shared carbohydrate structures and an array of endogenous lectins.

We have cited only a few examples among numerous antibodies that were originally raised to glycoprotein receptors, glycoprotein enzymes or to differentiation antigens of whole cells, and later found to recognize oligosaccharide determinants. We predict that immunosequencing procedures exploiting these extraordinary by-products of the hybridoma approach (Köhler & Milstein, 1975), as well as intentionally raised antibodies to glycoprotein oligosaccharides, will be used widely both for structural studies of glycoprotein oligosaccharides and for visualizing their subcellular organization. A new micro-immunochemical approach involving the construction of lipid conjugates of *O*- and *N*-linked glycoprotein oligosaccharides prior to antigenic analyses should greatly facilitate the identification of novel carbohydrate antigens on glycoproteins. The availability of monoclonal antibodies with well defined carbohydrate specificities is now of key importance to the biochemistry of glycoproteins.

We wish to acknowledge the work and helpful discussions of our colleagues supported by the Medical Research Council, Arthritis and Rheumatism Council, Cancer Research Campaign and the Leukaemia Research Fund which have contributed to many of the concepts in this article. We are grateful to Drs. E. F. Hounsell and P. Scudder for constructive criticisms, Miss Caroline Wetter for the illustrations and Mrs. Maureen Moriarty for the preparation of this manuscript.

References

- Alexander, S. & Elder, J. H. (1984) *Science* **226**, 1328–1330
- Baenziger, J. U. (1985) *Am. J. Pathol.* **121**, 382–391
- Bahl, O. P. & Kalyan, N. K. (1983) in *Role of Peptides and Proteins in Control of Reproduction* (McCann, S. M. & Dhindsa, D. S., eds.), pp. 293–327, Elsevier
- Barondes, S. J. (1984) *Science* **223**, 1259–1264
- Bird, J. M. & Kimber, S. J. (1984) *Dev. Biol.* **104**, 449–460
- Blanchard, D., Cartron, J. P., Fournet, B., Montreuil, J., van Halbeek, H. & Vliegthart, J. F. G. (1983) *J. Biol. Chem.* **258**, 7691–7695
- Bremer, E. G., Hakomori, S., Bowen-Pope, D. F., Raines, E. & Ross, R. (1984) *J. Biol. Chem.* **259**, 6818–6825
- Bremer, E. G., Schlessinger, J. & Hakomori, S. (1986) *J. Biol. Chem.* **261**, 2434–2440
- Brown, A., Feizi, T., Gooi, H. C., Embleton, M. J., Picard, J. K. & Baldwin, R. W. (1983) *Biosci. Rep.* **3**, 163–170
- Calvo, F. O. & Ryan, R. J. (1985) *Biochemistry* **24**, 1953–1959
- Carding, S. R., Thorpe, R., Childs, R. A., Spitz, M. & Feizi, T. (1984) *Biochem. J.* **220**, 253–260
- Carding, S. R., Thorpe, S. J., Thorpe, R. & Feizi, T. (1985) *Biochem. Biophys. Res. Commun.* **127**, 680–686
- Carlsson, S. R., Sasaki, H. & Fukuda, M. (1986) *J. Biol. Chem.* **261**, 12787–12795
- Childs, R. A. & Feizi, T. (1987) in *Handbook of Immunoblotting of Proteins* (Bjerrum, O. J. & Heegaard, N. H. H., eds.), CRC Press, Boca Raton, in the press
- Childs, R. A., Pennington, J., Uemura, K., Scudder, P., Goodfellow, P. N., Evans, M. J. & Feizi, T. (1983) *Biochem. J.* **215**, 491–503
- Childs, R. A., Gregoriou, M., Scudder, P., Thorpe, S. J., Rees, A. R. & Feizi, T. (1984) *EMBO J.* **3**, 2227–2233
- Childs, R. A., Berger, E. G., Thorpe, S. J., Aegerter, E. & Feizi, T. (1986) *Biochem. J.* **238**, 605–611
- Childs, R. A. & Feizi, T. (1987) in *Handbook of Immunoblotting of Proteins* (Bjerrum, O. J. & Heegaard, N. H. H., eds.), CRC Press, Boca Raton, in the press
- Chou, K. H., Ilyas, A. A., Evans, J. E., Quarles, R. H. & Jungalwala, F. B. (1985) *Biochem. Biophys. Res. Commun.* **128**, 383–388
- Cummings, R. D. & Kornfeld, S. (1982) *J. Biol. Chem.* **257**, 11235–11240
- Cummings, R. D., Soderquist, A. M. & Carpenter, G. (1985) *J. Biol. Chem.* **260**, 11944–11952
- Donald, A. S. R., Yates, A. D., Soh, C. P. C., Morgan, W. T. J. & Watkins, W. W. (1984) *Biochem. Soc. Trans.* **12**, 596–599
- Edwards, J., Dysart, J. & Hughes, R. C. (1976) *Nature (London)* **264**, 66–68
- Elder, J. H., McGee, J. S. & Alexander, S. (1986) *J. Virol.* **57**, 340–342
- Feizi, T. (1981a) *Immunol. Commun.* **10**, 127–156
- Feizi, T. (1981b) *Trends Biochem. Sci.* **6**, 333–335
- Feizi, T. (1983) *Biochem. Soc. Trans.* **11**, 263–265
- Feizi, T. (1984) *Biochem. Soc. Trans.* **12**, 545–549
- Feizi, T. (1985) *Nature (London)* **314**, 53–57
- Feizi, T. (1987) in *Molecular Immunology of Complex Carbohydrates* (Wu, A. M., ed.), Plenum Press, New York, in the press
- Feizi, T. & Childs, R. A. (1985) *Trends Biochem. Sci.* **10**, 24–29
- Feizi, T., Kabat, E. A., Vicari, G., Anderson, B. & Marsh, W. L. (1971) *J. Immunol.* **106**, 1578–1592
- Feizi, T., Gooi, H. C., Childs, R. A., Picard, J. K., Uemura, K., Loomes, L. M., Thorpe, S. J. & Hounsell, E. F. (1984) *Biochem. Soc. Trans.* **12**, 591–596
- Fenderson, B. A., Zehavi, U. & Hakomori, S. (1984) *J. Exp. Med.* **160**, 1591–1596
- Finne, J. (1983) *Biochem. Soc. Trans.* **11**, 269–270
- Finne, J., Krusius, T. & Järnefelt, J. (1980) *Proc. Int. Congr. Pure Appl. Chem. 17th (Varmavuori, A., ed.)*, pp. 147–159, Pergamon Press, Oxford
- Fox, N., Damjanov, I., Knowles, B. & Solter, D. (1983) *Cancer Res.* **43**, 669–678
- Freeze, H. H., Mierendorf, R. C., Wunderlich, R. & Dimond, R. L. (1984) *J. Biol. Chem.* **259**, 10641–10643
- Fujiki, H., Yamashita, K., Saganuma, M., Horiuchi, T., Taniguchi, N. & Makita, A. (1986) *Biochem. Biophys. Res. Commun.* **138**, 153–158
- Fukuda, M. N., Fukuda, M. & Hakomori, S. (1979) *J. Biol. Chem.* **254**, 5458–5465
- Gabius, H. J., Engelhardt, R., Rehm, S., Deerberg, F. & Cramer, F. (1986) *Tumour Biology* **7**, 71–81
- Gahmberg, C. G. & Hakomori, S. (1973) *J. Biol. Chem.* **248**, 4311–4317
- Gahmberg, C. G., Nilsson, K. & Andersson, L. C. (1979) *Proc. Natl. Acad. Sci. U.S.A.* **76**, 4087–4091
- Gardas, A. (1982) *Arch. Biochem. Biophys.* **216**, 440–448
- Goldstein, I. J. & Hayes, C. E. (1978) *Adv. Carbohydr. Chem. Biochem.* **35**, 127–340
- Gooi, H. C., Feizi, T., Kapadia, A., Knowles, B. B., Solter, D. & Evans, M. J. (1981) *Nature (London)* **292**, 156–158
- Gooi, H. C., Thorpe, S. J., Hounsell, E. F., Rumpold, H., Kraft, D., Förster, O. & Feizi, T. (1983a) *Eur. J. Immunol.* **13**, 306–312
- Gooi, H. C., Schlessinger, J., Lax, I., Yarden, Y., Libermann, T. A. & Feizi, T. (1983b) *Biosci. Rep.* **3**, 1045–1052
- Gooi, H. C., Uemura, K., Edwards, P. A. W., Foster, C. S., Pickering, N. & Feizi, T. (1983c) *Carbohydr. Res.* **120**, 293–302
- Gooi, H. C., Hounsell, E. F., Lax, I., Kris, R. M., Libermann, T. A., Schlessinger, J., Sato, J. D., Kawamoto, T., Mendelsohn, J. & Feizi, T. (1985a) *Biosci. Rep.* **5**, 83–94

- Gooi, H. C., Picard, J., Hounsell, E. F., Gregoriou, M., Rees, A. R. & Feizi, T. (1985b) *Mol. Immunol.* **22**, 689–693
- Hakomori, S. (1985) *Cancer Res.* **45**, 2405–2414
- Hanisch, F. G., Egge, H., Peter-Katalinic, J., Uhlenbruck, G., Dienst, C. & Fangmann, R. (1985) *Eur. J. Biochem.* **152**, 343–351
- Hirohashi, S., Clausen, H., Yamada, T., Shimosato, Y. & Hakomori, S.-I. (1985) *Proc. Natl. Acad. Sci. U.S.A.* **82**, 7039–7043
- Hounsell, E. F. (1987) *Chem. Soc. Rev.*, in the press
- Hounsell, E. F., Gooi, H. C. & Feizi, T. (1981) *FEBS Lett.* **131**, 279–282
- Jacob, F. (1979) *Curr. Top. Dev. Biol.* **13**, 117–137
- Kabat, E. A. (1976) in *Structural Concepts in Immunology and Immunochemistry* (Kabat, E. A., ed.), 2nd edn., pp. 143–197, Holt, Riehart and Wilson, New York
- Kabat, E. A. (1982) *Am. J. Clin. Pathol.* **78**, 281–292
- Kabat, E. A., Liao, J., Sherman, W. H. & Osserman, E. F. (1984) *Carbohydr. Res.* **130**, 289–297
- Kapadia, A., Feizi, T. & Evans, M. J. (1981) *Exp. Cell Res.* **131**, 185–195
- Kim, J. Y. H., Goldenring, J. R., DeLorenzo, R. J. & Yu, R. K. (1986) *J. Neurosci. Res.* **15**, 159–166
- Knapp, W., Majdic, O., Stockinger, H., Bettelheim, P., Liszka, K., Köller, U. & Peschel, C. (1984) *Med. Oncol. Tumor Pharmacother.* **1**, 257–262
- Knowles, B. B., Rappaport, J. & Solter, D. (1982) *Dev. Biol.* **93**, 54–58
- Kobata, A. (1984) in *Biology of Carbohydrates* (Ginsburg, V. & Robbins, P. W., eds.), pp. 87–161, Wiley, New York
- Kobata, A. & Yamashita, K. (1984) *Pure Appl. Chem.* **56**, 821–832
- Köhler, G. & Milstein, C. (1975) *Nature (London)* **256**, 495–497
- Kornfeld, R. & Kornfeld, S. (1985) *Annu. Rev. Biochem.* **54**, 631–664
- Kruse, J., Keilhauer, G., Faissner, A., Timpl, R. & Schachner, M. (1985) *Nature (London)* **316**, 146–150
- Lai, E. & Kabat, E. A. (1985) *Mol. Immunol.* **22**, 1021–1037
- Leffler, H. & Barondes, S. H. (1986) *J. Biol. Chem.* **261**, 10119–10126
- Lerner, R. A. (1984) *Adv. Immunol.* **36**, 1–45
- Le Pendu, J., Fredman, P., Richter, N. D., Magnani, J. L., Willingham, M. C., Pastan, I., Oriol, R. & Ginsburg, V. (1985) *Carbohydr. Res.* **141**, 347–349
- Lis, H. & Sharon, N. (1986) *Annu. Rev. Biochem.* **55**, 35–67
- Lisowska, E. (1981) *Rev. Fr. Transfus. Immuno-Hematol.* **24**, 75–84
- Magnani, J. L. (1984) *Biochem. Soc. Trans.* **12**, 543–545
- Martin, G. R. (1980) *Science* **209**, 768–776
- Mayes, E. L. V. & Waterfield, M. D. (1984) *EMBO J.* **3**, 531–537
- Mehmet, H., Scudder, P., Tang, P. W., Hounsell, E. F., Caterson, B. & Feizi, T. (1986) *Eur. J. Biochem.* **157**, 385–391
- Mizoguchi, A., Takasaki, S., Maeda, S. & Kobata, A. (1984) *J. Biol. Chem.* **259**, 11949–11957
- Mollicone, R., Bara, J., Le Pendu, J. & Oriol, R. (1985) *Lab. Invest.* **53**, 219–227
- Moore, S., Woodrow, C. F. & McClelland, D. B. L. (1982) *Nature (London)* **295**, 529–531
- Morgan, W. T. J. & Watkins, W. M. (1951) *Br. J. Exp. Pathol.* **32**, 34–48
- Mort, A. J. & Lampport, D. T. A. (1977) *Anal. Biochem.* **82**, 289–309
- Moskowitz, M. & Treffers, H. P. (1950) *Science* **111**, 717–719
- Muramatsu, T., Gachelin, G., Nicolas, J. F., Condamine, H., Jakaob, H. & Jacob, F. (1978) *Proc. Natl. Acad. Sci. U.S.A.* **75**, 2315–2319
- Muramatsu, T., Gachelin, G., Damonville, M., Delarbre, C. & Jacob, F. (1979) *Cell* **18**, 183–191
- Niemann, H., Watanabe, K., Hakomori, S., Childs, R. A. & Feizi, T. (1978) *Biochem. Biophys. Res. Commun.* **81**, 1286–1293
- Parker, P. J., Young, S., Gullick, W. J., Mayes, E. L. V., Bennett, P. & Waterfield, M. D. (1984) *J. Biol. Chem.* **259**, 9906–9912
- Pazur, J. H., DeHoff, D. K., Miskiel, F. J. & Baumrucker, C. R. (1986) *Carbohydr. Res.* **149**, 137–147
- Pennington, J. (1985) Ph.D. Thesis, Council for National Academic Awards, London
- Pennington, J. E., Rastan, S., Roelcke, D. & Feizi, T. (1985) *J. Embryol. Exp. Morphol.* **90**, 335–361
- Rahman, A. F. R. & Longenecker, B. M. (1982) *J. Immunol.* **129**, 2021–2024
- Rastan, S., Thorpe, S. J., Scudder, P., Brown, S., Gooi, H. C. & Feizi, T. (1985) *J. Embryol. Exp. Morphol.* **87**, 115–128
- Raz, A. & Lotan, R. (1986) *Cancer Res.* **46**, 3667–3672
- Rumpold, H., Obexerand, G. & Kraft, D. (1982) in *NK Cells and Other Natural Effector Cells* (Herbermann, R. B., ed.), pp. 47–52, Academic Press, New York and London
- Sairam, M. R. & Bhargavi, G. N. (1985) *Science* **229**, 65–67
- Scudder, P., Childs, R. A., Feizi, T., Joziassse, D. H., Schiphorst, W. E. C. M. & Van den Eijnden, D. H. (1982) *Biochem. Biophys. Res. Commun.* **104**, 272–279
- Scudder, P., Hanfland, P., Uemura, K. & Feizi, T. (1984) *J. Biol. Chem.* **259**, 6586–6592
- Scudder, P., Tang, P. W., Hounsell, E. F., Lawson, A. M., Mehmet, H. & Feizi, T. (1986) *Eur. J. Biochem.* **157**, 365–373
- Schreiber, A. B., Libermann, T. A., Lax, I., Yarden, Y. & Schlessinger, J. (1983) *J. Biol. Chem.* **258**, 846–853
- Sly, W. S. & Fischer, H. D. J. (1982) *Cell. Biochem.* **18**, 67–85
- Solter, D. & Knowles, B. B. (1978) *Proc. Natl. Acad. Sci. U.S.A.* **75**, 5565–5569
- Spooner, E., Fukuda, M., Klock, J. C., Oates, J. E. & Dell, A. (1984) *J. Biol. Chem.* **259**, 4792–4801
- Steuden, I., Duk, M., Czerwinski, M., Radzikowski, C. & Lisowska, E. (1985) *Glycoconjugate J.* **2**, 303–314
- Stockinger, H., Majdic, O., Liszka, K., Aberer, W., Bettelheim, P., Lutz, D. & Knapp, W. (1984) *J. Natl. Cancer Inst.* **73**, 7–11
- Stojanovic, D., Hughes, R. C., Feizi, T. & Childs, R. A. (1983) *J. Cell Biochem.* **21**, 119–127
- Struck, D. & Lennarz, W. (1980) in *The Biochemistry of Glycoproteins* (Lennarz, W. J., ed.), pp. 35–73, Plenum, New York
- Tang, P. W. & Feizi, T. (1987) *Carbohydr. Res.* **161**, 133–143
- Tang, P. W., Gooi, H. C., Hardy, M., Lee, Y. C. & Feizi, T. (1985) *Biochem. Biophys. Res. Commun.* **132**, 474–480
- Tang, P. W., Scudder, P., Mehmet, H., Hounsell, E. F. & Feizi, T. (1986) *Eur. J. Biochem.* **160**, 537–545
- Tetteroo, P. A. T., Mulder, A., Lansdorp, P. M., Zola, H., Baker, D. A., Visser, F. J. & von dem Borne, A. E. G. Kr. (1984) *Eur. J. Immunol.* **14**, 1089–1095
- Thorpe, S. J. & Feizi, T. (1984) *Biosci. Rep.* **4**, 673–685
- Triadou, N. (1984) *J. Immunol. Methods* **73**, 283–291
- Uemura, K., Childs, R. A., Hanfland, P. & Feizi, T. (1983) *Biosci. Rep.* **3**, 577–588
- Uemura, K., Macher, B. A., DeGregorio, M., Scudder, P., Buehler, J., Knapp, W. & Feizi, T. (1985) *Biochim. Biophys. Acta* **846**, 26–36
- Urdal, D. L., Brentnall, T. A., Bernstein, I. D. & Hakomori, S. (1983) *Blood* **62**, 1022–1026
- Voak, D., Lennox, B., Lowe, A. D., Downie, D. M., Merry, A. H., Jarvis, J. & Milstein, C. (1983) *Proc. Br. Blood Transfus. Soc.* **1**, 83–112
- Walter, G. (1986) *J. Immunol. Methods* **88**, 149–161
- Wasniowska, K., Reichert, C. M., McGinniss, M. H., Schroer, K. R., Zopf, D., Lisowska, E., Messeter, L. & Lundblad, A. (1985) *Glycoconjugate J.* **2**, 163–176
- Watanabe, K., Hakomori, S., Childs, R. A. & Feizi, T. (1979) *J. Biol. Chem.* **254**, 3221–3228
- Watkins, W. M. (1980) *Adv. Hum. Genet.* **10**, 1–136 and 379–385
- Woodward, M. P., Young, W. W., Jr. & Bloodgood, R. A. (1985) *J. Immunol. Methods* **78**, 143–153

Yarden, Y. & Schlessinger, J. (1985) Ciba Found. Symp. **116**, 23–45
Yamamoto, K., Tsuji, T., Tarutani, O. & Toshiaki, O. (1985) Biochim. Biophys. Acta **838**, 84–91

Young, W. W., Jr., Johnson, H. S., Tamura, Y., Karlsson, K.-Å., Larson, G., Parker, J. M. R., Khare, D. P., Spohr, U., Baker, D. A., Hindsgaul, O. & Lemieux, R. U. (1983) J. Biol. Chem. **258**, 4890–4894