Human *N*-acetylgalactosamine-4-sulphate sulphotase

Purification, monoclonal antibody production and native and subunit M_r values

Gary J. GIBSON,* Gino T. P. SACCONI, Doug A. BROOKS, Peter R. CLEMENTS and John J. HOPWOOD

Department of Chemical Pathology, Adelaide Children's Hospital, 72 King William Road, North Adelaide, South Australia 5006, Australia

Initial purification of *N*-acetylgalactosamine-4-sulphate sulphotase from human liver homogenates containing approx. 1 mg of enzyme in 26 g of soluble proteins was achieved by a six-column chromatography procedure and yielded approx. 40 μ g of a single major protein species. Enzyme thus prepared was used to produce *N*-acetylgalactosamine-4-sulphate sulphotase-specific monoclonal antibodies. The use of a monoclonal antibody linked to a solid support facilitated the purification of approx. 0.5 mg of *N*-acetylgalactosamine-4-sulphate sulphotase from a similar liver homogenate. Moreover the enzyme isolated contained a single protein species, shown by SDS/polyacrylamide-gel electrophoresis to have an M_r of 57 000, which dissociated into subunits of M_r 43 000 and 13 000 in the presence of reducing agents. Essentially identical enzyme preparations were isolated from homogenates of human kidney and lung and from concentrated human urine. The native protein M_r of enzyme from human liver and kidney was assessed by gel-permeation chromatography to be 43 000 on Ultrogel AcA and Bio-Gel P-150. The liver *N*-acetylgalactosamine-4-sulphate sulphotase was shown to have pH optima of approx. 4 and 5.5 with the oligosaccharide substrate (GalNAc4S-GlcA-GalitolNAc4S) and fluorogenic substrate (methylumbelliferyl sulphate) respectively. K_m values of 60 μ M and 4 mM and V_{max} values of 2 and 20 μ mol/min per mg were determined with the oligosaccharide and fluorogenic substrates respectively.

INTRODUCTION

N-Acetylgalactosamine-4-sulphate sulphotase (EC 3.1.6.1, 4-sulphate sulphotase, first termed arylsulphotase B before the natural substrate *in vivo* was discovered) is one of a group of enzymes responsible for the intralysosomal degradation of dermatan sulphate and chondroitin 4-sulphate. A deficiency in 4-sulphate sulphotase is responsible for the pathogenesis of the Maroteaux-Lamy syndrome (mucopolysaccharidosis type VI). This is an autosomal recessive disorder of glycosaminoglycan catabolism leading to storage and excretion of excessive amounts of dermatan sulphate and chondroitin 4-sulphate and a variety of clinical phenotypes, but in its severe form results in dwarfism, gross skeletal changes and cloudy corneas without mental deterioration (McKusick & Neufeld, 1983).

The purification of 4-sulphate sulphotase was initiated to investigate the structure, kinetics and interactions of this enzyme to gain a better understanding of the regulation of lysosomal glycosaminoglycan degradation and to provide sufficient pure enzyme to enable the study of cellular uptake. Conventional purification of 4-sulphate sulphotase has proven difficult and laborious, largely because of its low abundance. McGovern *et al.* (1982) have reported the isolation of 4-sulphate sulphotase from human liver with a purification of 8000-fold and yield of 17%. They obtained a single protein species of M_r 38 000 on polyacrylamide-gel electrophoresis and M_r 48 000 on gel filtration. By contrast, Steckel *et al.* (1983) isolated 4-sulphate sulphotase from human placenta in a yield of 1.5% and purification of 12 000-fold. They reported a single protein species of

M_r 57 000 that gave rise to a major species of M_r 47 000 when subjected to polyacrylamide-gel electrophoresis under reducing conditions.

We have described previously the development of techniques for the concurrent isolation, from human liver, of a number of lysosomal enzymes involved in the degradation of glycosaminoglycans (Mahuran *et al.*, 1983b; Clements *et al.*, 1983, 1985a; Freeman & Hopwood, 1986). In the present paper we report the extension of these techniques for the isolation of 4-sulphate sulphotase for monoclonal antibody production, the utilization of the monoclonal antibody bound to an agarose support for the purification of a single protein species in high yield and partial characterization of this enzyme from a variety of human tissues.

MATERIALS

Human livers, lungs and kidneys were obtained from autopsies of normal individuals with intervals *post mortem* ranging from 24 h to 72 h. Tissues were stored at -20°C for up to 8 weeks. Urine from eight male donors was collected daily and stored at -20°C . Pharmacia Fine Chemicals (Uppsala, Sweden) supplied concanavalin A-Sepharose, chelating Sepharose 6B and the M_r standard kits for both gel filtration and SDS/polyacrylamide-slab-gel-electrophoresis M_r determinations. Amicon (Danvers, MA, U.S.A.) supplied Blue A-agarose and Blue B-agarose (Matrex) gels, hollow-fibre concentrator (10 000- M_r exclusion limit) and YM10 filtration membranes. Bio-Rad Laboratories (Richmond, CA, U.S.A.) supplied Bio-Gel P-150, Affigel

* To whom correspondence should be addressed.

10, peroxidase-coupled goat anti-(mouse Ig) antibody, a silver stain kit and protein assay reagent. LKB (Bromma, Sweden) supplied Ultrogel HA and AcA 34. 4-Methylumbelliferyl sulphate was from Koch-Light (Haverhill, Suffolk, U.K.). The radiolabelled trisaccharide substrate, *N*-acetylgalactosamine 4-sulphate-(β 1-4)-glucuronic acid-(β 1-3)-*N*-acetyl[1- 3 H]galactosaminitol 4-sulphate (GalNAc4S-GlcA-GalitolNAc4S), was prepared as previously described (Hopwood *et al.*, 1986).

METHODS

Purification of 4-sulphate sulphatase: six-column procedure

All steps were performed at 4 °C and all buffers contained 0.1 mM-dithioerythritol, 0.02% NaN₃ and, with the exception of buffer A, 10% (v/v) glycerol.

Extraction. Human liver, kidney and lung (700–800 g) were homogenized in 3 vol. of buffer A (15 mM-dimethylglutarate buffer, pH 6, containing 500 mM-NaCl) and centrifuged as previously described (Mahuran *et al.*, 1983b). Urine (6–12 litres) was concentrated to approx. 500 ml by ultrafiltration in an Amicon hollow-fibre concentrator, adjusted to pH 6 and to 15 mM-dimethylglutarate/500 mM-NaCl/0.1 mM-dithioerythritol by addition of an equal volume of 2 $\times$ concentrated buffer A (30 mM-dimethylglutarate buffer, pH 6, containing 1.0 M-NaCl and 0.2 mM-dithioerythritol). Material precipitated during storage was removed by centrifugation at 1500 *g* for 20 min.

Concanavalin A-Sepharose/Blue A-agarose chromatography. The 27000 *g* supernatant of the homogenate was applied to a 300 ml column of concanavalin A-Sepharose and washed with sufficient buffer A to lower the eluate protein concentration to less than 5 μ g/ml (usually 7–10 litres). The concanavalin A-Sepharose column was connected to a 300 ml column of Blue A-agarose equilibrated with buffer A, and the two columns were equilibrated with 750 ml of 15% (w/w) methyl α -mannoside in buffer A as described elsewhere (Mahuran *et al.*, 1983b; Clements *et al.*, 1983). The recycle time was reduced to 24 h and the flow rate adjusted to 2 ml/min. The methyl α -mannoside was displaced from the columns with 1 litre of buffer A (concanavalin A eluate), and the concanavalin A-Sepharose column was separated from the Blue A-agarose column. The latter column was then washed with more buffer A until the eluate protein concentration was less than 5 μ g/ml. The Blue A-agarose column was then washed with 1 litre of buffer B (15 mM-dimethylglutarate buffer, pH 6, containing 50 mM-NaCl) followed by sufficient buffer C (30 mM-Tris/HCl buffer, pH 7.5, containing 50 mM-NaCl) to lower the eluate protein concentration to less than 5 μ g/ml. The 4-sulphate sulphatase activity was eluted by application of a 2 $\times$ 2-litre linear gradient of NaCl (50 mM–1 M) in buffer C (Blue A eluate). The fractions containing 4-sulphate sulphatase activity were pooled and concentrated to about 200 ml by ultrafiltration in an Amicon hollow-fibre concentrator and/or further concentrated in an Amicon stirred cell fitted with a YM-10 membrane. Any resulting precipitate was removed by centrifugation.

Second Blue A-agarose chromatography. The concentrated Blue A eluates from two preparations were combined, adjusted to pH 6 with 1 M-acetic acid, diluted to about 400 ml with buffer A and applied to a 300 ml Blue A-agarose column equilibrated with the same buffer. The column was then washed with buffer A followed by 500 ml of buffer B and 300 ml of buffer C. The 4-sulphate sulphatase activity was eluted with a 2 $\times$ 1-litre linear gradient (50 mM–1 M) of NaCl in buffer C (20 ml fractions), pooled and concentrated to 10–25 ml.

Hydroxyapatite chromatography. The concentrate was then diluted 3-fold with buffer D (5 mM-sodium phosphate buffer, pH 6.8, containing 10 mM-NaCl) and again concentrated by ultrafiltration to about 5–10 ml. This solution, diluted with buffer D without NaCl to a final Cl[−] ion concentration of approx. 10 mM (assayed with a Cl[−]-specific electrode), was applied to a 50 ml column of hydroxyapatite equilibrated with buffer D. The column was then washed with 500 ml of buffer D, 200 ml of 15 mM-sodium phosphate buffer, pH 6.8, containing 50 mM-NaCl and 100 ml of 30 mM-sodium phosphate buffer, pH 6.8, containing 100 mM-NaCl. The 4-sulphate sulphatase activity was eluted with 200 ml of 60 mM-sodium phosphate buffer, pH 6.8, containing 200 mM-NaCl. Fractions (20 ml) containing 4-sulphate sulphatase activity were pooled, concentrated, adjusted to pH 5 with 1 M-acetic acid and diluted with 2 vol. of 25 mM-sodium acetate buffer, pH 5.0.

Blue B-agarose chromatography. The 4-sulphate sulphatase eluate from the hydroxyapatite column was applied at about 1–2 ml/min to a 50 ml column of Blue B-agarose equilibrated with buffer E (25 mM-sodium acetate buffer, pH 5.0, containing 100 mM-NaCl). The column was then washed with 250 ml of buffer E and eluted with 200 ml of buffer F (25 mM-sodium acetate buffer, pH 5.0, containing 500 mM-NaCl). Fractions (20 ml) containing 4-sulphate sulphatase activity were pooled and concentrated.

Chelating Sepharose chromatography. The above concentrate was applied to a 5 ml column of chelating Sepharose charged with Cu²⁺ ions and equilibrated with buffer F. The column was then washed with 20 ml of buffer F, followed by 75 ml of 20 mM-dimethylglutarate buffer, pH 4.0, containing 500 mM-NaCl and then 75 ml of the same buffer at pH 3.5.

Dithioerythritol was added to the collecting vessel (usually 50 μ l of 10 mM stock solution). Fractions (5 ml) of the pH 3.5 eluate were collected, and those containing 4-sulphate sulphatase activity were immediately pooled and concentrated to a final volume of 2–3 ml. Any precipitate that developed upon overnight storage was removed by centrifugation, and supernatants were used to prepare monoclonal antibodies.

Preparation of monoclonal antibody

Preparations of partially purified 4-sulphate sulphatase (enzyme obtained after the Cu²⁺-chelating-Sepharose chromatography step) were used to immunize a single Balb C mouse by using approx. 10 μ g of protein per injection given in Freund's adjuvant at 4-weekly intervals. After four injections, a final boost was given without adjuvant 4 days before the fusion. The fusion was carried

out by using a previously described protocol (Clements *et al.*, 1985a). Fusion products were screened for antibody activity against partially purified enzyme preparations by enzyme-linked immunosorbent assay with a peroxidase-labelled goat anti-(mouse Ig) antibody detection system (Bio-Rad Laboratories). Briefly, 50 μ l of enzyme (10 μ g/ml) was attached to nitrocellulose paper, and additional reactive sites were blocked by 1% (w/v) albumin in 20 mM-Tris/HCl buffer, pH 7, containing 0.25 M-NaCl. Culture supernatants from hybrid colonies were incubated with the nitrocellulose paper for 30 min at room temperature and then washed twice with 20 mM-Tris/HCl buffer, pH 7, containing 0.25 M-NaCl and incubated with 50 μ l of peroxidase-labelled goat anti-(mouse Ig) antibody (30 min at room temperature). After a further three washes, the assay was developed by using horseradish peroxidase colour-development reagent (Bio-Rad Laboratories).

Reactive antibodies were further characterized by their capacity to react with active enzyme in an antibody immunoprecipitation assay. An affinity-purified goat anti-(mouse Ig) antibody was attached to Affigel 10 (Bio-Rad Laboratories) by using the manufacturer's directions (with approx. 1–5 mg of protein/ml of Affigel). The test antibody (culture supernatant) was then bound to the Affigel-coupled anti-Ig antibody and used to bind 4-sulphate sulphatase from either crude or partially purified enzyme extracts. Enzyme binding was carried out in 20 mM-Tris/HCl buffer, pH 7.0, containing 250 mM-NaCl at 4 °C for 2 h. After three washes in the same buffer, the Affigel-bound enzyme was assayed with the fluorogenic substrate.

Hybridoma antibodies of interest were immuno-purified from cell-culture supernatants by immunoaffinity chromatography with Affigel 10-bound goat anti-(mouse Ig) antibody. Approx. 1 litre of culture supernatant was bound to a 10 ml column of Affigel-coupled goat anti-(mouse Ig) [4 mg of affinity-purified goat anti-(mouse Ig) antibody coupled to 10 ml of Affigel 10] and the column was washed with 10 column volumes of 0.2 M-Tris/HCl buffer, pH 7.0, containing 250 mM-NaCl. The purified immunoglobulin was eluted from the goat anti-(mouse Ig) antibody with 1 M-NH₃. The eluate was neutralized to pH 7–8 with acetic acid, then dialysed extensively against 0.2 M-Tris/HCl buffer, pH 7.0, containing 250 mM-NaCl or 0.1 M-NaHCO₃, pH 8.5, for coupling to Affigel. One immunoglobulin, designated ASB 4.1, was attached to Affigel to generate an immunoaffinity column (20 cm $\times$ 1 cm) for the purification of 4-sulphate sulphatase. The heavy-chain isotype of the purified anti-(4-sulphate sulphatase) antibody was determined by a standard Ouchterlony double-immunodiffusion protocol.

Purification of 4-sulphate sulphatase: monoclonal-antibody affinity chromatography

Human liver, kidney and lung were extracted as described above, and human urine was concentrated using pressure dialysis in an Amicon stirred cell with YM10 membrane. Human liver 4-sulphate sulphatase was purified by concanavalin A-Sepharose/Blue A-agarose chromatography, as described above. Since only approx. 50% of the 4-sulphate sulphatase bound to the Blue A-agarose column under the conditions described, the two fractions (i.e. concanavalin A eluate and Blue A eluate) were processed separately. For the preparation of

kidney, lung and urine enzymes, and in some experiments liver enzyme, the use of Blue A-agarose column chromatography was omitted and tissue homogenate was partially purified by concanavalin A-Sepharose column chromatography only.

4-Sulphate sulphatase solutions thus prepared were concentrated to approx. 10 ml, adjusted to pH 6.8 and chloride concentration of 250 mM and applied to the ASB 4.1 affinity column previously equilibrated with 20 mM-Tris/HCl buffer, pH 6.8, containing 250 mM-NaCl, 10% (v/v) glycerol and 0.02% NaN₃. The column was clamped off for 2 h, then washed with equilibration buffer and eluted with 50 mM-sodium citrate buffer, pH 4.0, containing 2.0 M-NaCl, 10% (v/v) glycerol and 0.02% NaN₃. The flow rate was maintained at 20 ml/h, and all operations were performed at 4 °C. Eluted 4-sulphate sulphatase was concentrated to 2–5 ml. Only 10–20% loss of activity was observed after several weeks' storage at 4 °C.

Gel-permeation chromatography

Native protein M_r was determined by gel-permeation chromatography on a column (2.5 cm $\times$ 155 cm) of Ultrogel AcA 34 in 15 mM-dimethylglutarate buffer, pH 6.0, containing 500 mM-NaCl, 10% (v/v) glycerol and 0.1 mM-dithioerythritol or on a column (1 cm $\times$ 100 cm) of Bio-Gel P-150 in 20 mM-Tris/HCl buffer, pH 6.8, containing 200 mM-NaCl and 10% (v/v) glycerol. Columns were calibrated with protein standards ferritin (M_r 440 000), catalase (M_r 232 000), fructose-bisphosphate aldolase (M_r 158 000), bovine serum albumin (M_r 67 000), ovalbumin (M_r 43 000), chymotrypsinogen A (M_r 25 000) and ribonuclease A (M_r 13 700).

SDS/polyacrylamide-slab-gel electrophoresis

Samples were prepared by a deoxycholate/trichloroacetic acid co-precipitation method as previously described (Mahuran *et al.*, 1983a). Discontinuous SDS/polyacrylamide gels run according to the method of Laemmli (1970), with 12.5% acrylamide, were stained with Coomassie Brilliant Blue R250 and, where necessary, with silver by the method of Merril *et al.* (1981). A Pharmacia M_r calibration kit consisting of phosphorylase *b* (M_r 94 000), bovine serum albumin (M_r 67 000), ovalbumin (M_r 43 000), carbonic anhydrase (M_r 30 000), soya-bean trypsin inhibitor (M_r 20 100) and α -lactalbumin (M_r 14 400) was used.

Enzyme activity

Oligosaccharide substrate. Enzyme activity and kinetic data were obtained, as described previously (Hopwood *et al.*, 1986), with a 12 μ l assay mixture containing 10 μ l of enzyme in 50 mM-sodium formate buffer, pH 3.5, bovine serum albumin (0.5 mg/ml), 570 pmol of substrate (GalNAc4S-GlcA-GalitolNAc4S) and β -hexosaminidase inhibitor (2-acetamido-2-deoxy-D-gluconolactone, 3 mM final concentration). One unit of activity represents the amount of enzyme required to convert 1 nmol of substrate into product (GalNAc-GlcA-GalitolNAc4S)/min. The trisaccharide substrate (GalNAc4S-GlcA-GalitolNAc4S) was chemically derived from one of the natural substrates, chondroitin 4-sulphate, and activity assayed by this procedure is specific for N-acetylgalactosamine-4-sulphate sulphatase.

Table 1. Purification of 4-sulphate sulphatase by the six-column procedure

For experimental details see the text. Enzyme activity was determined with the oligosaccharide substrate GalNAc4S-GlcA-GalitolNAc4S.

Step	Total activity (units)	Total protein (mg)	Specific activity (units/mg)	Purification (fold)	Recovery (%)
Homogenate	919	32 600	0.028	1	100
Concanavalin A-Sepharose	489	194	2.5	90	53
Blue A-agarose I	519	51	9.39	340	56
Blue A-agarose II	401	15	26.8	960	43
Hydroxyapatite	148	4.9	29.6	1000	16
Blue B-agarose	86	1.6	53.3	1900	9.2
Cu ²⁺ -chelating Sepharose	41	0.14	282	10000	4.4

Fluorogenic substrate. The assay with 4-methylumbelliferyl sulphate as substrate was performed essentially as described by Chang *et al.* (1981), with the exception that AgNO₃ was omitted and bovine serum albumin (50 µg/ml) was included in the assay mixture. One unit represents the amount of enzyme needed to produce 1 nmol of methylumbelliferone/min. This assay is not specific for *N*-acetylgalactosamine-4-sulphate sulphatase. Galactose-3-sulphate sulphatase (aryl-sulphatase A), present in the tissues examined at activities similar to 4-sulphate sulphatase, also cleaves 4-methylumbelliferyl sulphate.

Other lysosomal enzymes. Galactose-3-sulphate sulphatase activity was assayed by the technique described by Poulos & Beckman (1978). Iduronate-2-sulphate sulphatase (Hopwood, 1979), *N*-acetylglucosamine-6-sulphate sulphatase (Freeman *et al.*, 1987) and sulphamate sulphohydrolase (Mahuran *et al.*, 1983b) activities were assayed with oligosaccharide substrates by the previously published techniques. α -L-Iduronidase activity was assayed with 4-methylumbelliferyl α -L-iduronide (Hopwood *et al.*, 1979).

RESULTS

Purification of human liver 4-sulphate sulphatase

The extraction conditions reported here for 4-sulphate sulphatase were sub-optimal but were employed to allow concurrent extraction and isolation of four other lysosomal hydrolases, α -L-iduronidase, iduronate-2-sulphate sulphatase, sulphamate sulphohydrolase and *N*-acetylglucosamine-6-sulphate sulphatase.

Six-column purification procedure. The purification of 4-sulphate sulphatase was greater than 10000-fold as determined by the 4-sulphate sulphatase-specific assay and represents a 4.4% yield (Table 1). About 90% of the 4-sulphate sulphatase activity bound to the concanavalin A-Sepharose column and was subsequently eluted with methyl α -mannoside together with the other four lysosomal hydrolases mentioned above. Approximately one-half of this 4-sulphate sulphatase activity bound to the Blue A-agarose column during the recycling step and the remainder appeared in the fraction not bound (concanavalin A-Sepharose eluate), together with

iduronate-2-sulphate sulphatase and *N*-acetylglucosamine-6-sulphate sulphatase activities.

Elution of the first Blue A-agarose column allowed the separation of 4-sulphate sulphatase activity from α -L-iduronidase activity. The combined concanavalin A-Sepharose/Blue A-agarose procedure was the major purification step in this procedure. 4-Sulphate sulphatase activity was eluted from Blue A-agarose at 300 mM-NaCl under the conditions employed and contained no galactose-3-sulphate sulphatase activity. Separation of 4-

Fig. 1. SDS/polyacrylamide-gel electrophoresis of 4-sulphate sulphatase prepared by the six-column procedure

The M_r values indicated for the component proteins were calculated by comparison with protein standards run on the same gel.

Table 2. Immunoprecipitation of 4-sulphate sulphatase

Test antibody was either bound to a goat anti-(mouse Ig) antibody-Affigel 10 affinity column or directly to Affigel 10. The ability of these substrates to bind 4-sulphate sulphatase was then examined as described in the Methods section and is expressed as the percentage of applied activity (determined by fluorogenic assay) bound to the substrate. Purified enzyme refers to 4-sulphate sulphatase isolated by monoclonal-antibody affinity chromatography.

	Enzyme activity bound (%)	
	Purified enzyme	Blue A eluate
Goat anti-(mouse Ig) only	11.4	14.0
Goat anti-(mouse Ig) + ASB 7.3	5.8	11.5
Goat anti-(mouse Ig) + ASB 17.2	5.1	7.9
Goat anti-(mouse Ig) + ASB 4.1	28.2	26.8
ASB 4.1-Affigel	45.5	46.8

sulphate sulphatase and sulphamate sulphohydrolase activities was achieved at the hydroxyapatite step, where sulphamate sulphohydrolase activity was eluted with 15 mM-sodium phosphate buffer, pH 6.8, containing 50 mM-NaCl, and 4-sulphate sulphatase was eluted with 60 mM-sodium phosphate buffer, pH 6.8, containing 200 mM-NaCl. The Blue B-agarose step increased the specific activity nearly 2-fold, and the subsequent Cu²⁺-chelating-Sepharose step resulted in a further 5-fold purification. The 4-sulphate sulphatase activity was eluted with pH 3.7 buffer.

4-Sulphate sulphatase isolated by this procedure was shown to contain a single major species of M_r 43000 together with minor species of M_r 64000, 36000, 29000 and 13000 by SDS/polyacrylamide-slab-gel electrophoresis run under reducing conditions (Fig. 1).

Monoclonal antibody characterization. Fifteen clones containing antibodies against a partially purified preparation of 4-sulphate sulphatase were detected by enzyme-linked immunosorbent assay. One clone, designated ASB 4.1, immunoprecipitated 4-sulphate sulphatase from both purified and relatively crude (Blue A eluate) preparations of enzyme, when assayed with the use of a second antibody attached to Affigel 10 (Table 2). The other 14 antibodies raised failed to bind active enzyme in this binding assay, as exemplified by ASB 7.3 and ASB 17.2 (see Table 2). The monoclonal antibody ASB 4.1 was characterized as being of the IgG₁ subclass. ASB 4.1 was purified by immunoaffinity chromatography and then bound to Affigel 10 to generate an immunoaffinity column for the purification of 4-sulphate sulphatase. The panel of 14 antibodies not immunoprecipitating 4-sulphate sulphatase enzyme activity were re-tested by enzyme-linked immunosorbent assay against ASB 4.1 immunopurified enzyme and showed no reactivity. These antibodies were therefore presumed to be against contaminants in the impure 4-sulphate sulphatase preparations.

The specificity of ASB 4.1 antibody was further assessed in an enzyme-linked immunosorbent assay against various glycoproteins. ASB 4.1 failed to react with albumin, gelatin, transferrin and casein. Similarly, ASB 4.1 failed to react with a panel of other partially purified lysosome-associated exohydrolases, including α -L-iduronidase, N-acetylglucosamine-6-sulphate sul-

Fig. 2. Monoclonal-antibody affinity purification of 4-sulphate sulphatase

(a) Blue A eluate obtained from human liver was applied to the ASB 4.1 monoclonal-antibody affinity column in 20 mM-Tris/HCl buffer, pH 6.8, containing 250 mM-NaCl, 10% glycerol, and 0.02% NaN₃, washed with this buffer and eluted with 50 mM-sodium citrate buffer, pH 4.0, containing 2.0 M-NaCl, 10% glycerol and 0.02% NaN₃. (b) Concanavalin A eluate from the same human liver preparation was applied to the monoclonal-antibody affinity column and eluted as described in (a). Enzyme activity was assayed by the non-specific technique with the methylumbelliferyl sulphate as substrate. The arrows indicate the start of elution. ●, Enzyme activity; ○, A₂₈₀. Note: the A₂₈₀ in fractions 2, 3 and 4 in (b) is off-scale.

Table 3. Purification of 4-sulphate sulphatase by monoclonal-antibody affinity chromatography

The enzyme bound to the Blue A column (2) was applied to the monoclonal-antibody (MAb) affinity column and fractions unretarded (3) and bound (4) were analysed separately. The enzyme that failed to bind to the Blue A column (5) was also applied to the monoclonal-antibody affinity column and fractions unretarded (6) and bound (7) were analysed. (8) Total bound enzyme was obtained by summation of the enzyme from the Blue A eluate that bound to the monoclonal-antibody affinity column with that from the concanavalin A eluate that bound to the monoclonal-antibody affinity column. Enzyme activity was determined with the oligosaccharide substrate GalNAc4S-GlcA-GalitoINAc4S.

Step		Total activity (units)	Total protein (mg)	Specific activity (units/mg)	Purification (fold)	Recovery (%)
Homogenate		1143	26400	0.043	1	100
Blue A eluate	(2)	531	25	21	490	47
Concanavalin A eluate	(5)	471	198	2.4	56	41
MAb affinity chromatography						
Blue A: not bound	(3)	8	15	0.5		0.7
Blue A: bound	(4)	334	0.24	1392	32000	29
Concanavalin A: not bound	(6)	6	170	0.04		0.5
Concanavalin A: bound	(7)	245	0.21	1167	27000	21
Total bound enzyme (concanavalin A + Blue A)	(8)	579	0.45	1287	30000	50

phatase, sulphamate sulphohydrolase and iduronate-2-sulphate sulphatase.

Monoclonal-antibody affinity purification procedure. 4-Sulphate sulphatase activity was efficiently separated by monoclonal-antibody affinity chromatography from the huge excess of contaminating proteins present in the Blue A or concanavalin A eluates (Fig. 2). This procedure has greatly shortened our 4-sulphate sulphatase preparation and facilitated the recovery of more than 50% of the enzyme activity present in the initial liver homogenate (Table 3). It should be noted that the large peak of activity shown with the fluorogenic substrate (Fig. 2b) that did not bind to the monoclonal-antibody affinity column contained only approx. 1% of the applied 4-sulphate sulphatase activity. The small amount of 4-sulphate sulphatase not bound to the column does not appear to differ significantly from the bound enzyme since, on re-application, almost all the activity bound to the monoclonal-antibody affinity column and was eluted in the high-salt low-pH step (results not shown).

The enzyme recovered from the monoclonal-antibody affinity column represents a 30000-fold purification over that found in the liver homogenate and 60- or 500-fold over that found in the Blue A or concanavalin A eluates respectively (Table 3). SDS/polyacrylamide-slab-gel electrophoresis (Fig. 3) revealed only a single protein species of M_r 57000 (lane 4), which gave rise to two proteins of M_r 43000 and M_r 13000 in the presence of dithioerythritol (lane 2). About 30 livers from donors of different age and sex, and after various periods of storage, yielded only this single species of 4-sulphate sulphatase. When the liver enzyme preparation was applied to the monoclonal-antibody affinity column in 200 mM-NaCl rather than 250 mM-NaCl, the bound fraction contained an additional protein species of M_r 29000 (lane 3), the mobility of which was unaffected by the presence of reducing agents (lane 5). Although variable, the 29000- M_r species frequently appeared as the major protein species present in these preparations.

Rechromatography at 250 mM-NaCl allowed the separation of the 57000- M_r and 29000- M_r species in the bound and unbound fractions respectively (lanes 7 and 6), and suggested that the lower- M_r species is not enzymically active 4-sulphate sulphatase, since the low level of activity present in this fraction could be accounted for by slight contamination with the 57000- M_r species. The possibility remains, however, that the 29000- M_r species is a form of 4-sulphate sulphatase with a low antibody affinity and low enzyme activity. Small amounts of an additional 36000- M_r protein found in fractions bound to the monoclonal-antibody affinity column in 200 mM-NaCl (Fig. 3, lane 6) were very rarely observed and are presumed to be another protein species loosely bound to the enzyme. Apparently identical 29000- M_r and 36000- M_r species were present in the preparation isolated by the six-column procedure (Fig. 1).

The 57000- M_r species of 4-sulphate sulphatase was also the major species isolated from kidney, lung and urine. However, particularly in enzyme prepared from lung, additional minor protein bands were observed. The lung preparation contained additional proteins of M_r 112000, 105000, 83000, 75000, 29000 and 27500. Kidney contained only the 75000- M_r minor species, and urine the 27500- M_r species (Fig. 3). Although present in small amounts, the minor protein species were consistently observed and in similar relative amounts in a number of preparations from each tissue. Separation from enzyme activity by gel-permeation chromatography or rechromatography on the monoclonal-antibody affinity column (results not shown) suggests that these proteins are not related to 4-sulphate sulphatase. The possibility remains, however, that they represent inactive enzyme.

Native molecular size

Gel filtration of the immunopurified 4-sulphate sulphatase isolated from liver and kidney resulted in a single symmetrical peak corresponding to an M_r of 43000 when run on Ultrogel AcA 34 in 500 mM-NaCl,

Fig. 3. Polyacrylamide-gel electrophoresis of 4-sulphate sulphatase prepared from liver, kidney, lung and urine by monoclonal-antibody affinity chromatography

Lanes 1–7, 8 and 9, 10–12 and 13–15 represent four separate gels. Lanes 1, 9, 12 and 15 contain standard protein mixtures. The M_r values of the component proteins are shown to the left of lane 1. Lanes 2–7, 11 and 14 contain 4-sulphate sulphatase isolated from human liver, lane 8 enzyme isolated from human kidney, lane 10 enzyme isolated from human lung, and lane 13 enzyme isolated from human urine. All samples were denatured before electrophoresis by incubation for 2 min at 100 °C in the presence of 0.1 M-dithioerythritol with the exception of samples in lanes 4 and 5, which were denatured in the absence of reducing agent. Enzyme in lanes 3 and 5 was purified from human liver by affinity chromatography in 20 mM-Tris/HCl buffer, pH 6.8, containing 200 mM-NaCl, 10% glycerol and 0.02% NaN_3 . All other enzyme samples were obtained after affinity chromatography in the same buffer but containing 250 mM-NaCl. Enzyme obtained by affinity chromatography in 200 mM-NaCl buffer (lane 3) was re-run over the monoclonal-antibody affinity column in 250 mM-NaCl buffer. The fraction not retarded after re-chromatography is shown in lane 6 and that bound to the column is shown in lane 7.

pH 6.0, or on Bio-Gel P-150 in 200 mM-NaCl, pH 6.8 (Fig. 4). The enzyme isolated from the Blue A eluate gave a profile identical with that of the enzyme isolated from the concanavalin A eluate. SDS/polyacrylamide-slab-gel electrophoresis of the enzyme activity peak eluted from the sizing columns demonstrated the presence of both the 43 000- M_r and 13 000- M_r components. No other peptides were observed in the enzyme activity peak, and the minor high- M_r species were eluted in the void volume fraction.

Kinetics

Human liver 4-sulphate sulphatase was shown to have a pH optimum of 5.5 with the fluorogenic substrate. This appeared to be unaffected by the type of buffer used (Fig. 5b). The pH optimum observed with the oligosaccharide substrate, however, was dependent on the type of buffer used. With formate buffer the pH optimum observed was 3.8, whereas with acetate or dimethylglutarate the pH optimum was approx. 4.4 (Fig. 5a). Our previous studies (Clements *et al.*, 1985b; Freeman & Hopwood, 1986) of the lysosomal enzyme degradation of substrates derived from glycosaminoglycans had similarly shown that the observed pH optima were susceptible to changes in the type of buffer ion used. One possible explanation is that buffer ions influence the interaction of substrate acidic groups with basic groups present in the enzyme. Arginine

residues have been implicated as components of substrate-binding sites for other enzymes acting on anionic substrates or cofactors (Riordan *et al.*, 1977; Gacesa *et al.*, 1981); however, other basic residues may also be involved (Rome *et al.*, 1978). The K_m was 4 mM and V_{max} 20 $\mu\text{mol/min}$ per mg with the fluorogenic substrate in 50 mM-acetate buffer, pH 5.5. The K_m was 60 μM and V_{max} 2 $\mu\text{mol/min}$ per mg with the oligosaccharide substrate in 50 mM-formate buffer, pH 3.5. Enzyme prepared by the six-column procedure and monoclonal-antibody affinity chromatography had similar kinetics.

DISCUSSION

We have described the purification of 4-sulphate sulphatase from human liver, the production of monoclonal antibodies to this enzyme, the use of monoclonal antibodies in the efficient preparation of pure enzyme and partial characterization of pure 4-sulphate sulphatase. The advantages of the combined concanavalin A-Sepharose/Blue A-agarose chromatography procedure for the concurrent preparation of several lysosomal enzymes has been discussed previously (Clements *et al.*, 1983). In the case of 4-sulphate sulphatase, this simple step achieved a 500-fold purification of the enzyme and facilitated subsequent

Fig. 4. Gel-permeation chromatography of 4-sulphate sulphatase

Chromatography of monoclonal-antibody affinity-purified 4-sulphate sulphatase was performed in 20 mM-Tris/HCl buffer, pH 6.8, containing 200 mM-NaCl on Bio-Gel P-150. Activity was assayed with both the oligosaccharide substrate (○) expressed as percentage breakdown into product and the fluorogenic substrate (●) in units/ml. SDS/polyacrylamide-gel electrophoresis of the fractions is also shown.

purification by hydroxyapatite and Cu^{2+} -chelating Sepharose chromatography for the initial preparation of 4-sulphate sulphatase antigen and, more recently, a partial purification before chromatography on the monoclonal-antibody affinity column.

The two-step procedure of concanavalin A-Sepharose/Blue A-agarose chromatography, followed by monoclonal-antibody affinity chromatography, allowed the isolation of approx. 0.5 mg of pure enzyme from an initial extract containing only approx. 1 mg of enzyme protein in 26 g of soluble liver proteins. Similar recoveries were obtained with preparations isolated from kidney, lung and urine. It seems likely, considering the high rate of recovery, that the major and possibly the only form of active 4-sulphate sulphatase present in human liver, kidney, lung and urine is a protein of M_r 57000, composed of peptides of M_r 43000 and 13000

linked by disulphide bonds. Although care was taken to minimize exposure of the 4-sulphate sulphatase to proteolytic enzymes, doubtless extracted from the liver during homogenization, and to minimize the time of liver storage, we cannot eliminate the possibility that the species characterized has undergone some enzyme processing either during extraction and purification or *in situ* after the death of the donor. However, the consistent M_r values of the component subunits of the enzyme (i.e. M_r 43000 and 13000) isolated from a number of different tissues and donors (which might be expected to contain a variety of proteolytic enzymes) suggests that proteolysis of the enzyme *in vitro* was not significant.

Approximately one-half the liver 4-sulphate sulphatase bound to the Blue A-agarose column under the conditions employed. This does not appear to reflect the presence of different species of enzyme, since 4-sulphate

Fig. 5. pH optimum of 4-sulphate sulphatase

(a) With the oligosaccharide substrate GalNAc4S-GlcA-GalitolNAc4S. (b) With the fluorogenic substrate 4-methylumbelliferyl sulphate. Enzyme was buffered to the appropriate pH in 50 mM-formate buffer (○), 50 mM-acetate buffer (□) or 50 mM-dimethylglutarate buffer (●).

sulphatase isolated from the concanavalin A eluate and 4-sulphate sulphatase isolated from the Blue A eluate produced identical patterns on SDS/polyacrylamide-gel electrophoresis and gel filtration and displayed the same pH optima and very similar K_m and V_{max} values with both the fluorogenic substrate and the oligosaccharide substrate. Furthermore, since all the 4-sulphate sulphatase in the concanavalin A eluate bound to the Blue A-agarose column on re-application (results not shown), it appears that the presence of 4-sulphate sulphatase in the concanavalin A eluate reflects only the capacity of the Blue A-agarose column to bind 4-sulphate sulphatase under the conditions employed. Although Blue A-agarose chromatography was a valuable preparative step in the purification of 4-sulphate sulphatase for antibody production, it was not essential before monoclonal-antibody affinity chromatography.

Steckel *et al.* (1983) have reported similar properties for 4-sulphate sulphatase isolated from human placenta, although recovery of the enzyme was only 1–2% of that present in their initial homogenate. They reported a 57000- M_r protein composed of a 47000- M_r polypeptide linked via disulphide bonds to two smaller polypeptides of M_r 11000 and 14500. The same authors also reported the isolation of an additional 57000- M_r form of 4-sulphate sulphatase from human fibroblasts by immunoprecipitation of radiolabelled proteins with a polyclonal antibody. This form was composed of 40000- M_r and 31000- M_r disulphide-linked polypeptides and apparently had very low enzyme activity. Although we found a 29000- M_r protein that displayed a lower affinity for the

monoclonal-antibody column, it does not appear to be the same as that observed by Steckel *et al.* (1983), since it was not affected by reducing agents. Furthermore, we failed to detect any accompanying 40000- M_r protein. It thus appears likely that human liver, kidney, lung and urine contain very little of this form of the enzyme, or that this form does not contain the epitope recognized by our monoclonal antibody, or that this protein species is unrelated to 4-sulphate sulphatase and was precipitated by contaminating antibodies present in the polyclonal preparation employed by Steckel *et al.* (1983). A small amount of 4-sulphate sulphatase failed to bind to the concanavalin A-Sepharose column, and a further fraction was lost during immunoaffinity chromatography. This probably reflects the binding equilibrium of the concanavalin A-Sepharose column and loss due to irreversible binding to containers and column matrix and enzyme denaturation during processing. However, it remains to be determined whether these fractions contain the 31000- M_r and 40000- M_r species described by Steckel *et al.* (1983).

McGovern *et al.* (1982) have also reported the isolation of 4-sulphate sulphatase from human liver, although the size of the protein on SDS/polyacrylamide-slab-gel electrophoresis appears to be slightly lower (M_r 38000) than that reported here or by Steckel *et al.* (1983). This may be due to the different techniques employed. The enzyme displayed a similar pH optimum and K_m with 4-nitrocatechol sulphate as substrate to those reported here with the fluorogenic substrate. In our hands the procedure described by McGovern *et al.* (1982) resulted in a very

low yield (less than 1 % recovery) of impure (many bands were observed on polyacrylamide-gel electrophoresis) enzyme. Gasa *et al.* (1987) have recently reported an isolation of 4-sulphate sulphatase from human liver using a modification of the techniques described by McGovern *et al.* (1982). This procedure incorporated an extra column purification step and resulted in the isolation of a protein of M_r 58 000 determined by SDS/polyacrylamide-gel electrophoresis run under non-reducing conditions with a yield of 17 % and purification of 13 000-fold.

We conclude that the 57 000- M_r species of 4-sulphate sulphatase is the major form found in human tissues and urine and have demonstrated that it can be simply and efficiently purified to homogeneity by the use of monoclonal-antibody affinity chromatography. Initial studies have indicated that it is taken up by fibroblasts from mucopolysaccharidosis VI patients in cell culture, but further studies will be required to investigate the mechanism of uptake and establish a protocol for enzyme replacement therapy for these patients.

This work was supported by grants from the Research Trust of the Adelaide Children's Hospital and the National Health and Medical Research Council of Australia. We thank Dr. A. Bourne of the Department of Histopathology, Adelaide Children's Hospital, and Dr. R. A. James, South Australian Forensic Science Division, for supplying autopsy material. The expert technical assistance of Merron Gelder in preparation of monoclonal antibodies, and Julie Hagedorn, Elaine Ravenscroft and Larissa Cox in enzyme preparation, is gratefully acknowledged.

REFERENCES

- Chang, P., Rosa, N. E. & Davidson, R. G. (1981) *Anal. Biochem.* **117**, 382–389
- Clements, P. R., Mahuran, D. J. & Hopwood, J. J. (1983) *J. Chromatogr.* **261**, 77–82
- Clements, P. R., Brooks, D. A., Saccone, G. T. P. & Hopwood, J. J. (1985a) *Eur. J. Biochem.* **152**, 21–28
- Clements, P. R., Muller, V. & Hopwood, J. J. (1985b) *Eur. J. Biochem.* **152**, 29–34
- Freeman, C. & Hopwood, J. J. (1986) *Biochem. J.* **234**, 83–92
- Freeman, C., Clements, P. R. & Hopwood, J. J. (1987) *Biochem. J.* **246**, 347–354
- Gacesa, P., Savitsky, M. J., Dodgson, K. S. & Olavesen, A. H. (1981) *Biochim. Biophys. Acta* **661**, 205–212
- Gasa, S., Balbaa, M., Nakamura, M., Yonemori, H. & Makita, A. (1987) *J. Biol. Chem.* **262**, 1230–1238
- Hopwood, J. J. (1979) *Carbohydr. Res.* **69**, 203–216
- Hopwood, J. J., Muller, V., Smithson, A. & Baggett, N. (1979) *Clin. Chim. Acta* **92**, 257–265
- Hopwood, J. J., Elliott, H., Muller, V. J. & Saccone, G. T. P. (1986) *Biochem. J.* **234**, 507–514
- Laemmli, V. K. (1970) *Nature (London)* **227**, 680–685
- Mahuran, D. J., Clements, P. R., Carella, M. & Strasberg, P. M. (1983a) *Anal. Biochem.* **129**, 513–516
- Mahuran, D. J., Clements, P. R. & Hopwood, J. J. (1983b) *Biochim. Biophys. Acta* **747**, 359–365
- McGovern, M. M., Vine, D. T., Haskins, M. E. & Desnick, R. J. (1982) *J. Biol. Chem.* **257**, 12605–12610
- McKusick, V. & Neufeld, E. (1983) in *The Metabolic Basis of Inherited Disease* (Stanbury, J. B., Wyngaarden, J. B., Fredrickson, D. S., Goldstein, J. L. & Brown, M. S., eds.), 5th edn., pp. 751–771, McGraw-Hill, New York
- Merril, C. R., Goldman, D., Sedman, S. A. & Ebert, M. (1981) *Science* **211**, 1437–1438
- Poulos, A. & Beckman, K. (1978) *Clin. Chim. Acta* **89**, 417–420
- Riordan, J. F., McElvany, K. D. & Borders, C. L. (1977) *Science* **195**, 884–886
- Rome, L. H., Garvin, A. J. & Neufeld, E. F. (1978) *Arch. Biochem. Biophys.* **189**, 344–353
- Steckel, F., Hasilik, A. & von Figura, K. (1983) *J. Biol. Chem.* **258**, 14322–14326

Received 31 March 1987/23 June 1987; accepted 21 August 1987