Cadaveric Dissections Based on Observations of Injuries to the Temporal Bone Structures following Head Trauma

Jarosław Wysocki, M.D., Ph.D.¹

ABSTRACT

One hundred temporal bones obtained from forensic autopsies were dissected to expose injured structures. Longitudinal fractures were present in 82%, transverse fractures in 11%, and mixed fractures in 7% of the cases. Facial canal injuries were present in almost half of the bones with longitudinal fractures (36/82), although cuts of the facial nerve stem were rarely encountered. Damages to the facial canal associated with longitudinal fractures were most frequently seen in the region of the geniculum. However, transverse fractures with facial canal involvement (7/11) most frequently occurred in the labyrinthine portion, causing a complete cut of the facial nerve. Injuries to the jugular bulb were also common (21/100) and associated with all types of temporal bone fractures. Observed damages to the auditory ossicles included disconnection of their joints or fractures of the malleus or stapes. Fractures of the incus were not observed. Injuries to the carotid canal were common (52/100), although an injury to the arterial wall was observed in only one specimen. The frequency and nature of damage in temporal bone fractures strictly reflect the type of fracture, especially in terms of facial nerve disorders: the most serious damage is observed with fractures that involve the otic capsule.

KEYWORDS: Temporal bone, fractures, cadavers, anatomy

Temporal bone fractures can adversely affect the quality of life when the vestibulocochlear organ or facial nerve is damaged. Temporal bone fractures are common. Thirty percent of head injuries involve a fracture of the cranial base, and 18% also affect the temporal bone.¹ The common causes of temporal

bone fractures are road accidents, falls, beatings, and gunshot wounds.²⁻⁶

The characteristic direction of the fracture line was used to classify temporal bone fractures as longitudinal, transverse, mixed, and oblique.^{4,7-15} In longitudinal fractures, the fracture line courses

Skull Base, volume 15, number 2, 2005. Address for correspondence and reprint requests: Jarosław Wysocki, M.D., Ph.D., Institute of Physiology and Pathology of Hearing, Pstrowskiego Str. 1, 01-943 Warsaw, Poland. E-mail: jwysocki@ib.amwaw.edu.pl. ¹Institute of Physiology and Pathology of Hearing, Warsaw, Poland. Copyright © 2005 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel: +1(212) 584-4662. 1531-5010,p;2005,15,02,99,108,ftx,en;sbs00440x.

parallel to the long axis of the pyramid, through the external auditory canal and tegmen tympani, passes the labyrinthine capsule, and reaches the middle cranial fossa.^{1-4,6-8,10,11} In transverse fractures, the line of fracture is perpendicular to the long axis of the pyramid or petrous bone,^{1,4,6,7,9} and its course is variable.

In terms of the labyrinth, two main groups of transverse fractures can be distinguished: internal fractures (anteromedial) and external fractures (posterolateral).^{16,17} Internal fractures run along the internal auditory canal, traverse the cochlea and geniculate ganglion, and reach the anterior edge of the pyramid. External fractures run through the entire internal ear (the cochlea and vestibule) along the facial canal to the geniculate ganglion. The middle ear, tympanic membrane, and external auditory canal are lateral to the fracture line and typically intact. Structures of the middle and internal ear can be involved with mixed fractures of the temporal bone.^{18,19} Mixed fractures compose less than 10% of all temporal bone fractures. Longitudinal fractures are regarded as the most common (70 to 90%) followed by transverse fractures (10 to 30%).^{1,3,12,18-20} Sometimes longitudinal fractures occur bilaterally.²¹ Oblique fractures run across descriptive anatomical planes and are a variant of mixed fractures.⁴

Damage to the facial nerve is a relatively common complication of temporal bone fractures, resulting in paralysis or paresis (10 to 50%).^{1-3,12,15-17,22-26} The pathological damage to the facial nerve can vary. In transverse fractures, the continuity of the facial nerve tends to be lost, while it is rare in longitudinal fractures. Paralysis of the facial nerve associated with longitudinal fractures can also have other causes: a piece of bone impinging on the nerve, pressure exerted on the nerve by a dislocated bone fragment, an intraneural hematoma or nerve contusion.^{1,22,25} There are other possible serious consequences of temporal bone fractures: a fistula associated with cerebrospinal fluid leakage and the possibility of meningitis,^{1-3,5,27} a traumatic perilymph fistula,^{28,29} cholesteatoma,¹⁶ hearing loss,³⁰⁻³² balance disturbances,^{16,33,34} and carotid injury.^{35,36} Data on temporal bone fractures ob-

tained from postmortem observations are sparse. Clinical reports yield only partial information about the temporal bone obtained from computed tomography (CT) and intraoperative observations. We therefore conducted a cadaveric study to characterize temporal bone fractures as fully as possible.

MATERIALS AND METHODS

One hundred cadaveric temporal bones obtained from the forensic autopsies of victims of head trauma were examined. Specimens were collected consecutively, without additional criteria of qualification with the exception of data availability (age, mechanism of trauma, management). The age of victims varied between 15 and 87 years (mean 38 years). Damage to the temporal bones was estimated under an operating microscope and standard set of otosurgical instruments. The course of the fracture line was used to classify the injuries as longitudinal, transverse, or mixed.

The studied materials were diverse. They consisted of extremely severe head trauma (including skull defragmentation) and general trauma associated with moderate skull damage. In the latter, the head injury was only one of several injuries and was not the direct cause of death. Thus, the data were analyzed according to the cause of death. In Group I, head trauma was a direct cause of death (62 bones). In Group II, head trauma was one of several injuries encountered during autopsy (38 bones).

RESULTS

In the examined materials, 82 fractures qualified as longitudinal, 11 as transverse, and 7 as mixed. All temporal bone structures could be involved in the fracture line and totally or partially damaged (Table 1). As a rule, longitudinal fractures did not involve the labyrinth, and transverse fractures left

Table 1 Localization of Damage to Temporal Bone Structures in 100 Fractures

Structure Destroyed	No. of Longitudinal Fractures	No. of Transverse Fractures	No. of Mixed Fractures
Vestibule*		7	5
Cochlea*		8	4
Semicircular canals*		5	5
Internal acoustic meatus*		5	1
Carotid canal	42	6	4
Eustachian tube	57	7	3
Jugular fossa	9	8	4
Sigmoid groove	13	2	2
Tympanic membrane*	54		3
External acoustic meatus	59		4

*Internal ear structures.

the external meatus intact. There were no significant differences in the pattern of the fracture line in the two groups. Therefore, the results for the two groups are presented jointly (Tables 1 and 2). The only statistically significant difference between the two groups was the frequency of facial canal fractures (Table 3), which were most frequent in Group I. These results are presented separately for both groups (Table 4). Also, see Figures 1, 2, and 3.

The following structures were involved in longitudinal fractures (Table 1): the osseous portion of the Eustachian tube, external acoustic meatus, tympanic membrane, sigmoid groove, jugular fossa, and the facial and carotid canals. Dislocations of auditory ossicles and their fractures were also

common (Table 2). According to the definition, internal ear structures were not involved with longitudinal fractures. In most longitudinal fractures, the tympanic membrane was ruptured, particularly when the external auditory meatus was involved.

Transverse fractures were associated with injuries to the internal, and, sometimes, also to the middle ear structures. Damage to the tympanic membrane or external auditory meatus was not observed, consistent with the definition of this fracture (Table 1). The fracture line was most often located in the anterior part of the labyrinth. Less frequently it ran through its posterior region (semicircular canals). In almost all cases (8/11), transverse fractures were associated with injury to the jugular fossa due to its vicinity to the labyrinth. If the fracture line ran through the oval window, the base of the stapes was also destroyed (Table 2).

Mixed fractures involving the structures of the middle, inner, and external ears reflected the combined features of longitudinal and transverse fractures. Damage to the auditory ossicles produced dislocated joints, fractures, or both (Table 2). Interestingly, fractures of the incus were not observed. The fracture line in the malleus ran through its neck and seldom through its head. Fractures of the stapes involved both its branches or its plate. Fractures of the facial canal were common and localized in the various topographical segments of the canal.

Table 2 Damage to the Auditory Ossicles in 100 Temporal Bone Fractures

Destroyed Structure and Type of Injury	No. of Longitudinal Fractures	No. of Transverse Fractures	No. of Mixed Fractures
Fracture			
Malleus	7	0	1
Incus	0	0	0
Stapes	11	2	4
Luxation			
Incudo-malleal joint	48	0	3
Incudo-stapedial joint	51	2	4

Table 3 Location of Facial Canal Injury in 100 Temporal Bone Fractures

Segment of Facial Canal	No. of Longitudinal Fractures		No. of Transverse Fractures		No. of Mixed Fractures	
	Nerve Stem Intact	Nerve Stem Cut	Nerve Stem Intact	Nerve stem Cut	Nerve Stem Intact	Nerve Stem Cut
Internal acoustic meatus	0	0	0	1	0	0
Labyrinthine	0	0	0	4	1	0
Region of genu	27	4	0	0	2	2
Tympanic	0	0	0	2	0	0
Mastoid	3	2	0	0	0	1

DISCUSSION

Most fracture lines did not run in a perpendicular plane; rather they tended to run more or less obliquely. This finding is consistent with other researchers' observations that oblique fractures are the most common.^{4,8} In our opinion, however, the course of a fracture plane described as oblique does not contradict the classification of fractures as longitudinal, transversal, or mixed types, as introduced by Ulrich.⁷ First, this classification is based on the three main parts of the ear, not on the geometry of the fracture plane, and has considerable clinical merit. As defined, longitudinal fractures avoid the labyrinthine structures, while transverse ones include them; mixed fractures run through both regions of the ear.^{1,16,17}

Fractures of the sigmoid groove, jugular fossa, or both were common injuries (2/5 of all studied fractures). Their frequency was highest in transverse fractures (Table 1). In all cases sigmoid groove or jugular fossa fractures were accompanied by interruption of the walls of the adjacent veins. This finding is consistent with clinical observations that fractures of the temporal bone may manifest with massive venous hemorrhage.^{3,6,10,11,16} However, fractures of the carotid canal, which were also often observed (half of all studied bones) were rarely associated with injury to the arterial wall (1 case, transverse fracture). This finding is consistent with some clinical observations, but CT, magnetic resonance imaging, and angiographic data suggest that small angiographic abnormalities due to carotid canal fracture are common.^{35,36}

Table 4 Location of Facial Canal Injury in Groups 1 (62 Patients) and 2 (38 Patients)

Segment of Facial Canal	No. of Longitudinal Fractures		No. of Transverse Fractures		No. of Mixed Fractures	
	Nerve Stem Intact	Nerve Stem Cut	Nerve Stem Intact	Nerve Stem Cut	Nerve Stem Intact	Nerve Stem Cut
GROUP 1						
Meatal	0	0	0	1	0	0
Labyrinthine	0	0	0	3	1	0
Region of genu	22	3	0	0	1	1
Tympanic	0	0	0	1	0	0
Mastoid	3	2	0	0	0	1
GROUP 2						
Meatal	0	0	0	0	0	0
Labyrinthine	0	0	0	1	0	0
Region of genu	5	1	0	0	1	1
Tympanic	0	0	0	1	0	0
Mastoid	0	0	0	0	0	0

Figure 1 Longitudinal fracture of the right temporal bone. The fracture line is parallel to the axis of the pyramid. The malleus, incus, and stapes are disconnected, the incus is luxated, and the malleus is fixed in the insertion of the tensor tympani. The fracture line passes through the region of the genu of the facial nerve and the first (initial) segment of the greater petrosal nerve. Both structures are interrupted. The facial nerve hangs over the interior of the tympanic cavity. The fracture includes the carotid canal housing the internal carotid artery. The plane of the fracture is oblique to the anatomical plane, dividing the temporal bone into superolateral and inferomedial portions (shown here). 1 Interior mastoid antrum, 2 body of incus, 3 tegmentum of mastoid antrum, 4 arcuate eminence, 5 eminence of lateral semicircular canal, 6 facial nerve interrupted in region of genu, 7 tendon of tensor tympani, 8 internal carotid artery, 9 head of malleus, 10 lower wall of the external acoustic meatus, and 11 stapes (head).

The tympanic membrane was injured in longitudinal and mixed fractures. When the fracture line did not include the external acoustic meatus, the membrane was intact, as occurred in 28 of 82 longitudinal fractures (34%) and in 4 of 7 mixed fractures. Clinically, it is observed as a hemotympanum phenomenon. Our findings comply with Tos's data that the frequency of this phenomenon is about 40%.⁶ In these cases, the fracture passed through the mastoid process and often through the sigmoid sinus.

The facial canal was injured in almost all mixed fractures (6/7), in almost half of the longitudinal ones (36/82), and in more than half of the transversal fractures (7/11) (Table 2). Therefore, in our material, the facial canal was injured in 49 of 100 fractures, although 20 fractures were associated with sectioning of the facial nerve (20% of fractures, 40% of fractures with facial canal involvement).

Data available in the literature indicate that the mean frequency of facial nerve dysfunction associated with pyramid fractures is about 16%. However, this percentage depends on the type of fracture. In longitudinal fractures, facial nerve paralysis is observed in 10 to 18% of cases (about 25% associated with the loss of facial nerve continuity) and in 38 to 50% of transverse fractures (100% with a cut nerve stem).^{1,3,6,17,22-26}

In the 82 longitudinal fractures observed in our material, the facial canal was injured in 36 cases (44%), and continuity of the facial nerve was lost in 6 of these 36 cases (7.3%). The facial canal was involved in 63% of the transverse fractures, all of which were associated with a severed nerve stem. Longitudinal fractures in Group II (multiorgan damage, head trauma not the main cause of death) were associated with injury to the facial canal in 6 of 31 cases (19%), and a complete cut of the facial

Figure 2 Transverse fracture of the pyramid in the right temporal bone. The fracture line passes the vestibule and cochlea in front of the round and oval windows. The fracture shows the interior of the vestibule, exposing the outlets of the semicircular canal and footplate from inside. The facial canal is severed at the level of the labyrinthine segment. The facial nerve is interrupted and its end hangs over the labyrinth. The tympanic membrane is intact with light reflex (an optic event visible during the normal otoscopy of the healthy ear). 1 Tympanic membrane, 2 handle of malleus, 3 long process of incus, 4 footplate of stapes, 5 interior of vestibule, 6 facial nerve severed (labyrinthine portion), 7 chorda tympani, 8 belly of tensor tympani muscle.

nerve was observed in 1 of 6 cases (17%). In transverse and mixed fractures, the differences between Groups I and II failed to reach statistical significance.

In our longitudinal fractures, the facial canal was injured only in the genu region or its mastoid portion. Based on the literature, the fracture line in relation to the facial canal in longitudinal fractures is located as follows: 4% in the labyrinthine segment, 64% in the tympanic segment, 25% in the genu region, and 7% in the mastoid segment. Meanwhile, in transverse fractures, the fracture line runs through the facial canal in the meatal segment in 10% of cases, in the labyrinthine

segment in 80%, and in the genu region in 10%.²² Our data support other authors' observations that most facial canal injuries (80 to 93%) involve the genu region.^{1,15,17,21-25}

According to the literature, fractures of the facial canal localized in the labyrinth or tympanic segment always interrupt continuity of the nerve, which is consistent with our findings (Table 3). In contrast to clinical reports, we observed no involvement of the tympanic region of the facial canal with longitudinal fractures.^{15,21,22,25} Injuries to the mastoid region were relatively uncommon (7% of cases with facial canal damage), consistent with other authors' data.^{1,22,23,25} According to many authors,

Figure 3 A mixed fracture of the temporal bone pyramid. The fracture line is perpendicular to the axis of the pyramid, breaking away its apex and showing the carotid canal. It then courses posteriorly, parallel to the axis of the pyramid along the middle ear. The plane of the fracture is first parallel to the frontal plane and then oblique to this plane. In the region of the facial nerve genu an osseous splinter is in the interior of the labyrinth. 1 Interior of the external acoustic meatus, 2 mastoid air cells, 3 interior of lateral semicircular canal, 4 interior of anterior semicircular canal, 5 facial nerve (labyrinthine segment), 6 head of malleus, 7 body of incus, and 8 chorda tympani.

transverse fractures are associated with the highest frequency of facial canal injuries. In contrast, in our study there was no significant difference in the frequency of these injuries associated with either longitudinal or transverse fractures. This finding probably reflects the high number of injuries in the genu region associated with longitudinal fractures, which may not have clinical manifestations. This observation seems to be particularly useful from a clinical perspective, justifying full exploration of the facial canal (up to the genu region) in decompression procedures.

CONCLUSIONS

Fractures of the sigmoid groove and/or jugular fossa, quite often injuries, always interrupt the walls of these vessels. Fractures of the carotid canal, although also frequent, seldom lead to the break of arterial wall. Thus, almost all serious ear hemorrhage is of venous origin. The facial canal in the

genu region is the most common location for injury. In transverse fractures, the nerve trunk is interrupted; in longitudinal and mixed fractures, the continuity of the facial nerve trunk may be preserved.

REFERENCES

1. Nosan DK, Benecke JE, Murr AH. Current perspective on temporal bone trauma. *Otolaryngol Head Neck Surg* 1997;117:67–71
2. Brodie HA, Thompson TC. Management of complications from 820 temporal bone fractures. *Am J Otol* 1997;18:188–197
3. Cannon CR, Jahrsdoefer RA. Temporal bone fractures: review of 90 cases. *Arch Otolaryngol* 1983;109:285–288
4. Ghorayeb BY, Yeakley JW. Temporal bone fractures: longitudinal or oblique. The case for oblique temporal bone fractures. *Laryngoscope* 1992;102:129–134
5. Liu-Shindo M, Hawkins DB. Basilar skull fractures in children. *Int J Pediatr Otolaryngol* 1989;17:109–117
6. Tos M. Course of and sequelae of 248 petrosal fractures. *Acta Otolaryngol* 1973;75:353–354
7. Ulrich K. Verletzungen des Gehörorgans bei Schädelbasisfrakturen. *Acta Otolaryngol Suppl* 1926;6:1–150

8. Aguilar EA. High resolution CT scan of temporal bone fractures: association of facial nerve paralysis with temporal bone fractures. *Head Neck Surg* 1987;9:162-166
9. Fredrickson JM, Griffith AW, Lindsay JR. Transverse fractures of the temporal bone. *Arch Otolaryngol* 1963;78:770-784
10. Hough JVD, Stuart WD. Middle ear injuries in skull trauma. *Laryngoscope* 1968;78:899-937
11. Hough JVD, McGee M. Otologic trauma. In: Paparella MM, Shumrick DA, Gluckman JL, Meyerhoff WL, eds. *Otolaryngology*. Philadelphia, PA: WB Saunders; 1991:1137-1160
12. McHough HE. The surgical treatment of facial paralysis and traumatic conductive deafness in fractures of the temporal bone. *Ann Otol Rhinol Laryngol* 1959;68:855-889
13. Travis LW, Stalnakier RL, Melvin JW. Impact trauma of the human temporal bone. *J Trauma* 1977;17:761-766
14. Voss O. Die Chirurgie der Schädelbasisfrakturen auf Grund. 25 jähriger Erfahrungen. Leipzig, Germany: Johann Ambrosius Barth; 1936:1-87
15. Yanagihara N, Murakami S, Nishihara S. Temporal bone fractures inducing facial nerve paralysis: a new classification and its clinical significance. *ENT J* 1997;76:79-86
16. Kelly KE, Tami TA. Temporal bone and skull base trauma. In: Jackler RK, Brackmann DE, eds. *Neurotology*. St. Louis, MO: CV Mosby; 1994:340-360
17. Kettel K. Peripheral facial paralysis in fractures of the temporal bone. *Arch Otolaryngol* 1950;51:25-41
18. Khan AA, Marion M, Hinojosa R. Temporal bone fractures: a histopathologic study. *Otolaryngol Head Neck Surg* 1985;93:177-186
19. Goodwin WJ Jr. Temporal bone fractures. *Otolaryngol Clin North Am* 1983;16:651-659
20. Kinney SE. Violence and the ear and temporal bone. *Arch Otolaryngol Head Neck Surg* 1992;118:581-583
21. Griffin JE, Altenau MM, Schaffer SD. Bilateral longitudinal temporal bone fractures: a retrospective review of seventeen cases. *Laryngoscope* 1979;89:1432-1435
22. Fisch U. Facial paralysis in fractures of the petrous bone. *Laryngoscope* 1974;84:2141-2154
23. Coker NJ, Kendall KA, Jenkins HA, Alford BR. Traumatic intratemporal facial nerve injury: management rationale for preservation of function. *Otolaryngol Head Neck Surg* 1987;97:262-269
24. Kamerer DB. Intratemporal facial nerve injuries. *Otolaryngol Head Neck Surg* 1982;90:612-615
25. Lambert PR, Brackmann DE. Facial paralysis in longitudinal temporal bone fractures: a review of 26 cases. *Laryngoscope* 1984;94:1022-1026
26. Brawley BW, Kelly WA. Treatment of basal skull fractures with and without cerebrospinal fluid fistulae. *J Neurosurg* 1967;26:57-61
27. Glasscock ME 3rd, McKennan KX, Levine SC. Persistent traumatic perilymph fistulas. *Laryngoscope* 1987;97:860-864
28. Emmett JR, Shea JJ. Traumatic perilymph fistula. *Laryngoscope* 1980;90:1513-1520
29. Hall JW, Huang-Fu M, Gennarelli TA. Auditory function in acute severe head injury. *Laryngoscope* 1982;92:883-889
30. Tos M. Prognosis of hearing loss in temporal bone fractures. *J Laryngol Otol* 1971;85:1147-1159
31. Lyos AT, Marsh MA, Jenkins HA, Coker NJ. Progressive hearing loss after transverse temporal bone fracture. *Arch Otolaryngol Head Neck Surg* 1995;121:795-799
32. Makishima K, Sobel SF, Snow JB. Histopathologic correlates of otoneurologic manifestations following head trauma. *Laryngoscope* 1976;86:1303-1314
33. Griffiths MV. The incidence of auditory and vestibular concussion following minor head injury. *J Laryngol Otol* 1979;93:253-265
34. Goodwin JR, Johnson MH. Carotid injury secondary to blunt head trauma: case report. *J Trauma* 1994;37:119-122
35. May M. Trauma to the facial nerve. *Otolaryngol Clin N Am* 1983;16:661-670
36. Resnick DK, Subach BR, Marion DW. The significance of carotid canal involvement in basilar cranial fracture. *Neurosurgery* 1997;40:1177-1181

Commentary

Dr. Wysocki presents a well-done, careful, and systematic microanatomical study on 100 consecutive temporal bone fractures associated with fatal head injury. Autopsy studies with these kind of numbers are becoming increasingly rare, particularly in the United States, yet a tremendous amount of information can come from them. This study is a good example.

He found that 82% of fractures were longitudinal, 11% transverse, and 7% mixed, which is not unexpected. He found a surprising incidence of facial canal disruption (37%) associated with longitudinal fractures, but nerve disruption occurred in only 7% versus 63% of cases of transverse fracture. He also noted a surprising incidence of ossicular disruption, fracture, or both, as well as jugular bulb injury. Both of these latter findings seem significantly higher than clinical experience would suggest and may reflect the severity of temporal bone fractures associated with fatal head injury, as opposed to basilar skull fractures associated with lesser degrees of survivable closed head injury.

The incidence of carotid canal fracture (52%) is very interesting given Dr. Wysocki's finding of

carotid artery injury in only 1 case. Clinical experience suggests that angiographic abnormalities will be present in 42% of cases studied for the presence of a basal skull fracture involving the carotid canal,¹ and as many as 18% of patients with carotid canal fractures may develop clinically symptomatic vascular complications.² The likely explanation for this discrepancy is that basal skull fractures through the carotid canal probably lead to intimal dissection as the most common injury, subsequently resulting in thrombosis or embolization rather than transmural disruption. I suspect that Dr. Wysocki's autopsy studies are less sensitive to detecting intimal injury as opposed to transmural disruption through the adventitia. From a clinical standpoint, we need to maintain heightened vigilance for vascular sequelae when carotid canal fractures are identified. Whether

or not routine further screening (angiography, computed tomography angiography, magnetic resonance angiography) is a cost-effective approach remains the subject of ongoing debate.^{1,3}

Mark E. Linskey, M.D.¹

REFERENCES

1. Kerwin AJ, Bynoe RP, Murray J, et al. Liberalized screening for blunt carotid and vertebral artery injuries is justified. *J Trauma* 2001;51:308–314
2. Resnick DK, Subach BR, Marion DW. The significance of carotid canal involvement in basilar cranial fracture. *Neurosurgery* 1997;40:1177–1181
3. Mayberry JC, Brown CV, Mullins RJ, Velmahos GC. Blunt carotid artery injury: the futility of aggressive screening and diagnosis. *Arch Surg* 2004;139:609–612; discussion 612–613